Published in final edited form as:

Qual Life Res. 2020 January ; 29(1): 191–199. doi:10.1007/s11136-019-02266-x.

Reliability and Validity of PROMIS® Pediatric Family Relationships Short Form in Children 8-17 Years of Age with Chronic Disease

Elizabeth D. Cox, MD PhD,

Department of Pediatrics, University of Wisconsin School of Medicine and Public Health, H6/558 Clinical Science Center, 600 Highland Avenue, Madison, WI, USA 53792-4108,

Jennifer R. Connolly, MS,

Department of Pediatrics, University of Wisconsin School of Medicine and Public Health, H6/577 Clinical Science Center, 600 Highland Avenue, Madison, WI, USA 53792-4108

Mari Palta, PhD,

Department of Population Health Sciences, University of Wisconsin School of Medicine and Public Health, 689 Wisconsin Alumni Research Foundation, 610 Walnut Street, Madison, WI, USA 53726-2336

Victoria P. Rajamanickam, MS,

Department of Biostatistics and Medical Informatics, University of Wisconsin School of Medicine and Public Health, H6/526 Clinical Science Center, 600 Highland Avenue, Madison, WI, USA 53792-4108

Kathryn E. Flynn, PhD

Department of Medicine, Medical College of Wisconsin, 9200 W. Wisconsin Ave, Milwaukee, WI, USA 53226-3522

Introduction

Children with chronic illnesses rely on their families in many ways: Parents and other family members provide key support for managing a child's disease, and gradually they support their children in taking over that management themselves. Further, many studies have shown that family support is strongly linked to both improved health as well as better psychosocial outcomes for chronically ill children [1–8]. Family relationships and family functioning often change over time, as children go through different developmental stages and eventually achieve autonomy from their parents [9]. Families of children with chronic illness are thus

ecox@wisc.edu, Phone: (608) 263-9104.

Compliance with ethical standards

Ethical approval: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Informed consent: All participating parents provided informed consent. Participating children ages 14 to 17 years also gave informed consent, while children ages 8 to 13 years provided assent.

Conflict of interest: The authors declare that they have no conflict of interest.

faced with the challenge of balancing the child's autonomy and ongoing development on the one hand, while also making sure that disease management behaviors occur [10–15].

Despite the important role that families play for chronically ill children, there are few validated measures available that assess a child's perceptions of family relationships [16–18]. Acknowledging the important role of family relationships and the lack of assessment tools, the National Institutes of Health (NIH) supported the development and validation of an item bank measuring children's experience with their families, the Pediatric Family Relationships measure, as part of the Patient-Reported Outcomes Measurement Information System® (PROMIS®). The measure was developed for self-report by children ages 8 to 17 years and report by parents of children ages 5 to 17 years. In accordance with PROMIS methodological standards for measure development and validation, the items were psychometrically evaluated in two large field studies in general populations in the United States [19]. However, evidence of the reliability and validity of this new measure for use in clinical settings with chronically ill children is needed.

This paper presents the findings from our investigation as to whether the Family Relationships measure demonstrates adequate reliability and validity when applied in a clinical population of chronically ill children and their parents. For this evaluation, we selected three chronic illnesses—asthma, type 1 diabetes, and sickle cell disease. About 6.5 million US children, or approximately 1/3 of all US children with a chronic illness, have been diagnosed with one of these diseases [20–23]. Our hypotheses were: 1) the error of measurement will vary across the range of PROMIS Family Relationships scores, 2) the measure will demonstrate adequate internal consistency and reliability, and 3) the measure will display higher correlations with PROMIS social and mental health domains, and smaller correlations with physical health domains.

Methods

Subjects and procedures

Children with asthma, type 1 diabetes, or sickle cell disease (N=275) were recruited from the pediatric specialty clinics of two Midwestern tertiary care centers from March 2016 to May 2018. During the recruitment period, care team members asked each potentially eligible family about interest in the study. Eligibility criteria for the study included being an English-speaking child who was 8 to 17 years of age with one of the three chronic diseases and with an English-speaking parent or legal guardian, since, at study initiation, the PROMIS Family Relationships measure was only available in English. Assessments were completed on paper or electronically (REDCap). The Institutional Review Board at the primary study site approved the study. All parents provided informed consent. Children ages 14 to 17 years also gave informed consent, while children ages 8 to 13 years provided assent. We did not collect any data on families who declined to participate.

PROMIS Measures

We administered PROMIS measures to all child and parent participants. For children, we administered the PROMIS Pediatric Short Form v1.0 - Family Relationships 8a, which

includes eight items using a 4-week recall period and 1–5-point response options from never to always. The items cover multiple facets of family relationships, such as sense of family (e.g., “I had a strong relationship with my family”), trust, dependability, and support, (e.g., “I got all the help I needed from my family”), love and caring, value and acceptance, enjoyment, and communication. Children also received the PROMIS Pediatric-25 Profile v2.0, which includes six 4-item short forms covering social (Peer Relationships), mental (Anxiety, Depressive Symptoms), and physical health (Fatigue, Pain Interference, and Physical Function-Mobility) plus a single Pain Intensity item. The Profile-25 domains use a 7-day recall period. All items use standard 1–5-point response options except Pain Intensity, which is 0–10. For parents, we included the corresponding PROMIS Parent Short Form v1.0 - Family Relationships 8a and the PROMIS Parent Proxy-25 Profile v2.0. Item Response Theory (IRT)-based T-scores were calculated for all measures except Pain Intensity (single item on 0–10 numerical rating scale). For the PROMIS measures included in our study, a mean T-score of 50 and standard deviation of 10 represents the general US population, including both healthy and chronically ill children [24]. Higher scores indicate greater levels of the construct. For example, for Family Relationships and Peer Relationships, higher scores are more favorable, but for Fatigue and Anxiety, higher scores are less favorable.

Other measures

Participant Characteristics—Children completed a standard, single item reflecting overall health status (five categories: poor, fair, good, very good, excellent). Parents completed items reflecting child age (continuous; 8–17 years), child gender (male, female, or other), child race (Black or African-American, White, American-Indian or Alaskan Native, Asian, Native Hawaiian or other Pacific Islander, or other; aggregated to Black or African-American, White, and Multi-race or other), and child ethnicity (Hispanic versus all other), as well as parent age (continuous; 27–64 years), parent education (less than high school, high school diploma, some college, bachelor’s degree, or post-graduate degree), and relationship to the child (12 options such as biological mother, biological father, step-mother, step-father, guardian, grandmother, grandfather, or other; which were aggregated to biological, adoptive, or step-mother versus all other).

Analyses—We used means (M) with standard deviations (SD) and proportions to describe participants. Within each chronic disease, the standard error of measurement (SEM) for individual scores was calculated in SAS using algorithms provided by PROMIS. We assessed the possibility that the scores have different levels of certainty at different levels of the score by graphing the SEM of the T-score over the entire range of scores within each disease type. We also assessed the percentage of scores with SEM >4, corresponding to a true variance of 10 and reliability of 0.85. For each chronic disease, IRT-estimated marginal reliability and Cronbach’s alpha reflecting internal consistency were calculated for child and parent reports. Convergent and discriminant validity were assessed from Spearman correlations of the Family Relationships scores with PROMIS-25 domains within each chronic disease. Results are reported as the correlation coefficients and boot-strapped 95% confidence intervals (CI). We regarded a two-tailed $p < 0.05$ as significant for this analysis, and correlation coefficients of < 0.30 as small, $r = 0.30$ to $r = 0.49$ as medium, and 0.50 as large.

Results

Participant characteristics

A total of 275 families contributed data for this work. Children's mean ages were 12.2 (asthma), 13.1 (type 1 diabetes), and 12.0 (sickle cell disease) years within the three disease groups respectively (Table 1). In asthma and sickle cell disease, about 60% of participants were 8 to 12 years of age, but in type 1 diabetes participants were nearly evenly split between 8 to 12 year and 13 to 17 year olds. With regard to race, the majority of children with type 1 diabetes were White, while the majority of children with asthma or sickle cell disease were either Black, multi-race or other. Most children reported their health as good, very good, or excellent. Most parent respondents were mothers (84%), with varying levels of educational attainment.

Family Relationships scores by chronic disease type

Across the three diseases, mean children's Family Relationships scores were similar ($M=53.0$ among children with asthma, $M=53.2$ in type 1 diabetes, and $M=54.6$ in sickle cell disease; Table 2). Parent scores differed significantly between the three diseases, with higher scores reported by parents of children with asthma ($M=56.2$) and sickle cell disease ($M=56.2$), than by parents of children with type 1 diabetes ($M=51.1$, $p<0.001$).

Standard error of measurement of individual PROMIS Family Relationships scores

Figure 1 shows SEMs across the range of estimated true scores for all three diseases, for children and parents. The maximum Family Relationships score was 63.9 for both children and parents, while the minimum was 17.8 for children and 25.8 for parents. The SEMs were similar across diseases. SEMs for those estimated to have true values below 50 were fairly constant and small, especially for parents. However, SEMs were larger for estimated scores above the normative mean of 50.

Among parent reports of Family Relationships, 54%, 35% and 64% of scores have an SEM above four points for asthma, type 1 diabetes, and sickle cell, respectively (Table 3). Corresponding values for children are 23%, 27%, and 38%, respectively.

IRT-estimated marginal reliability and internal consistency

For children, IRT-estimated marginal reliability was 0.84 in each of the three disease groups (Table 4). For parents, IRT-estimated marginal reliability was highest among parents of children with type 1 diabetes (0.85) and lowest for parents of children with asthma (0.75). For sickle cell disease, the IRT-estimated reliability for parents was 0.80. Within each of the three chronic illnesses, internal consistency for both child and parent responses was good with Cronbach's alpha 0.88.

Correlations with other PROMIS-25 domains

Within each of the three disease groups, children's Family Relationships scores correlated significantly and in the expected direction with scores on the PROMIS social health and mental health domains (Table 5). Specifically, within each of the three disease groups, moderate positive correlations with Peer Relationships, moderate negative correlations with

Depressive Symptoms, and small to moderate negative correlations with Anxiety were found for child respondents. Similar patterns were seen among parent responses.

For children with type 1 diabetes and sickle cell disease, Family Relationships scores also correlated significantly and in the expected direction, but to a lesser degree, with PROMIS physical health domains. For example, Fatigue exhibited a moderate negative correlation with Family Relationships among children with type 1 diabetes and a small negative correlation among children with sickle cell disease. Among children with type 1 diabetes, Family Relationships exhibited small negative correlations with Pain Interference and Pain Intensity, as well as small positive correlation with Mobility. However, this was not the case for children with asthma. For parents, similar results were seen for these PROMIS domains.

Discussion

We sought to understand whether the PROMIS Family Relationships measure demonstrates adequate reliability and validity when applied in a clinical population of chronically ill children and their parents. Findings from this work extend evidence for the reliability and validity of the PROMIS Family Relationships v1.0 8-item short forms for use with the 6.5 million US children who are impacted by one of these three chronic diseases. In analyses of data from chronically ill children and their parents, we found that children's assessments of their family relationships were similar across the diseases, while parent scores were significantly higher in asthma and sickle cell disease than in type 1 diabetes. Standard errors of measurement for Family Relationships scores were smallest for low scores, suggesting greater certainty for assessments that would raise concern (i.e., worse relationships). Across all three diseases, the measure demonstrated adequate reliability for both children and their parents. In addition, convergent and discriminant validity were supported, as the measure displayed expected correlations with other PROMIS health domains. For children with one of the three chronic diseases included in our study, the PROMIS Family Relationships short form offers a reliable and valid approach to assessing family relationships, from both the child and parent perspective.

In general, the distribution of PROMIS Family Relationships scores in our sample of chronically ill children is similar to those of the general population in which the measure was developed, although mean values for the scores are higher in our sample [19]. Specifically, both populations had maximum Family Relationships scores of 63 to 64. Standard deviations for the scores within each of the three disease groups were similar to those of the general population (10) for both children (9.8) and parents (9.0). Among the general population, positive skew and ceiling effects were noted, similar to those in our population. As with the general population, the PROMIS Family Relationships measure appears to be more useful for discriminating among scores for children with poorer relationships than for those with better relationships [19]. Hence, individual scores are estimated most precisely in the range of most clinical concern.

Despite these similarities, for each of the three diseases in our study, children's mean Family Relationships scores were about 3 points (0.3 SD) higher than the general population mean (51.6). Mean parent Family Relationships scores were also higher by as much as 6 points

(0.6 SD) in sickle cell disease and asthma, compared to the general population (51.2), but similar to the general population mean for parents of children with type 1 diabetes. This is in contrast to findings from prior work in the general population, in which parents of children with chronic illness reported lower Family Relationships scores with a significant effect size of 0.2 [19]. However, in this same study, having a chronic illness was not associated with children's Family Relationships scores. Also, the specific chronic conditions that children in the general population sample had are unknown, and the severity and impact of the chronic conditions may differ substantially. In the general population study, parents reported whether their child had a chronic condition that "lasted or was expected to last at least 12 months AND interferes with his/her activities." Mean Family Relationships scores for children and parents in our sample were also higher than those reported in a study focused on children with cancer, both on and off therapy [25].

Other possible explanations for the higher Family Relationships scores in this work may lie in differences in the characteristics of the samples, beyond just the presence and type of chronic illness. For example, we recruited only children who presented for care of their chronic illness at a clinic, whereas other studies in general populations did not limit recruitment to clinical sites. Thus, children in this study had families who ensured they received healthcare for their chronic disease. A second potential explanation may be differences in the socio-demographic make-up of the two samples. Specifically, our sample included more children 8 to 12 years of age than in the general population on which the measure was developed. Younger children and their parents have been noted to have higher Family Relationships scores. In addition, Black or African-American children, whose parents within the general population reported better family relationships than parents of White children, made up 44% of our sample. However, in the general population, Black or African American children did not report better family relationships than their White peers [19].

Among children with asthma, type 1 diabetes, or sickle cell disease, the reliability of the PROMIS Family Relationships 8a short form, while certainly adequate for group comparisons, was lower than that seen in the general population. Specifically, among the general population, internal consistency assessed with Cronbach's alpha was 0.94 for children and 0.91 for parents [19]. To date, the published literature does not contain IRT marginal reliabilities for the measure within the general population, but our evaluation of this parameter suggests adequate reliability among the three disease types for both children and parents. Further, our results suggest that in some but not all of our respondent groups, the measure also achieves the higher reliability threshold (>0.9) recommended for individual patient assessment [26].

Convergent and discriminant validity were supported through expected correlations with other PROMIS domains. Specifically, for both children and parents, moderate, positive correlations were consistently seen with aspects of social or mental health, across all three diseases. This was as expected, given family relationships are an element of social health, which "encompasses the ways in which individuals connect with important others, including communication, companionship and understanding, and the quality, reciprocity and size of an individual's social network" [27]. Further, these findings are consistent with prior concept

elicitation work to develop the PROMIS Family Relationships measure, in which children themselves articulated how their family relationships influence peer relationships, through friend selection [19]. In addition, Family Relationships scores displayed small to moderate negative correlations with both anxiety and depression, as reported by children and parents. This is consistent with accepted diagnostic criteria for these entities, which requires functional impairment in social or family settings [28].

Our study has several limitations. For this evaluation, we included only English-speaking families whose child had at least one of three chronic diseases—asthma, type 1 diabetes, or sickle cell disease. Although these diseases represent over 1/3 of children with pediatric chronic illness, generalizing our findings to the entire population of children with chronic conditions should be done with caution. Further, because we recruited participants from a tertiary care facility, the children with asthma may well have more severe disease than those with asthma who are cared for predominantly by primary care physicians. Future work could evaluate reliability and validity of the Spanish-language version of the instrument, which is now available, and also validate the measure among children with other pediatric chronic illnesses. As expected within the three disease groups included in our sample, participants were predominantly White or African American. The geographic diversity of our sample was limited, though we recruited participants from two large centers that capture both rural and urban families. Families were aware of study participation, potentially leading to higher scores for Family Relationships, as a result of socially desirable survey responses. Lastly, we included only children 8 to 17 years of age and their parents, aligning with the age group for which pediatric PROMIS measures have been designed for child self-report.

In summary, this study's findings add to the evidence for reliability and validity of the PROMIS Pediatric and Parent Short Form v1.0 Family Relationships 8a for use with children who have a chronic disease. Our results suggest adequate psychometric properties for use among three chronic diseases that encompass 1/3 of the children with a chronic condition. In addition, the measure's properties appear consistent between children with chronic disease in this study and children from the general population on which the measure's initial rigorous development occurred. These conclusions complement qualitative work demonstrating the content validity of the PROMIS Family Relationships short form among children with chronic diseases and their parents [29]. Evidence of the measure's reliability and validity in multiple contexts furthers the case for its widespread use. Ultimately, the measure could be used to evaluate outcomes from interventions that focus on improving family interactions and also could support healthcare organizations and researchers in identifying families who may need additional support.

Acknowledgments

Funding: Research reported in this publication was supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health under Award Number U19 AR069519. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Perez-Marin M, Gomez-Rico I, Montoya-Castilla I. Type 1 diabetes mellitus: psychosocial factors and adjustment of pediatric patient and his/her family. Review. Arch Argent Pediatr 2015; 113(2): 158–162. [PubMed: 25727829]
2. Moreira H, Frontini R, Bullinger M, Canavarro MC. Caring for a child with type 1 diabetes: links between family cohesion, perceived impact, and parental adjustment. J Fam Psychol 2013; 27(5): 731–742. [PubMed: 24015704]
3. Raphael JL, Butler AM, Rattler TL, Kowalkowski MA, Mueller BU, Giordano TP. Parental information, motivation, and adherence behaviors among children with sickle cell disease. Pediatr Blood Cancer 2013; 60(7): 1204–1210. [PubMed: 23303618]
4. Logan DE, Radcliffe J, Smith-Whitley K. Parent factors and adolescent sickle cell disease: associations with patterns of health service use. J Pediatr Psychol 2002; 27(5): 475–484. [PubMed: 12058011]
5. Hildenbrand AK, Barakat LP, Alderfer MA, Marsac ML. Coping and coping assistance among children with sickle cell disease and their parents. J Pediatr Hematol Oncol 2015; 37(1): 25–34. [PubMed: 24327131]
6. La Greca AM, Auslander WF, Greco P, Spetter D, Fisher EB, Santiago JV. I get by with a little help from my family and friends: adolescents' support for diabetes care. J Pediatr Psychol 1995; 20(4): 449–476. [PubMed: 7666288]
7. Christiaanse ME, Lavigne JV, Lerner CV. Psychosocial aspects of compliance in children and adolescents with asthma. J Dev Behav Pediatr 1989; 10(2): 75–80. [PubMed: 2708540]
8. Cutfield SW, Derraik JGB, Reed PW, Hofman PL, Jefferies C, Cutfield WS. Early markers of glycaemic control in children with type 1 diabetes mellitus. PLoS One 2011; 6(9): e25251.
9. Hazen E, Schlozman S, Beresin E. Adolescent psychological development: a review. Pediatr Rev 2008; 29(5): 161–167. [PubMed: 18450837]
10. Hanson CL, Henggeler SW, Harris MA, Burghen GA, Moore M. Family system variables and the health status of adolescents with insulin-dependent diabetes mellitus. Health Psychol 1989; 8(2): 239–253. [PubMed: 2737175]
11. Hauser ST, Jacobson AM, Lavori P, Wolfsdorf JI, Herskowitz RD, Milley JE et al. Adherence among children and adolescents with insulin-dependent diabetes mellitus over a four-year longitudinal follow-up: II. immediate and long-term linkages with the family milieu. J Pediatr Psychol 1990; 15(4): 527–542. [PubMed: 2258799]
12. Hanson CL. Developing systemic models of the adaptation of youths with diabetes In: La Greca AM, Siegel LJ, Wallander JL, Walker CE, editors. Advances in Pediatric Psychology: Stress and Coping in Child Health. New York, NY: Guilford Press; 1992 p. 212–241.
13. Walders N, Drotar D, Kerckmar C. The allocation of family responsibility for asthma management tasks in African-American adolescents. J Asthma 2000; 37(1): 89–99. [PubMed: 10724302]
14. Helgeson VS, Palladino DK, Reynolds KA, Becker DJ, Escobar O, Siminerio L. Relationships and health among emerging adults with and without type 1 diabetes. Health Psychol 2014; 33(10): 1125–1133. [PubMed: 23914816]
15. Rosland AM, Heisler M, Piette JD. The impact of family behaviors and communication patterns on chronic illness outcomes: a systematic review. J Behav Med 2012; 35(2): 221–239. [PubMed: 21691845]
16. Olson D FACES IV and the Circumplex Model: validation study. J Marital Fam Ther 2011; 37(1): 64–80. [PubMed: 21198689]
17. Moos RH, Moos BS. Family environment scale manual. Consulting Psychologists Press; 1994.
18. Furman W, Giberson RS. Identifying the links between parents and their children's sibling relationships Close relationships and socioemotional development. Westport, CT, US: Ablex Publishing; 1995 p. 95–108.
19. Bevans KB, Riley AW, Landgraf JM, Carle AC, Teneralli RE, Fiese BH et al. Children's family experiences: development of the PROMIS® pediatric family relationships measures. Qual Life Res 2017; 26(11): 3011–3023. [PubMed: 28643117]

20. Van Cleave J, Gortmaker SL, Perrin JM. Dynamics of obesity and chronic health conditions among children and youth. *JAMA* 2010; 303(7): 623–630. [PubMed: 20159870]
21. Centers for Disease Control and Prevention. Most Recent National Asthma Data. 2019; Available from: https://www.cdc.gov/asthma/most_recent_national_asthma_data.htm.
22. Centers for Disease Control and Prevention. Data & Statistics on Sickle Cell Disease. 2017; Available from: <https://www.cdc.gov/ncbddd/sicklecell/data.html>.
23. Menke A, Orchard TJ, Imperatore G, Bullard KM, Mayer-Davis E, Cowie CC. The prevalence of type 1 diabetes in the United States. *Epidemiology* 2013; 24(5): 773–774. [PubMed: 23903880]
24. DeWalt DA, Gross HE, Gipson DS, Selewski DT, DeWitt EM, Dampier CD et al. PROMIS® pediatric self-report scales distinguish subgroups of children within and across six common pediatric chronic health conditions. *Qual Life Res* 2015; 24(9): 2195–2208. [PubMed: 25715946]
25. Erker C, Yan K, Zhang LY, Bingen K, Flynn KE, Panepinto J. Impact of pediatric cancer on family relationships. *Cancer Medicine* 2018; 7(5): 1680–1688. [PubMed: 29577633]
26. Fayers PM, Machin D. *Quality of Life: The Assessment, Analysis and Interpretation of Patient-reported Outcomes*. 3 ed: John Wiley & Sons, Ltd; 2016 p. 139.
27. Hahn EA, Devellis RF, Bode RK, Garcia SF, Castel LD, Eisen SV et al. Measuring social health in the patient-reported outcomes measurement information system (PROMIS): item bank development and testing. *Qual Life Res* 2010; 19(7): 1035–1044. [PubMed: 20419503]
28. American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 5th ed. Arlington, VA: American Psychiatric Publishing; 2013.
29. Flynn KE, Kliems H, Saoji N, Svenson J, Cox ED. Content validity of the PROMIS® pediatric family relationships measure for children with chronic illness. *Health Qual Life Outcomes* 2018; 16(1): 203. [PubMed: 30340500]

Figure 1.
Standard Error of Measurement by Family Relationships Score for Children and Parents by Chronic Disease

Table 1.Family characteristics by disease type^a (N=275)

Child characteristics	Asthma (n=73)	Diabetes (n=122)	Sickle Cell (n=80)
Age, years (mean (SD))	12.2 (2.6)	13.1 (2.7)	12.0 (2.5)
8–12 years, % (n)	61.6% (45)	48.4% (59)	63.8% (51)
13–17 years, % (n)	38.4% (28)	51.6% (63)	36.3% (29)
Female, % (n)	35.6% (26)	47.5% (58)	52.5% (42)
Race, % (n)			
Black or African-American	54.8% (40)	7.4% (9)	91.3% (73)
White	28.8% (21)	83.6% (102)	0.0% (0)
Multi-race or other	9.6% (7)	8.2% (10)	3.8% (3)
Ethnicity, % (n)			
Hispanic or Latino	17.8% (13)	9.8% (12)	5.0% (4)
Health status, child report, % (n)			
Poor	1.4% (1)	0.8% (1)	2.5% (2)
Fair	6.8% (5)	4.1% (5)	11.3% (9)
Good	45.2% (33)	33.6% (41)	30.0% (24)
Very good	32.9% (24)	41.8% (51)	36.3% (29)
Excellent	12.3% (9)	17.2% (21)	18.8% (15)
Parent characteristics			
Age, years (mean (SD))	41.3 (7.4)	41.1 (6.1)	37.3 (7.4)
Relationship to child, mother, % (n)	82.2% (60)	86.9% (106)	82.5% (66)
Education, % (n)			
Less than high school	11.0% (8)	3.3% (4)	15.0% (12)
High school diploma	21.9% (16)	3.3% (4)	16.3% (13)
Some college	28.8% (21)	38.5% (47)	32.5% (26)
Bachelor's degree	15.1% (11)	32.8% (40)	10.0% (8)
Post-graduate degree	11.0% (8)	12.3% (15)	7.5% (6)

^aValues may not add to 100% due to rounding or non-response.

Table 2.

Pediatric PROMIS domain scores

PROMS Domain	Child Report ^a			Parent Report ^a		
	Asthma (n=71)	Diabetes (n=120)	Sickle (n=79)	Asthma (n=70)	Diabetes (n=120)	Sickle Cell (n=80)
	mean (SD)	mean (SD)	mean (SD)	mean (SD)	mean (SD)	mean (SD)
Social Health						
Family Relationships ^b	53.0 (9.2)	53.2 (9.0)	54.6 (9.1)	56.2 (7.9)	51.1 (9.2)	56.2 (9.6)
Peer Relationships ^b	49.6 (9.4)	52.4 (9.7)	50.1 (10.9)	49.8 (11.3)	51.2 (9.8)	50.3 (11.3)
Mental Health						
Anxiety	48.6 (11.3)	45.8 (10.3)	47.3 (11.7)	44.2 (9.4)	45.7 (11.0)	44.5 (11.2)
Depressr/e Symptoms	48.8 (10.0)	45.1 (9.5)	46.3 (11.0)	45.1 (9.8)	45.6 (10.2)	44.3 (10.5)
Physical Health						
Fatigue	49.0 (10.1)	45.3 (10.8)	48.2 (11.0)	48.6 (9.9)	45.2 (10.5)	48.9 (12.2)
Pain Interference	51.1 (10.6)	44.9 (9.7)	52.0 (12.0)	51.2 (9.6)	45.4 (8.6)	52.0 (12.5)
Physical Function - Mobility ^b	48.7 (8.7)	53.1 (6.2)	49.5 (8.9)	47.1 (8.5)	53.5 (5.3)	47.9 (9.4)
Pain Intensity ^c	3.1 (2.3)	1.9 (2.2)	3.8 (3.6)	2.4 (2.6)	1.1 (1.6)	3.0 (3.4)

^aSample sizes for children and parents differ due to non-response.^bHigher scores indicate better health.^cOn a scale of 0–10, where 10 is worst pain.

Table 3.

Overall standard errors of measurement (SEM) of PROMIS Family Relationships short form scores for chronically ill children 8–17 years old

	Child Report					Parent Report				
	n	Overall SEM	Minimum SEM	Maximum SEM	Percentage SEM above 4	n	Overall SEM	Minimum SEM	Maximum SEM	Percentage SEM above 4
Asthma	71	3.53	2.0	5.5	22.5	70	4.58	2.2	5.8	54.3
Diabetes	120	3.63	2.0	5.6	26.7	119	3.84	2.2	5.8	34.5
Sickle Cell	79	4.00	2.0	5.6	38.0	80	4.81	2.2	5.8	63.8

Table 4.

Reliability of PROMIS Family Relationships short form scores for chronically ill children 8–17 years old

	Child Report			Parent Report		
	n	IRT estimated marginal reliability	Cronbach's Alpha (95% CI)	n	IRT estimated marginal reliability	Cronbach's Alpha (95% CI)
Asthma	71	0.87	0.92 (0.84, 0.96)	70	0.75	0.88 (0.75, 0.94)
Diabetes	120	0.86	0.93 (0.89, 0.95)	119	0.85	0.91 (0.88, 0.93)
Sickle Cell	79	0.84	0.89 (0.82, 0.93)	80	0.80	0.94 (0.89, 0.96)

Table 5.

spearman correlation and 95% confidence interval of PROMIS Family Relationship with PROMIS 25 domains for chronically ill children

PROMIS Measure	Asthma		Diabetes		Sickle Cell	
	Child Report (n=70)	Parent Report (n=69)	Child Report (n= 120)	Parent Report (n=119)	Child Report (n=78)	Parent Report (n=80)
Social Health						
Peer Relationships	0.32 (0.15, 0.50)	0.25 (0.07, 0.43)	0.47 (0.35, 0.59)	0.56 (0.45, 0.67)	0.39 (0.22, 0.57)	0.40 (0.23, 0.55)
Mental Health						
Anxiety	-0.27 (-0.45, -0.09)	-0.26 (-0.43, -0.10)	-0.39 (-0.50, -0.26)	-0.29 (-0.41, -0.16)	-0.26 (-0.42, -0.10)	-0.38 (-0.54, -0.22)
Depressive Symptoms	-0.30 (-0.48, -0.11)	-0.32 (-0.50, -0.13)	-0.40 (-0.51, -0.25)	-0.58 (-0.68, -0.48)	-0.39 (-0.56, -0.23)	-0.60 (-0.74, -0.48)
Physical Health						
Fatigue	-0.11 (-0.30, 0.08)	-0.06 (-0.25, 0.13)	-0.39 (-0.51, -0.26)	-0.42 (-0.54, -0.30)	-0.22 (-0.38, -0.05)	-0.19 (-0.37, -0.01)
Pain Interference	-0.06 (-0.23, 0.12)	0.06 (-0.13, 0.25)	-0.25 (-0.38, -0.13)	-0.20 (-0.34, -0.07)	0.03 (-0.15, 0.21)	-0.11 (-0.27, 0.06)
Mobility	0.05 (-0.14, 0.24)	-0.08 (-0.28, 0.11)	0.20 (0.06, 0.33)	0.29 (0.15, 0.42)	0.11 (-0.06, 0.27)	0.18 (0.01, 0.36)
Pain Intensity	0.04 (-0.13, 0.23)	0.03 (-0.16, 0.22)	-0.25 (-0.38, -0.12)	-0.27 (-0.41, -0.13)	-0.01 (-0.19, 0.16)	-0.22 (-0.39, -0.04)