Published in final edited form as:

Free Radic Biol Med. 2019 February 01; 131: 18–26. doi:10.1016/j.freeradbiomed.2018.11.013.

Synthesis and evaluation of ^{13}C -labeled 5-5-dimethyl-1-pyrroline-N-oxide aimed at in vivo detection of reactive oxygen species using hyperpolarized ^{13}C -MRI

Keita Saito^{#a}, Deepak Sail^{#b}, Kazutoshi Yamamoto^a, Shingo Matsumoto^c, Burchelle Blackman^b, Shun Kishimoto^a, Jeffery Brender^a, Rolf E. Swenson^b, James B. Mitchell^a, Murali C. Krishna^a

^aRadiation Biology Branch, National Cancer Institute, Bethesda, MD, USA

^bImaging Probe Development Center, National Heart, Lung, and Blood Institute, Rockville, MD, USA

^cGraduate School of Information Science and Technology, Hokkaido University, Sapporo, Hokkaido, Japan

[#] These authors contributed equally to this work.

Abstract

Effective means to identify the role of reactive oxygen species (ROS) mediating several diseases including cancer, ischemic heart disease, stroke, Alzheimer's and other inflammatory conditions in in vivo models would be useful. The cyclic nitrene 5,5-Dimethyl-1-pyrroline-N-oxide (DMPO) is a spin trap frequently used to detect free radicals in vitro using Electron Paramagnetic Resonance (EPR) spectroscopy. In this study, we synthesized ^{13}C -labeled DMPO for hyperpolarization by dynamic nuclear polarization, in which ^{13}C NMR signal increases more than 10000-fold. This allows in vivo ^{13}C MRI to investigate the feasibility of in vivo ROS detection by the ^{13}C -MRI. DMPO was ^{13}C -labeled at C5 position, and deuterated to prolong the T_1 relaxation time. The overall yield achieved for 5- ^{13}C -DMPO- d_9 was 15%. Hyperpolarized 5- ^{13}C -DMPO- d_9 provided a single peak at 76 ppm in the ^{13}C -spectrum, and the T_1 was 60 s in phosphate buffer making it optimal for in vivo ^{13}C MRI. The buffered solution of hyperpolarized 5- ^{13}C -DMPO- d_9 was injected into a mouse placed in a 3 T scanner, and ^{13}C -spectra were acquired every 1 s. In vivo studies showed the signal of 5- ^{13}C -DMPO- d_9 was detected in the mouse, and the T_1 decay of ^{13}C signal of hyperpolarized 5- ^{13}C -DMPO- d_9 was 29 s. ^{13}C -chemical shift imaging revealed that 5- ^{13}C -DMPO- d_9 was distributed throughout the body in a minute after the intravenous injection. A strong signal of 5- ^{13}C -DMPO- d_9 was detected in heart/lung and kidney, whereas the signal in liver was small compared to other organs. The results indicate hyperpolarized 5- ^{13}C -DMPO- d_9

*Corresponding author: Murali C. Krishna, Radiation Biology Branch, Center for Cancer Research, National Cancer Institute, Building 10, Room B3B69, NIH, 9000 Rockville Pike, Bethesda, MD 20892-1002. Phone: 301-496-7511; Fax: 301-480-2238 murali@helix.nih.gov.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

provided sufficient ^{13}C signal to be detected in the mouse in several organs, and can be used to detect ROS in vivo.

Graphical Abstract:

Keywords

hyperpolarized ^{13}C -MRI; spin trapping; DMPO: reactive oxygen species

INTRODUCTION:

Nitrone and nitroso based spin traps have been used to trap short lived free radicals with life times in the order of nanoseconds to stabilize as radical spin-adducts with longer half-lives (minutes) which can be detected by Electron Spin Resonance (EPR) spectroscopy (1). From the EPR spectra of the spin-adducts which accumulate to detectable levels, the identity of the short-lived intermediates can be confirmed (2). The EPR-spin trapping methodology has been used to detect, characterize and quantify short-lived intermediates generated in various radiolytic, photochemical, sonochemical, and enzymatic reactions (3–6). Nitron based spin traps such as DMPO, PBN (*N-tert*-Butyl- α -phenylnitrone) and POBN (α -(4-Pyridyl *N*-oxide)-*N-tert*-butylnitrone) have been widely used in chemical and cellular studies to detect free radical adducts using EPR spectroscopy (1,7–9). Newer derivatives of DMPO such as 5-(diethoxyphosphoryl)-5-methyl-1-pyrroline-*N*-oxide (DEPMPO), 2-ethoxycarbonyl-2-methyl-3,4-dihydro-2H-pyrrole-1-oxide (EMPO), 5-*tert*-butoxycarbonyl-5-methyl-1-pyrroline-*N*-oxide (BMPO) have been developed for enhanced sensitivity in detecting radicals where the spin adducts are longer lived (4,10–16). Nitrones have been shown to have minimal to no toxicity in vitro and in vivo (~200 mg/kg in mice, rats) and were effective as antioxidants in vitro and in vivo against various types of oxidative stress (5,17–20). Free radical adducts of nitrones such as PBN and POBN were detected after in vivo administration in mice which were exposed to whole body ionizing radiation and the protective effects were attributed to their radical scavenging ability (18,19,21). While EPR spectroscopy has been widely used to detect free radical intermediates, NMR based techniques and immune-spin trapping techniques have enabled the use of nitrones using NMR (22–24), immunology and MRI methods for the detection of free radical based end products in vitro and in vivo (23–28). Of the nitrones developed for spin trapping applications, the PBN derivative NXY-059 has been evaluated for clinical use in protecting against post ischemic reperfusion injury in ischemic stroke in brain (21,29–31).

Carmichael et al., conducted a systematic study to quantify the EPR-detectable radical adducts formed when a known quantity of free radicals were generated using ionizing radiation (32). Free radicals with poor reactivity such as superoxide were trapped with > 90% efficiency generating the corresponding spin-adduct (4,14) (33). Superoxide radical reactions occur primarily at the nitron bond of DMPO yielding the DMPO-O_2^- spin adducts which themselves are short lived and are converted to the DMPO-OH radical which is a more persistent radical (Eq 1, 2).

Conversely, nitron traps yielded radical adducts of the hydroxyl radical in the range of 14 – 35% in spite of diffusion limited rates of reaction (Eq 3).

The low yield of the spin adducts of these highly reactive radicals, regardless of the efficient rates of reaction were attributed to non-selective reaction on the spin trap other than at the nitron bond, which are not detectable by EPR (Eq 4).

Greater than 60% of the reactions of short lived radicals with nitrones result in intermediates which are also transient and ultimately form stable diamagnetic end products (Eq 5) (32).

To monitor such transient intermediates of DMPO reaction with highly reactive radicals, which are rapidly converted to diamagnetic end products, hyperpolarized MRI methods may be applied since the hyperpolarized spin states in DMPO may survive the brief paramagnetic state of the intermediates. While the paramagnetic intermediates cannot be detected, the diamagnetic end-products produced in reactive oxygen species (ROS) can be detected by ^{13}C MRI. The injected 5- ^{13}C -DMPO and the ROS mediated diamagnetic end-products will have distinct ^{13}C chemical shifts from which are distinguishable.

Hyperpolarization of the nuclear spin states using dynamic nuclear polarization (DNP) in ^{13}C labeled at sites with $T_1 \sim 30$ s provided > 10,000 fold enhancement in sensitivity for ^{13}C MRI and allowed metabolic studies of the conversion of these substrates to their products in vivo in pre-clinical and clinical settings. Requirements for a candidate molecule suitable for in vivo hyperpolarized MRI include: a) to be non-toxic at high bolus concentrations for in vivo administration; b) have a carbon atom in the molecule with T_1 longer than 20 s; c) high aqueous solubility (> 5 M); d) form a glass when mixed with the paramagnetic trityl radicals and frozen to enable DNP. Molecules which satisfy these conditions can be detected with

high sensitivity after DNP by in vivo ^{13}C MRI. Endogenous molecules such as pyruvate, fumarate, ketoglutarate etc., are examples of molecules where the carbonyl carbon atoms have been enriched with ^{13}C and polarized to provide 4 orders of magnitude increase in sensitivity. This enhanced sensitivity allowed detection and imaging of these substrates and their metabolic products in vivo. With the 4–5 orders of sensitivity enhancement provided by DNP, it is possible in practice to detect and image $\sim 5\text{ }\mu\text{M}$ of ROS-induced diamagnetic end products in vivo.

DMPO is liquid at room temperature and it forms glass when it is frozen with trityl radicals, making it suitable for DNP. In this study, we identified C-5 as a site in the molecule with a long T_1 for C atom, synthesized ^{13}C -labeled DMPO, and investigated feasibility of hyperpolarized ^{13}C -MRI and established conditions to image its biodistribution in vivo in a mouse. In vivo models of oxidative stress where high levels of reactive oxygen species (ROS) are generated focally, hyperpolarized MRI methods may become practical for the detection and imaging of free radical processes.

MATERIALS AND METHODS

Reagents

DMPO was purchased from Dojindo Laboratories. OX063 and the symmetric trityl radical (commonly known as the “Finland radical”) were purchased from GE Healthcare. Tris, ethylenediaminetetraacetic acid (EDTA), hypoxanthine, Fe-EDTA, SOD, and catalase were purchased from Sigma-Aldrich. Xanthine oxidase (XO) was purchased from Roche Molecular Biochemicals (Indianapolis, IN, USA). Isotopically labeled compounds were obtained from Cambridge Isotope Laboratories, Inc., MA. All other reagents and solvents were obtained from Sigma Aldrich, Inc. and used as received unless otherwise noted. Flash chromatography was performed on a Teledyne Isco CombiFlash Companion instrument with UV detection at 220 and 254nm. Analytical HPLC analysis was performed on an Agilent 1200 Series instruments equipped with multi wavelength detectors using an Agilent SB C18 column ($4.6 \times 50\text{mm}$, $3.5\text{ }\mu\text{m}$) with a flow rate of 1 mL/min . Solvent A was 0.05 % TFA in water, solvent B was 0.05 % TFA in ACN, and a linear gradient of 5% B to 95% B over 10 min was used. APCI mass spectrometry was performed on 6130 Quadrupole LC/MS Agilent Technologies instrument equipped with diode array detector. HRMS data were acquired using Waters XEVO G2-XS Q-ToF in ESI positive mode. ^1H and ^{13}C NMR spectra were recorded with a Varian spectrometer operating at 400MHz and 101 MHz respectively. Chemical shifts are reported in parts per million (δ) and are referenced to tetramethylsilane (TMS).

Synthesis of 5- ^{13}C -DMPO- d_9

Schemes of 5- ^{13}C -DMPO- d_9 synthesis are summarized in Figure 1.

Acetone-[2- ^{13}C , d_6] (3): A 3N LiOD solution in D_2O (0.625 mL) was added to Acetone-[2- ^{13}C] (10.0 g, 170 mmol) and D_2O (20.0 mL). The reaction mixture was stirred for 72h at RT. The acetone was distilled via short path distillation. The distillate was subjected to the same reaction conditions except that the stirring was for 48 h. The process was repeated after

second and third distillation. At the end of four cycles, 98.6 % deuteration of compound 3 (9.87 g, 152 mmol) was obtained in a 89% yield. ^{13}C NMR (101 MHz, D_2O) δ 8215.5 (C–2)

Isopropanol-[2- ^{13}C , d_7] (4): Ruthenium on carbon (5%, 4.8 g) was added to a solution of acetone-[2- ^{13}C , d_6] compound 3 (9.8 g, 150.6 mmol) in D_2O (40.0 mL) in a Parr reactor vessel. Air from the vessel was evacuated and replaced with deuterium gas at 45 PSI pressure. The reaction vessel was shaken for 16 h keeping the pressure of deuterium at 45 PSI. A small aliquot was analyzed by ^{13}C NMR and the reaction was found to be complete. The reaction mixture was filtered over a celite pad and the residue washed with deionized water. The combined filtrate was distilled via a short path distillation. The distillate obtained was the pure desired product 4 (9.11 g, 136 mmol) in an 90% yield. ^{13}C NMR (101 MHz, D_2O) δ 63.51 (C–2, t, J = 22.2 Hz)

2-Bromopropane-[2- ^{13}C , d_7] (5): A solution of 48 % DBr in D_2O (50.0 mL) was slowly added to Isopropanol-[2- ^{13}C , d_7] compound 4 (9.10 g, 136 mmol) at 0°C. The reaction mixture was stirred for 16h at RT. After a short path distillation the product 5 (13.2 g, 103 mmol) was obtained in a 76% yield. ^{13}C NMR (101 MHz, CDCl_3) δ 44.85 (C–2, t, J = 23.3 Hz)

2-Nitropropane-[2- ^{13}C , d_7] (6): Silver nitrite (16.20 g, 105.3 mmol) was added to a mixture of molecular sieves (4Å, 5g) and 2-bromopropane-[2- ^{13}C , d_7] compound 5, (11.5 g, 88.4 mmol) in 1,4-dioxane (100.0 mL). The mixture was stirred in the dark for 48 h. It was then analyzed by ^{13}C NMR and found to be complete. The reaction mixture was filtered and the residue washed with ether. The filtrates were combined and the solvent concentrated at reduced pressure, while keeping the temperature of water bath below 20°C. The dried product 6 (5.10 g, 53.6 mmol) was obtained in a 60% yield. ^{13}C NMR (101 MHz, DMSO) δ 78.21 (C–2, t, J = 22.8 Hz)

Methyl-4-methyl-4-nitro pentanoate-[2- ^{13}C , d_9] (8): Methyl acrylate-2,3,3- d_3 compound 7 (5.0 g; 53.6 mmol) was added to a solution of 2-nitropropane-[2- ^{13}C , d_7] compound 6, (5.1 g; 52.5 mmol) and Triton-B (40% in H_2O , 2.0 mL) in 1,4-dioxane (100.0 mL) at 70°C for 60 min. The temperature of the reaction was then raised to 85°C and maintained for 3 h. The solution was cooled to RT, diluted with 200 mL DCM and washed with 1 N HCl (300ml). The aqueous phase was then washed with DCM (3×100 mL). All organic fractions were combined, dried with magnesium sulfate and concentrated. The oil 8 (5.39g, 29.1 mmol) was obtained in a 52% yield. $\delta^{13}\text{C}$ NMR (101 MHz, CDCl_3)¹⁵ δ 86.8 (C–4), 51.72(–OCH₃), 28.4 (C–3), 24.8 (C–2)

4-Methyl-4-nitro pentanal-[2- ^{13}C , d_9] (9): DIBAL-H (1M in DCM, 32 mL, 32 mmol) was added over 30 min. to a mixture of 4 Å molecular sieves (2.0 g), and nitropentanoate 8 (4.74 g; 25.6 mmol) in DCM (45 mL) at –78°C. The reaction mixture was stirred for 1h at –78°C under an atmosphere of Ar. It was then quenched with addition of wet silica (20 g silica + 10 g water). The temperature was raised to RT. The reaction mixture was then filtered and the residue washed with 3 × 50 mL DCM. All DCM fractions were combined and washed with 300 mL 1N HCl solution. Aqueous phases were extracted with 2 × 100 mL

DCM. Organic fractions were combined, dried with MgSO_4 , and concentrated provide product 9 (3.59 g, 23.1 mmol) in an 90% yield. ^{13}C NMR (101 MHz, CDCl_3) δ 199.8 (–CHO), 86.70 (C–4); Lit¹⁶: ^{13}C NMR (75.47 MHz, CDCl_3) δ 25.61 (– CH_3), 32.09 (C–3), 38.60 (C–2), 87.09 (C–4), 199.68 (–CHO).

Dimethyl pyrroline-N-oxide-[5- ^{13}C , d_9] (10): Acetic acid- d_4 (5.98 g, 93.5 mmol) was added over 1 h to a mixture of activated zinc dust (3.05 g; 47.7 mmol, 2.1 eq.) and pentanal (3.45 g, 22.2 mmol, 1.0 eq.) in ethanol- d (50 mL) at 0°C under Ar. The reaction mixture was then stirred at 4°C for 44 h. It was then diluted with cold ethanol (20 mL) and filtered. The residue was washed with 4×75 mL cold ethanol. The ethanol phases were combined and concentrated. The crude syrup obtained was chromatographed twice on a normal phase silica (100 % DCM to 9:1 DCM-MeOH) to provide DMPO 10 (0.891 g, 7.23 mmol) in a 32% yield. ^{13}C NMR (101 MHz, CDCl_3) δ 73.1 (C–5); Lit¹⁷: ^{13}C NMR (101 MHz, CDCl_3): δ 23.9 (C–3), 25.2 (– CH_3), δ 33.8 (C–4), δ 72.8 (C–5), 129.0 (*nitronyl*). ESI-MS: 124.1 (M⁺).

The NMR data is as follows: ^1H NMR (400 MHz) CDCl_3 δ 6.79 (dd, $J = 6.7, 2.7$ Hz, 1H), 2.55 (d, $J = 4.4$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ 73.1 (C–5); Lit¹⁷: ^{13}C NMR (101 MHz, CDCl_3): δ 23.9 (C–3), 25.2 (– CH_3), δ 33.8 (C–4), δ 72.8 (C–5), 129.0 (*nitronyl*). HRMS (ESI/Q-TOF) m/z : $[\text{M}+\text{H}]^+$ Calcd for $\text{C}_5^{13}\text{CH}_3\text{D}_9\text{NO}^+$ 124.1512; Found 124.1513

Animal studies

All animal experiments were carried out in compliance with the Guide for the care and use of laboratory animal resources (National Research Council, 1996) and approved by the National Cancer Institute Animal Care and Use Committee (NCI-CCR-ACUC (Bethesda). Female C3H/Hen mice were supplied by the Frederick Cancer Research Center, Animal Production (Frederick, MD).

The mice were anesthetized with isoflurane (4% for induction and 1.5–2.5 % for maintaining anesthesia) in medical air (500 mL/min) and were positioned prone in a MRI scanner. During MRI scans, the breathing rate of the mouse was monitored with a pressure transducer (SA Instruments Inc., Stony Brook, NY) and was maintained at 70 ± 15 breaths per minute. Core body temperature was also monitored with a non-magnetic rectal temperature probe (FISO Technologies, INC., Quebec, Canada) and was maintained at $36 \pm 1^\circ\text{C}$ with a flow of warm water. For administration of 5- ^{13}C -DMPO- d_9 , a 30-gauge needle was cannulated into the tail vein and was extended using polyethylene tubing (PE-10).

Hyperpolarized ^{13}C -MRI studies

MRI scans were performed on a 3 T scanner controlled with EasyScan and PowerScan (MR Solutions, Acton, MA), a 4.7 T scanner controlled with ParaVision 5.1 (Bruker Bio-Spin MRI GmbH), or a 3 T scanner (Achieva 3T-TX, Philips Healthcare, Best, Netherland). A 17 mm home-built ^{13}C solenoid coil placed inside of a saddle coil for ^1H and a 35 mm ^{13}C - ^1H saddle coil was used for phantom and a mouse, respectively. In the MRI scans of mice body, T_2 -weighted anatomical images were obtained using a fast spin echo sequence (FSET2) with TE of 13 ms, TR of 2,500 ms, 8 slices, 2 mm thickness, resolution of 0.25×0.25 mm.

5- ^{13}C -DMPO- d_9 (30 μL) containing 15 mM symmetric trityl and 2.5 mM gadolinium chelate ProHance (Bracco Diagnostics, Milano, Italy) was hyperpolarized using the Hypersense DNP polarizer (Oxford Instruments, Abingdon, UK) as described previously. After 60–90 min, the hyperpolarized sample was rapidly dissolved in 4.5 mL of PBS containing 100 mg/L EDTA. The hyperpolarized 5- ^{13}C -DMPO- d_9 solution (60 mM) was intravenously injected through a catheter placed in the tail vein of the mouse (12 $\mu\text{L/g}$ body weight). For the dynamic study, ^{13}C spectra in mouse body were acquired every 1 s for 240 s. ^{13}C two-dimensional spectroscopic images were also acquired 25–47 s after the beginning of the DMPO injection, with a 32×32 mm, field of view in a 8 mm coronal slice through the body, a matrix size of 16×16 , spectral width of 3330 Hz, repetition time of 85 ms, and excitation pulse with a flip angle of 10° . The total time required to acquire each image was 23 s. In the MRI scans of mice head, 5- ^{13}C -DMPO- d_9 was hyperpolarized using the SPINlab (GE Healthcare) for 3–4 hours, and the scans were performed using the Philips 3 T scanner. ^{13}C two-dimensional spectroscopic images were acquired 30 s after the beginning of the DMPO injection, with a 28×28 mm, field of view in a 10 mm axial slice through the head, a matrix size of 14×14 , spectral width of 3333 Hz, repetition time of 86 ms, and excitation pulse with a flip angle of 3° .

RESULTS

Synthesis of ^{13}C -labeled DMPO (Deepak)

Preliminary ^{13}C NMR spectroscopy studies conducted using a 4.7 T MRI scanner identified that of all carbon atoms, the carbon atom at the 5-position of DMPO (Figure 2A) with a chemical shift of 76 ppm (Figure 2B) had the longest T_1 (Figure 2C). Since DMPO is amphipathic, we tested the polarization efficiency using the Finland-COOH radical, which is also an amphipathic paramagnetic probe. A frozen glass mixture of the DMPO and the Finland-COOH radical was polarized for 60 minutes and rapidly dissolved to room temperature and measuring its NMR spectrum immediately. A significant enhancement of the signal intensity of the ^{13}C NMR signal was observed (Figure 2D) compared to the signals from DMPO polarized with OX063 (Figure 2B). Based on these observations, DMPO enriched with ^{13}C -labeled at C5 position and deuterated was synthesized to further prolong the T_1 relaxation time.

Since the initial synthesis of unlabeled DMPO (34,35) several other researchers have synthesized DMPO and its analogs using various routes (36–42). Based on works of Beth et al. (43) and Janzen et al. (38), synthesis of labeled DMPO as a spin trap agent was first explored by Pou et al. (37). In order to prepare DMPO with ^{13}C at position C5 and deuterated protons on neighboring carbons (to prolong the T_1 relaxation times) we followed the procedure of Le et al. (41) and Leinisch et al. (42). We found that the ^{13}C -C2 deuterated three carbon starting material could be effectively prepared from; the h-d exchange on acetone ^{13}C -C2 as described in the literature (44) under basic conditions. Per-deuterated acetone- ^{13}C -C2 was then reduced to isopropanol- ^{13}C -C2 with Pd/C under D_2 atmosphere. Labeled isopropanol was brominated with DBr, converted to 2-nitropropane with silver nitrite and then alkylated with methyl acrylate-2,3,3- d_3 .

The resulting nitropentanoate was reduced to aldehyde with DIBAL-H. This was followed by treatment with activated zinc and acetic acid to yield the cyclic nitron, DMPO. However, the resulting DMPO was not fully deuterated due to the DIBAL reduction. The overall yield achieved for DMPO-d₉ was 15%. Figure 1 shows the overall synthetic scheme followed.

Hyperpolarization of ¹³C-DMPO

For successful polarization, the molecule to be polarized should dissolve at high concentrations (~ 5 M) and form a homogenous glass with the trityl radical when frozen. Of the four trityl radical analogs for hyperpolarization of ¹³C-DMPO, namely OX063, Finland trityl, OX063-HCl, Finland-COOH, Finland-COOH was found to be the most suited for glass formation and polarization of DMPO. Figure 3A shows the polarization build-up of DMPO with natural abundance ¹³C (open circles) and 5-¹³C-DMPO-d₉ (closed circles) with 15 mM Finland-COOH and 2.5 mM Gd chelate ProHance. The time constant of polarization build-up in the hyperpolarizer was found to be ~ 23 min. After the polarization was maximal, the hyperpolarized samples were dissolved in PBS containing EDTA, and ¹³C-spectra were measured every 1 s using the 4.7 T scanner with 10 deg flip angle. The T₁ relaxation time of non-enriched/non-deuterated DMPO and 5-¹³C-DMPO-d₉ was 26 s and 60 s, respectively, suggesting that deuteration of DMPO extended the T₁ significantly (Figure 3B). The T₁ of 5-¹³C-DMPO-d₉ was similar to that of 1-¹³C-pyruvic acid, which is the most frequently used probe in hyperpolarized ¹³C-MRI in vivo, suggesting ¹³C-DMPO has sufficiently long T₁ to permit in vivo studies. With the known toxicological data of DMPO from prior studies where large bolus doses have no adverse effects (20) and with the long T₁ of the 5-C atom, the results suggest that imaging of hyperpolarized DMPO in vivo monitoring its biodistribution and its transformation is feasible.

In Vivo ¹³C MRI of hyperpolarized 5-¹³C-DMPO-d₉

The in vivo behavior of DMPO was tested in a mouse where the DMPO after dissolution of the hyperpolarized mixture was administered through a tail vein cannula of a mouse placed in a double tuned ¹H/¹³C coil and the ¹³C signal monitored every 1 s using the 3 T scanner with 10 deg flip angle pulse. The signal of 5-¹³C-DMPO-d₉ was detected in the mouse body immediately after its injection and detectable for more than 150 s (Figure 4A). Figure 4B shows the signal intensity curve of 5-¹³C-DMPO-d₉ after iv administration of the hyperpolarized sample in the mouse. From the signal intensity curve, the T₁ decay of 5-¹³C-DMPO-d₉ in mouse was estimated to be 29s. The persistence of the hyperpolarized signal in vivo for this duration points to the feasibility of imaging DMPO biodistribution and possible use for monitoring its transformation either enzymatically or through reactive oxygen species.

We carried out ¹³C-chemical shift imaging in a mouse body after iv administration of hyperpolarized 5-¹³C-DMPO-d₉. Figure 5A shows consecutive anatomical slices (2 mm thickness each) of the mouse body. Figure 5B shows the ¹³C-CSI overlaid on the anatomic image obtained from 8mm slice thickness of the mouse body 25–47 s after hyperpolarized 5-¹³C-DMPO-d₉ injection. The DMPO map obtained from the chemical shift image (Figure 5C) revealed that ¹³C-DMPO was distributed through the mouse body within a minute of its injection. The signal was higher in the heart/lung region and kidney, whereas the signal from

liver was relatively low compared to the other region (Figure 5D, E). To obtain a higher signal in the mouse liver, we did slice selected ^{13}C -MRS by averaging 100 transients (Figure 5 F). Signal to noise ratio was much better compared to ^{13}C -CSI. The relative levels of DMPO in the heart, kidney and liver regions are shown in Figure 5G. No metabolites were detected in the normal mouse liver suggesting that enzymatic conversion of DMPO was minimal.

Several studies showed that DMPO exhibits anti-oxidative and anti-inflammatory effects in vivo (45,46). A recent study showed that DMPO injected directly into the intranigral region in rats afforded protection against 6-OH-Dopamine mediated neurotoxicity (10). With the enhanced sensitivity provided by hyperpolarization, it is possible to examine the biodistribution of DMPO in the brain following systemic administration intravenously. A tail vein cannula was placed in an anesthetized mouse placed in a doubly tuned resonator encompassing the head region. Figure 6A shows the ^{13}C -CSI overlayed on the anatomic image in an axial profile. DMPO biodistribution in the brain was monitored immediately after hyperpolarized DMPO was administered (Figure 6B). The ^{13}C image shows evidence of DMPO accumulation in the brain region suggesting that it penetrates the blood-brain barrier rapidly and accumulates at effective concentrations while retaining polarization. This image represents the first imaging evidence for the localization of DMPO in the brain and may explain the observed antioxidant effects it exerts.

DISCUSSION:

DNP of low γ nuclei in molecules enriched with ^{13}C , ^{15}N has allowed imaging of endogenous molecules and their biochemical transformations in vivo. ^{13}C enriched pyruvate, fumarate, urea, glucose, gluconolactate, bicarbonate, have been successfully imaged in vivo in various animal models to obtain information pertaining to metabolism, perfusion and pH. Exogenous molecules such as ascorbate, glycerate, methyl glutamine were also developed and tested for probe for perfusion, pH. Most probes report on biochemical, physiologic, and microenvironmental features. Dehydroascorbate has been used to probe the levels of ROS by monitoring its transformation to ascorbate and determining the ration of DHA/AA (47). However, this transformation being a 2-electron process, the ratio may also be significantly influenced by the tissue redox status. The search for hyperpolarizable antioxidants which can report on oxidative stress and provide noninvasive capabilities to monitor these reactions led to considering nitrones as candidate probes for the following reasons: a) Nitrones can have carbon atoms which can have potentially long ^{13}C T1s. b) Nitrones are non-toxic even at high bolus doses typically used for HP MRI; c) NXY-059, a nitron derivative has been characterized to have anti-oxidant activity and has been developed and used for human applications to limit post-ischemic reperfusion injury (21). DMPO shares many of these features and has been successfully polarized. DMPO has long been used as a spin trap to scavenge reactive oxygen species, identify the intermediates and quantify them in simple chemical systems using EPR spectroscopy. However, in cellular studies, it was of limited use since the spin adducts undergo rapid reduction to diamagnetic products which are EPR invisible. NMR-Spin trapping overcame some of these limitations. In vivo detection of ROS was demonstrated by EPR by administering nitron spin traps and analyzing the tissue extracts (22,24). Direct in vivo detection of spin adducts was also demonstrated using POBN

by using low-frequency EPR by exposing tumors to high radiation doses (19). These approaches however, cannot be translated to in vivo imaging because of poor sensitivity.

HP-MRI with 5-¹³C-DMPO-d₉ has the potential to provide imaging assessment of reactive oxygens species because: a) the signals of the original molecule can be significantly enhanced by the DNP method; b) the molecule biodistribution in vivo is effective to have adequate global concentrations in organs/regions of interest; c) appropriate models of ROS mediated conversion of the DMPO to stable diamagnetic products proceeds with sufficient efficiency to have detectable levels of products with characteristic chemical shifts. The hyperpolarized DMPO biodistribution monitored by ¹³C MRI revealed that significant levels were attained in heart, kidney, brain with detectable levels in the liver and muscle. The demonstration of the in vivo biodistribution of DMPO, by ¹³C MRI using hyperpolarization techniques sets the stage for oxidative stress evaluation in vivo. With suitable in vivo models of focal ROS generation at high levels such as post-ischemic reperfusion injury or systemic ROS generation such as with lipopolysaccharide challenge, this method can be applied for in vivo imaging of oxidative stress.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

REFERENCES:

1. Janzen EG. Spin Trapping. *Accounts Chem Res* 1971;4(1):31-& 10.1021/ar50037a005.
2. Buettner GR. Spin trapping: ESR parameters of spin adducts. *Free Radic Biol Med* 1987;3(4):259–303. [PubMed: 2826304]
3. Villamena FA, Zweier JL. Detection of reactive oxygen and nitrogen species by EPR spin trapping. *Antioxid Redox Signal* 2004;6(3):619–29 10.1089/152308604773934387. [PubMed: 15130289]
4. Zhang H, Joseph J, Vasquez-Vivar J, Karoui H, Nsanzumuhire C, Martasek P, et al. Detection of superoxide anion using an isotopically labeled nitron spin trap: potential biological applications. *FEBS Lett* 2000;473(1):58–62. [PubMed: 10802059]
5. Halpern HJ, Yu C, Barth E, Peric M, Rosen GM. In situ detection, by spin trapping, of hydroxyl radical markers produced from ionizing radiation in the tumor of a living mouse. *Proc Natl Acad Sci U S A* 1995;92(3):796–800. [PubMed: 7846054]
6. Riesz P, Kondo T, Carmichael AJ. Sonochemistry of acetone and acetonitrile in aqueous solutions. A spin trapping study. *Free Radic Res Commun* 1993;19 Suppl 1:S45–53. [PubMed: 8282231]
7. McCormick ML, Buettner GR, Britigan BE. The spin trap alpha-(4-pyridyl-1-oxide)-N-tert-butyl nitron stimulates peroxidase-mediated oxidation of deferoxamine. Implications for pharmacological use of spin-trapping agents. *J Biol Chem* 1995;270(49):29265–9. [PubMed: 7493957]
8. Buettner GR, Britigan BE. The spin trapping of superoxide with M4PO (3,3,5,5-tetramethylpyrroline-N-oxide). *Free Radic Biol Med* 1990;8(1):57–60. [PubMed: 2157635]
9. Janzen EG. ESR Detection and Identification of Short-Lived Free Radicals by Spin Trapping. *Appl Spectrosc* 1971;25(1):139-&.
10. Kitamura Y, Kamibayashi M, Inden M, Yanagida T, Shibaike T, Takata K, et al. Detoxification of 6-hydroxydopamine-induced Parkinsonian neurodegeneration by G-CYPMPO, a novel radical trapper. *Neurochem Int* 2011;58(6):721–7 10.1016/j.neuint.2011.02.013. [PubMed: 21333707]
11. Xu Y, Kalyanaraman B. Synthesis and ESR studies of a novel cyclic nitron spin trap attached to a phosphonium group-a suitable trap for mitochondria-generated ROS? *Free Radic Res* 2007;41(1):1–7 10.1080/10715760600911147. [PubMed: 17164173]

12. Kamibayashi M, Oowada S, Kameda H, Okada T, Inanami O, Ohta S, et al. Synthesis and characterization of a practically better DEPMPO-type spin trap, 5-(2,2-dimethyl-1,3-propoxy cyclophosphoryl)-5-methyl-1-pyrroline N-oxide (CYPMPO). *Free Radic Res* 2006;40(11):1166–72. 10.1080/10715760600883254. [PubMed: 17050170]
13. Clement JL, Finet JP, Frejaville C, Tordo P. Deuterated analogues of the free radical trap DEPMPO: synthesis and EPR studies. *Org Biomol Chem* 2003;1(9):1591–7. [PubMed: 12926292]
14. Zhao H, Joseph J, Zhang H, Karoui H, Kalyanaraman B. Synthesis and biochemical applications of a solid cyclic nitron spin trap: a relatively superior trap for detecting superoxide anions and glutathyl radicals. *Free Radic Biol Med* 2001;31(5):599–606. [PubMed: 11522444]
15. Timmins GS, Liu KJ, Bechara EJ, Kotake Y, Swartz HM. Trapping of free radicals with direct in vivo EPR detection: a comparison of 5,5-dimethyl-1-pyrroline-N-oxide and 5-diethoxyphosphoryl-5-methyl-1-pyrroline-N-oxide as spin traps for HO* and SO4*. *Free Radic Biol Med* 1999;27(3–4):329–33. [PubMed: 10468206]
16. Frejaville C, Karoui H, Tuccio B, Le Moigne F, Culcasi M, Pietri S, et al. 5-(Diethoxyphosphoryl)-5-methyl-1-pyrroline N-oxide: a new efficient phosphorylated nitron for the in vitro and in vivo spin trapping of oxygen-centered radicals. *J Med Chem* 1995;38(2):258–65. [PubMed: 7830268]
17. Villamena FA, Das A, Nash KM. Potential implication of the chemical properties and bioactivity of nitron spin traps for therapeutics. *Future Med Chem* 2012;4(9):1171–207. 10.4155/fmc.12.74. [PubMed: 22709256]
18. Young HK, Floyd RA, Maidt ML, Dynlacht JR. Evaluation of nitron spin-trapping agents as radioprotectors. *Radiat Res* 1996;146(2):227–31. [PubMed: 8693073]
19. Lai EK, Crossley C, Sridhar R, Misra HP, Janzen EG, McCay PB. In vivo spin trapping of free radicals generated in brain, spleen, and liver during gamma radiation of mice. *Arch Biochem Biophys* 1986;244(1):156–60. [PubMed: 3004337]
20. Khan N, Wilmot CM, Rosen GM, Demidenko E, Sun J, Joseph J, et al. Spin traps: in vitro toxicity and stability of radical adducts. *Free Radic Biol Med* 2003;34(11):1473–81. [PubMed: 12757857]
21. Floyd RA, Kopke RD, Choi CH, Foster SB, Doblas S, Towner RA. Nitrones as therapeutics. *Free Radic Biol Med* 2008;45(10):1361–74. 10.1016/j.freeradbiomed.2008.08.017. [PubMed: 18793715]
22. Khramtsov VV, Reznikov VA, Berliner LJ, Litkin AK, Grigor'ev IA, Clanton TL. NMR spin trapping: detection of free radical reactions with a new fluorinated DMPO analog. *Free Radic Biol Med* 2001;30(10):1099–107. [PubMed: 11369499]
23. Berliner LJ, Khramtsov V, Fujii H, Clanton TL. Unique in vivo applications of spin traps. *Free Radic Biol Med* 2001;30(5):489–99. [PubMed: 11182519]
24. Khramtsov V, Berliner LJ, Clanton TL. NMR spin trapping: detection of free radical reactions using a phosphorus-containing nitron spin trap. *Magn Reson Med* 1999;42(2):228–34. [PubMed: 10440946]
25. Towner RA, Saunders D, Smith N, Towler W, Cruz M, Do S, et al. Assessing long-term neuroinflammatory responses to encephalopathy using MRI approaches in a rat endotoxemia model. *Geroscience* 2018;40(1):49–60. 10.1007/s11357-018-0009-z. [PubMed: 29417380]
26. Mason RP. Imaging free radicals in organelles, cells, tissue, and in vivo with immuno-spin trapping. *Redox Biol* 2016;8:422–9. 10.1016/j.redox.2016.04.003. [PubMed: 27203617]
27. Towner RA, Smith N, Saunders D, Lupu F, Silasi-Mansat R, West M, et al. In vivo detection of free radicals using molecular MRI and immuno-spin trapping in a mouse model for amyotrophic lateral sclerosis. *Free Radic Biol Med* 2013;63:351–60. 10.1016/j.freeradbiomed.2013.05.026. [PubMed: 23722162]
28. Towner RA, Smith N, Saunders D, De Souza PC, Henry L, Lupu F, et al. Combined molecular MRI and immuno-spin-trapping for in vivo detection of free radicals in orthotopic mouse GL261 gliomas. *Biochim Biophys Acta* 2013;1832(12):2153–61. 10.1016/j.bbadis.2013.08.004. [PubMed: 23959048]
29. de Souza PC, Smith N, Pody R, He T, Njoku C, Silasi-Mansat R, et al. OKN-007 decreases VEGFR-2 levels in a preclinical GL261 mouse glioma model. *Am J Nucl Med Mol Imaging* 2015;5(4):363–78. [PubMed: 26269774]

30. de Souza PC, Balasubramanian K, Njoku C, Smith N, Gillespie DL, Schwager A, et al. OKN-007 decreases tumor necrosis and tumor cell proliferation and increases apoptosis in a preclinical F98 rat glioma model. *J Magn Reson Imaging* 2015;42(6):1582–91 10.1002/jmri.24935. [PubMed: 25920494]
31. Coutinho de Souza P, Smith N, Atolagbe O, Ziegler J, Njoku C, Lerner M, et al. OKN-007 decreases free radical levels in a preclinical F98 rat glioma model. *Free Radic Biol Med* 2015;87:157–68 10.1016/j.freeradbiomed.2015.06.026. [PubMed: 26119786]
32. Carmichael AJ, Makino K, Riesz P. Quantitative aspects of ESR and spin trapping of hydroxyl radicals and hydrogen atoms in gamma-irradiated aqueous solutions. *Radiat Res* 1984;100(2):222–34. [PubMed: 6093187]
33. Ben-Hur E, Carmichael A, Riesz P, Rosenthal I. Photochemical generation of superoxide radical and the cytotoxicity of phthalocyanines. *Int J Radiat Biol Relat Stud Phys Chem Med* 1985;48(5): 837–46. [PubMed: 2997064]
34. Brown RFC, Clark VM, Todd A. DELTA-1-PYRROLINE N-OXIDES. *Proceedings of the Chemical Society of London* 1957(3):97–8.
35. Bonnett R, Clark VM, Giddey A, Todd A. EXPERIMENTS TOWARDS THE SYNTHESIS OF CORRINS .1. THE PREPARATION AND REACTIONS OF SOME DELTA-1-PYRROLINES - A NOVEL PROLINE SYNTHESIS. *Journal of the Chemical Society* 1959(JUN):2087–93 10.1039/jr9590002087.
36. Janzen EG, Zhang YK, Arimura M. SYNTHESIS AND SPIN-TRAPPING CHEMISTRY OF 5,5-DIMETHYL-2-(TRIFLUOROMETHYL)-1-PYRROLINE N-OXIDE. *Journal of Organic Chemistry* 1995;60(17):5434–40 10.1021/jo00122a021.
37. Pou S, Rosen GM, Wu YX, Keana JFW. SYNTHESIS OF DEUTERIUM-CONTAINING AND N-15-CONTAINING PYRROLINE 1-OXIDES - A SPIN TRAPPING STUDY. *Journal of Organic Chemistry* 1990;55(14):4438–43 10.1021/jo00301a042.
38. Janzen EG, Zhang YK, Haire DL. SYNTHESIS OF A NOVEL NITRONE, 2-PHENYL-5,5-DIMETHYL-1-PYRROLINE N-OXIDE (NITRONYL-C-13), FOR ENHANCED RADICAL ADDEND RECOGNITION AND SPIN ADDUCT PERSISTENCE. *Journal of the American Chemical Society* 1994;116(9):3738–43 10.1021/ja00088a009.
39. Haire DL, Hilborn JW, Janzen EG. A MORE EFFICIENT SYNTHESIS OF DMPO-TYPE (NITRONE) SPIN TRAPS. *Journal of Organic Chemistry* 1986;51(22):4298–300 10.1021/jo00372a039.
40. Golubev VA, Shilov GV, Sen VD. Synthesis of new functionalized 5,5-dimethyl-1-pyrroline 1-oxides and their investigation as spin traps. *Russian Chemical Bulletin* 2010;59(11):2081–5 10.1007/s11172-010-0358-y.
41. Le DD, Zhang YS, Nguyen D, Moravek J. Synthesis of 5,5-dimethyl-1-pyrroline-N-oxide-2-C-14. *Journal of Labelled Compounds & Radiopharmaceuticals* 2000;43(11):1107–11 10.1002/1099-1344(20001015)43:11<1107::aid-jlcr396>3.3.co;2-f
42. Leinisch F, Jiang J, Deterding LJ, Mason RP. Simplified synthesis of isotopically labeled 5,5-dimethyl-pyrroline N-oxide. *Molecules* 2011;16(10):8428–36 10.3390/molecules16108428. [PubMed: 21986521]
43. Beth AH, Venkataramu SD, Balasubramanian K, Dalton LR, Robinson BH, Pearson DE, et al. N-15-SUBSTITUTED AND H-2-SUBSTITUTED MALEIMIDE SPIN LABELS - IMPROVED SENSITIVITY AND RESOLUTION FOR BIOLOGICAL ELECTRON-PARAMAGNETIC-RES STUDIES. *Proceedings of the National Academy of Sciences of the United States of America-Biological Sciences* 1981;78(2):967–71 10.1073/pnas.78.2.967.
44. Paulsen PJ, Cooke WD. PREPARATION OF DEUTERATED SOLVENTS FOR NUCLEAR MAGNETIC RESONANCE SPECTROMETRY. *Analytical Chemistry* 1963;35(10):1560-& 10.1021/ac60203a072.
45. Munoz MD, Della Vedova MC, Bushel PR, Ganini da Silva D, Mason RP, Zhai Z, et al. The nitron spin trap 5,5-dimethyl-1-pyrroline N-oxide dampens lipopolysaccharide-induced transcriptomic changes in macrophages. *Inflamm Res* 2018;67(6):515–30 10.1007/s00011-018-1141-z. [PubMed: 29589052]

46. Zhai Z, Gomez-Mejiba SE, Ramirez DC. The nitron spin trap 5,5-dimethyl-1-pyrroline N-oxide affects stress response and fate of lipopolysaccharide-primed RAW 264.7 macrophage cells. *Inflammation* 2013;36(2):346–54 10.1007/s10753-012-9552-4. [PubMed: 23053730]
47. Keshari KR, Sai V, Wang ZJ, Vanbrocklin HF, Kurhanewicz J, Wilson DM. Hyperpolarized [1–¹³C]dehydroascorbate MR spectroscopy in a murine model of prostate cancer: comparison with ¹⁸F-FDG PET. *J Nucl Med* 2013;54(6):922–8 10.2967/jnumed.112.115402. [PubMed: 23575993]

Highlights

- Spin traps/radical scavengers cannot be currently imaged by MRI
- ^{13}C labeled spin trap DMPO can be imaged in vivo after hyperpolarization
- Its biodistribution was followed in vivo after hyperpolarization.
- ^{13}C MRI with hyperpolarized DMPO can be used for in vivo free radical detection/imaging.

Scheme 1: Synthesis of 2-Nitropropane-[2- ^{13}C ; d_7].

Scheme 2: Synthesis of Methyl 4-methyl-4-nitropentanoate-[2- ^{13}C ; 2,3,3- d_3]

Scheme 3: Synthesis of DMPO-[5- ^{13}C , d_9]

Figure 1:
Summary of schemes of 5- ^{13}C -DMPO- d_9 synthesis.

Figure 2:

(A) Position of ^{13}C -labeled in DMPO which has the longest T_1 . (B) ^{13}C -MR spectrum of unlabeled DMPO hyperpolarized with OX063. Single peak was detected at 76 ppm which was from ^{13}C atom at the 5-position of DMPO. (C) The signal decay after dissolution in PBS. T_1 was calculated to be 26 s from the decay curve. (D) ^{13}C -MR spectrum of unlabeled DMPO with natural abundance ^{13}C , hyperpolarized with Finland-COOH. The signal to noise ratio of DMPO peak was remarkably increased by using Finland-COOH.

Figure 3:

(A) Hyperpolarization build-up of unlabeled DMPO (○) and ^{13}C -labeled DMPO- d_9 (●) with 15 mM Finland-COOH and 2.5 mM Gd chelate ProHance. The build-up was monitored by an NMR spectroscopy in the hyperpolarizer. (B) The decay of ^{13}C -signal at 76 ppm of unlabeled DMPO (○) and ^{13}C -labeled DMPO- d_9 (●). Deuteration of DMPO extended the T_1 of the ^{13}C signal from 26 s to 60 s.

Figure 4:
(A) Dynamic ^{13}C -MR spectra in a mouse body after hyperpolarized 5- ^{13}C DMPO- d_9 injection. The spectra were acquired every 1 s for 240 s. DMPO peak (76 ppm) was observed. (B) Signal intensity curve of DMPO calculated from the spectra. The T_1 was estimated to be 29 s from the curve.

Figure 5:

(A) T_2 -weighted anatomical images of a mouse body (4 adjacent slices). The thickness of each slice was 2 mm. (B) ^{13}C -chemical shift image (CSI) in the mouse body acquired 25 s after hyperpolarized 5- ^{13}C -DMPO- d_9 injection. The ^{13}C -CSI was obtained from 8 mm thickness slice, and the size of each pixel was 2 mm $\times$ 2 mm. (C) Distribution of DMPO in the mouse body calculated from the ^{13}C -CSI. The positions of heart/lung, liver, and kidney were obtained from the T_2 -weighted anatomy. (D, E) ^{13}C -MR spectrum of heart and left kidney obtained from the ^{13}C CSI. (F) ^{13}C -MR spectrum from a mouse liver with 100 times

averaging. (G) Averaged signal intensity of ^{13}C -DMPO in each organ calculated from the DMPO distribution map.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 6:
(A) ^{13}C -CSI in the mouse head acquired 30 s after hyperpolarized 5- ^{13}C -DMPO- d_9 injection. The CSI was obtained from 10 mm thickness slice, and overlaid on a 2 mm thickness of T_2 -weighted anatomical image. (B) Distribution of DMPO in the mouse head calculated from the ^{13}C -CSI.