Published in final edited form as:

J Am Chem Soc. 2019 February 13; 141(6): 2268–2273. doi:10.1021/jacs.9b00237.

Ir-Catalyzed Intermolecular Branch-Selective Allylic C-H Amidation of Unactivated Terminal Olefins

Honghui Lei, Tomislav Rovis*

Department of Chemistry, Columbia University, New York, New York 10027, United States

Abstract

An efficient method for intermolecular branch-selective allylic C-H amidation has been accomplished via Ir(III) catalysis. The reaction proceeds through initial allylic C-H activation, supported by the isolation and crystallographic characterization of an allyl-Ir(III) intermediate, followed by a subsequent oxidative amidation with readily available dioxazolones as nitrenoid precursors. A diverse range of amides are successfully installed at the branched position of terminal alkenes in good yields and regioselectivities. Importantly, the reaction allows the use of amide-derived nitrenoid precursors avoiding problematic Curtius-type rearrangements.

Graphical Abstract

Due to the ubiquity of nitrogen-containing functionalities in both natural and synthetic bioactive molecules,¹ C-N bond formation reactions are some of the most frequently used transformations in medicinal chemistry.² In addition to classic strategies which usually require prefunctionalization, direct amination of C-H bonds could dramatically simplify synthetic routes, providing more straightforward disconnections for amine synthesis.³ A number of reports have recently demonstrated chelate-assisted C-H amination; however, these methods usually employ pre-installed coordinating directing groups that limit their general application.⁴ In contrast, allylic C-H amination only involves a simple alkene as the “directing group” which is not only prevalent in a variety of compounds but is also easily manipulated with a diverse range of robust methods.

Current strategies in allylic C-H amination utilize either C-H insertion by a metal nitrenoid⁵ or nucleophilic amination of allylmetal species generated via C-H activation.⁶ Despite these

*Corresponding Author tr2504@columbia.edu.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website. Detailed experimental procedures, characterization data, copies of ¹H NMR and ¹³C NMR spectra for all isolated compounds (PDF)

The authors declare no competing financial interest.

advances, controlling chemoselectivity and regioselectivity remains challenging in intermolecular allylic C-H amination reactions. For instance, terminal olefins that react with a metal-nitrenoid species suffer from poor chemoselectivity between aziridination of the alkene and desired C-H amination.^{5h,7} (Figure 1a(i)). An alternative strategy was independently reported by the White^{6d} and Liu^{6e} groups in 2008, demonstrating that linear amination could be achieved under Pd(II) catalysis (Figure 1a(ii)). The reaction is proposed to proceed through an allyl-Pd(II) intermediate that is generated via allylic C-H activation, followed by reductive quenching with a nitrogen nucleophile at the less hindered terminal position. Based on these initial reports, many modifications and improvements have been developed in recent years,⁸ offering various choices for the synthesis of linear allylic amines. On the other hand, the branch-selective allylic C-H amination of terminal olefins has been achieved through an intramolecular manner.^{5c,5e,5g,5i, 6b, 6g,9} Two important exceptions are Tambar's work employing a sulfurdiiimide reagent followed by a Pd-catalyzed asymmetric [2,3]-rearrangement, and Hartwig's Pd-catalyzed oxidation followed by asymmetric Ir-catalyzed amination to deliver the branched amination product.¹⁰ Moreover, the majority of these examples involve the use of a nitrogen bearing a sulfonyl or carbamate group; amide-derived nitrenoids are more challenging due to the potential intervention of a Curtius-type rearrangement.¹¹ Herein, we report an intermolecular Ir(III)-catalyzed branch-selective allylic C-H amidation of unactivated terminal olefins.

We have recently reported the intramolecular Ir-catalyzed diamination of alkenyl hydroxamate esters wherein the transformation was proposed to partially proceed via Ir-nitrenoid species.¹² We wondered whether Ir nitrenoids might have sufficient lifetime to allow intermolecular aziridination or amination of feedstock α -olefins. Inspired by previous reports of group 9 metal-catalyzed C-H amidations,^{3f} together with our long-term interests in C-H functionalizations,¹³ the hypothesis was initially tested with $[\text{Cp}^*\text{MCl}_2]_2$ as the pre-catalysts (M=Rh, Co, Ir), and either sulfonyl azides¹⁴ or dioxazolones¹⁵ as nitrene precursors. Cossy and coworkers have reported that $\text{Cp}^*\text{Rh(III)}$ promotes an intramolecular allylic C-H amidation reaction proceeding through an allyl-Rh(III) intermediate.^{6g} We previously demonstrated that the allylic activation chemistry could be coupled with a C-C bond formation.¹⁶ More recently, Blakey and coworkers reported an intermolecular allylic C-H amidation of β -substituted styrenes that exhibits exclusive linear selectivity with allylbenzene.⁶ⁱ In sharp contrast, the allylic C-H amidation of **1a** with $[\text{Cp}^*\text{RhCl}_2]_2$ in combination with methyl dioxazolone **2a** gives the branched amidation product **3a** as the major isomer, albeit with a moderate yield and regioselectivity (Table 1, entry 1). We speculated that this was due to the oxidative nitrene formation and insertion process which favors the more electron-rich position. Both yield and regioselectivity were found to be significantly improved with $[\text{Cp}^*\text{IrCl}_2]_2$ while $[\text{Cp}^*\text{CoCl}_2]_2$ failed to produce any amidation products (entries 2, 3).

The higher reactivity of $\text{Cp}^*\text{Ir(III)}$ is consistent with the observations and computational arguments advanced by Baik and Chang.¹⁷ Additionally, the optimized conditions include a catalytic amount of AgNTf_2 and LiOAc . We also found that the use of Li_2CO_3 instead of LiOAc produces **3a** in 62% yield with only 10 mol% of AgNTf_2 (see the Supporting Information). $\text{Cp}^*\text{Ir(OAc)}_2$, which was believed to be the reactive catalyst generated in situ,

was ineffective without adding additional AgNTf₂ (entry 5, 6). Taken together with the above observations, we suggest that AgNTf₂ probably helps the dissociation of the acetate ligand (or the amide product) from iridium, which promotes the alkene coordination and further activation. It is also worth noting that the yield is dramatically reduced when CsOAc is used instead of LiOAc (entry 7). Given that CsOAc is much more soluble than LiOAc in DCE, this further suggests that excess acetate inhibits the reaction, supporting our assertion that a monoacetato-Ir is required. Moreover, tosyl azide was also tested. It was found to afford moderate yield of branched amidation product with excellent regioselectivity, when the reaction was heated to 80 °C (entry 11).

Having established the optimal conditions for branch-selective allylic amidation, we next sought to examine the scope of this transformation with various terminal alkenes. The reaction tolerates a broad range of functional groups, giving rise to various branched amidation products. As shown in Scheme 2, several commonly used oxygen, nitrogen or arene containing functional groups are compatible, affording corresponding amidation products in good yields and regioselectivities (**3b-3h**). Even substrates with susceptible functionalities, like -Br, -CN and -CO₂H, participated smoothly with the standard condition to provide the desired products (**3i-3k**). Although a lower yield was observed with the alkene bearing a carboxylic acid group, the regioselectivity remains unaffected, giving the branched product **3k** predominately. Despite precedent for Ir-catalyzed dehydrogenation of alkenes bearing homoallylic carbonyl groups under oxidative conditions,¹⁸ exclusive amidation of 4-phenyl-1-butene and ethyl 4-pentenoate was observed with no evidence of diene formation (**3l, 3m**). Although yield and regioselectivity are moderate with ethyl 4-pentenoate, the alternative conditions with TsN₃ as the nitrene precursor leads to moderate yield and excellent regioselectivity (**3m**). Furthermore, substrates with homoallylic heteroatoms were found to decrease the reactivity. Moderate conversions and yields were obtained even with elevated temperature (**3n, 3o**). Additionally, sterically hindered allyl-cyclohexane was remarkably well-tolerated (**3p**). Allylarenes with electron- rich (-OMe) and electron-poor (-CF₃) substituents were also examined. It was found that the regioselectivity decreased from >20:1 to 2.6:1 by changing the para- substituent from -OMe to -CF₃ (**3q-3s**). Excellent regioselectivity could be restored with TsN₃ as the nitrene precursor.

The generality of dioxazolone coupling partners were next explored. Primary, secondary and tertiary alkyl groups on the dioxazolone are tolerated with no significant decrease in reactivity (**4b-4f**). Additionally, the cyclopropane dioxazolone could also be utilized without detectable ring-opening (**4d**). Moreover, electron-donating and electron-withdrawing substituted phenyl dioxazolones are well tolerated (**4g-4j**). Even an alkenyl substituted dioxazolone can participate to afford **4k** in moderated yield. In general, these aryl and alkenyl dioxazolones are slightly less reactive and require elevated temperature for full conversion. Under all these reaction conditions, the Curtius/Lossen-type rearrangement is not competitive^{5i,19} presumably because of a change in mechanism (*vide infra*).

Further mechanistic insight was gained from isotopic labeling studies and stoichiometric reactions. When subjecting **1aa-d₂** to the optimized reaction conditions using acetic acid as the proton source, no deuterium leaching was observed at the allylic position of the amidation product. This indicates that the allylic C-H activation is irreversible under the

standard reaction condition. Furthermore, a smaller kinetic isotopic effect from intermolecular competition reaction compared to the one from intramolecular reaction was observed ($KIE = 1.9$ vs 2.6). We speculate that the allylic C–H bond activation is probably not the sole rate-determining step.²⁰ Stoichiometric studies were also performed. An allyl-Ir(III) complex **5** was trapped by an additional *p*-toluenesulfonamide ligand which stabilizes the intermediate for isolation. Interestingly, the subsequent amidation is only achieved with the assistance of AgNTf₂, which delivers the desired product in high yield and regioselectivity. In this case, AgNTf₂ might also act as a Lewis acid promoting the dissociation of the *p*-toluenesulfonamide ligand to regenerate the active species with an empty coordination site. Finally, stoichiometric reactions with **1ab-d₁** were also conducted, leading to the same distribution between deuterated and protonated products of either the allyl-Ir(III) complex ($P_D/P_H = 2.6$) or the amidation product ($P_D/P_H = 2.7$) (see SI).

Based on the above observations, we hypothesize the following catalytic cycle (Scheme 5). The active catalyst is presumably the coordinatively unsaturated cationic monoacetato Cp*Ir(III) **I** which is generated from [Cp*IrCl₂]₂, AgNTf₂, and LiOAc. Alkene coordination and irreversible metalation form allyl-Ir(III) species **III**. The oxidation of Ir(III) via N–O bond cleavage and CO₂ extrusion produces the key allyl-Ir-nitrenoid species **IV**. Subsequent migratory insertion would install the desired amide bond at the internal position. Finally proto-demetalation regenerates the active catalyst and produces the amide product. Importantly, the allyl Ir intermediate **III** is more oxidizable than **I** or **II** and only undergoes oxidation/nitrene formation once the allyl moiety is in place. Thus, the Ir nitrenoid undergoes reductive elimination to form product faster than Curtius/Lossen-type rearrangement. This is consistent with Chang's observations in intramolecular C–H insertion with related Ir nitrenoids.⁵¹

In summary, we have developed an intermolecular branch-selective allylic C–H amidation reaction via Ir(III) catalysis. The inner-sphere Ir-nitrenoid insertion after allylic C–H activation is key for prevention of undesired aziridination and achieving branched-selectivity. This opens a new pathway for branched functionalizations of terminal olefins. Further efforts at elucidating the mechanism and extending this chemistry are currently underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

We thank NIGMS (GM80442) for support. We thank Daniel Paley (Columbia) for solving the structure of **5**.

REFERENCES

1. Vitaku E; Smith DT; Njardarson JT Analysis of the structural diversity, substitution patterns, and frequency of nitrogen heterocycles among U.S. FDA approved pharmaceuticals. *J. Med. Chem* 2014, 57, 10257. [PubMed: 25255204]
2. For selected reviews, see: (a) Roughley SD; Jordan AM The medicinal chemist's toolbox: an analysis of reactions used in the pursuit of drug candidates. *J. Med. Chem* 2011, 54, 3451.

[PubMed: 21504168] (b) Brown DG; Boström J Analysis of Past and Present Synthetic Methodologies on Medicinal Chemistry: Where Have All the New Reactions Gone? *J. Med. Chem* 2016, 59, 4443. [PubMed: 26571338]

3. (a) For selected reviews, see: Collet F; Lescot C; Dauban P Catalytic C-H amination: the stereoselectivity issue. *Chem. Soc. Rev* 2011, 40, 1926. [PubMed: 21234469] (b) Ramirez TA; Zhao B; Shi Y Recent advances in transition metal-catalyzed sp³ C-H amination adjacent to double bonds and carbonyl groups. *Chem. Soc. Rev* 2012, 41, 931. [PubMed: 21842094] (c) Jeffrey JL; Sarpong R Intramolecular C(sp³)-H amination. *Chem. Sci* 2013, 4, 4092. (d) Jiao J; Murakami K; Itami K Catalytic Methods for Aromatic C-H Amination: An Ideal Strategy for Nitrogen-Based Functional Molecules. *ACS Catal.* 2016, 6, 610. (e) Kim H; Chang S Transition-Metal-Mediated Direct C-H Amination of Hydrocarbons with Amine Reactants: The Most Desirable but Challenging C-N Bond-Formation Approach. *ACS Catal.* 2016, 6, 2341. (f) Park Y; Kim Y; Chang S Transition Metal-Catalyzed C-H Amination: Scope, Mechanism, and Applications. *Chem. Rev* 2017, 117, 9247. [PubMed: 28051855]
4. (a) Yoo EJ; Ma S; Mei TS; Chan KS; Yu JQ Pd-catalyzed intermolecular C-H amination with alkylamines. *J. Am. Chem. Soc* 2011, 133, 7652. [PubMed: 21520961] (b) He G; Zhao Y; Zhang S; Lu C; Chen G Highly efficient syntheses of azetidines, pyrrolidines, and indolines via palladium catalyzed intramolecular amination of C(sp³)-H and C(sp²)-H bonds at gamma and delta positions. *J. Am. Chem. Soc* 2012, 134, 3. [PubMed: 22191666] (c) Kim JY; Park SH; Ryu J; Cho SH; Kim SH; Chang S Rhodium-catalyzed intermolecular amidation of arenes with sulfonyl azides via chelation-assisted C-H bond activation. *J. Am. Chem. Soc* 2012, 134, 9110. [PubMed: 22624801] (d) Ryu J; Kwak J; Shin K; Lee D; Chang S Ir(III)-catalyzed mild C-H amidation of arenes and alkenes: an efficient usage of acyl azides as the nitrogen source. *J. Am. Chem. Soc* 2013, 135, 12861. [PubMed: 23957845] (e) Matsubara T; Asako S; Ilies L; Nakamura E Synthesis of anthranilic acid derivatives through iron-catalyzed ortho amination of aromatic carboxamides with N-chloroamines. *J. Am. Chem. Soc* 2014, 136, 646. [PubMed: 24380435] (f) Shang M; Sun SZ; Dai HX; Yu JQ Cu(II)-mediated C-H amidation and amination of arenes: exceptional compatibility with heterocycles. *J. Am. Chem. Soc* 2014, 136, 3354. [PubMed: 24527701] (g) Yan Q; Chen Z; Yu W; Yin H; Liu Z; Zhang Y Nickel-catalyzed direct amination of arenes with alkylamines. *Org. Lett* 2015, 17, 2482. [PubMed: 25942045]
5. (a) For selected references, see: Yu XQ; Huang JS; Zhou XG; Che CM Amidation of saturated C-H bonds catalyzed by electron-deficient ruthenium and manganese porphyrins. A highly catalytic nitrogen atom transfer process. *Org. Lett* 2000, 2, 2233. [PubMed: 10930251] (b) Liang C; Collet F; Robert-Peillard F; Müller P; Dodd RH; Dauban P Toward a Synthetically Useful Stereoselective C-H Amination of Hydrocarbons. *J. Am. Chem. Soc* 2008, 130, 343. [PubMed: 18072775] (c) Lu H; Jiang H; Hu Y; Wojtas L; Zhang XP Chemoselective intramolecular allylic C-H amination versus C=C aziridination through Co(II)-based metalloradical catalysis. *Chem. Sci* 2011, 2, 2361. (d) Gephart RT; Warren TH, Copper-Catalyzed sp³ C-H Amination. *Organometallics* 2012, 31, 7728. (e) Paradine SM; White MC Iron-catalyzed intramolecular allylic C-H amination. *J. Am. Chem. Soc* 2012, 134, 2036. [PubMed: 22260649] (f) Roizen JL; Harvey ME; Du Bois J Metal-Catalyzed Nitrogen-Atom Transfer Methods for the Oxidation of Aliphatic C-H Bonds. *Acc. Chem. Res* 2012, 45, 911. [PubMed: 22546004] (g) Paradine SM; Griffin JR; Zhao J; Petronico AL; Miller SM; Christina White M A manganese catalyst for highly reactive yet chemoselective intramolecular C(sp³)-H amination. *Nat. Chem* 2015, 7, 987. [PubMed: 26587714] (h) Dolan NS; Scamp RJ; Yang T; Berry JF; Schomaker JM Catalyst-Controlled and Tunable, Chemoselective Silver-Catalyzed Intermolecular Nitrene Transfer: Experimental and Computational Studies. *J. Am. Chem. Soc* 2016, 138, 14658. [PubMed: 27726353] (i) Hong SY; Park Y; Hwang Y; Kim YB; Baik MH; Chang S Selective formation of γ -lactams via C-H amidation enabled by tailored iridium catalysts. *Science* 2018, 359, 1016. [PubMed: 29496875]
6. (a) For selected references, see: Beccalli EM; Broggini G; Paladino G; Penoni A; Zoni C Regioselective Formation of Six- and Seven-Membered Ring by Intramolecular Pd-Catalyzed Amination of N-Allyl-anthranilamides. *J. Org. Chem* 2004, 69, 5627. [PubMed: 15307732] (b) Fraunhoffer KJ; White MC syn-1,2-Amino alcohols via diastereoselective allylic C-H amination. *J. Am. Chem. Soc* 2007, 129, 7274. [PubMed: 17516648] (c) Srivastava RS; Tarver NR; Nicholas KM Mechanistic Studies of Copper(I)-Catalyzed Allylic Amination. *J. Am. Chem. Soc* 2007, 129, 15250. [PubMed: 18004850] (d) Reed SA; White MC Catalytic Intermolecular Linear Allylic C-H Amination via Heterobimetallic Catalysis. *J. Am. Chem. Soc* 2008, 130, 3316. [PubMed:

18302379] (e)Liu G; Yin G; Wu L Palladium-catalyzed intermolecular aerobic oxidative amination of terminal alkenes: efficient synthesis of linear allylamine derivatives. *Angew. Chem. Int. Ed* 2008, 47, 4733.(f)Luzung MR; Lewis CA; Baran PS Direct, chemoselective N-tert-prenylation of indoles by C-H functionalization. *Angew. Chem. Int. Ed* 2009, 48, 7025.(g)Cochet T; Bellosa V; Roche D; Ortholand JY; Greiner A; Cossy J Rhodium(III)-catalyzed allylic C-H bond amination. Synthesis of cyclic amines from omega-unsaturated N-sulfonylamines. *Chem. Commun* 2012, 48, 10745.
(h)Shibata Y; Kudo E; Sugiyama H; Uekusa H; Tanaka K Facile Generation and Isolation of π -Allyl Complexes from Aliphatic Alkenes and an Electron-Deficient Rh(III) Complex: Key Intermediates of Allylic C-H Functionalization. *Organometallics* 2016, 35, 1547.(i)Burman JS; Blakey SB Regioselective Intermolecular Allylic C-H Amination of Disubstituted Olefins via Rhodium/ π -Allyl Intermediates. *Angew. Chem. Int. Ed* 2017, 56, 13666.

- 7 (a). Mahy JP; Bedi G; Battioni P; Mansuy D Allylic amination of alkenes by tosyliminoiodobenzene: manganese porphyrins as suitable catalysts. *Tetrahedron Lett.* 1988, 29, 1927.(b)Wehn PM; Lee J; Du Bois J Stereochemical Models for Rh-Catalyzed Amination Reactions of Chiral Sulfamates. *Org. Lett* 2003, 5, 4823. [PubMed: 14653683] (c)Barman DN; Nicholas KM Copper-Catalyzed Intramolecular C-H Amination. *Eur. J. Org. Chem* 2011, 2011, 908;(d)Cramer SA; Jenkins DM Synthesis of aziridines from alkenes and aryl azides with a reusable macrocyclic tetracarbene iron catalyst. *J. Am. Chem. Soc* 2011, 133, 19342. [PubMed: 22081884]
- 8 (a). Reed SA; Mazzotti AR; White MC A catalytic, Bronsted base strategy for intermolecular allylic C-H amination. *J. Am. Chem. Soc.* 2009, 131, 11701. [PubMed: 19645492] (b)Shimizu Y; Obora Y; Ishii Y Intermolecular Aerobic Oxidative Allylic Amination of Simple Alkenes with Diarylamines Catalyzed by the Pd(OCOCF₃)₂/NPMoV/O₂ System. *Org. Lett* 2010, 12, 1372. [PubMed: 20158264] (c)Yin G; Wu Y; Liu G Scope and Mechanism of Allylic C-H Amination of Terminal Alkenes by the Palladium/PhI(OPiv)₂ Catalyst System: Insights into the Effect of Naphthoquinone. *J. Am. Chem. Soc* 2010, 132, 11978. [PubMed: 20690676] (d)Xiong T; Li Y; Mao L; Zhang Q; Zhang Q Palladium-catalyzed allylic C-H amination of alkenes with N-fluorodibenzene-sulfonimide: water plays an important role. *Chem. Commun* 2012, 48, 2246.
(e)Pattillo CC; Strambeanu II; Calleja P; Vermeulen NA; Mizuno T; White MC Aerobic Linear Allylic C-H Amination: Overcoming Benzoquinone Inhibition. *J. Am. Chem. Soc* 2016, 138, 1265. [PubMed: 26730458] (f)Vemula SR; Kumar D; Cook GR Palladium-Catalyzed Allylic Amidation with N-Heterocycles via sp³ C-H Oxidation. *ACS Catal.* 2016, 6, 5295.(g)Ma R; White MC C-H to C-N Cross-Coupling of Sulfonamides with Olefins. *J. Am. Chem. Soc* 2018, 140, 3202. [PubMed: 29432000]
- 9 (a). Milczek E; Boudet N; Blakey S Enantioselective C-H amination using cationic ruthenium(II)-pybox catalysts. *Angew. Chem. Int. Ed* 2008, 47, 6825.(b)Zalatan DN; Du Bois J A Chiral Rhodium Carboxamidate Catalyst for Enantioselective C-H Amination. *J. Am. Chem. Soc* 2008, 130, 9220. [PubMed: 18582043] (c)Wu L; Qiu S; Liu G Bronsted base-modulated regioselective Pd-catalyzed intramolecular aerobic oxidative amination of alkenes: formation of seven-membered amides and evidence for allylic C-H activation. *Org. Lett* 2009, 11, 2707. [PubMed: 19456146] (d)Nahra F; Liron F; Prestat G; Mealli C; Messaoudi A; Poli G Striking AcOH acceleration in direct intramolecular allylic amination reactions. *Chem. - Eur. J* 2009, 15, 11078. [PubMed: 19795438]
- 10 (a). Bao H; Tambar UK Catalytic Enantioselective Allylic Amination of Unactivated Terminal Olefins via an Ene Reaction/[2,3]-Rearrangement. *J. Am. Chem. Soc* 2012, 134, 18495. [PubMed: 23106555] (b)Sharma A; Hartwig JF Enantioselective Functionalization of Allylic C-H Bonds Following a Strategy of Functionalization and Diversification. *J. Am. Chem. Soc* 2013, 135, 17983. [PubMed: 24156776]
- 11 (a). Dube P; Nathel NFF; Vetelino M; Couturier M; Aboussafy CL; Pichette S; Jorgensen ML; Hardink M Carbonyldiimidazole-Mediated Lossen Rearrangement. *Org. Lett* 2009, 11, 5622. [PubMed: 19908883] (b)Li D; Wu T; Liang K; Xia C Curtius-like Rearrangement of an Iron-Nitrenoid Complex and Application in Biomimetic Synthesis of Bisindolylmethanes. *Org. Lett* 2016, 18, 2228. [PubMed: 27116426]
12. Conway JH Jr.; Rovis T Regiodivergent Iridium(III)-Catalyzed Diamination of Alkenyl Amides with Secondary Amines: Complementary Access to γ - or δ -Lactams. *J. Am. Chem. Soc* 2018, 140, 135. [PubMed: 29278330]

13. Piou T; Rovis T Electronic and Steric Tuning of a Prototypical Piano Stool Complex: Rh(III) Catalysis for C-H Functionalization. *Acc. Chem. Res* 2018, 51, 170. [PubMed: 29272106]
- 14 (a). Intrieri D; Zardi P; Caselli A; Gallo E Organic azides: "energetic reagents" for the intermolecular amination of C-H bonds. *Chem Commun* 2014, 50, 11440.(b)Shin K; Kim H; Chang S Transitionmetal-catalyzed C-N bond forming reactions using organic azides as the nitrogen source: a journey for the mild and versatile C-H amination. *Acc. Chem. Res* 2015, 48, 1040. [PubMed: 25821998]
15. Park Y; Park KT; Kim JG; Chang S Mechanistic Studies on the Rh(III)-Mediated Amido Transfer Process Leading to Robust C-H Amination with a New Type of Amidating Reagent. *J. Am. Chem. Soc* 2015, 137, 4534. [PubMed: 25789561]
16. Archambeau A; Rovis T Rhodium(III)-Catalyzed Allylic C(sp³)-H Activation of Alkenyl Sulfonamides: Unexpected Formation of Azabicycles *Angew. Chem. Int. Ed* 2015, 54, 13337.
17. Park Y; Heo J; Baik M-H; Chang S Why is the Ir(III)-Mediated Amido Transfer Much Faster Than the Rh(III)-Mediated Reaction? A Combined Experimental and Computational Study. *J. Am. Chem. Soc* 2016, 138, 14020. [PubMed: 27690406]
18. Wang Z; He Z; Zhang L; Huang Y Iridium-Catalyzed Aerobic α,β -Dehydrogenation of γ,δ -Unsaturated Amides and Acids: Activation of Both α - and β -C-H bonds through an Allyl-Iridium Intermediate. *J. Am. Chem. Soc* 2018, 140, 735. [PubMed: 29251918]
19. No Curtius or Lossen-type rearrangement byproducts were observed from the reaction of 1-decene (**1a**) and 3-methyl-1,4,2-dioxazol-5-one (**2a**) under the standard condition in DCM-*d*₂. See Supporting Information for more details.
20. Simmons EM; Hartwig JF On the interpretation of deuterium kinetic isotope effects in C-H bond functionalizations by transition-metal complexes. *Angew. Chem. Int. Ed* 2012, 51, 3066.

a. Previous work:

i) Metal nitrenoid insertion (outer-sphere C-H insertion)

ii) Pd(II)-catalyzed linear allylic amination

b. This work: branch-selective

Scheme 1. Catalytic Allylic C-H Amidation

Scheme 2. Alkene Scope

^a60 °C was used. ^b80 °C was used. ^cTsN₃ was used instead of **2a**. ^dNMR yield. ^eYield based on recovered starting material.

Scheme 3. Dioxazolone Scope

^a60 °C was used. ^b80 °C was used.

Scheme 4. Mechanistic Studies

Scheme 5. Proposed Mechanism

Table 1.

Reaction Development^a

entry	Catalyst	Base	yield ^b	B:L ^c
1	[Cp*RhCl ₂] ₂	LiOAc	40%	2.9:1
2	[Cp*CoCl ₂] ₂	LiOAc	N.D.	--
3	[Cp*IrCl ₂] ₂	LiOAc	75% (73%) ^d	>20:1
4 ^e	[Cp*IrCl ₂] ₂	LiOAc	24%	>20:1
5 ^f	Cp*Ir(OAc) ₂	--	75%	>20:1
6 ^{f,g}	Cp*Ir(OAc) ₂	--	N.R.	--
7	[Cp*IrCl ₂] ₂	CsOAc	8%	--
8	--	LiOAc	N.R.	--
9	[Cp*IrCl ₂] ₂	--	15%	--
10 ^g	[Cp*IrCl ₂] ₂	LiOAc	N.R.	--
11 ^h	[Cp*IrCl ₂] ₂	LiOAc	39% ⁱ	>20:1

^aReactions were conducted on a 0.1 mmol scale using **1a** (1.0 equiv), **2a** (1.5 equiv), [Cp*IrCl₂]₂ (2.5 mol%), AgNTf₂ (15 mol%) and LiOAc (20 mol%).

^bYield of dominant isomer as determined by ¹H NMR.

^cDetermined by analysis of ¹H NMR of the unpurified reaction mixture.

^dIsolated yield.

^eAgNTf₂ (10 mol%) was used.

^fCp*Ir(OAc)₂ (5 mol%) was used.

^gWithout AgNTf₂.

^hTsN₃ instead of **2a** was applied at 80 °C.

ⁱYield of the corresponding allylic tosyl amide.