Published in final edited form as:

J Hosp Palliat Nurs. 2020 February ; 22(1): 5–11. doi:10.1097/NJH.0000000000000608.

On Best Interests: A Case for Clinical Ethics Consultation

Karen O. Moss, PhD, RN, CNL [Assistant Professor],

Center on Healthy Aging, Self-Management and Complex Care, #368 Newton Hall | 1585 Neil Avenue, Columbus, OH 43210, College of Nursing, The Ohio State University

Robert Guerin, PhD,

Clinical Ethicist, University Hospitals Cleveland Medical Center, Clinical Assistant Professor, Department of Bioethics, School of Medicine, Case Western Reserve University

Olubukunola Mary Dwyer, JD, MA,

Clinical Ethicist, University Hospitals Cleveland Medical Center, Clinical Assistant Professor, Department of Bioethics, School of Medicine, Case Western Reserve University

Celia E. Wills, PhD, RN [Associate Professor],

Center on Healthy Aging, Self-Management and Complex Care, College of Nursing | The Ohio State University

Barbara Daly, PhD, RN, FAAN

Clinical Ethicist, University Hospitals Cleveland Medical Center, The Gertrude Perkins Oliva Professor in Oncology Nursing, Department of Bioethics, School of Medicine, Frances Payne Bolton School of Nursing | Case Western Reserve University

Abstract

Surrogate healthcare decision-making is often a challenge for everyone involved. In the case of incapacitated patients, family members, nurses, healthcare providers, and other members of the healthcare team often grapple with determining the most appropriate clinical course of action. For these difficult patient scenarios, the expertise of clinical ethics consultants are sought to assist with complex healthcare decision-making. Clinical ethics consultation is designed to provide a more objective “outside” opinion and offer advice to the patient, family, and entire care team to support and guide decisions. Nurses are well positioned to initiate assistance from Clinical Ethics Consult Services in support of patient and family advocacy. This article presents a case analysis based on the Stakeholder, Facts, Norms, and Options Framework to analyze the best interest course of action for Mr. K, a patient diagnosed with abdominal pain due to end-stage liver cirrhosis and lacks decisional capacity in regard to his own treatment decision-making. The case analysis highlights specific examples of how nurses can provide information, facilitate discussion, and otherwise support patients and families to achieve best interest outcomes.

Corresponding author: Phone: 614-688-3100, moss.391@osu.edu.

This paper was developed as a result of the first author’s participation in the Clinical Ethics Immersion Program (CLEIP) through the Cleveland Clinic Center for Bioethics.

Conflict of Interest: The authors declare no conflicts of interest with respect to the research, authorship, and/or publication of this article.

Keywords

Best interests; clinical ethics; consultation; shared decision-making

The Case

Mr. K is a young adult male admitted to the intensive care unit 12 days ago because of hypotension and intractable abdominal pain due to end-stage liver cirrhosis. He has a history of alcohol and other substance abuse, as well as a seizure disorder. He is currently dependent on vasopressors and daily blood transfusions for hemodynamic stability. Mr. K has bleeding esophageal varices and other gastric bleeding. He is obtunded and therefore non-responsive, but breathing spontaneously. His code status is “full code.” He is not married and has four children, all under the age of 15 years. On admission to the hospital, he identified his father as his surrogate decision-maker, but he has no written advance directive. On the few occasions when his father visited, he would not discuss anything related to his son’s code status or end of life, simply saying, “Do everything.” His father has not been in to see him recently. However, Mr. K’s mother has been visiting him in the hospital often.

The Clinical Ethics Consult Service (CECS) team visited Mr. K, and after an assessment, including a discussion with the healthcare team, confirmed his current lack of decision-making capacity. Since Mr. K’s father was not visiting frequently and reportedly did not wish to discuss his code status or end of life, the CECS team agreed to contact his mother. Mr. K’s mother was asked about what she knew of his son’s current condition. She said that she believed it was his alcoholism that led to his liver failure. She was informed that the healthcare team believed he was most likely dying and required daily blood transfusions to maintain his life at this point. While she was accepting of his poor prognosis, she was saddened. She mentioned that she only recently learned of his hospitalization and had been in contact with Mr. K’s father via text messages. When asked about Mr. K’s goals, his mother said that he wanted to be home with his kids and provided no additional details.

The following day Mr. K became further hemodynamically unstable, requiring several blood transfusions during the night. A Computerized Tomography (CT) scan was completed to determine if there was an identifiable source of bleeding. It was agreed beforehand by the healthcare team that if the bleeding source was identified and an intervention, such as banding, resection, or cautery, was feasible, they would do so. The healthcare team was scheduled to hold a family meeting with Mr. K’s parents that evening, and the nurse asked the CECS if it would be ethically justified to impose treatment limitations without surrogate permission. Specifically, if the course of bleeding could not be corrected, was the team obligated to continue to provide blood transfusions every day and were they obligated to attempt resuscitation when the inevitable arrest occurred?

Ethical Analysis

This case presents a number of complex ethical issues. Among these are: 1) Given this patient’s condition and prognosis, is it ethically justified to limit continued use of blood transfusions? 2) Who should be the decision maker? With regard to the latter, when parents

are the surrogates, but they disagree with each other, who should the healthcare team consider to be the decision maker? And, once there is clarification about who has authority to make decisions, is the team obligated to follow the wishes of that person regardless of the team's assessment of Mr. K's best interest? Resolving this dilemma requires understanding of surrogacy in general and the best interest standard specifically as it relates to healthcare decisions.

Shared Decision-Making

Methods and approaches of healthcare decision-making have swung from being based upon paternalism, which is healthcare provider directed, to autonomy, which is patient driven. More recently, both of these approaches have lost favor and shared decision-making has become a preferred model of decision-making. This change occurred as the result of the recognition that healthcare decisions need to be medically appropriate as well as consistent with the patient's values. The acceptance of this change has been recognized by both western society and its healthcare community.¹ There is a great deal of literature discussing this method of decision-making, ranging from the ideal of shared decision-making and the principles involved, to how to accomplish this approach in various clinical settings, including the intensive care unit.²⁻⁴

In its usual form, shared decision-making is a division of labor and responsibility that occurs, at a minimum, between the patient (or their agent or surrogate if the patient lacks decision-making capacity) and a healthcare provider. The patient's duty in this dyad is to provide his/her values and goals, while the healthcare provider contributes knowledge and experience.⁵ Not to be excluded from this equation is the ever important role of the nurse in this process.⁶

Although this division of labor seems simple, the burdens involved are not always equally shared by those involved, and its meaning can change depending on specific aspects of the patient-provider relationship. For example, a patient hospitalized in an internal medicine unit for ten days, who has informed the inpatient healthcare team of his/her goals might choose to follow the recommended care plan without explicitly being involved in care plan development. In this circumstance, the primary medical care provider along with the nurse (and input from the remainder of the healthcare team) determines the appropriate treatment plan in context of the patient's goals and preferences. The patient controls whether and how much to work together with the healthcare provider to make treatment decisions and retains the ability to either reject or acquiesce to the recommendation that the healthcare provider makes. This collaborative process allows patients, surrogates, and clinicians to reach the most appropriate decisions while considering the best scientific evidence and the patient's values, goals, and preferences.⁴

An Alternative Case

In an alternate example, the patient who has decision-making capacity might choose to work directly with the current members of the healthcare team as partners in decision-making. The difference between the scenarios depends largely on how involved the patient or the

agent (proxy) chooses to be involved in the healthcare decision-making process. How much a patient wants to be involved can vary depending on various factors such as individual preference, comfort with the care provider, the type of decisions to be made, potential benefits and risks of alternative courses of action, severity of illness, capacity to make decisions, and when decisions need to take place. Clinicians should remain open to tailoring the decision-making process based on patient and surrogate preferences.⁴

On Best Interests

The moral authority of surrogate decision-makers is not absolute. As Buchanan and Brock state in their seminal publication, the legal standard of substituted judgment first emerged in probate court rulings regarding financial decisions on behalf of persons deemed incompetent of making decisions for themselves.⁷ Appreciating the vulnerability of such situations, and the speculative nature of substituted judgment, the rulings noted that “a gift may be made to another from an incompetent’s estate if the ward would have made this gift if competent, but *only if* the gift is not so large that it would deprive the ward of basic necessities of life, even if the ward, if competent, would have been more generous.”⁷ In this context, the courts recognized that surrogates are liable to make decisions for others that more likely track the surrogate’s interests rather than the ward’s interests. In order to protect the ward against such misdeeds, the “basic necessities” limitation was set.

Certainly, in the context of healthcare surrogate decision-making, there are cases in which treatment or use of healthcare technology is withheld or withdrawn such that the incapacitated patient is not only deprived of the basic necessities of life but is deprived of life itself. And since such cases exist with clear ethical support—such as for a patient in a permanent vegetative state who has an advance directive indicating a preference to withhold or withdraw life-sustaining interventions in this situation. Differentiation of the limitations of financial and healthcare surrogate decision-making is needed. A look at the latter suggests that healthcare surrogate decision-makers’ moral authority should be limited when the evidence for patient preferences is minimal or nonexistent. For example, imagine if family members at the bedside are asked whether their loved one ever spoke about living permanently on a ventilator and/or with a feeding tube. “We never spoke about such things,” they reply. “Who talks about that?” Here, and in similar cases, the limitation of surrogates’ moral authority stems from the lack of evidence regarding what the patient would be likely to decide if s/he were considered competent to do so.

The Best Interest Standard

The best interest standard for surrogate healthcare decision-making is designed to facilitate decision-making in cases in which there is minimal or nonexistent evidence regarding a patient’s preferences toward the decisions at hand. Generally speaking, the best interest standard instructs surrogate healthcare decision-makers to determine a treatment option that generates the most good or benefit for the patient. What this treatment option is will depend on the patient’s interests, the degree of value s/he assigns to interests such as avoiding pain or retaining the ability to speak and interact with family.

This standard is not an abstract judgment, such as “An ideal, rational being would choose X.” It attempts to take into account what Ronald Dworkin calls “critical” interests.⁸ These critical interests are integral to who the person is: critical interests seem tied to commitments and choices that define a person’s narrative or are part of a consistent character that is the mark of integrity. For example, a patient who has a long-held preference to avoid life-sustaining treatment in the context of a severely painful terminal illness for which there is inadequate symptom relief available, would be representative of a critical interest in the value of quality of life over quantity of life. In this sense, the best interest standard maintains respect for the patient as a person even in the absence of good evidence about what the patient would decide in this particular instance.

Like the basic necessities limitation in financial decision-making, the best interest standard is a built-in institutional safeguard to protect patients against surrogate decision-makers (as well as healthcare providers) *even if surrogates are accurately reflecting a patient’s preferences*. Consider a patient who is incapacitated, lacks an advance directive, and is suffering from complications of end-stage dementia. The healthcare team tells the family that the patient is dying and recommends hospice care. Moreover, and until the patient enrolls in hospice, the healthcare team notes that it will not perform cardiopulmonary resuscitation (CPR) in the event of a cardiac arrest. Now imagine that the family disagrees with the do-not-resuscitate (DNR) order. They state that the patient is a fighter and would want to go out with a fight. The important point to note here is that even if the patient prefers CPR, the vulnerability of this patient, the lack of evidence supporting the surrogates’ claim, and the degree of harm that would likely occur in the setting of resuscitation, overrides the surrogates’ authority. Surrogates cannot place patients in vulnerable situations with a high likelihood of harm, even if the patient would agree. This is one example of how preference-sensitive these decisions can be and stresses the clinicians’ fiduciary responsibility to use evidence-based practice and experience in order to implement appropriate treatment and testing.⁴

Stakeholders, Facts, Norms, and Options (SFNO) Framework

Also known as the *So Far No Objections Framework*, the SFNO framework method considers Stakeholders, Facts, Norms, and Options as a basis for analyzing ethical cases. The SFNO method uses root cause analysis to examine three major sources of uncertainty or disagreement regarding decisions when: 1) persons involved have competing interests, 2) uncertainty or disagreement exists about relevant facts, or, 3) disagreement, or uncertainty exists regarding ethical norms. This approach uses specific probing questions for identifying ethically appropriate solutions.⁹ The SFNO approach is applied here to the case of Mr. K.

Stakeholders

Mr. K currently lacks capacity to participate in his healthcare decision-making. Although the patient is always the central stakeholder for treatment decision-making, and his preferences and values and goals (insofar as known) therefore carry the most weight, Mr. K’s cognitive status hinders the ability to learn of these directly from him. When asked about Mr. K’s goals, his mother said that he wanted to be with his children and provided no additional

details. This was an indicator for the CECS team to further explore the life story and personal identity of the patient (and his parents). This could provide information that could help identify patient and family values that shape healthcare decision-making. Respecting the values of all stakeholders requires investing the time and energy needed to fully appreciate and attempt to incorporate these values in treatment planning.

Because Mr. K's father was not present and was assessed on multiple occasions as unwilling to speak about Mr. K's prognosis, the CECS reached out to his mother. This was an important step as it allowed the CECS to consult with another person who could potentially make decisions on Mr. K's behalf. There was also a concern for Mr. K's dependent children and their wellbeing. If Mr. K's father is unavailable to serve as surrogate, then his mother could fulfill this role; as noted above, both have equal authority as next of kin. His father may have been struggling with coming to terms with his son's morbidity and mortality, might have been uncomfortable with the burden of decision-making, or might have had other competing responsibilities that prevented him from being present.

Facts

Explicitly reviewing and considering the facts relevant to the ethical question is an important step. It sometimes reveals differences among the team in what is known or assumed, and it reveals what is not known. For example, there was not an adequate understanding of the reasoning of Mr. K's father and his adamant insistence on full code status and "doing everything." The role that Mr. K's history of seizure disorder and more recent alcohol abuse played in his father's reasoning was not understood. It could be that Mr. K's father just could not bear to make a decision that might hasten his son's death, leaving Mr. K's young children without their biological father for the rest of their lives. He also might have simply been overburdened by his other obligations at the time. Similarly, it was not known at the time of this dilemma how involved Mr. K's mom had been with her son and what her potential role as caregiver for Mr. K's children would be. While obtaining more complete facts about these issues might not, in themselves, resolve the dilemma, they could be very relevant to reaching a sound ethical response in regard to understanding best interests of the patient and family members.

It also was unclear how much of Mr. K's *current condition* had been discussed with him and his family to date at this point, particularly given his father's resistance to having the conversation. Apart from Mr. K's mother's expression that he would have wanted to be home with his children, there seemed to be little known about his values and belief system.

Equally important to identify are the healthcare facts. Mr. K's advanced liver disease in itself carried a very poor prognosis. In the absence of curative interventions, it was likely that his bleeding would continue, and this soon would lead to his death. On the other hand, if a bleeding source could be identified and addressed, although he would still eventually succumb to his liver failure, there would be a chance of survival for weeks to months.

The facts regarding the justification for using CPR in the event of an arrest are relatively well known. In a recent national study conducted by American Heart Association® Get with the Guidelines® resuscitation investigators, the average survival to discharge after CPR for

adults receiving assisted/mechanical ventilation or vasoactive agents in intensive care units or telemetry units was 18.6%.¹⁰ When considering significant added risk factors for Mr. K (coagulopathy, liver failure, substance abuse, and vasopressor dependence), there is no meaningful likelihood of survival without concomitant anoxic damage, but a very high probability of inflicting further damage.

Norms

The principle of autonomy through consideration of previously known patient wishes and consideration of surrogates' viewpoints to establish best interest for Mr. K are applicable to this case. Also, nonmaleficence and beneficence are key principles establishing the obligation to help and to refrain from causing harm by carrying out actions such as CPR. During CPR, chest compressions can result in broken ribs and possible anoxic injury, further complicating Mr. K's condition and worsening his already poor prognosis.

According to the American Medical Association's Code of Ethics, if a patient lacks decision-making capacity, the surrogate may decline an intervention or ask for an intervention to be discontinued in keeping with ethical standards for surrogate decision-making.¹¹ This Code of Ethics states that while there may be an emotional difference between not initiating an intervention and discontinuing it, there is no ethical difference between withholding and withdrawing treatment.¹¹ When an intervention, such as the administration of blood transfusions for Mr. K., no longer helps to achieve the patient's desired quality of life or goals for care, it is ethically appropriate for healthcare providers to withdraw or to withhold it. Consultation with a CECS should be sought whenever the surrogate's decision regarding treatment appears inconsistent with the healthcare provider's best professional judgement, and thus, potentially not in the patient's healthcare interest, or violates the patient's previously expressed goals for care.¹¹ In Mr. K's case there was a lack of evidence regarding previously expressed goals for care, but his assigned surrogate decision maker wanted everything done to preserve his life. Both the medical and nursing codes of ethics offer the use of palliative care for symptom management even when cure-directed medical interventions are being withdrawn or withheld.^{11,12} Both also support the use of advance directives and early establishment of goals of care for treatments that are in keeping with the known patient values and belief systems.^{11,12}

The American Nurses Association Code of Ethics advises nurses to carefully consider decisions regarding resuscitation status and withholding and withdrawal of life-sustaining therapies.¹² Care delivery for an incapacitated patient should be consistent with the patient's previously expressed wishes and known values. It is understood that a surrogate would make decisions for a patient such as Mr. K with these in mind; however, in the absence of an appropriate decision maker, decisions will be made by the healthcare team that are in the best interests of the patient.¹² The American Academy of Hospice and Palliative Care Medicine (AAHPM) and the Hospice and Palliative Nurses Association (HPNA) endorse that withholding and withdrawing nonbeneficial treatment are ethically and morally indistinguishable and are considered acceptable through the course of progressive, life-limiting illness.^{13,14} Both HPNA and AAHPM believe that surrogate decision makers are to be included in this process. However, patient's preferences and best interests, rather than the

surrogate's beliefs and wishes alone, should direct the care plan, which always includes relief of suffering.^{13,14} It is essential for nurses to be familiar with these position statements and their responsibility for applying them to their clinical practice, particularly as it relates to surrogate healthcare decision-making.¹⁵

Options

The options for Mr. K include the decision to have the interventional radiology (IR) team perform a procedure to attempt to repair the cause of the bleeding or to withhold this intervention. Along with the decision to perform a CT scan of Mr. K's abdomen, the option to intervene was presented based on the premise that if the results of the CT scan indicated that there was a cause for the bleed that could be repaired, then this intervention would be justified in that it could extend his survival for a short time. Whether or not to limit transfusions was another decision at hand for Mr. K. Part of this decision had to be the intent of the transfusions. Based on the ability to identify and repair the source of the bleed, continuing transfusions could be part of a plan to continue life support in order to allow for some meaningful period of survival. Or, even if no repairable source of bleeding could be identified, transfusions could be continued for a brief time in order to give family more opportunity to be with the patient, or simply to have a few more days to evaluate the patient's clinical condition. If the latter were the case, it would be appropriate to determine a preset time for reconsideration of the plan. Briefly postponing final decisions about withdrawing interventions can be useful practice in healthcare decision-making.¹¹

The decision whether or not to perform CPR on Mr. K in the event of a cardiopulmonary arrest was also considered. As noted above, the decision to withhold CPR should rest on assessment of likelihood of harms versus benefits in this particular case, given known probabilities of outcomes of CPR and Mr. K's specific clinical situation. Table 1 presents a summary of probing questions and considerations.

Case Discussion

In further considering Mr. K's status, his bleeding is in the context of end-stage liver cirrhosis. He complained of pain when awake, requires blood transfusions for hemodynamic support and a radiological procedure that might lead to control of his bleeding if a source can be identified. The ethical question is whether there is an obligation to continue blood transfusions (if the source of the bleed cannot be identified) and whether or not to attempt resuscitation if Mr. K arrests. Little is known about Mr. K's values and preferences. According to his mother he would like to be home with his children. It is possible that if more information could be gathered, it could be determined that Mr. K would appreciate hospice care at home so he may be with his children. This would satisfy both the obligation to keep him comfortable at the end of life (support his well-being) and the obligation to respect his preferences (support his autonomy).

Mr. K's father—who the patient has identified as his surrogate—is stating, “do everything.” As discussed above, even if Mr. K's father is accurately reflecting his son's preferences, which is very uncertain, the best interest standard acts as a safeguard against surrogates who place patients at high risk for preventable suffering and harm. Resuscitation for Mr. K would

be likely to cause harm with a high probability of little to no benefit. He would likely incur bruising and broken ribs, as well as a possible anoxic injury. All things considered, his condition likely would be worse after the arrest than it is at baseline. In addition, there is ethical justification for withholding blood transfusions if the source of the bleeding cannot be identified. If neither Mr. K nor his surrogate decision-maker can provide evidence that there are specific short-term goals that blood transfusions can support and that he can appreciate (e.g., an anniversary, birthday, family visit, etc.), the transfusions should be considered nonbeneficial. There is therefore ethical support for not only writing a DNAR (Do Not Attempt Resuscitation) order but for also discontinuing blood transfusions.

Case Conclusion

Mr. K was taken to the gastroenterology suite, after obtaining permission from his mother for the procedure. Several areas of bleeding were identified and cauterized. However, upon arrival back to the unit, he experienced cardiopulmonary arrest. Cardiopulmonary resuscitation was withheld, and he died within minutes.

Discussion

Surrogate shared decision making rests on the availability of appropriate surrogates, their willingness to engage in discussion, and clarity on the part of the healthcare team about decisions (e.g. treatment options) that are available for consideration and those that are not. In the case presented, Mr. K is critically ill, and it has been determined that he lacks decision-making capacity. This determination eliminates his ability to determine whether and how much of a partner he wants to be in the shared decision-making process. Mr. K has identified his father as his surrogate, but the question arises of whether his father is an appropriate surrogate decision maker. As described above, the patient or agent's role in the dyad requires that they identify the patient's values and goals. Mr. K's father has thus far not been in regular contact with providers and has declined to discuss treatment decisions with the healthcare team. He has not provided the healthcare team with information about Mr. K's values or goals, although he has given the directive to "do everything." Given the father's limited involvement in shared decision-making with the healthcare team, an argument can be made that decision-making authority should pass to Mr. K's mother. She has been willing to speak with the team about one of Mr. K's goals and has been present and willing to be involved in decision-making.

Summary and Role of the Nurse

The significance of the nurse on CECS committees is recognized both nationally and internationally,^{16,17} including the importance of their ethics education, training, and practice.^{18,19} The current case presented was initiated by the bedside nurse. In this instance, as often true for clinical ethics cases, the issues at hand are not always clear, making the need for ethics consultation warranted. Nurses in various nursing roles request and participate in clinical ethics consultation, with patient advocacy being a top reason for this decision.²⁰ In these instances, it is imperative for nurses involved to gather necessary information to help provide needed data to the CECS to aid in their organization of facts in order to support the

most appropriate decision. The nurse should be present at the team/family meetings to answer questions and clarify misunderstandings, and serve as a liaison between the CECS team, other members of the healthcare team, and the family. As an important member of the healthcare team, the nurse's perspective is crucial to the discussion with the CECS team in ensuring that information is fully presented. Also essential is the nurses' obligation to help engage the family. For example, in the case presented, Mr. K's father may have been intimidated by the healthcare providers. Part of the nurses' role is to help to identify and address family concerns. This could be through clarification of healthcare jargon and answering and anticipating questions or concerns that the family may have. By doing this, the nurse can help family members to find their voice while building their self-efficacy for surrogate decision making.

Nurses and other healthcare team members also have an obligation to refrain from causing harm. Nurses often identify cardiopulmonary arrest and initiate CPR. Therefore, it is not uncommon for the nurse to be an assertive member of the healthcare team in addressing a patient's resuscitation status and ensuring that a DNAR order is initiated when appropriate. For the case at hand, it was important for the nurse to evaluate whether or not initiating CPR is harmful or not in advance of the sudden deterioration in Mr. K's condition. The emotional harm that cases such as Mr. K's present for the family is significant, and support for the family must be prioritized as well.

The case presented in this paper is an example of a common clinical scenario in which the role of the nurse is highlighted as an important part of patient and family centered care. The nurse not only cares for the physical but also the psychological well-being of the patient and family. However, the benefits of engaging the unbiased support of the CECS is invaluable to care planning and helps to ensure that the patient and family needs are being addressed in the most morally sound way, consistent with ethical standards. These collaborative partnerships help to ensure that the highest quality care is delivered and to ensure the best patient and family outcomes, even in the midst of unexpected or unfortunate patient circumstances such as the one presented in this article.

Acknowledgements:

Dr. Moss was funded by a T32 fellowship from the National Institutes of Health in Symptom Management and Palliative Care Research in Adults with Advanced Disease (T32NR01421304) in the Frances Payne Bolton School of Nursing at Case Western Reserve University when this manuscript idea was conceptualized.

References

1. President's Commission for the Study of Ethical Problems in Medicine and Biomedical and Behavioral Research. Making health care decisions: A report on the ethical and legal implications of informed consent in the patient-practitioner relationship. 1982;1
2. AUTHOR BLINDED.
3. Bibas L, Peretz-Larochelle M, Adhikari NK, et al. Association of surrogate decision-making interventions for critically ill adults with patient, family, and resource use outcomes: A systematic review and meta-analysis. *JAMA Network Open*. 2019;2(7):e197229–e197229. Doi: doi:10.1001/jamanetworkopen.2019.7229 [PubMed: 31322688]

4. Kon AA, Davidson JE, Morrison W, et al. Shared decision making in ICUs: An American College of Critical Care Medicine and American Thoracic Society policy statement. *Critical Care Medicine*. 2016;44(1):188–201. Doi: 10.1097/CCM.0000000000001396 [PubMed: 26509317]
5. Brock DW. The ideal of shared decision making between physicians and patients. *Kennedy Institute of Ethics Journal*. 1991;1(1):28–47. Doi: 10.1353/ken.0.0084 [PubMed: 10113819]
6. Truglio-Londrigan M, Slyer JT. Shared decision-making for nursing practice: An integrative review. *The Open Nursing Journal*. 2018;12:1–14. Doi: 10.2174/1874434601812010001 [PubMed: 29456779]
7. Buchanan AE, Brock DW. *Deciding for Others: The Ethics of Surrogate Decision Making*. Cambridge: Cambridge University Press; 1990.
8. Dworkin R *Life's Dominion: An Argument About Abortion and Euthanasia*. New York: Random House; 1993.
9. DuBois JM. *Ethics in Mental Health Research: Principles, Cases, and Guidance*. New York: Oxford University Press; 2008.
10. Ofoma UR, Basnet S, Berger A, et al. Trends in Survival After In-Hospital Cardiac Arrest During Nights and Weekends. *Journal of the American College of Cardiology*. 2018;71(4):402–411. 10.1016/j.jacc.2017.11.043 [PubMed: 29389356]
11. American Medical Association. *AMA Principles of Medical Ethics: I, IV*. 2016 <https://www.ama-assn.org/sites/ama-assn.org/files/corp/media-browser/principles-of-medical-ethics.pdf>
12. American Nurses Association. *Code of Ethics for Nurses*. 2015 <https://www.nursingworld.org/practice-policy/nursing-excellence/ethics/code-of-ethics-for-nurses/>
13. Hospice and Palliative Nurses Association. *HPNA Position Statement Withholding/Withdrawing Life-Sustaining Therapies*. 2016 https://advancingexpertcare.org/HPNA/Leadership/Position_Statements/HPNA/Leadership_Advocacy/Position_Statements.aspx?hkey=a794c17f-c88f-42d9-a09f-caf413e42aa3
14. American Academy of Hospice and Palliative Medicine. *Statement on Withholding and Withdrawing Nonbeneficial medical Interventions*. 2011 <http://aahpm.org/positions/withholding-nonbeneficial-interventions>
15. Cheon J, Coyle N, Wiegand DL, Welsh S. Ethical Issues Experienced by Hospice and Palliative Nurses. *Journal of Hospice & Palliative Nursing*. 2015;17(1):7–13. Doi: 10.1097/NJH.0000000000000129
16. Hajibabae F, Joolae S, Cheraghi MA, Salari P, Rodney P. Hospital/clinical ethics committees' notion: An overview. *J Med Ethics Hist Med*. 2016;9:17–17. [PubMed: 28523118]
17. de Brito GMG, de Oliveira Santa Rosa D. Nurses performance in clinical ethics committees and commissions: An integrative review. *Nursing Ethics*. 2017;26(3):688–699. Doi: 10.1177/0969733017724611 [PubMed: 28933257]
18. Hoskins K, Grady C, & Ulrich C. Ethics education in nursing: Instruction for future generations of nurses. *The Online Journal of Issues in Nursing*. 2018;23(1). Doi: 10.3912/OJIN.Vol23No01Man03
19. McCabe MS. CE Ethics Series. Reenergizing the Hospital Ethics Committee. *Journal of hospice and palliative nursing*. 2015;17(6):480–489. Doi: 10.1097/NJH.0000000000000190
20. Bartlett VL, Finder SG. Lessons learned from nurses' requests for ethics consultation: Why did they call and what did they value? *Nursing Ethics*. 2018;25(5):601–617. Doi: 10.1177/0969733016660879 [PubMed: 27514740]

Table 1:

Use of SFNO for Ethical Decision Making

Component	Questions to Consider	Possible Responses
Stakeholder	<ul style="list-style-type: none"> Who has a stake in the decisions being made? Should one stakeholder be prioritized over the other(s)? 	<ul style="list-style-type: none"> Mr. K., his parents, and healthcare team
Facts	<ul style="list-style-type: none"> What are the pertinent facts? 	<ul style="list-style-type: none"> End stage liver disease with no curative options No written advance directive Mr. K. identified his father as his decision maker on admission Father says “do everything” but does not visit often, and will not engage in discussions with the healthcare team Mother says Mr. K. wants to be home with his kids and nothing more Mr. K requires daily blood transfusions for hemodynamic stability Full code status
Norms	<ul style="list-style-type: none"> What are the relevant ethical principles, norms, and values? What is the conflict at hand? 	<ul style="list-style-type: none"> Principles of autonomy, beneficence, and nonmaleficence Uncertainty exists about whether or not blood transfusions should continue, Mr. K should be taken to IR, and if CPR should be performed in the event of cardiopulmonary arrest
Options	<ul style="list-style-type: none"> Which actions should be considered? Is there a compromise? 	<ul style="list-style-type: none"> Perform the IR procedure or do not perform Continue blood transfusions indefinitely, continue for a short time, or discontinue now Perform CPR or withhold CPR in event of arrest