Identification of eukaryotic translation initiation factors and the temperature-dependent nature of *Turnip mosaic virus* epidemics in allopolyploid *Brassica juncea*

Jannat Shopan¹ · Chang Liu¹ · Zhongyuan Hu¹ · Mingfang Zhang^{1,2} · Jinghua Yang^{1,2}

Received: 27 May 2019 / Accepted: 6 January 2020 / Published online: 28 January 2020
© King Abdulaziz City for Science and Technology 2020

Abstract

Eukaryotic translation initiation factors (*eIFs*) are essential protein complexes involved in the translation of mRNA into proteins. These initiation factors are generally used as targets in the control of plant RNA virus infections. In the present study, we identified a total 190 *eIFs*, clustered phylogenetically into 40 distinct subfamilies in the allopolyploid *Brassica juncea*. Extensive evolutionary duplications of the *eIFs* in *B. juncea* suggest their increased genetic diversity and wide adaptability. The induction of expressions in some of the *eIFs* after inoculation against *Turnip mosaic virus* (TuMV) provided candidate targets to be used in the control of viral infections. In addition, the expression profiles of *eIFs* under different temperatures suggested that the TuMV epidemic was temperature dependent. The *eIFs* expressions suggested that the systemic viral infections were more acute in plants grown between 20 °C and 28 °C. In addition, our results revealed that new subgroups of *eIFs*, *eIF2β*, *eIF2α*, *eIF2Bβ*, *EF1A*, and *PABP* could be represented as targets for antiviral strategies in *B. juncea*. In summary, our findings would be helpful in studying the complex mechanisms of *eIF*-mediated, temperature-dependent RNA virus control in *B. juncea*.

Keywords Eukaryotic translation initiation factors · Phylogenetic evolution · Virus epidemic · Duplication

Introduction

Eukaryotic translation initiation factors (*eIFs*) consist of several phylogenetic subfamilies that are required for the initiation of protein synthesis through multiphase processes (Jackson et al. 2010). The translation of mRNA into protein is a complex process, which requires the 43S and 48S initiation complexes. The 43S complex consists of the 40S ribosomal subunit, some of the *eIFs* (*eIF4E*, *eIF4G*, *eIF4A*,

PABP and *eIF3*), and the initiator tRNA bound to the capped mRNA formed by the *eIF4E* (cap-binding) subunit at the 5'-end. The recognition of the start codon influences the formation of the 48S complex; the 48S and 60S subunits together form the 80S ribosome, following which the protein synthesis begins at the ribosomal P site (Jackson et al. 2010).

Over 40 *eIF* subfamilies have been identified in plant and animal species; however, only few of them have been characterized functionally (Li et al. 2019). The *eIFs* are involved in various aspects of plant growth and development as well as disease resistance. For example, *eIF2Ba* is required for the translation suppression of the vesicular stomatitis virus, which suggests its regulatory role in viral infections in mammalian cells (Elsby et al. 2011). Moreover, *eIF2B* is a key regulator of the guanine nucleotide exchange factor (GEF), which is responsible for the recycling of the heterotrimeric G-protein, *eIF2*, to permit translation initiation (Wortham and Proud 2015). Similarly, *eEF1A*, *eEF1B*, *eIF4B*, and poly(A)-binding proteins (*PABPs*) were reported to be related with viral infections in many plants (Jackson et al. 2010; Li et al. 2019; Hwang et al. 2013). While *eIF2Bβ*, *eIF4E*, *eIF(iso)4E*, *eIF4G*, and *eIF(iso)4G* have been seen to

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s13205-020-2058-0>) contains supplementary material, which is available to authorized users.

✉ Jinghua Yang
yangjinghua@zju.edu.cn

¹ Laboratory of Germplasm Innovation and Molecular Breeding, Institute of Vegetable Science, Zhejiang University, Hangzhou 310058, China

² Key Laboratory of Horticultural Plant Growth, Development and Quality Improvement, Ministry of Agriculture, Hangzhou 310058, China

establish recessive resistance to plant viral infections (Shopan et al. 2017), they have also been found to help translate viral RNA, regulate their replication, and aid in their systemic movement with the plant cells (Hwang et al. 2013).

Turnip mosaic virus (TuMV) is a member of the potyvirus genus affecting many plant species and is a limiting factor in *Brassica* production. The recessive resistance genes of potyvirus as well as of several other genera have been mapped from the *eIFs* of the *eIF4* family (Shopan et al. 2017). For example, *eIFiso4E* (*lsp1* locus) is responsible for TuMV RNA translation and the systemic movement of the virus in *Arabidopsis* (Duprat et al. 2002). Two important, similar TuMV resistance genes, *retr01* and *retr02*, have been isolated from *eIFiso4E* in *Brassica rapa* (Qian et al. 2013). A few other potyvirus resistance loci, such as *Clover yellow vein virus*, *Pepper vein mottle virus*, *Chilli vein mottle virus*, *Potato virus Y*, *Tobacco etch virus*, *Pepper mottle virus*, *Lettuce mosaic virus*, *Pea seed-borne mosaic virus*, and *Bean yellow mosaic virus*, were also isolated from *eIF4E* or *eIF(iso)4E* (Sato et al. 2005; Truniger and Aranda 2009; Duprat et al. 2002). In *Poaceae* species, *eIF(iso)4G* is correlated with *Sobemovirus* resistance in rice, with the *tsv1* and *rymv-1* loci controlling the systemic movement of *Rice yellow mottle virus* (Truniger and Aranda 2009). Similarly, *eIF4E* has been found to be resistant to virus genera such as *Carmovirus* in melon, with the *nsv* locus controlling the *Melon necrotic spot virus* and *Bymovirus* in barley, with the *rym4*, *rym5*, and *rym6* loci controlling the *Barley mild mosaic virus* (BaMMV) (Truniger and Aranda 2009). In addition, recently, a dominant TuMV resistance gene *RTM3* (Restricted TEV Movement 3) has been cloned from the *eIF3b* subfamily in *Arabidopsis thaliana* (Rubio et al. 2019).

In this study, we identified the eukaryotic translation initiation factor genes in the allopolyploid *B. juncea* genome. We investigated the effects of temperature on TuMV systemic infections and their symptoms in *B. juncea*. In addition, we also investigated the expression profiles of *BjuEIFs* in response to TuMV inoculation, which provided useful clues for choosing the suitable TuMV resistance responsive candidates for further functional analyses.

Materials and methods

Plant materials, TuMV inoculation, and treatment

Brassica juncea var. STZ plants were grown in a growth chamber with the day and night temperatures maintained at 22 °C and 20 °C, respectively, and with a 16-h photoperiod. The TuMV_ZJ isolate was prepared for inoculation. The inoculation was done according to the method described by Walsh et al. (1999) and Shopan et al. (2017). Twenty plants were included for each temperature treatment. The

inoculated plants were monitored regularly, for a period of 4 weeks, for symptom development. To study the effects of temperature on the systemic infection by TuMV, the inoculated plants were initially maintained at 22 °C. After 20 days, they were transferred to growth chambers maintained at temperatures 10 °C, 15 °C, 20 °C, 28 °C, and 35 °C, respectively, for 5 days. The upper, newly developed, uninoculated leaves were collected at different times for the quantitative reverse transcription polymerase chain reaction (qRT-PCR) measurements. The experiments were replicated three times.

Identification of *eIFs* in *B. juncea*

The coding sequences of 95 *A. thaliana* *eIFs* from the TAIR database (<https://arabidopsis.org/index.jsp>) were used as the set of reference genes (Swarbreck et al. 2007) (Table S1). The homolog sequences of *B. rapa* *eIFs* were taken from the BRAD database (<https://brassicadb.org/brad/>) (Cheng et al. 2011). The sequences of *B. juncea* and *Brassica nigra* were obtained from the BRAD database (<https://brassicadb.org/brad/>) (Yang et al. 2018). Homologs of *eIFs* in other species were taken from Phytozome v10.3 (<https://www.phytozome.net/>) (Goodstein et al. 2012). *Brassica oleracea* and *Brassica napus* *eIFs* were obtained from the Bolbase (<https://ocri-genomics.org/bolbase/>) (Liu et al. 2014) and Genoscope (<https://www.genoscope.cns.fr/blat-server/cgi-bin/colza/webBlat>) databases, respectively. The existence of the *eIF* core sequences in candidate genes was confirmed by the Pfam (<https://pfam.sanger.ac.uk/search>) and SMART (<https://smart.embl-heidelberg.de/>) web-based resources. In addition, we also checked the conserved domains of the identified *eIF* proteins with the help of the National Center for Biotechnology Information (NCBI) database (<https://www.ncbi.nlm.nih.gov/Structure/cdd/cgi>).

Bioinformatics analysis of *eIFs*

For phylogenetic analysis, the protein sequences of the *eIFs* were classified using the Clustal W software using default parameters (Thompson et al. 2003). The resulting sequences were used to construct a phylogenetic tree using the neighbor-joining (NJ) method, and the bootstrap values were deliberated with 1000 replications using MEGA 5.0 (Tamura et al. 2011). Furthermore, the online tool ProtComp was used to predict the subcellular localization (ProtComp 9.0: <https://linux1.softberry.com/berry.phtml?topic=protcomppl&group=programs&subgroup=proloc>) of *eIFs* in *B. juncea*. Following this, the chromosomal locations of the *eIF* genes in *A. thaliana* and *B. juncea* were determined from the TAIR and BRAD genomics databases, respectively. The CIRCOS software was used for comparative analysis of the multiple genomes by adopting the circular layouts and

minimizing the inherent difficulties in visualizing complex genomic data (Krzywinski et al. 2009). Besides, to identify the conserved motifs of the *eIFs* of *B. juncea*, the online program Multiple Expectation Maximization for Motif Elicitation (MEME, <https://meme-suite.org/V4.9.0>) was used using default parameters. Pfam and SMART were used to check the conserved domains (https://smart.embl-heidelberg.de/smart/set_mode.cgi?NORMAL=1). The DOG 1.0 software (<https://dog.biocuckoo.org/>) was used to construct the protein structures of *BjuAeIFs*. The exon–intron organization of *B. juncea eIFs* was determined by comparing the predicted coding sequences with their corresponding full-length sequences using the online program Gene Structure Display Server (GSDS, <https://gsds.cbi.pku.edu.cn>). The information on protein phosphorylation sites and signal peptides were manifested based on the data collected from the NetPhos 3.1 Server (<https://www.cbs.dtu.dk/services/NetPhos/>) and the SignalP online program (<https://www.cbs.dtu.dk/services/SignalP/>), respectively. In addition, the *cis*-regulatory elements were predicted using the PlantCARE (<https://bioinformatics.psb.ugent.be/webtools/plantcare/html/>) database (Lescot et al. 2002). The graphic alignment tool for the comparative sequence analysis (GATA 0.7) program was used to perform a comparative analysis of the promoter regions (Nix and Eisen 2005).

K_s value analysis

All *A. thaliana eIF* coding sequences were aligned with subgenomes A and B of *B. juncea* using Clustal W (Thompson et al. 2003). The Nei and Gojobori method was used to calculate the K_a and K_s values, as previously described (Zhang et al. 2006). During K_s value analysis, the alignments of the coding sequences were regulated by applying an in-house Perl script. The K_s values of *B. juncea* A and B subgenome *eIFs* were also analyzed and plotted as histograms.

Expression profiling analysis

The expression profiles of the *eIFs* were evaluated under TuMV infection using qRT-PCR. Total RNA was extracted from the TuMV-infected [0, 3, 6, 9, and 12 days post-inoculation (DPI)] tissue samples using the Trizol reagent (Invitrogen, Carlsbad, CA, USA). The first-strand cDNAs were synthesized using the Transcriptor First-Strand cDNA Synthesis kit (Roche, USA). Quantification of the RT-PCR reactions was performed with the ABI PRISM 7900HT sequence detection system (Applied Biosystems, USA) using FastStart Universal SYBR Green Master (Roche), following the manufacturer's instructions. The AF128094/qRT-25S gene was used as the reference in all qRT-PCR analyses. Each treatment was conducted independently and in triplicate. The relative expression levels of the *eIFs* were calculated

according to the method discussed previously (Livak and Schmittgen 2001). The primers used in this analysis are listed in Table S8. The expression profiles were analyzed and visualized using the MeV4.9.0 software (The Institute for Genomic Research, USA). The primers (Table S8) of *BjuEIFs* were designed by GenScript (<https://www.genscript.com/ssl-bin/app/primer>).

Results

Genome-wide identification of *eIFs* in *B. juncea*

In the present study, we obtained a total of 190 *eIFs* from the *B. juncea* genome, 99 and 91 from subgenomes A and B, respectively (Table S1). In addition, we studied the characteristics of the *eIF* genes in *B. juncea*, including coding sequence length (CDS), amino acid (aa) residue length, chromosome distribution, and subcellular localization (Fig. S1, Tables S2 and S4). Based on the subcellular localization predictions, we conclude that most of the *eIFs* in *B. juncea* were probably located in the cytoplasm and the rest in various organelles (Table S4).

Gene duplication plays a very important role in gene family expansion and protein functional diversification (Lynch and Conery 2000). To explore the contribution of duplication events in *eIFs*, we analyzed the occurrence of tandem duplications during the evolution of this *eIFs* gene family. From phylogenetic analysis and chromosomal localization of *BjuEIFs*, we identified a total of 33 duplicate gene pairs for the *B. juncea* subgenome A (Fig. 2) and 4 gene pairs for the tandem duplicate (Table S3). Similarly, 35 pairs of duplicate/triplicate genes were identified for the *B. juncea* subgenome B (Figs. S1 and S2, Table S3). Almost all of the *BjuEIF* duplicate genes contained similar domain and motifs (Figs. 2, S2, Table S3). These findings suggested that gene duplication events contributed to the expansion of the *eIF* gene family in *B. juncea*.

To explore the evolutionary relationship among the *eIFs*, a total of 543 *eIF* domain sequences were predicted from *A. thaliana*, *B. rapa*, *B. nigra*, and *B. juncea* using alignments of full-length *eIF* protein sequences (Fig. 1, Table S1). They were then classified into 40 distinct groups based on their phylogenetic trees. The 190 *BjuEIFs* were distributed throughout most of the groups, indicating the various functions of *eIFs* in *B. juncea* (Table S1).

Gene structure, conserved domains, and motif analysis of *eIFs* in *B. juncea*

Phylogenetic analysis was carried out using the full-length conserved aa sequences of the 190 *B. juncea eIFs* (Figs. 2a, S2a). Based on the NJ phylogenetic tree, they were divided

Fig. 1 Phylogenetic tree of *eIF* proteins from *A. thaliana*, *B. rapa*, *B. nigra*, and *B. juncea* (a, b). The phylogenetic tree is based on a sequence alignment of 543 *eIF* protein sequences from 5 genomes.

The unrooted tree was generated with Clustal W using the NJ method. Bootstrap values from 1000 replicated at each node. The *eIF* proteins are grouped into 40 variant clades

into 39 subfamilies for *BjuAeIFs* and 38 subfamilies for *BjuBeIFs*. No member was found corresponding to the subfamily *eIF4G* in the *B. juncea* subgenome A. Similarly, no members corresponding to subfamilies *eIF3e* and *eIF3h* were found in the *B. juncea* subgenome B, suggesting that they may have been lost during the process of evolution. The exon/intron structures of *BjuAeIFs* and *BjuBeIFs* were examined to gain further insight into their possible structural evolution (Figs. 2b, S2B). Our results indicated that there is a strong correlation between phylogeny and the exon/intron structure, suggesting that genes clustering together generally possessed similar structures. In brief, all members of subfamily *eIF2Bβ* (A and B subgenomes) had nine exons and eight introns. On the other hand, members of subfamily *eIF4E* and those of its isoform *eIFiso4E* showed two

categories, one having five exons and four introns and the other having four exons and three introns. The observation suggests that those genes may have fully or partially lost their coding sequence during evolution.

To provide further confirmation of the evolutionary relationships among the *BjuAeIFs*, we determined the distribution of their conserved domains (Fig. 2c) and observed the diversity of *BjuBeIFs*. The results showed that 30 corresponding distinct motifs exist in the *B. juncea* subgenome B (Fig. S2C, Tables 1, S6). By consensus, an *eIF* protein has the following basic structure: a low-complexity domain and an *eIF/EF/IF/eRF/MIF/MA* domain with active sites. Among the 99 *BjuAeIF* proteins, 51 contain at least 1 low-complexity domain. No signal peptide was found in both *BjuAeIF* and *BjuBeIF*

Fig. 2 The *eIF* genes in the *B. juncea* subgenome A. **a** Phylogenetic analysis of *B. juncea* eIF proteins. Numbers above or below branches of the tree indicate bootstrap values. **b** Exon-intron structures of *B.*

juncea eIF genes. **c** The distribution of conserved domains in *B. juncea* subgenome A eIF proteins. The relative positions of each domain within each protein are shown in color

Table 1 Number of different *cis*-elements presented in the promoter regions of the temperature-responsive *eIF* genes

Gene name	AAGAA-motif	MBS	LTR	ABRE	TCA-element	HSE	ERE	ARE	TGACG-motif	CGTCA-motif	GCN4_motif	MRE	G-Box	Skn-1_motif	GARE-motif	GATA-motif
<i>BjuA020776</i>		2	2		1	1		3	2	3				3		1
<i>BjuA028636</i>	1	4	1	1	2			2				1	1	1		
<i>BjuA022587</i>		1	1	1			1	1	1	1		1	2			1
<i>BjuA037803</i>	1	1	1	1	1	1	1	1					2			
<i>BjuA009417</i>	1		1	1	2			1	1	1			2		1	1
<i>BjuA037087</i>			1	1					1			1	2			1
<i>BjuB016629</i>		1	1	3	1				1				1			
<i>BjuB013529</i>	2	3	3		1			1	3	3			2	2	1	1
<i>BjuB022871</i>	1	3	2					3	1	1	1			4	1	1
<i>BjuB035362</i>	1	3	1			1		3				1	1	1		1
<i>BjuB032683</i>			1	3			1	1				1	1			1
<i>BjuB005408</i>	1		1	1	1			1	1	1			2			

(Figs. S7, S8). In addition, ten *BjuAeIF* proteins had an RRM domain and ten others had a PolyA domain. The protein of the members of a typical subfamily *eIF4A* has a DEXDc domain and a HELICc domain. Thirteen *BjuAeIF* proteins were identified in the PINT domain and three in the transmembrane domain. We identified the motif sequences of *BjuBeIFs*; of these, a total of 30 distinct motifs were identified (Figs. S2C, S3). All members of the *eIF2Ba* subfamily have their specific motif serials such as 27, 23, 11, 16, 19, and 9. Overall, our findings revealed that the specific functions of each subfamily might be distinct for specific motifs (Fig. S3).

***eIF2B* (α , β , δ), *eIF4E*, *eIFiso4E* and *eIF4G* subfamily genes**

Based on our previous studies, we have chosen the *eIF2B* (α , β , δ), *eIF4E*, *eIFiso4E*, *eIF4G* subfamily genes, which have induced responses after TuMV infections (Shopan et al. 2017; Qian et al. 2013; Truniger and Aranda 2009). The genomic data of *eIF2B* (α , β , δ), *eIF4E*, *eIFiso4E*, and *eIF4G* from the angiosperm were searched from the NCBI and Phytozome databases using the BLASTn program (Benson et al. 2013; Goodstein et al. 2012). Based on the phylogenetic analysis, we found that *eIFs* present in the same group are likely to possess similar functions (Figs. 3, 4). In addition, we also investigate the evolutionary relationships among these group genes from different *Brassicaceae* species (Fig. 4). In this result we have confirmed the duplication of genes and their specializations in *Brassicaceae* species (Bowers et al. 2003).

K_s value analysis

We have calculated the K_s (synonymous substitution rates) and K_a (nonsynonymous substitution rates) values using the orthologs identified for *B. juncea* and *A. thaliana* *eIFs*. The results indicated that the K_a/K_s ratios of a majority of *eIF* orthologs were less than 1, representing purifying selection on the *eIFs* (Fig. 5a) (Duan et al. 2015). The K_s values of the *B. juncea* *eIFs* ranged from 0.4 to 0.6 and averaged 0.5 (~24 Myr). The homologous members of a single gene showed different evolutionary rates, but the divergence of the homologous members was mainly concurrent with the *Brassica*-specific whole-genome triplication (WGT) event. Simultaneously, we also investigated the divergence of the *eIFs* and found that the synonymous substitution rate (K_s) was less than 1 for all others except those from the *eIF1* and *EF2* subgroup genes (Fig. 5b).

Fig. 3 Cladogram of *Brassicaceae* *eIF2B* (α , β , δ), *eIF4E*, *eIFiso4E*, and *eIF4G*-like proteins in relation to the conserved *eIF2B* (α , β , δ), *eIF4E*, *eIFiso4E*, and *eIF4G* proteins of angiosperms/flowering plants, which have the same homologous genes

Transcriptional expression of *eIFs* in *B. juncea*

The 5'UTR region (the promoter and enhancer region) of a gene contains a *cis*-regulatory element that controls the gene expression patterns (Zhang et al. 2005; Lv et al. 2016). We observed the putative enhancer *cis*-elements in the promoter regions of some of the *eIF* genes; a few of the subfamily genes among them were found to have positive responses for TuMV infections (Tables 1, S6). Based on the *cis*-regulatory elements in the 2 kb promoter regions of *eIFs* in *B. juncea*, we showed that all *eIF2β*, *eIF2α*, *eIF2Bβ*, *poly(A)*, *EF1A*, *eIF4E*, and *eIFiso4E* subfamily genes contain at least one *cis*-element type in their promoter sequences (Tables 1, S6, S7), suggesting that these *eIFs* might be involved in TuMV

resistance. Interestingly, the promoter regions of all the duplicate gene pairs of *eIF2Bβ*, *eIF4E*, and *eIFiso4E* subfamily genes showed their divergence, though observed in conserved regions (Fig. S4), as seen in our previous studies (Shopan et al. 2017; Qian et al. 2013). Furthermore, a protein's subcellular localization plays an extremely important role in understanding specific functions and biological processes in cells (Wallbach et al. 2016). On the basis of localization prediction, we speculate that *eIF2Bβ* might be located in the plasma membrane; *eIF4E* and *eIFiso4E* in the extracellular matrix/peroxisome, and *eIF4G* in the nucleus. In addition, protein phosphorylation is the most abundant post-translational modification and a key regulatory step in eukaryotic signal transduction (Kumer and

Fig. 4 Copy number variation in the *eIF2B* (α , β , δ , and γ), *eIF4E*, *eIFiso4E*, and *eIF4G* genes in *Brassicaceae* crops. The phylogenetic trees of the *AteIF* genes is shown in the left and the species topology tree at the top. The phylogenetic tree of the *AteIF* genes was constructed using protein sequences with MEGA5 using the NJ method. Copy numbers of each gene in *A. thaliana* (Ath), *Arabidopsis lyrata* (Aly), *B. rapa* (Bra), *B. nigra* (Bni), *B. oleracea* (Bol), *B. juncea* (Bju), *B. napus* (Bna), *Boechera stricta* (Bst), *Capsella grandiflora* (Cgr), *Capsella rubella* (Cru), and *Eutrema salsugineum* (Esa)

Fig. 5 Pairwise comparisons of K_s values of *eIF*-related genes. **a** The distribution of K_s values of *eIF*-related genes between *A. thaliana* vs. BjuA and *A. thaliana* vs. BjuB. **b** The distribution of K_s values of *eIF*-related genes between BjuA vs. BjuB. The blue mark indicates

the main concentrated area of the K_s value (0.4–0.5). The K_s values of these *eIF*-related genes ranged from 0.4 to 0.6 and averaged 0.5 (~24 Myr)

Vrana 1996). Based on *eIF* protein sequence phosphorylation site prediction, results showed that every *eIF* protein sequence contains three types of phosphorylation sites such

as Serine (S), Threonine (T), and Tyrosine (Y) residues, with S phosphorylation being the most common (Figs. S5 and S6, Table S5). These phosphorylation sites might be involved in

the regulation of protein activities when plants are subject to stress.

Temperature-dependent systemic infection of TuMV in *B. juncea*

Based on the reported results, a number of recessive resistance genes of TuMV are shown to encode *eIFs*, which include *eIF4E* and *eIF4G* and their isoforms, *eIF(iso)4E*, and *eIF(iso)4G* (Nicaise 2014; Revers and Nicaise 2014). On the other hand, the *eIF2B α* subunit has been shown to be required for the translation suppression of vesicular stomatitis virus, suggesting its regulatory role in viral infection in mammals (Elsby et al. 2011). In our present study, we chose *eIF4E*, *eIF4G*, and their isoforms *eIF(iso)4E* and *eIF(iso)4G*, *eIF2B* (α , β , δ , and ϵ), and *RF-S* subfamily genes to analyze the expression profiles at intervals of different days after TuMV infection (Fig. 6). Five *eIFs*, namely *eIF2B β* (*BjuA006209*), *eIF4E* (*BjuB006445*), two *eIF(iso)4E* (*BjuA018845*, *BjuA014379*), and *eIF4G* (*BjuB023035*), showed positive responses quickly and persistently. Among them, the gene expressions of *BjuA006209* and *BjuA018845* increased gradually until a maximum rate

was reached at 12 days. In contrast, we found that other *eIFs* had no response to TuMV infections. We also compared the expressions of the segmentally duplicated genes (*BjuA006209* duplicate with *BjuB006522*) and found that they diverged significantly in their expression patterns (Fig. 6). This implies the association of *eIF2B β* , *eIF4E*, and *eIF4G* and their isoforms, *eIF(iso)4E*, and *eIF(iso)4G* with TuMV resistance; it may be noted that *eIF2B β* is the first gene that acted as a resistance gene to a plant virus.

To study the effects of temperature on the systemic infections of TuMV in *Brassicaceae*, we inoculated the plants with TuMV at 22 °C for 20 days. The plants were then transferred to a growth chamber with different temperature settings (10 °C, 15 °C, 20 °C, 28 °C, and 35 °C) for 1 week and were moved back to 22 °C (Chung et al. 2015, 2017; Parizipour et al. 2018). The results suggested that 20–28 °C is the most favorable temperature range for TuMV virus proliferation in plants. The results indicated that the systemic infection rate of TuMV is slow below 15 °C, and that the viral RNA levels began to gradually fall at 35 °C. We thus inferred that RNA silencing increased at 35 °C, consistent with the data of previous reports (Chung et al. 2017; Velazquez et al. 2010). Accordingly,

Fig. 6 Expression profiles of *B. juncea* *eIF2B α* , *eIF2B β* , *eIF2B δ* , *eIF2B ϵ* , *eIF4G*, *eIFiso4G*, *eIF4E*, and *eIFiso4E* subfamily genes after TuMV infection in different intervals. *RF-S* subgroup-related genes were used as the control. Upper leaf samples were collected at 0, 3, 6, 9, and 12 DPI

Fig. 7 The effects of temperature on the transcript abundance of *eIF* family genes after 2 weeks of TuMV_ZJ inoculation

the observed reduction in systemic infections at 35 °C may be attributable to the increased host resistance mediated by enhanced RNA silencing at higher temperatures (Chung et al. 2017). Under temperature treatment, most of the detected *eIFs* showed positive responses at 28 °C (Fig. 7), which may be induced by the systemic infection of TuMV of *B. juncea*. Among the detected *eIFs*, *BjuB032683* (*poly(A)*), *BjuB005408* (*poly(A)*), *BjuA037803* (*poly(A)*), *BjuA022587* (*eIF2β*), *BjuA009417* (*EF1A*), *BjuB016629*

(*eIF2α*), and *BjuB035362* (*eIFiso4E*) exhibited a quicker response in terms of their expression levels to temperature than the others. Furthermore, the LTR *cis*-regulatory element, which is responsible for low-temperature inducibility, could only be found in the promoters of some of the *eIFs* (Fig. 7, Tables 1, S6). All these *eIFs* were found to be upregulated by TuMV infection under 28 °C. The results suggested that such *eIFs* have a positive involvement in the biotic and abiotic stress signal pathways.

Discussion

eIFs are characterized by the initiation of eukaryotic protein synthesis, which is a multistep process requiring a large number of initiation factors, and constitutes one of the largest protein families. The functions of these factors are to recruit mRNAs to the 40S ribosomal subunit by scanning ribosomes along with the recruited mRNA to identify the appropriate start codon. Hence, *eIFs* facilitate the recruitment of the 60S ribosomal subunit to allow translation to commence (Jackson et al. 2010). In this study, we showed that the evolutionary meaning of extensive duplication of *eIFs* indicates their genetic complexity, novelty, diversity, and wide adaptation (Yang et al. 2018). The *eIFs* are highly conserved in eukaryotes, with nearly 60% identity in *A. thaliana* and *B. juncea* orthologs. The orthologs did not differ significantly between the subgenomes A and B, as confirmed by the K_s value analysis. Segmental duplication, tandem duplication, whole genome duplication (WGD), and WGT events were the main causes for gene family expansion (Edger and Pires 2009; Duan et al. 2015; You et al. 2015). In our study, the tandem duplication suggested that gene duplications are the main contributors to the expansion of the *B. juncea eIFs* family. After duplication events, some gene copies sustain, because they have important functions, while those that are functionally redundant may be lost (Qian et al. 2010). *B. juncea* has undergone WGT events with both of its ancestors, *B. rapa* and *B. nigra*, providing a resource for studying the evolution of polyploid genomes (Wang et al. 2011; Yang et al. 2018).

eIFs are simultaneously involved in different aspects of plant growth and development, including pathogen and virus responses (Truniger and Aranda 2009). In this study, we were especially interested in identifying the *eIFs* responsible for TuMV resistance in *B. juncea*. In addition, *eIFs* in same group are likely to perform similar functions. For example, groups *eIF4E*, *eIFiso4E*, and *eIF4G* were involved in potyvirus resistance (Duprat et al. 2002; Revers and Nicaise 2014; Nicaise 2014), while *eIF4B*, *eEF1A*, *eEF1B*, and *PABPs* were found to be involved in other viral infections in many plant species (Li et al. 2019; Kato et al. 2010). Our results also found some new subgroups of *eIFs*, namely, *eIF2β*, *eIF2α*, *eIF2Bβ*, *EF1A*, and *poly(A)*, representing new targets and mechanisms for antiviral strategies in *B. juncea* (Figs. 6, 7).

Eukaryotic translation initiation is a particularly complicated stage of gene expression, which is required for some initiation factors. Moreover, *cis*-elements and expression profile analysis of *BjueIFs* provided further evidences for their involvement in TuMV resistance. At least one of stress-related *cis*-elements was detected

in the promoter regions of the *BjueIFs*. Our qRT-PCR results showed that the expressions of the *BjuA006209* and *BjuA018845* genes were highly inducible by TuMV among the 30 putative TuMV-related *BjueIFs* (Fig. 6). Furthermore, the increase in temperature was found to negatively affect the expression of symptoms caused by certain plant viruses. For example, the systemic symptoms of TuMV in Chinese cabbage increased with increasing temperature from 23 to 28 °C (Chung et al. 2015). Potato leafroll virus caused more severe symptoms in *Physalis floridana* kept between 20 °C and 25 °C (Chung et al. 2017). Similarly, our study under different temperature treatments showed that the expressions of some of the *eIFs* were significantly upregulated at 28 °C (Fig. 7). In addition, the promoter of *BjuA006209* contains stress-responsive *cis*-elements such as HSE, and, the promoter of some other *eIFs* contains stress-responsive *cis*-elements such as LTR, which suggested that the presence of such *cis*-elements plays an important role in regulating the gene expression levels in response to TuMV infections (Zhang et al. 2005; Lv et al. 2016). Moreover, many of *BjueIFs* were downregulated against TuMV infection, indicating that the proteins might act as negative regulators in resistance responses. The functions of the *BjueIFs* in TuMV resistance will be further characterized in subsequent work. The dynamics of plant virus epidemics are likely to be influenced by the direct consequences of climate change, because increased temperature could directly influence the abundance and activity of transmission vectors (Trezbicki et al. 2015). Some recent studies have reported the impact of high temperature, water deficit, amount of light, and CO₂ concentration changes on both the rates of infection and transmission efficiencies of plant viruses (Hily et al. 2016; Van Munster et al. 2017; Trębicki et al. 2017; Yvon et al. 2017; Chung et al. 2017). Therefore, it is worth investigating the mechanism of these *eIFs* in response to environmental changes to find a more stable and broad-spectrum resistance in future.

Conclusions

In conclusion, we demonstrated the expression profile of *eIFs* in response to TuMV infection in *B. juncea*. The *B. juncea eIFs* from different subfamilies exhibited diverse responsive patterns to environmental changes. The results also uncovered the candidate-resistant genes to TuMV. Further studies are required to elucidate the role of *eIFs* in the regulation of potyvirus resistance. Our findings provide valuable clues for advance functional research on the *eIF* family in *Brassicaceae* crops and its improvement in adaptation to TuMV resistance via molecular approaches.

Acknowledgements This work was supported by grants from the National Natural Science Foundation of Zhejiang Province (Grant No. Z20C150008) and the National Natural Science Foundation of China (Grant No. 31872095). We thank LetPub (<https://www.letpub.com>) for its linguistic assistance during the preparation of this manuscript.

Compliance with ethical standards

Conflict of interest There is no conflict of interest.

References

- Benson D, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman D, Ostell J, Sayers E (2013) GenBank. *Nucleic Acids Res* 41(D1):D36–D42
- Bowers JE, Chapman BA, Rong JK, Paterson AH (2003) Unravelling angiosperm genome evolution by phylogenetic analysis of chromosomal duplication events. *Nature* 422(6930):433–438
- Cheng F, Liu SY, Wu J, Fang L, Sun SL, Liu B, Li PX, Hua W, Wang XW (2011) BRAD, the genetics and genomics database for *Brassica* plants. *BMC Plant Biol* 11:136
- Chung BN, Choi KS, Ahn JJ, Joa JH, Do KS, Park KS (2015) Effects of temperature on systemic infection and symptom expression of Turnip mosaic virus in Chinese cabbage (*Brassica campestris*). *Plant Pathol J* 31(4):363–370
- Chung BN, Canto T, Tenllado F, Choi KS, Joa JH, Ahn JJ, Kim CH, Do KS (2017) The effects of high temperature on infection by potato virus Y, potato virus A, and Potato leafroll virus (vol 32, pg 321, 2016). *Plant Pathol J* 33(1):101–101
- Duan WK, Song XM, Liu TK, Huang ZN, Ren J, Hou XL, Du JC, Li Y (2015) Patterns of evolutionary conservation of ascorbic acid-related genes following whole-genome triplication in *Brassica rapa*. *Genome Biol Evol* 7(1):299–313
- Duprat A, Caranta C, Revers F, Menand B, Browning KS, Robaglia C (2002) The Arabidopsis eukaryotic initiation factor (iso)4E is dispensable for plant growth but required for susceptibility to potyviruses. *Plant J* 32(6):927–934
- Edger PP, Pires JC (2009) Gene and genome duplications: the impact of dosage-sensitivity on the fate of nuclear genes. *Chromosome Res* 17(5):699–717
- Elsby R, Heiber JF, Reid P, Kimball SR, Pavitt GD, Barber GN (2011) The alpha subunit of eukaryotic initiation factor 2B (*eIF2B*) is required for *eIF2*-mediated translational suppression of vesicular stomatitis virus. *J Virol* 85(19):9716–9725
- Goodstein DM, Shu SQ, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U, Putnam N, Rokhsar DS (2012) Phytozome: a comparative platform for green plant genomics. *Nucleic Acids Res* 40(D1):D1178–D1186
- Hily JM, Poulicard N, Mora MÁ, Pagán I, García-Arenal F (2016) Environment and host genotype determine the outcome of a plant–virus interaction: from antagonism to mutualism. *New Phytol* 209(2):812–822
- Hwang J, Oh CS, Kang BC (2013) Translation elongation factor 1B (*eEF1B*) is an essential host factor for Tobacco mosaic virus infection in plants. *Virology* 439(2):105–114
- Jackson RJ, Hellen CUT, Pestova TV (2010) The mechanism of eukaryotic translation initiation and principles of its regulation. *Nat Rev Mol Cell Biol* 11(2):113–127
- Kato Y, Konishi M, Shigyo M, Yoneyama T, Yanagisawa S (2010) Characterization of plant eukaryotic translation initiation factor 6 (*eIF6*) genes: the essential role in embryogenesis and their differential expression in Arabidopsis and rice. *Biochem Biophys Res Commun* 397(4):673–678
- Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA (2009) Circos: an information aesthetic for comparative genomics. *Genome Res* 19(9):1639–1645
- Kumer SC, Vrana KE (1996) Intricate regulation of tyrosine hydroxylase activity and gene expression. *J Neurochem* 67(2):443–462
- Lescot M, Dehais P, Thijs G, Marchal K, Moreau Y, Van de Peer Y, Rouze P, Rombauts S (2002) PlantCARE, a database of plant *cis*-acting regulatory elements and a portal to tools for in silico analysis of promoter sequences. *Nucleic Acids Res* 30(1):325–327
- Li LS, Luo C, Huang F, Liu ZL, An ZY, Dong L, He XH (2019) Identification and characterization of the mango eIF gene family reveals *MieIF1A-a*, which confers tolerance to salt stress in transgenic Arabidopsis. *Sci Hortic* 248:274–281
- Liu SY, Liu YM, Yang XH, Tong CB, Edwards D, Parkin IAP, Zhao MX, Ma JX, Yu JY, Huang SM, Wang XY, Wang JY, Lu K, Fang ZY, Bancroft I, Yang TJ, Hu Q, Wang XF, Yue Z, Li HJ, Yang LF, Wu J, Zhou Q, Wang WX, King GJ, Pires JC, Lu CX, Wu ZY, Sampath P, Wang Z, Guo H, Pan SK, Yang LM, Min JM, Zhang D, Jin DC, Li WS, Belcram H, Tu JX, Guan M, Qi CK, Du DZ, Li JN, Jiang LC, Batley J, Sharpe AG, Park BS, Ruperao P, Cheng F, Waminal NE, Huang Y, Dong CH, Wang L, Li JP, Hu ZY, Zhuang M, Huang Y, Huang JY, Shi JQ, Mei DS, Liu J, Lee TH, Wang JP, Jin HZ, Li ZY, Li X, Zhang JF, Xiao L, Zhou YM, Liu ZS, Liu XQ, Qin R, Tang X, Liu WB, Wang YP, Zhang YY, Lee J, Kim HH, Denoeud F, Xu X, Liang XM, Hua W, Wang XW, Wang J, Chalhoub B, Paterson AH (2014) The *Brassica oleracea* genome reveals the asymmetrical evolution of polyploid genomes. *Nat Commun* 5:3930
- Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* 25(4):402–408
- Lv X, Lan S, Guy KM, Yang J, Zhang M, Hu Z (2016) Global expressions landscape of NAC transcription factor family and their responses to abiotic stresses in *Citrullus lanatus*. *Sci Rep* 6:30574
- Lynch M, Conery JS (2000) The evolutionary fate and consequences of duplicate genes. *Science* 290(5494):1151–1155
- Nicaise V (2014) Crop immunity against viruses: outcomes and future challenges. *Front Plant Sci* 5:660
- Nix DA, Eisen MB (2005) GATA: a graphic alignment tool for comparative sequence analysis. *BMC Bioinform* 6:9
- Parizipour MHG, Ramazani L, Sardrood BP (2018) Temperature affected transmission, symptom development and accumulation of Wheat dwarf virus. *Plant Prot Sci* 54(4):222–233
- Qian WF, Liao BY, Chang AYF, Zhang JZ (2010) Maintenance of duplicate genes and their functional redundancy by reduced expression. *Trends Genet* 26(10):425–430
- Qian W, Zhang SJ, Zhang SF, Li F, Zhang H, Wu J, Wang XW, Walsh JA, Sun RF (2013) Mapping and candidate-gene screening of the novel Turnip mosaic virus resistance gene *retro2* in Chinese cabbage (*Brassica rapa* L.). *Theor Appl Genet* 126(1):179–188
- Revers F, Nicaise V (2014) Plant resistance to infection by viruses. In: eLS. Wiley, Chichester. <https://doi.org/10.1002/9780470015902.a0000757.pub3>
- Rubio B, Cosson P, Caballero M, Revers F, Bergelson J, Roux F, Schurdi-Levraud V (2019) Genome-wide association study reveals new loci involved in *Arabidopsis thaliana* and Turnip mosaic virus (Tu MV) interactions in the field. *New Phytol* 221(4):2026–2038
- Sato M, Nakahara K, Yoshii M, Ishikawa M, Uyeda I (2005) Selective involvement of members of the eukaryotic initiation factor 4E family in the infection of *Arabidopsis thaliana* by potyviruses. *FEBS Lett* 579(5):1167–1171
- Shopan J, Mou HP, Zhang LL, Zhang CT, Ma WW, Walsh JA, Hu ZY, Yang JH, Zhang MF (2017) Eukaryotic translation initiation factor 2B-beta (*eIF2B β*), a new class of plant virus resistance gene. *Plant J* 90(5):929–940

- Swarbreck D, Wilks C, Lamesch P, Berardini TZ, Garcia-Hernandez M, Foerster H, Li D, Meyer T, Muller R, Ploetz L (2007) The Arabidopsis information resource (TAIR): gene structure and function annotation. *Nucleic Acids Res* 36(suppl_1):D1009–D1014
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28(10):2731–2739
- Thompson JD, Gibson TJ, Higgins DG (2003) Multiple sequence alignment using ClustalW and ClustalX. *Curr Protoc Bioinform* 1:2.3. 1–2.3. 22
- Trebicki P, Nancarrow N, Bosque-Pérez NA, Rodoni B, Aftab M, Freeman A, Yen A, Fitzgerald GJ (2017) Virus incidence in wheat increases under elevated CO₂: a 4-year study of yellow dwarf viruses from a free air carbon dioxide facility. *Virus Res* 241:137–144
- Trebicki P, Nancarrow N, Cole E, Bosque-Perez NA, Constable FE, Freeman AJ, Rodoni B, Yen AL, Luck JE, Fitzgerald GJ (2015) Virus disease in wheat predicted to increase with a changing climate. *Glob Change Biol* 21(9):3511–3519
- Truniger V, Aranda MA (2009) Recessive resistance to plant viruses. *Adv Virus Res* 75:119
- Van Munster M, Yvon M, Vile D, Dader B, Fereres A, Blanc S (2017) Water deficit enhances the transmission of plant viruses by insect vectors. *PLoS ONE* 12(5):e0174398
- Velazquez K, Renovell A, Comellas M, Serra P, Garcia ML, Pina JA, Navarro L, Moreno P, Guerri J (2010) Effect of temperature on RNA silencing of a negative-stranded RNA plant virus: *Citrus psorosis* virus. *Plant Pathol* 59(5):982–990
- Wallbach M, Escobar JD, Babaeikishomi R, Stahnke MJ, Blume R, Schroder S, Kruegel J, Maedler K, Kluth O, Kehlenbach RH, Miosge N, Oetjen E (2016) Distinct functions of the dual leucine zipper kinase depending on its subcellular localization. *Cell Signal* 28(4):272–283
- Walsh JA, Sharpe AG, Jenner CE, Lydiate DJ (1999) Characterisation of resistance to turnip mosaic virus in oilseed rape (*Brassica napus*) and genetic mapping of *TuRB01*. *Theor Appl Genet* 99(7–8):1149–1154
- Wang XW, Wang HZ, Wang J, Sun RF, Wu J, Liu SY, Bai YQ, Mun JH, Bancroft I, Cheng F, Huang SW, Li XX, Hua W, Wang JY, Wang XY, Freeling M, Pires JC, Paterson AH, Chalhoub B, Wang B, Hayward A, Sharpe AG, Park BS, Weisshaar B, Liu BH, Li B, Liu B, Tong CB, Song C, Duran C, Peng CF, Geng CY, Koh CS, Lin CY, Edwards D, Mu DS, Shen D, Soumpourou E, Li F, Fraser F, Conant G, Lassalle G, King GJ, Bonnema G, Tang HB, Wang HP, Belcram H, Zhou HL, Hirakawa H, Abe H, Guo H, Wang H, Jin HZ, Parkin IAP, Batley J, Kim JS, Just J, Li JW, Xu JH, Deng J, Kim JA, Li JP, Yu JY, Meng JL, Wang JP, Min JM, Poulain J, Wang J, Hatakeyama K, Wu K, Wang L, Fang L, Trick M, Links MG, Zhao MX, Jin MN, Ramchiary N, Drou N, Berkman PJ, Cai QL, Huang QF, Li RQ, Tabata S, Cheng SF, Zhang S, Zhang SJ, Huang SM, Sato S, Sun SL, Kwon SJ, Choi SR, Lee TH, Fan W, Zhao X, Tan X, Xu X, Wang Y, Qiu Y, Yin Y, Li YR, Du YC, Liao YC, Lim Y, Narusaka Y, Wang YP, Wang ZY, Li ZY, Wang ZW, Xiong ZY, Zhang ZH (2011) The genome of the mesopolyploid crop species *Brassica rapa*. *Nat Genet* 43(10):1035–U1157
- Wortham NC, Proud CG (2015) eIF2B: recent structural and functional insights into a key regulator of translation. *Biochem Soc Trans* 43:1234–1240
- Yang JH, Liu DY, Wang XW, Ji CM, Cheng F, Liu BN, Hu ZY, Chen S, Pental D, Ju YH, Yao P, Li XM, Xie K, Zhang JH, Wang JL, Liu F, Ma WW, Shopan J, Zheng HK, Mackenzie SA, Zhang MF (2018) The genome sequence of allopolyploid *Brassica juncea* and analysis of differential homoeolog gene expression influencing selection (vol 48, pg 1225, 2016). *Nat Genet* 50(11):1616–1616
- You J, Zhang L, Song B, Qi X, Chan Z (2015) Systematic analysis and identification of stress-responsive genes of the NAC gene family in *Brachypodium distachyon*. *PLoS ONE* 10(3):e0122027
- Yvon M, Vile D, Brault V, Blanc S, Van Munster M (2017) Drought reduces transmission of Turnip yellows virus, an insect-vectored circulative virus. *Virus Res* 241:131–136
- Zhang W, Ruan J, T-hD Ho, You Y, Yu T, Quatrano RS (2005) Cis-regulatory element based targeted gene finding: genome-wide identification of abscisic acid-and abiotic stress-responsive genes in *Arabidopsis thaliana*. *Bioinformatics* 21(14):3074–3081
- Zhang Z, Li J, Zhao X-Q, Wang J, Wong GK-S, Yu J (2006) KaKs_Calculator: calculating Ka and Ks through model selection and model averaging. *Genom Proteom Bioinform* 4(4):259–263