Published in final edited form as:

Obesity (Silver Spring). 2020 January ; 28(1): 171–177. doi:10.1002/oby.22662.

It's absolutely relative: The effect of age on the body mass index-mortality relationship in postmenopausal women

Hailey R. Banack¹, Jennifer W. Bea², Andrew Stokes³, Candyce H. Kroenke⁴, Marcia L. Stefanick⁵, Shirley A. Beresford⁶, Chloe E. Bird⁷, Lorena Garcia⁸, Robert Wallace⁹, Robert A. Wild¹⁰, Bette Caan¹¹, Jean Wactawski-Wende¹

¹Department of Epidemiology and Environmental Health, School of Public Health and Health Professions, University at Buffalo, The State University of New York

²Department of Nutrition Sciences and University of Arizona Cancer Center, University of Arizona

³Department of Global Health and Center for Global Health and Development, Boston University School of Public Health

⁴Division of Research, Kaiser Permanente Northern California, Oakland, CA

⁵Division of Cardiovascular Medicine, Department of Medicine, Stanford, University School of Medicine, Stanford, CA, United States.

⁶Department of Epidemiology, School of Public Health, University of Washington, Seattle, Washington.

⁷RAND Corporation, Santa Monica, California.

⁸Department of Public Health Sciences, University of California Davis, Davis, CA, USA.

⁹Department of Epidemiology, College of Public Health, University of Iowa

¹⁰Department of Obstetrics and Gynecology, University of Oklahoma Health Sciences Center

¹¹Division of Research, Kaiser Permanente Northern California, Oakland, CA

Abstract

Objective: The use of relative and absolute effect estimates has important implications for the interpretation of study findings. Likewise, examining additive and multiplicative interaction can lead to differing conclusions about the joint effects of two exposure variables. The aim of this paper is to examine the relationship between BMI and mortality on the relative and absolute scales and investigate interaction between BMI and age.

Methods: We used data from 68,132 participants in the Women's Health Initiative (WHI). We estimated the risk ratio and risk difference of BMI on mortality. We also included a product term to examine interaction between BMI and age on the multiplicative scale and calculated the relative excess risk of interaction to measure additive interaction.

Corresponding Author: Hailey Banack, Department of Epidemiology and Environmental Health, School of Public Health and Health Professions, University at Buffalo, The State University of New York, 270 Farber Hall, Buffalo, NY 14214-8001, hrbanack@buffalo.edu.

Conflicts of Interest: The authors declare no conflict of interest

Results: Results demonstrated that the mortality risk ratio decreased as women aged but the mortality risk difference increased as women aged. We found evidence of additive and multiplicative interaction between age and BMI.

Conclusions: In postmenopausal women, the relative mortality risk associated with high BMI decreased with increasing age, but the absolute risk of high BMI increased with increasing age. This indicates the importance of considering the interaction between age and BMI to understand mortality risk in older women.

Keywords

obesity; aging; risk ratio; risk difference; interaction

Introduction

There is a substantial body of evidence demonstrating that the mortality risk due to obesity changes with aging.(1–7) The consistent change in associations between risk factors and outcomes with aging has long been recognized (2, 8). Understanding the changing association of obesity and mortality with aging is extremely relevant today given the increasing population prevalence of obesity (9) and rapid aging of our population(10). Over time, the proportion of older adults in the population has grown considerably: by 2030, more than 20% of American adults are expected to be over 65 years of age, while only 13% were above 65 years in 2010 and 9.8% in 1970.(10) At the same time, the prevalence of obesity among older adults (>60 years) has risen steadily; recent estimates indicate that 38.5% of men and 43.1% of women are obese in the United States.(9) This is a particular concern for older women as the prevalence of obesity is 3.7% higher for older men but 6.6% higher for older women compared to middle age men and women (40–59 years old)(9).

The effect of obesity on health outcomes in older adults is frequently of interest to researchers, clinician-scientists, and public health practitioners. By definition, the fact that the relationship between BMI and mortality differs across age categories is evidence of interaction between age and BMI. (3) Interaction can be assessed on either the additive or multiplicative scale.(11–13) Several authors have argued that additive interaction is more relevant than statistical or multiplicative interaction to understand disease etiology and for public health purposes.(12, 14–16) VanderWeele and Knol provide several clear examples of why additive interaction is the more relevant public health measure: in the real-world, additive interaction provides direct insight into the number of deaths that can be avoided by treating a high-risk group, but multiplicative interaction does not.(15) A null multiplicative effect between obesity and old age should not be used as evidence that weight management and intervention strategies are less important in older adults.(13, 17)

The aim of this paper is to examine the relationship between BMI and mortality on both the relative (risk ratio) and absolute (risk difference) scale in a large cohort of postmenopausal women. Additionally, we will investigate the presence of additive and multiplicative interaction between BMI and age. Understanding the interaction between BMI and age is important to identify groups at particularly high risk for mortality. Taken together, these

analyses will provide obesity researchers with a comprehensive understanding of age-related differences in the BMI- mortality relationship in older women.

Methods

We used data from the Women's Health Initiative (WHI) clinical trials in this analysis. The WHI is a large longitudinal study of postmenopausal women across the United States focused on preventing heart disease, cancer, and osteoporotic fractures in older women.(18–20) Recruitment started in 1993 and the trials were all completed by 2005. Outcome ascertainment is still ongoing through the WHI extension studies I and II (2005–2010, 2010–2020). Information about recruitment and eligibility criteria have been published in detail elsewhere.(18) Briefly, postmenopausal women aged 50–79 were recruited to take part in partially overlapping randomized clinical trials: two hormone therapy (HT) trials testing either a combination of estrogen and progestin versus placebo or estrogen alone compared to a placebo, a dietary modification (DM) trial, and a vitamin (calcium and vitamin D; CaD) supplementation trial.(19, 21) Participants of the vitamin supplementation trial were recruited from the participants of the dietary modification trial and/or the HT trials; to be part of the CaD trial participants would have already consented to participate in the DM trial or HT trials at annual visits 1 or 2.(22)

The analytic cohort for this study consists of 68,132 women. In total, there were 27,347 women in the HT trials, 48,836 women in the DM trial, and 36,282 women in the CaD trial. (19, 21) In order to enhance recruitment effort, women were allowed to enroll in more than one clinical trial: for instance, of the participants in the HT trial, 29.4% also participated in the DM and 58.8% were part of the CaD trial.(23) After recruitment and randomization, clinic visits were required annually, and consisted of questionnaires, anthropometric measures, and clinical examinations.(18) The study protocol was reviewed by institutional review boards at each of the 40 WHI clinical centers as well as the WHI coordinating center and each participant provided informed consent.(22)

Outcome ascertainment

All-cause mortality is the primary outcome of interest in this analysis, with follow-up until June 2017. The WHI coordinating center collects information on vital status through mail, telephone, searches of medical record and death certificates, and the National Death Index. (19, 22)

Exposure variable and covariates

The exposure variable of interest is body mass index (BMI), an index of weight-for-height calculated as weight in kilograms (kg) divided by the square of height in meters (m²). Height and weight were measured annually at clinic visits by trained examiners using standard procedures. For this analysis, we will categorize participants into standard BMI groups: BMI <18.5 kg/m², 18.5–24.9 kg/m², 25–29.9 kg/m², 30–34.9 kg/m², 35–39.9 kg/m², and ≥40 kg/m². Information on relevant confounding variables was collected through questionnaires or in-clinic measurements, and includes information on demographic

characteristics, personal and medical history, lifestyle habits, medication use, and physical measurements.

Statistical Analysis

To illustrate the crude probability of survival according to baseline BMI category, the first step in this analysis was to create a Kaplan-Meier survival curve. We then estimated the relative and absolute effects of BMI on all-cause mortality, comparing the risk difference and risk ratio of mortality for each BMI category. Finally, we quantified interaction on the additive and multiplicative scales. To examine the influence of age at baseline, we present the main results stratified by baseline age group (50–59, 60–69, and 70–79).

To calculate the crude and multivariate adjusted relative risk (risk ratio) of mortality comparing women with a BMI of 25–29.9 kg/m² to women with BMIs of <18.5, 18.5–24.9, 30–34.9 kg/m², 35–39.9 kg/m², and 40 kg/m², we used a generalized linear model with a log link and Poisson distribution.⁽¹⁶⁾ BMI 25–29.9 kg/m² was used as the referent group because it is the group associated with the lowest risk of mortality and less likely to be affected by illness-related weight loss than BMI 18.5–24.9.^(24–26)

To calculate the risk difference by BMI category, we used a two-step process that involved obtaining the predicted probabilities from the generalized linear model and then using marginal standardization to obtain average marginal effects.⁽²⁷⁾ Average marginal effects are an absolute measure of effect, and can be interpreted similar to a risk difference (i.e., the difference in outcome risk comparing each BMI category to the referent group).⁽²⁷⁾ We adjusted for relevant confounding variables including: trial(s) and trial arm(s), race/ethnicity, education, marital status, health insurance status, self-rated health, smoking status and smoking history, personal hormone therapy use, baseline age, and alcohol use. In addition, in Supplementary Table 1, we present the rate ratio and rate difference results, also estimated from Poisson regression models but with an offset term to account for the log of the number of days of follow-up.

To estimate interaction on the additive and multiplicative scales, we included product interaction terms between BMI categories, as dummy variables, and age as a continuous variable (age per 5 year increase) in a generalized linear model with log link and Poisson distribution. Age was scaled to 5 year increments by dividing continuous age by 5.⁽¹²⁾ In this analysis, we first created dummy variables for BMI categories compared to the referent group. We then created product terms by multiplying the BMI dummy variables by age, and included them in a Poisson regression model:

$$\log(E(Y|X,Z)) = \beta_0 + \beta_1 \text{BMI}_1 + \beta_2 \text{BMI}_2 + \beta_3 \text{BMI}_4 + \beta_4 \text{BMI}_5 + \beta_5 \text{BMI}_6 + \beta_6 \text{Age} + \beta_7 \text{BMI}_1 \times \text{Age} + \beta_8 \text{BMI}_2 \times \text{Age} + \beta_9 \text{BMI}_4 \times \text{Age} + \beta_{10} \text{BMI}_5 \times \text{Age} + \beta_{11} \text{BMI}_6 \times \text{Age} + \beta_k Z$$

Where BMI₁ represents the dummy variable for BMI<18.5 compared to the referent group, 25–29.9, BMI₂ represents the dummy variable for BMI 18.5–24.9 compared to the referent group, BMI₄ represents the dummy variable for BMI 30–34.9 compared to the referent group, BMI₅ represents the dummy variable for BMI 35–39.9 compared to the referent group, and BMI₆ represents the dummy variable for BMI>40 compared to the referent

group. We again controlled for all of the aforementioned confounding variables, noted by the vector Z .

The value of the coefficients and 95% confidence intervals from the product terms between BMI category and age (β_7 to β_{11}) provide information about multiplicative interaction. In a logistic regression model, or other exponential models, a non-zero value for the coefficient of the product term indicates deviation from exact multiplicativity (i.e., the presence of sub- or super-multiplicative interaction).

To quantify the magnitude of additive interaction, we used the results from the regression model to calculate the relative excess risk of interaction (RERI)(12, 28) for each BMI category:

$$\text{RERI for BMI}_{<18.5} = \exp(\beta_1 \text{BMI}_1 + \beta_6 \text{Age} + \beta_7 \text{BMI}_1 \times \text{Age}) - \exp(\beta_1 \text{BMI}_1) - \exp(\beta_6 \text{Age}) + 1$$

$$\text{RERI for BMI}_{18.5-24.9} = \exp(\beta_2 \text{BMI}_2 + \beta_6 \text{Age} + \beta_8 \text{BMI}_2 \times \text{Age}) - \exp(\beta_2 \text{BMI}_2) - \exp(\beta_6 \text{Age}) + 1$$

$$\text{RERI for BMI}_{30-34.9} = \exp(\beta_3 \text{BMI}_4 + \beta_6 \text{Age} + \beta_9 \text{BMI}_4 \times \text{Age}) - \exp(\beta_3 \text{BMI}_4) - \exp(\beta_6 \text{Age}) + 1$$

$$\text{RERI for BMI}_{35-39.9} = \exp(\beta_4 \text{BMI}_5 + \beta_6 \text{Age} + \beta_{10} \text{BMI}_5 \times \text{Age}) - \exp(\beta_4 \text{BMI}_5) - \exp(\beta_6 \text{Age}) + 1$$

$$\text{RERI for BMI}_{40} = \exp(\beta_5 \text{BMI}_6 + \beta_6 \text{Age} + \beta_{11} \text{BMI}_6 \times \text{Age}) - \exp(\beta_5 \text{BMI}_6) - \exp(\beta_6 \text{Age}) + 1$$

An RERI value of 0 indicates the joint effects of age and BMI are exactly additive, meaning the combined effect of BMI and age on mortality is exactly equal to the sum of the individual effects and there is no additive interaction. An RERI value less than 0 indicates sub-additivity, when the combined effect of BMI and mortality are less than the sum of the individual effects on mortality. An RERI value greater than 0 indicates the combined effects of BMI and age are larger than the sum of the individual effects, or super additivity.(28) We estimated 95% confidence intervals for the RERI using bootstrapping.(12)

Results

Demographic characteristics of the study population ($n=68,132$) at baseline are presented in Table 1. Over the total WHI study follow-up, 17,785 deaths occurred. At baseline, the average age of study participants was 62.2 years (± 6.9). The majority of women had BMI 25–29.9 kg/m^2 at baseline (36%), followed by 18.5–24.9 kg/m^2 (27%), 30–34.9 kg/m^2 (22%), 35–39.9 kg/m^2 (10%), 40 kg/m^2 (4%), and <18.5 kg/m^2 (0.4%). The mean BMI of the study sample was 28.9 kg/m^2 ($\text{SD}=5.9$).

Figure 1 depicts a Kaplan Meier (unadjusted) survival curve by baseline BMI. In the early years of follow-up, having very low BMI (<18.5 kg/m^2) was associated with the lowest survival probability. However, over the entire study period, having a BMI 40 kg/m^2 or 35–39.9 kg/m^2 was associated with the lowest survival. The cumulative probability of survival

for women with BMI 34–39.9 kg/m² was 0.60 (0.55, 0.63) and 0.57 (0.53, 0.61) for the >40 category. The probability of survival was very similar among women who have BMI 25–29.9 kg/m² and 18.5–24.9 kg/m². In Supplementary Figure 1, we present Kaplan Meier survival curves stratified by age group at baseline (50–59, 60–69, and 70–79).

Table 2 contains relative (risk ratio) and absolute (risk difference) estimates and 95% confidence intervals for the effect of BMI category on mortality, stratified by age at baseline. Consistent with previous research, examination of the stratified relative risk results demonstrates that the relative risks of high BMI are attenuated in the oldest group of women (70–79) compared with the youngest group (50–59). In the youngest women, the mortality risk ratios are 1.08 (0.98, 1.18) for BMI 30–34.9 kg/m², 1.38 (1.24, 1.53) for BMI 35–39.9 kg/m², and 1.74 (1.54, 1.96) for women with BMI >40 kg/m², whereas in the oldest women, the mortality rate ratios were 1.02 (0.98, 1.07) for BMI 30–34.9 kg/m², 1.08 (1.02, 1.15) for BMI 35–39.9 kg/m², and 1.18 (1.07, 1.30) for women with BMI greater than 40 kg/m². These results represent a decrease in the risk ratio of 5.9% in the 30–34.9 kg/m² category, 27.8% in the 34–39.9 category, and 47% in the >40 category comparing the oldest (70–79) and youngest women (50–59). Rate ratio results demonstrate a similar trend (presented in Supplemental Table 1).

The risk difference results presented in Table 2 demonstrate differences in the BMI-mortality relationship by baseline age cohort, but in the opposite direction from the rate ratio results. In contrast to the risk ratio results, the mortality risk difference increased in women who were older at baseline. In 50–59 year old women, the risk difference (per 10,000) was 74 (38, 111), 212 (159, 264), and 432 (349, 515) for women who had BMI of 30–34.9, 35–39.9, or >40 kg/m², compared to the referent group. In 60–69 year old women, the risk difference results for the same categories were 177 (91, 264), 504 (379, 1227), and 1030 (832, 1364), and in 70–79 year old women, the results were 384 (197, 572), 1093 (362, 1226) and 2231 (1799, 2663). These results strongly suggest that the absolute risk of mortality increases with increasing age and BMI. Rate difference results are presented in Supplemental Table 1. The difference between relative and absolute effect estimates is illustrated in Figure 2 comparing the risk ratio and risk difference results for the youngest (50–59) and oldest (70–79) year old age groups.

Interaction results are presented in Table 3. The coefficients (95% CI) for the product terms for BMI <18.5, 30–34.9, 35–39.9, and >40 kg/m² per 5 year age increase were −0.09 (−0.24, 0.05), −0.03 (−0.06, 0.003), −0.08 (−0.12, −0.04) and −0.13 (−0.19, −0.08), suggesting negative interaction on the multiplicative scale, also called sub-multiplicative interaction. The relative excess risk of interaction results (RERI) and 95% confidence intervals for the high BMI categories (30–34.9, 35–39.9, and >40 kg/m²) strongly indicated the presence of super-additive interaction between age and BMI. The RERI values for each of the high BMI categories per 5 year age increase were 0.26 (0.02, 0.50), 0.99 (0.46, 1.52), and 2.51 (1.17, 3.85). Results for the interaction between BMI 18.5–24.9 and age were qualitatively different than for the other BMI categories, suggesting sub-additive interaction (RERI = −0.02; 95% CI: −0.17, 0.14) and super-multiplicative interaction (β = 0.004; 95% CI: −0.03, 0.03). However, the confidence intervals for both the RERI and interaction term are quite wide, making it difficult to draw a definitive conclusion for this category.

Discussion

Using prospectively collected data from the WHI clinical trials, we examined the joint effects of BMI and age on mortality in a cohort of postmenopausal women. We compared risk difference and risk ratio results and quantified interaction between age and BMI on the additive and multiplicative scales. We found evidence to support the frequently reported pattern of attenuation of obesity-related mortality risk across age cohorts, but only on the relative scale. The opposite pattern was seen when considering the absolute scale (risk differences); there was a clear increase in the number of excess deaths due to high BMI in older compared to younger women. These conclusions were also supported by the results of the interaction analysis, we found evidence of a sub-multiplicative and super-additive interaction between increasing age and BMI category.

Prior work examining the relationship between BMI and mortality in older adults in the general population and in the WHI has consistently reported that the relative risks associated with high BMI decline with age.(1, 4, 13, 29, 30) For example, in a study of 13,451 male and female participants in the Leisure World Cohort Study (mean age=73.5 years), Corrada and colleagues reported that obese participants only had increased mortality up to age 75. (29) A similar pattern of attenuation has been reported in other studies from the WHI cohort: compared to BMI 18.5–24.9, for women with BMI of 30–34.9, Chen and colleagues reported hazard ratios (HR) of 1.08 (0.95 –1.23) for 50–59 year old women and 0.98 (0.92, 1.06) for women aged 70–79, for women with BMI of 35–39.9 they reported HRs of 1.61 (1.39, 1.87) for 50–59 year olds and 1.11 (1.00, 1.23) for 70–79 year olds, and for women with BMI 40, the HRs were 1.82 (1.55, 2.15) for 50–59 year olds and 1.08 (0.92, 1.26) for 70–79 year olds.(31) Using data from a sub-cohort of WHI women, Bea et al. also reported a consistent pattern of age-attenuation in analyses stratified by age group at baseline.(3) These results support previous work on the differing effects of relative and absolute estimates when studying obesity in older men and women (8). Similar to the results presented herein, Stevens and colleagues (1999) demonstrated that the relative age-related mortality risks of obesity (BMI 30 kg/m²) decrease while the absolute age-related mortality risk increased. (8)

Relative risks are often reported in the literature because they are familiar to researchers and easily interpreted. However, it is important to keep in mind that relative risks can potentially obscure actual differences in the risk between groups.(2, 8, 32–35). For instance, using an example of a study looking at the relationship between obesity and mortality, reporting a risk ratio (RR) of 2.0 would be interpreted as a two-fold increase in mortality risk in the exposed group (obese) relative to the unexposed group (non-obese).(36) This may appear to be a meaningful increase in mortality risk, but it could result from having exceedingly low risks in both the exposed and unexposed group, such as 0.0002% in the exposed (obese) group and 0.0001% in the unexposed (non-obese) group ($RR = \text{Risk}_{\text{exposed}} / \text{Risk}_{\text{unexposed}}$). Reporting an absolute risk difference (RD) of 0.0001% ($RD = \text{Risk}_{\text{exposed}} - \text{Risk}_{\text{unexposed}}$) is less likely to be interpreted as a meaningful difference, either clinically or from a population health perspective, but gives a true indication of the low mortality risk due to exposure.(32, 36)

In studies of older adults, the mortality risk difference comparing obese and non-obese may still be the same (or greater) in older individuals compared to younger individuals owing to the increased mortality risk associated with aging.(8, 33) If the risk in the unexposed group increases with time, the same observed risk ratio in different age groups would correspond with a larger change in absolute risk in older individuals than younger individuals.(8, 33, 37) Consider this example of a study exploring the relationship between BMI and heart failure: if the hazard ratio for a 5-unit change in BMI was 1.20 in 40 year old individuals, but 1.10 in 80 year old individuals, it might appear as though the increase in risk associated with a 5-unit change in BMI is 50% greater (20% risk increase vs 10% risk increase) in 40-year olds compared with 80-year olds. However, it is necessary to also consider the risk of the outcome in each group. If the incidence of heart failure is 3 per 100,000 in younger adults (40 year olds) but 19 per 100,000 in older adults (80 year olds), the absolute difference of a 5-unit reduction in BMI would be approximately 3 times greater at age 80 than age 40.(35)

It is important to note that both relative and absolute estimates can be valid measures of effects in cohort studies, choosing one over the other has important implications for the interpretation of study findings.(32, 33) The same is true of additive and multiplicative interaction.(13) The absolute effect estimates directly indicate the excess number of deaths that can be attributed to each BMI category, whereas relative effect estimates can only inform how much more (or less) likely an individual is to experience the outcome, relative to those in the referent group. Since the baseline risks are different in younger and older individuals, the risk ratios are difficult to compare. Similarly, it has long been recognized that additive interaction is more relevant for clinicians, public health practitioners, and policymakers because it highlights the individuals in the population who would benefit most from an intervention.(15, 38) Based on the interpretation of super additive interaction from our results, the number of deaths in aging women who have high BMI values is greater than the number of deaths expected from either high BMI or increasing age independently.(13, 28, 39) Ultimately, whether additive or multiplicative interaction is most relevant depends on the question that the researchers are trying to answer. Some authors have suggested it is good practice to examine both types of interaction in an analysis.(15, 40)

Although there are many articles focused on the relationship between obesity and mortality in older adults, several pertinent questions remain unanswered. We have presented an analysis in a large, well-characterized cohort, but this type of analysis does not answer questions about the effect of specific interventions (e.g., diet, exercise, pharmacotherapy (41)) on mortality risk among older women with high BMI. These are questions best answered in the context of a rigorously conducted randomized controlled trial. We used baseline BMI to simplify the analysis and interpretation for the purpose of demonstrating the differences between relative and absolute effect estimates. Another limitation of this analysis is the use of BMI categories as the primary exposure, as it may not adequately represent body composition in postmenopausal women(42). We considered the BMI-mortality relationship in our analyses, however it would be interesting to examine the interaction between age and other anthropometric indices (e.g., waist circumference, body surface area) in future work. Finally, it is important to note that all of the participants in the WHI are women, which has important implications for the generalizability of our study findings.

Conclusion

The objective of this manuscript was to investigate the effect of BMI on mortality in postmenopausal women on the relative and absolute scales and examine interaction between aging and BMI on the additive and multiplicative scales. Our results provide an important reminder of the different interpretations of relative and absolute effect estimates and highlight the need for future research exploring obesity in older adults.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements:

We thank Dr. Noel Weiss for his thoughtful comments on an initial version of this analysis.

We also acknowledge the Women's Health Initiative (WHI): Program Office: (National Heart, Lung, and Blood Institute, Bethesda, Maryland) Jacques Rossouw, Shari Ludlam, Joan McGowan, Leslie Ford, and Nancy Geller Clinical Coordinating Center: (Fred Hutchinson Cancer Research Center, Seattle, WA) Garnet Anderson, Ross Prentice, Andrea LaCroix, and Charles Kooperberg Investigators and Academic Centers: (Brigham and Women's Hospital, Harvard Medical School, Boston, MA) JoAnn E. Manson; (MedStar Health Research Institute/Howard University, Washington, DC) Barbara V. Howard; (Stanford Prevention Research Center, Stanford, CA) Marcia L. Stefanick; (The Ohio State University, Columbus, OH) Rebecca Jackson; (University of Arizona, Tucson/Phoenix, AZ) Cynthia A. Thomson; (University at Buffalo, Buffalo, NY) Jean Wactawski-Wende; (University of Florida, Gainesville/Jacksonville, FL) Marian Limacher; (University of Iowa, Iowa City/Davenport, IA) Jennifer Robinson; (University of Pittsburgh, Pittsburgh, PA) Lewis Kuller; (Wake Forest University School of Medicine, Winston-Salem, NC) Sally Shumaker; (University of Nevada, Reno, NV) Robert Brunner Women's Health Initiative Memory Study: (Wake Forest University School of Medicine, Winston-Salem, NC) Mark Espeland.

Sources of Funding: The WHI program is funded by the National Heart, Lung, and Blood Institute, National Institutes of Health, U.S. Department of Health and Human Services through contracts HHSN268201600018C, HHSN268201600001C, HHSN268201600002C, HHSN268201600003C, and HHSN268201600004C. HRB is supported by the Banting Postdoctoral Fellowship Program from the Canadian Institutes of Health Research (CIHR)

References

1. Stevens J, Cai J, Pamuk ER, Williamson DF, Thun MJ, Wood JL. The Effect of Age on the Association between Body-Mass Index and Mortality. *New England Journal of Medicine* 1998;338(1):1–7. [PubMed: 9414324]
2. Kaplan GA, Haan MN, Wallace RB. Understanding Changing Risk Factor Associations With Increasing Age in Adults. *Annual Review of Public Health* 1999;20(1):89–108.
3. Bea JW, Thomson CA, Wertheim BC, Nicholas JS, Ernst KC, Hu C, et al. Risk of Mortality According to Body Mass Index and Body Composition Among Postmenopausal Women. *American journal of epidemiology* 2015;182(7):585–96. [PubMed: 26350478]
4. Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA* 2013;309(1):71–82. [PubMed: 23280227]
5. Heiat A, Vaccarino V, Krumholz HM. An evidence-based assessment of federal guidelines for overweight and obesity as they apply to elderly persons. *Archives of Internal Medicine* 2001;161(9):1194–203. [PubMed: 11343442]
6. Adams KF, Schatzkin A, Harris TB, Kipnis V, Mouw T, Ballard-Barbash R, et al. Overweight, obesity, and mortality in a large prospective cohort of persons 50 to 71 years old. *N Engl J Med* 2006;355(8):763–78. [PubMed: 16926275]

7. Berrington de Gonzalez A, Hartge P, Cerhan JR, Flint AJ, Hannan L, MacInnis RJ, et al. Body-mass index and mortality among 1.46 million white adults. *N Engl J Med* 2010;363(23):2211–9. [PubMed: 21121834]
8. Stevens J, Cai J, Juhaeri, Thun MJ, Williamson DF, Wood JL. Consequences of the Use of Different Measures of Effect to Determine the Impact of Age on the Association between Obesity and Mortality. *American journal of epidemiology* 1999;150(4):399–407. [PubMed: 10453816]
9. Hales C, Carroll M, Fryar C, Ogden C. Prevalence of obesity among adults and youth: United States, 2015–2016. Hyattsville, MD 2017.
10. Ortman J, Velkoff V, Hogan H. An aging nation: The older population in the United States: U.S. Census Bureau 2014 [Available from: <http://www.census.gov/prod/2014pubs/p25-1140.pdf>.
11. VanderWeele TJ. On the distinction between interaction and effect modification. *Epidemiology* 2009;20(6):863–71. [PubMed: 19806059]
12. Knol MJ, van der Tweel I, Grobbee DE, Numans ME, Geerlings MI. Estimating interaction on an additive scale between continuous determinants in a logistic regression model. *International journal of epidemiology* 2007;36(5):1111–8. [PubMed: 17726040]
13. Mehta N, Zheng H. Do the effects of major risk factors for mortality rise or fall with age?; 2018 Contract No.: 18–888.
14. Greenland S, Rothman KJ. Concepts of Interaction In: Rothman KJ, Greenland S, editors. *Modern Epidemiology* Philadelphia, Pa.: Lippincott-Raven; 1998.
15. VanderWeele Tyler J, Knol Mirjam J. A Tutorial on Interaction. *Epidemiologic Methods* 2014 p. 33.
16. McNutt L-A, Wu C, Xue X, Hafner JP. Estimating the Relative Risk in Cohort Studies and Clinical Trials of Common Outcomes. *American journal of epidemiology* 2003;157(10):940–3. [PubMed: 12746247]
17. Mehta N, Preston S. Are major behavioral and sociodemographic risk factors for mortality additive or multiplicative in their effects? *Social Science & Medicine* 2016;154:93–9. [PubMed: 26950393]
18. Design of the Women’s Health Initiative Clinical Trial and Observational Study. *Controlled Clinical Trials* 1998;19(1):61–109. [PubMed: 9492970]
19. Rossouw J, Anderson G, Prentice R, LaCroix A, Kooperberg C, Stefanick M, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: Principal results from the women’s health initiative randomized controlled trial. *JAMA* 2002;288(3):321–33. [PubMed: 12117397]
20. Manson JE, Chlebowski RT, Stefanick ML, et al. Menopausal hormone therapy and health outcomes during the intervention and extended poststopping phases of the women’s health initiative randomized trials. *JAMA* 2013;310(13):1353–68. [PubMed: 24084921]
21. Ritenbaugh C, Patterson RE, Chlebowski RT, Caan B, Fels-Tinker L, Howard B, et al. The women’s health initiative dietary modification trial: overview and baseline characteristics of participants. *Annals of epidemiology* 2003;13(9, Supplement):S87–S97. [PubMed: 14575941]
22. Neuhouser ML, Aragaki AK, Prentice RL, et al. Overweight, obesity, and postmenopausal invasive breast cancer risk: A secondary analysis of the women’s health initiative randomized clinical trials. *JAMA Oncology* 2015;1(5):611–21. [PubMed: 26182172]
23. Hays J, Hunt JR, Hubbell FA, Anderson GL, Limacher M, Allen C, et al. The women’s health initiative recruitment methods and results. *Annals of epidemiology* 2003;13(9):S18–S77. [PubMed: 14575939]
24. Durazo-Arvizu R, McGee D, Li Z, Cooper R. Establishing the Nadir of the Body Mass Index-Mortality Relationship: A Case Study. *Journal of the American Statistical Association* 1997;92(440):1312–9.
25. Wong ES, Wang BC, Garrison LP, Alfonso-Cristancho R, Flum DR, Arterburn DE, et al. Examining the BMI-mortality relationship using fractional polynomials. *BMC Medical Research Methodology* 2011;11(1):175. [PubMed: 22204699]
26. Winter JE, MacInnis RJ, Wattanapenpaiboon N, Nowson CA. BMI and all-cause mortality in older adults: a meta-analysis. *The American journal of clinical nutrition* 2014;99(4):875–90. [PubMed: 24452240]

27. Muller CJ, MacLehose RF. Estimating predicted probabilities from logistic regression: different methods correspond to different target populations. *Int J Epidemiol* 2014;43(3):962–70. [PubMed: 24603316]
28. Richardson DB, Kaufman JS. Estimation of the relative excess risk due to interaction and associated confidence bounds. *Am J Epidemiol* 2009;169(6):756–60. [PubMed: 19211620]
29. Corrada MM, Kawas CH, Mozaffar F, Paganini-Hill A. Association of Body Mass Index and Weight Change with All-Cause Mortality in the Elderly. *American journal of epidemiology* 2006;163(10):938–49. [PubMed: 16641311]
30. Janssen I. Morbidity and Mortality Risk Associated With an Overweight BMI in Older Men and Women. *Obesity* 2007;15(7):1827–40. [PubMed: 17636102]
31. Chen Z, Klimentidis YC, Bea JW, Ernst KC, Hu C, Jackson R, et al. Body Mass Index, Waist Circumference, and Mortality in a Large Multiethnic Postmenopausal Cohort—Results from the Women’s Health Initiative. *Journal of the American Geriatrics Society* 2017;65(9):1907–15. [PubMed: 28229456]
32. Poole C. On the Origin of Risk Relativism. *Epidemiology* 2010;21(1):3–9 [PubMed: 20010205]
33. Weiss NS, Treggiari MM. Postmenopausal hormone use and skeletal fracture: does the size of the benefit decrease with increasing age? *Obstetrics & Gynecology* 2002;100(2):364–8. [PubMed: 12151165]
34. King NB, Harper S, Young ME. Use of relative and absolute effect measures in reporting health inequalities: structured review. *The BMJ* 2012;345:e5774.
35. Howard G, C GD. A call for caution in the interpretation of the observed smaller relative importance of risk factors in the elderly. *Annals of epidemiology* 1998;8(7):411–4. [PubMed: 9738686]
36. Noordzij M, van Diepen M, Caskey FC, Jager KJ. Relative risk versus absolute risk: one cannot be interpreted without the other. *Nephrology Dialysis Transplantation* 2017;32(suppl_2):ii13–ii8.
37. Mayeda ER, Banack HR, Bibbins-Domingo K, Zeki Al Hazzouri A, Marden JR, Whitmer RA, et al. Can Survival Bias Explain the Age Attenuation of Racial Inequalities in Stroke Incidence?: A Simulation Study. *Epidemiology* 2018;29(4):525–32. [PubMed: 29621058]
38. Rothman KJ, Greenland S, Walker AM. Concepts of interaction. *Am J Epidemiol* 1980;112(4):467–70. [PubMed: 7424895]
39. Bally M, Nadeau L, Brophy JM. Studying additive interaction in a healthcare database: Case study of NSAIDs, cardiovascular profiles, and acute myocardial infarction. *PLOS ONE* 2018;13(8):e0201884.
40. Knol M, Egger M, Scott P, Mirjam G, Vandenbroucke J. When one depends on the other: reporting of interaction in case-control and cohort studies. *Epidemiology* 2009;20(2):161–6. [PubMed: 19034025]
41. LeBlanc ES, Patnode CD, Webber EM, Redmond N, Rushkin M, O’Connor EA. Behavioral and Pharmacotherapy Weight Loss Interventions to Prevent Obesity-Related Morbidity and Mortality in Adults: Updated Evidence Report and Systematic Review for the US Preventive Services Task Force USPSTF Evidence Report: Behavioral Interventions to Prevent Adult Obesity-Related Outcomes USPSTF Evidence Report: Behavioral Interventions to Prevent Adult Obesity-Related Outcomes. *JAMA* 2018;320(11):1172–91. [PubMed: 30326501]
42. Banack HR, Stokes A, Fox MP, Hovey KM, Cespedes Feliciano EM, LeBlanc ES, et al. Stratified Probabilistic Bias Analysis for Body Mass Index-related Exposure Misclassification in Postmenopausal Women. *Epidemiology* 2018;29(5):604–13. [PubMed: 29864084]

Study Importance Questions

- Using data from a large sample of postmenopausal women in the Women's Health Initiative, in this paper we examine relative (e.g., risk ratio, rate ratio) and absolute (e.g., risk difference, rate difference) effect estimates for the BMI-mortality relationship.
- We report on interaction between age and BMI on mortality risk on both the additive and multiplicative scales. We found evidence of super-additive and sub-multiplicative interaction between age and high BMI. Super-additive interaction implies that the number of deaths due to the combination of obesity and aging is greater than the sum number of deaths that would be due to either exposure independently, whereas sub-multiplicative interaction indicates that the product of the joint effects of obesity in aging is less than the product of the two exposures individually.
- Obesity in older adults is an important topic; as life expectancies continue to rise and given the high population prevalence of obesity, there is a need for clinical, epidemiologic and public health research on effective policies and intervention programs specifically aimed at older adults with high BMI

Figure 1.
Kaplan Meier survival curve by BMI categories

Figure 2.
Graphical comparison of risk ratio and risk difference results for the effect of BMI on mortality among women who were 50–59 at baseline and 70–79 at baseline

Table 1.

Demographic characteristics of the WHI clinical trial cohort at baseline (n=68132)

	WHI clinical trial cohort (n=68132)
Age, years (mean $\pm$ SD)	62.1 $\pm$ 7.0
Age category at baseline (n; %)	
50-59	27,408 (40.2)
60-69	30,193 (44.3)
70-79	10,531 (15.5)
Race/ethnicity (n; %)	
Non-Hispanic White	55,631 (81.7)
Non-Hispanic Black	7,000 (10.3)
Hispanic	2,876 (4.2)
Other	2,601 (3.8)
Education (n; %)	
Some high school	3,795 (5.6)
High school diploma or GED	12,502 (18.5)
Post-secondary school	33,871 (50.0)
Post-graduate school	17,515 (25.9)
Smoking status (n; %)	
Current smoker	5,352 (7.9)
Former smoker	27,596 (41.0)
Never smoker	34,409 (51.1)
Marital status (n; %)	
Married	41,340 (60.9)
Widowed	11,588 (17.1)
Divorced	11,076 (16.3)
Has health insurance (n; %)	63,320 (93.8)
Employed (n; %)	23,822 (39.4)
History of CVD (%)	12,959 (19.0)
BMI categories (kg/m ² ; n; %)	
<18.5	294 (0.43)
18.5-24.9	18,333 (26.9)
25-29.9	24,310 (35.7)
30-34.9	15,240 (22.3)
35-39.9	6,757 (9.9)
>40	3,171 (4.7)

	WHI clinical trial cohort (n=68132)
Body mass index (mean $\pm$ SD)	28.9 $\pm$ 5.9
Years since menopause (mean $\pm$ SD)	14.4 $\pm$ 9.0

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2.

Comparison of relative (risk ratio) and absolute (risk difference) estimates from multivariate adjusted models stratified by baseline age cohort (50–59, 60–69, 70–79)

	Mortality Risk Ratio (95% CI)	Mortality Risk Difference (95% CI) per 10,000
Age 50–59		
<18.5 kg/m ²	1.34 (0.86, 2.08)	91 (–78, 261)
18.5–24.9 kg/m ²	0.99 (0.91, 1.09)	12 (–22, 46)
25–29.9 kg/m ²	-ref-	-ref-
30–34.9 kg/m ²	1.08 (0.98, 1.18)	74 (38, 111)
35–39.9 kg/m ²	1.38 (1.24, 1.53)	212 (159, 264)
>40 kg/m ²	1.74 (1.54, 1.96)	432 (349, 515)
Age 60–69		
<18.5 kg/m ²	1.13 (0.91, 1.40)	218 (–187, 622)
18.5–24.9 kg/m ²	1.01 (0.96, 1.05)	29 (–52, 109)
25–29.9 kg/m ²	-ref-	-ref-
30–34.9 kg/m ²	1.09 (1.04, 1.15)	177 (91, 264)
35–39.9 kg/m ²	1.18 (1.11, 1.25)	504 (379, 1227)
>40 kg/m ²	1.29 (1.19, 1.40)	1030 (832, 1364)
Age 70–79		
<18.5 kg/m ²	1.01 (0.84, 1.21)	471 (–404, 1347)
18.5–24.9 kg/m ²	1.03 (0.99, 1.07)	62 (–112, 236)
25–29.9 kg/m ²	-ref-	-ref-
30–34.9 kg/m ²	1.02 (0.98, 1.07)	384 (197, 572)
35–39.9 kg/m ²	1.08 (1.02, 1.15)	1093 (362, 1226)
>40 kg/m ²	1.18 (1.07, 1.30)	2231 (1799, 2663)

Table 3.

Interaction results on the additive and multiplicative scale for the effect of BMI category on mortality per 5 year age increase

	Relative Excess Risk of Interaction (95% CI)	Additive Interaction	Interaction (β) Coefficient	Multiplicative Interaction
Per 5-year age increase				
<18.5 kg/m ²	1.15 (−1.37, 3.67)	Super-additive	−0.09 (−0.24, 0.05)	Sub-multiplicative
18.5–24.9 kg/m ²	−0.02 (−0.17, 0.14)	Sub-additive	0.004 (−0.03, 0.03)	Super-multiplicative
25–29.9 kg/m ²	---	--	--	--
30–34.9 kg/m ²	0.26 (0.02, 0.50)	Super-additive	−0.03 (−0.06, 0.003)	Sub-multiplicative
35–39.9 kg/m ²	0.99 (0.46, 1.52)	Super-additive	−0.08 (−0.12, −0.04)	Sub-multiplicative
>40 kg/m ²	2.51 (1.17, 3.85)	Super-additive	−0.13 (−0.19, −0.08)	Sub-multiplicative