ORIGINAL RESEARCH

Seven cervical sensorimotor control tests measure different skills in individuals with chronic idiopathic neck pain

Rutger M.J. de Zoete^{a,b,c,d,*}, Peter G. Osmotherly^{a,b,c}, Darren A. Rivett^{a,b,c}, Suzanne J. Snodgrass^{a,b,c}

^a School of Health Sciences, Faculty of Health and Medicine, The University of Newcastle, Callaghan, NSW, Australia

^b Center for Brain and Mental Health Research, The University of Newcastle, Callaghan, NSW, Australia

^c Hunter Medical Research Institute, Newcastle, NSW, Australia

^d Recover Injury Research Centre, NHMRC Centre of Research Excellence in Recovery Following Road Traffic Injuries, The University of Queensland, Herston, Australia

Received 23 August 2018; received in revised form 24 October 2018; accepted 26 October 2018

Available online 11 November 2018

KEYWORDS

Neck;
Neck pain;
Proprioception;
Feedback;
Sensory

Abstract

Background: Sensorimotor control is commonly reported in neck pain research and rapidly gaining interest in clinical practice. Joint position error (conventional and torsion), postural balance, subjective visual vertical, head tilt response, The Fly[®], smooth pursuit neck torsion and head steadiness are tests that have been reported to assess cervical sensorimotor control. However, it is unknown whether clinicians could use one test, or a test battery, to appropriately assess cervical sensorimotor control and improve efficiency. Our main research question is: Do seven cervical sensorimotor control tests measure unique or similar characteristics of sensorimotor control in individuals with chronic idiopathic neck pain?

Methods: Principle components factor analysis. Data from seven cervical sensorimotor control tests of 50 participants with chronic idiopathic neck pain were included. Individual factors, potentially related to sensorimotor control, were determined by Eigen values >1.00 and inspection of a loading plot. Items with loadings ≥ 0.40 were considered satisfactory for inclusion in a factor.

Results: All cervical sensorimotor control tests were found to measure unique skills. Four factors were isolated with two, postural balance and head steadiness, accounting for most of the variance across tests. The remaining two factors, continuous movement accuracy and perceived verticality, contributed less to the observed variance.

* Corresponding author at: Recover Injury Research Centre, The University of Queensland, Level 7, Oral Health Centre, 288 Herston Road, Herston, QLD 4006, Australia.

E-mail: r.dezoete@uq.edu.au (R.M. de Zoete).

Conclusion: Postural balance and head steadiness were the major underlying factors explaining cervical sensorimotor control in the current sample. However, our results imply that all seven tests are independent and measure different skills. It is not possible to recommend a test battery for clinical practice, as all tests measure unique skills which appear to be independent of each other.

© 2018 Associação Brasileira de Pesquisa e Pós-Graduação em Fisioterapia. Published by Elsevier Editora Ltda. All rights reserved.

Introduction

Sensorimotor control (SMC) has been defined as the afferent and efferent information streams, as well as the central processing of these two, contributing to joint stability.¹ Cervical sensorimotor control (CSMC), which comprises of head and neck movement control, eye movement control, and postural stability, is commonly considered important in neck pain research.²

Specifically, in individuals with idiopathic neck pain, it has been suggested that CSMC may be affected.^{3–5} To date, seven CSMC tests have been suggested useful in the assessment of individuals with idiopathic neck pain⁶: the joint position error (JPE),⁷ postural balance,^{7,8} subjective visual vertical,⁹ head tilt response,¹⁰ smooth pursuit neck torsion,^{11,12} head steadiness,¹³ and The Fly[®] test.¹⁴

The most commonly used test is the JPE test, which has been reported in a large number of studies and included in several reviews and meta-analyses.^{6,15,16} Despite its widespread use, the clinical significance of the JPE test remains unclear due to the lack of a gold standard for comparison. This also applies to the other suggested tests that have similarly not been validated, as the clinical meaningfulness of these tests is based on minimal research, conducted in varying populations.^{12,17–29}

Despite the minimal understanding of their meaningfulness, the clinical use of CSMC tests in individuals with neck pain is rapidly gaining interest.³⁰ However, the availability of multiple tests purportedly assessing CSMC may introduce a challenge for clinicians in trying to select an appropriate test. Furthermore, clinicians may believe they have a valid measurement of SMC by just using one test, or conversely believe they might need to use all available tests to properly assess SMC, as it is currently unclear how many and which tests provide a clinically meaningful understanding of SMC. This lack in understanding may possibly lead to invalid clinical assessments, and potentially unnecessary use of valuable resources.

SMC comprises multiple subsystems: proprioception and kinesthesia (both responsible for the sensation of head and neck orientation relative to the body), the visuomotor system (the processing of visual information for the purpose of movement) and the vestibular system (located in the inner ear, predominantly responsible for spatial orientation of the head and neck, and postural balance).¹ It might be expected that different tests would measure the performance of one or more of these subsystems. However, available evidence

does not allow for an easy comparison of reported SMC outcomes in order to determine whether a particular subsystem is being assessed by each of these tests.³¹ Individual studies report different outcomes in different neck pain populations^{21,28,32} and source populations.^{22,24,27}

If groups of tests can be identified that address the different subsystems of CSMC,⁴ a test battery could potentially be developed for clinical practice. This may lead to a reduction in the amount of time and resources needed to assess CSMC in clinical practice.

Presently, based solely on face validity, it is proposed that different tests measure different subsystems of CSMC.^{1,33} The JPE and the head steadiness tests have been suggested to assess primarily cervical proprioceptive or kinaesthetic afferent input.^{7,13} The postural balance test is proposed to evaluate the vestibular and visual systems, but is thought to also incorporate cervical proprioceptive information.²⁴ Both the subjective visual vertical test and the head tilt response test assess the visual perception of verticality linked to a motor output, potentially reflecting the visuomotor system.^{10,34} The Fly[®] test combines a visual input with a continuous movement control exercise, and is proposed to address the visuomotor system as well as proprioception/kinesthesia.³⁵ Finally, the smooth pursuit neck torsion test is considered to assess oculomotor control.^{11,12}

Even though similarities in the assessed subsystems are to be expected across these tests based on their face validity, to date, any overlap between different tests has not been investigated and it remains unclear whether tests can actually be grouped together to assess a particular characteristic of SMC.³⁶ Tests are often used interchangeably as it is unclear which and how many tests are needed to get a clinically meaningful measurement. Until it is possible to validate cervical sensorimotor control tests using a gold standard, assessing whether tests measure the same or different subsystems of sensorimotor control is an important question in order to use time and resources efficiently. The present study aims to determine whether CSMC tests measure the same, or different, subsystems of SMC in individuals with chronic idiopathic neck pain. It is hypothesized that conditions of the same test, as well as different tests, should cluster together. This would identify clusters of tests that would assess different subsystems of CSMC. Subsequently, clustering of tests could be used clinically in determining what tests should be used in CSMC assessment.

Methods

Design

For this factor analysis, that follows the STROBE guidelines,³⁷ data were collected between July 2016 and June 2017. This study complies with the Declaration of Helsinki and was approved by the Human Research Ethics Committee of The University of Newcastle, Australia (H2016-0099). All participants included in this study provided written informed consent.

Participants

Participants with chronic idiopathic neck pain were included, recruited from The University of Newcastle Callaghan campus, the Hunter Medical Research Institute Research Register, and physiotherapy practices in Newcastle, NSW, Australia. Local advertising and social media were also used to recruit participants. Adults with chronic idiopathic neck pain were eligible to participate if they rated their pain at least 4/10 on a numeric rating scale. Exclusion criteria were limited vision (after correction), current or past history of diabetes, current or past history of migraine headaches, history of neck trauma or surgery, and other neurological or musculoskeletal conditions that may be considered to affect SMC.

Sample size

For factor analyses, there is no definitive method for sample size calculation. A sample-to-variables ratio between 2:1 and 5:1 is most common for factor analyses according to a recent systematic review.³⁸ With seven main outcome variables (the seven SMC tests), a sample size of 35 participants with idiopathic neck pain was considered sufficient.³⁹ As this study was part of a larger study with follow-up measurements, a sample of 50 participants was recruited to allow for withdrawals.

Procedure

Participant characteristics were recorded at the start of each session. Body mass index (BMI) was calculated from height and weight measures, using a stadiometer to the nearest cm (Surgical and Medical Supplies Pty. Ltd., Rose Park, SA, Australia) and a standard analogue scale to the nearest kg (model 762 1019009, seca gmbh & co., Hamburg, Germany) respectively. As subjective pain response varies with different recall periods in individuals with neck pain, the 100mm visual analogue scale (VAS), anchored by no pain on the left and worst pain imaginable on the right, was used to assess current pain, average pain over the last seven days, and average pain over the last four weeks.⁴⁰ The neck disability index (NDI, scored 0–50) was used to assess perceived disability.⁴¹ Physical activity levels were assessed using the Godin-Shephard Leisure-Time Physical Activity Questionnaire.⁴² Cervical range of motion (ROM) was assessed with the cervical range of motion instrument (CROM, Performance Attainment Associates, Lindstrom, MN,

U.S.) for rotation (left and right), flexion and extension.⁴³ From three measurements for each direction, range of motion was recorded as the mean total range in the transverse (rotation) and sagittal (flexion/extension) plane.

Sensorimotor control tests

Seven SMC tests, as identified by a recent systematic review,⁴⁴ were included in this factor analysis: JPE, postural balance, subjective visual vertical, head tilt response, the Fly, smooth pursuit neck torsion, and head steadiness. [Table 1](#) provides a detailed description of each of the tests.

For some tests multiple test conditions were utilized: JPE was tested conventionally and with torsion, postural balance was tested in four conditions with eyes open and eyes closed, and neutral and torsion stance, the Fly test used three movement patterns differing in difficulty, and the head steadiness test was performed in a low load and high load condition. In total, seven tests were included, consisting of a total of 14 test conditions. A custom script written in MATLAB (MATLAB R2015a, The MathWorks, Inc., Natick, MA, U.S.) was used to randomize the order of range of motion tests and SMC test conditions.

Data analysis

All cervical sensorimotor test variables were measured on an ordinal scale and assumptions for performing the principal components factor analysis were checked prior to analysis including linearity, multicollinearity, and homoscedasticity. The correlation between test results was examined using Spearman's rho to assess the strength of any association between test results. A principle components factor analysis was then conducted to identify any relationships between the results of individual SMC tests and to expose the number of underlying factors present in the data. The optimum number of factors present was determined by the number of Eigen values greater than 1.00 and visual interpretation of the factor structure using a loading plot. In order to further clarify the factors, a Promax oblique rotation was performed. An oblique rotation was selected as we postulated that the factors identified as skills of SMC in people with chronic idiopathic neck pain would not necessarily be independent of each other and may influence other factors present in this population. Item loadings on each factor were greater than or equal to 0.40 for inclusion in that factor. Elements within each identified factor were subsequently analyzed and each factor was characterized. All statistical analyses were performed using Stata 13 (StataCorp. 2013. *Stata Statistical Software: Release 13*. College Station, TX: StataCorp LP).

Results

The sample of individuals with chronic idiopathic neck pain consisted of 30 females and 20 males with a mean age of 41.61 (SD: 17.52) years. Participant demographics are presented in [Table 2](#). Strong correlations (Spearman's rho higher than 0.60) were found between similar conditions of the same test, but not between different tests ([Table 3](#)).

Table 1 Cervical sensorimotor control measures.*Joint position error (JPE)*

In the conventional JPE test, the participant was seated on a chair facing a target on the wall 90 cm away. A standard laser pointer (Electus Distribution Pty. Ltd, Rydalmere, NSW, Australia) was placed on top of the head and the participant was blindfolded. While starting with the laser pointer exactly in the centre of the target, the participant was instructed to move the head away from the target. Upon returning to the centre, the difference between starting position and final position was assessed. This was repeated three times for right rotation, left rotation, flexion, and extension, and the mean error for each direction was calculated for analyses. The head was repositioned to the centre of the target after each attempt. To minimize sensory reference information, a foam cushion was placed under the feet.⁸

For the JPE torsion test, a laser pointer (Electus Distribution Pty. Ltd, Rydalmere, NSW, Australia) was attached to the mid-sternum region using a strap around the chest. While the participant was seated in a neutral position, the laser pointer was aligned with the centre of the target on the wall 90 cm away. While the examiner gently held the head stationary, the participant was instructed to rotate the trunk away from the target and return to the starting position. Similar to the conventional JPE, the difference between starting position and final position was measured. This was repeated three times for rotation to the right and to the left, and the mean value was used for each direction. For both tests, movements were performed through comfortable range of motion.⁸

Postural balance

In the assessment of postural balance, a Kistler force platform (type 9260AA6, Kistler Instrumente AG, Switzerland) was used. While the participant was asked to take a comfortable stance on the platform, the centre of pressure was recorded for 30 s. This was done with the eyes both open and closed in a neutral head position. This was then repeated with neck torsion to the left and to the right (both eyes open and eyes closed). In order to achieve the torsion stance, the examiner gently held the participant's head while they rotated their body 45 degrees, including their feet, to either the right or left. The reference position on the platform was marked with pieces of visible electrical tape. For the collection and analysis of postural sway data (total sway path in mm), MARS (S2P Ltd, Slovenia) software was used.

Subjective visual vertical

An Oculus Rift (SDK 2.0, Oculus VR, CA, U.S.) virtual reality device was used for the subjective visual vertical test. Custom made software as described by Treleaven and Takasaki⁹ was displayed in the virtual reality device. Two dots representing the end points of a 20 degree tilted imaginary line were presented within a square that was tilted 18 degrees. Both the tilted square and dots could be positioned either clockwise or anticlockwise, which was randomized by the software. The computer mouse could be used to rotate a button, allowing rotation of the dots by a minimum of 0.01 degree. The participant was instructed to reposition the dots so that the imaginary line was positioned vertically (i.e., dots directly over each other). The deviation from the true vertical was recorded as the average of the deviation of the two dots. This procedure was repeated 20 times and the mean deviation from these repetitions was used for analyses.³⁴

Head tilt response

For the head tilt response test, the Oculus Rift virtual reality device was used with custom software (Libra@Home, Israel) as described by Geisinger, Ferreira, and Suarez.¹⁰ In the virtual reality device, a tilted white line was presented. The line was tilted either clockwise or anticlockwise, and the participant was instructed to tilt the head laterally so that the line would be positioned vertically. The Oculus Rift's head tracking sensors recorded the head tilt angle, which was compared to the line that was presented in the virtual reality device.¹⁰

The Fly[®]

The original Fly test software was used as introduced by Kristjansson et al.,¹⁴ together with an electromagnetic tracking device (Polhemus Liberty, FasTrak, Colchester, VT, U.S.) to record head movement. On a computer screen, a representation of the head position was displayed with a crosshair, as well as a moving target (the "Fly"). By moving the head and neck, the target could be followed and movement accuracy (distance between crosshair and target) was measured. This was repeated for three different movement patterns – easy, medium, and difficult – which differed in length, time, and complexity. For the duration of each movement pattern, the mean distance between crosshair and target was used for analyses. A fourth movement pattern was used to familiarize participants with the test.

From the principle components factor analysis, four factors with Eigen values over 1.00 emerged. Table 4 presents these factors, as well as the constituent items that were included in factors 1–4.

Factor one consisted of all four postural balance test conditions; eyes open, eyes closed, torsion eyes open, torsion eyes closed, and was thus labelled as 'postural balance'. Factor two consisted of both low and high load conditions of the head steadiness test and was labelled 'head steadiness'.

Factor three consisted of the medium and difficult movement patterns of the Fly test and was labelled 'continuous movement accuracy'. Factor four consisted of the subjective visual vertical and head tilt response tests and was labelled 'perception of verticality'. Through observation of the factor loadings plot (Fig. 1) it was determined that factor one (postural balance) and factor two (head steadiness) formed clusters visually distinct from the other SMC tests.

Table 1 (Continued)

Smooth pursuit neck torsion

The smooth pursuit neck torsion test was adapted from Tjell and Rosenhall¹¹ and Treleaven, Jull, and LowChoy.¹² Testing was performed with the participant, seated facing a wall 100 cm away. Custom made equipment which consisted of a laser pointer attached to a motor was positioned behind the participant with the laser pointer just above the head. The laser pointer, which projected a red dot (the target) on the wall, followed a sinusoidal pattern from left to right with a visual angle of 40 degrees. The target moved with a consistent speed of 20 degrees per second and a frequency of 0.2 Hz. The participant was asked to follow the moving target with their eyes, while keeping their head still. The target moved from left to right and back ten times, and eye movements were recorded using surface electrooculography. This was done with the body straight ahead, and also in a torsion position in which the body was rotated 45 degrees on the chair but the head kept stationary. Eye movement was recorded using Ag/AgCl surface electrodes (Cleartrace, ConMed, Utica, NY, U.S.) positioned bilaterally just lateral to the eyes, and with one ground electrode on the forehead. This method has been previously utilized to record the corneo-retinal potential for the horizontal assessment of binocular electrooculography.⁴⁵ The electrooculography signal was filtered through a 70 Hz low pass filter and the data were de-identified prior to analysis. Data were analyzed by graphing the average velocity of eye movement and identifying the proportion of saccades for the sixth to ninth sequence. The difference between neutral and the average of the left and right torsion positions was calculated and used as outcome measure.

Head steadiness

For head steadiness testing, an electromagnetic motion tracking device (Polhemus Liberty, FasTrak, Colchester, VT, U.S.) was used to record head movement. Two conditions were investigated; low load with the participant seated on a treatment table with the back/head support at a 60 degree angle, and high load, with the participant lying supine on a horizontal treatment table. The participant was asked to lift their head approximately 1 cm off the table and hold it still for 40 s. Head movement (i.e., head angular velocity), was recorded continuously during the tasks.¹³ Participants that were not able to hold the lifted position for a minimum of 10 s were excluded from the analysis. Raw data from sensor recordings were used to calculate average displacement and velocity over 40 s.¹³

Table 4 Cervical sensorimotor control test item loadings for each of the four factors.

Sensorimotor control test	Postural balance (Eigen value = 3.15)	Head steadiness (Eigen value = 2.06)	Continuous movement accuracy (Eigen value = 1.55)	Perceived verticality (Eigen value = 1.49)
Subjective visual vertical				0.466
Head tilt response				0.750
Conventional joint position error				
Joint position error torsion				
Balance EO	0.574			
Balance EC	0.790			
Balance torsion EO	0.996			
Balance torsion EC	1.018			
Fly (easy)				
Fly (medium)			0.733	
Fly (difficult)			0.550	
Head steadiness – low load		0.818		
Head steadiness – high load		0.982		
Smooth pursuit neck torsion				

Item loadings <0.40 are omitted from this table, as these did not account for a large proportion of the variance within each factor. EC, eyes closed; EO, eyes open.

Discussion

The present study used factor analysis to investigate whether existing CSMC tests measure the same, or different, subsystems of SMC. Two factors, postural balance and head steadiness, explained the majority of the variance in measured SMC. However, the results of this study demonstrate that most CSMC tests measure unique skills, and it is not possible to group them according to the different

subsystems of SMC. For clinical practice, this means that different tests measure unique skills, thus a clinician cannot simply select one test. It is also not possible to recommend a specific battery of tests which would suffice in the assessment of CSMC, as all tests are unique.

Two factors, postural balance (factor one) and head steadiness (factor two), had relatively high Eigen values of 3.15 and 2.06 respectively, demonstrating that they are responsible for a large proportion of the variance across the

Table 2 Demographics of participants ($n = 50$) with descriptive data for each test and test condition (mean, SD or median, IQR).

Characteristic	
Gender (n female)	30 (60%)*
Age (years)	35.5 (IQR 24.0–55.3)
Body mass index	24.8 (IQR 21.6–29.3)
Physical activity	Insufficient: 11 (22%)*
(n per group)	Moderate: 10 (20%)*
	Active: 29 (58%)*
VAS current (0–100 mm)	30.16 ($\pm$ 18.77)
VAS 7 days (0–100 mm)	49 (IQR 26–62)
VAS 4 weeks (0–100 mm)	50.78 ($\pm$ 20.06)
Pain duration	130 ($\pm$ 148)
(in months)	
NDI (0–50)	14.96 ($\pm$ 6.19)
Range of motion (deg)	
Total rotation	124.33 (IQR 97.83–137.67)
Total sagittal	125.33 (IQR 103.67–134.17)
(flexion/extension)	
SVV (deg)	3.74 (IQR 1.92–6.08)
HTR (deg)	2.90 (IQR 2.01–4.42)
JPE (deg)	
Conventional	3.85 (IQR 3.16–4.58)
Torsion	4.88 (IQR 3.54–6.83)
Balance (mm)	
Eyes open	227.5 (IQR 176.5–298.5)
Eyes closed	288.0 (IQR 208.8–391.3)
Torsion EO	261.0 (IQR 211.5–311.9)
Torsion EC	314.5 (IQR 260.3–401.3)
Fly (mm)	
Easy	2.10 (IQR 1.86–2.50)
Medium	2.15 (IQR 1.94–2.66)
Difficult	2.82 (IQR 2.42–3.53)
Head steadiness (mm/s)	
Low load	0.41 (IQR 0.35–0.62)
High load	0.61 (IQR 0.50–0.78)
SPNT (difference)	0.02 (IQR 0.01–0.03)

Physical activity measured by The Godin-Shephard Leisure-Time Physical Activity Questionnaire. EC, eyes closed; EO, eyes open; HTR, head tilt response; JPE, joint position error; NDI, neck disability index; SPNT, smooth pursuit neck torsion; SVV, subjective visual vertical; VAS, visual analogue scale.

* N (%) for categorical data.

tests. Factors three and four were less distinct clusters and accounted for less of the overall variance observed. From inspection of the factor loading plot, it is clear that postural balance and head steadiness groupings are distinct from the other tests, which implies that these two factors measure different characteristics of SMC, which do not appear to be captured by other SMC tests.

It has been suggested that postural balance is an important measure of SMC in individuals with neck pain.^{3,46} Cervical proprioceptors, the vestibular system and the visuomotor system are important in providing relevant

afferent information for postural balance, and this may be affected in individuals with neck pain.³³ Even though little consistency exists in the reported discriminating abilities of postural balance testing between individuals with idiopathic neck pain and healthy individuals,^{3,6,24,46} postural balance may be a unique characteristic of SMC, not captured by the other tests that are all performed seated. Alternatively, balance tests may rely on other systems outside of CSMC and thus may be testing a different construct to the other tests.

Factor two consisted of the two test conditions of the head steadiness test (i.e., low load and high load), and did not include other tests. The head steadiness test has been used in the sensorimotor assessment of individuals with neck pain in only one identified study that found differences between individuals with idiopathic neck pain and healthy controls.¹³ However, based on the test's characteristics, it might be argued that the head steadiness test is predominantly a muscle endurance test and not a true SMC test. Rather than being the result of differences in SMC skills, the discriminating capabilities of this test may illustrate a difference in neck muscle endurance and activation which has been reported in other literature.^{47,48} Other tests examined in this study do not require high levels of muscle strength, which is probably why the head steadiness test was found to be distinct from the other tests.

Factor three was labelled continuous movement accuracy and consisted of two conditions of the Fly test; the medium and difficult movement pattern. The Fly test is a continuous movement accuracy test suggested to involve sensory (visual) information processing and a motor output specific for the cervical spine (head movement).^{35,49} These results suggest the medium and difficult Fly test conditions measure unique skills that are not being assessed in the other tests. However, limited evidence^{14,35} is available for the Fly test and outcomes have only been examined cross-sectionally.

In the present factor analysis, not all Fly test conditions cluster together in one factor: the easy movement pattern failed to contribute to factor three. This was unexpected as the different test conditions appear to be very similar and the three conditions are together considered as one test. The only notable differences between the three conditions are the test duration and the number of changes in movement direction. The easy pattern takes 25 s, the medium takes 40 s and the difficult pattern takes 50 s, and the number of direction changes increases with each level of difficulty. A potential reason for the easy condition not clustering with the two more demanding conditions (and also with none of the other tests) is its difficulty: it may simply be too easy to test continuous movement accuracy. However, previous studies did not find a difference in the discriminative ability of the three conditions of the Fly test in both whiplash and idiopathic neck pain populations.^{14,35}

The subjective visual vertical test and the head tilt response test clustered in factor four. The two tests are both spatial orientation tests in which perceived verticality is assessed.^{9,10} The head tilt response test has been introduced as supplementary to the subjective visual vertical test¹⁰ as it introduces a motor output more specific to the cervical spine (i.e., head tilt, compared to arm/hand movement to

Table 3 Correlations between cervical sensorimotor control tests and test conditions in individuals with chronic idiopathic neck pain.

	SVV	HTR	JPE conven- tional	JPE torsion	Balance EO	Balance EC	Balance torsion EO	Balance torsion EC	Fly Easy	Fly Medium	Fly Difficult	Head steadiness Low	Head steadiness High
HTR HTR	0.370*												
JPE Conventional	0.311	0.038											
JPE torsion	0.160	-0.087	0.372										
Balance EO	-0.250	0.181	0.010	-0.102									
Balance EC	-0.396*	0.051	-0.185	-0.007	0.727*								
Balance torsion EO	-0.361*	0.167	-0.097	-0.101	0.750*	0.781*							
Balance torsion EC	-0.439*	0.064	-0.156	-0.092	0.715*	0.805*	0.903*						
Fly Easy	0.261	0.107	0.235	0.233	-0.213	-0.188	-0.106	-0.192					
Fly Medium	0.332*	0.174	0.074	0.099	-0.227	-0.181	0.011	-0.069	0.544*				
Fly Difficult	0.366*	0.175	0.083	0.295*	-0.106	-0.136	-0.097	-0.135	0.624*	0.665*			
Head steadiness Low	-0.034	-0.028	-0.125	0.097	0.183	0.258	0.087	0.091	0.127	-0.055	0.208		
Head steadiness High	-0.039	-0.017	0.034	0.057	0.210	0.120	0.043	-0.056	0.212	-0.074	0.265	0.702*	
SPNT	0.144	-0.006	0.003	-0.095	0.000	-0.046	0.035	-0.067	-0.144	-0.102	-0.031	0.020	0.103

EC, eyes closed; EO, eyes open; HTR, head tilt response; JPE, joint position error; SPNT, smooth pursuit neck torsion; SVV, subjective visual vertical. Correlations provided are Spearman's rho, p -value between brackets, bold and * denotes statistically significant ($\alpha = 0.05$).

Table 4 Cervical sensorimotor control test item loadings for each of the four factors.

Sensorimotor control test	Postural balance (Eigen value = 3.15)	Head steadiness (Eigen value = 2.06)	Continuous movement accuracy (Eigen value = 1.55)	Perceived verticality (Eigen value = 1.49)
Subjective visual vertical				0.466
Head tilt response				0.750
Conventional joint position error				
Joint position error torsion				
Balance EO	0.574			
Balance EC	0.790			
Balance torsion EO	0.996			
Balance torsion EC	1.018			
Fly (easy)				
Fly (medium)			0.733	
Fly (difficult)			0.550	
Head steadiness – low load		0.818		
Head steadiness – high load		0.982		
Smooth pursuit neck torsion				

Item loadings <0.40 are omitted from this table, as these did not account for a large proportion of the variance within each factor. EC, eyes closed; EO, eyes open.

Figure 1 Factor loading plot of all cervical sensorimotor control tests. Postural balance and head steadiness account for most of the variance across tests.

EC, eyes closed; EO, eyes open; HTR, head tilt response; JPE, joint position error; SPNT, smooth pursuit neck torsion; SVV, subjective visual vertical.

control a computer mouse in the subjective visual vertical test). However, as the two tests cluster together in factor four, it can be argued that they both measure spatial orientation.

Based on the assumption that all tests measure SMC, stronger associations between different tests were expected. From the current findings, it can be concluded that each of these CSMC tests measure different skills that have been suggested relevant in SMC assessment. If multiple CSMC tests were found to cluster together and contribute to one factor, this would have supported the assumption that SMC is a single entity that might be measured using a single test or a small number of different tests. However, the present outcomes lead to the more likely explanation that each of these tests measure their own individual skill, which may potentially all be different characteristics of SMC.

A noteworthy observation is the fact that the most commonly used cervical SMC test, the JPE test, failed to contribute to any of the factors in this study. This suggests that this test must assess a unique skill, different to all of the other tests we investigated. Although the two test conditions of JPE (conventional and torsion) might have been expected to strongly correlate based on their face validity, Table 3 demonstrates they do not ($\rho = .37$). Possibly, the subsystems of CSMC assessed by the two test conditions may be different. The conventional test condition appears to assess both the proprioceptive and vestibular subsystems, whereas the torsion condition, in which the head is kept still, the vestibular system may not contribute as much. The findings from the current study cannot determine whether the skill assessed by JPE relates to a specific characteristic of SMC, only that it must be unique to the other tests evaluated in the current study.

Limitations and strengths

Required sample size in factor analysis remains contentious. Despite the lack of clear sample size requirements for factor analyses, a participant:variable ratio of 5:1 was considered sufficient.³⁹ Based on available literature, however, it was assumed that all test conditions that are part of the same test, for example the three conditions of The Fly[®] test, could be considered as one factor. As outlined above, the results show that the three conditions of the Fly test do not cluster together in one factor, which results in the creation of an extra factor and thus a reduction in statistical power.

A strength of the current study is the use of factor analysis to explore commonalities between the available tests. Unlike related literature to date which solely reports the group-comparison findings of cross-sectional studies, this study has allowed us to address whether available CSMC tests measure similar skills. As this issue has not been addressed before, the present findings contribute to our understanding of available CSMC tests in both clinical and research settings.

A methodological strength of the current study is that test methods similar to the original test descriptions were used for all tests. By closely collaborating with other research groups, we were able to replicate test methods and even use the original test equipment and software for most of the tests. This results in the methods of the present study being highly comparable to earlier studies and strengthens its external validity. Furthermore, as most tests used computerized recordings, the likelihood that the examiner influenced the recording of test outcomes is negligible, which reduces assessor bias.

Conclusions

All CSMC tests measure unique skills and cannot be grouped together under SMC subsystems such as proprioception, visuomotor control and the vestibular system. Only very similar tests, or test conditions that are part of the same test, were found to cluster together. This indicates that the available CSMC tests do not measure similar skills. Clinicians assessing CSMC therefore arguably have to perform each test, as each provides unique information. The clinical relevance of this information, in terms of the concept known as CSMC, remains unknown.

Acknowledgements

The researchers wish to acknowledge the support of the HMRI Research Volunteer Register for their assistance in participant recruitment.

References

1. Riemann BL, Lephart SM. The sensorimotor system. Part I: The physiologic basis of functional joint stability. *J Athl Train*. 2002;37(1):71–79.
2. Treleaven J. Sensorimotor disturbances in neck disorders affecting postural stability, head and eye movement control. *Man Ther*. 2008;13(1):2–11.
3. Jørgensen MB, Skotte JH, Holtermann A, Sjøgaard G, Petersen NC, Søgaard K. Neck pain and postural balance among workers with high postural demands – a cross-sectional study. *BMC Musculoskelet Disord*. 2011;12:176.
4. Meisingset I, Woodhouse A, Stensdotter AK, et al. Evidence for a general stiffening motor control pattern in neck pain: a cross sectional study pathophysiology of musculoskeletal disorders. *BMC Musculoskelet Disord*. 2015;16(1).
5. Rudolfsson T, Djupsjöbacka M, Hager C, Björklund M. Effects of neck coordination exercise on sensorimotor function in chronic neck pain: a randomized controlled trial. *J Rehabil Med*. 2014;46(9):908–914.
6. de Zoete RM, Osmotherly PG, Rivett DA, Farrell SF, Snodgrass SJ. Sensorimotor control in individuals with idiopathic neck pain and healthy individuals: a systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2016.
7. Revel M, Andre-Deshays C, Minguet M. Cervicocephalic kinesthetic sensibility in patients with cervical pain. *Arch Phys Med Rehabil*. 1991;72:288–291.
8. Chen X, Treleaven J. The effect of neck torsion on joint position error in subjects with chronic neck pain. *Man Ther*. 2013;18(6):562–567.
9. Treleaven J, Takasaki H. High variability of the subjective visual vertical test of vertical perception, in some people with neck pain – should this be a standard measure of cervical proprioception? *Man Ther*. 2015;20(1):183–188.
10. Geisinger D, Ferreira E, Suarez A, Suarez H. Head tilt response: a complementary test to the subjective visual vertical. *J Vestib Res*. 2010;20(5):381–389.
11. Tjell C, Rosenhall U. Smooth pursuit neck torsion test: a specific test for cervical dizziness. *Am J Otol*. 1998;19(1):76–81.
12. Treleaven J, Jull G, LowChoy N. Smooth pursuit neck torsion test in whiplash-associated disorders: relationship to self-reports of neck pain and disability, dizziness and anxiety. *J Rehabil Med*. 2005;37(4):219–223.
13. Woodhouse A, Liljebäck P, Vasseljen O. Reduced head steadiness in whiplash compared with non-traumatic neck pain. *J Rehabil Med*. 2010;42(1):35–41.
14. Kristjansson E, Oddsdóttir GL. ‘‘The Fly’’: a new clinical assessment and treatment method for deficits of movement control in the cervical spine. *Spine (Phila Pa 1976)*. 2010;35(23):E1298–E1305.
15. de Vries J, Ischebeck BK, Voogt LP, et al. Joint position sense error in people with neck pain: a systematic review. *Man Ther*. 2015;20(6):736–744.
16. Stanton TR, Leake HB, Chalmers KJ, Moseley GL. Evidence of impaired proprioception in chronic, idiopathic neck pain: systematic review and meta-analysis. *Phys Ther*. 2016;96(6):876–887.
17. Elsig S, Luomajoki H, Sattelmayer M, Taeymans J, Tal-Akabi A, Hilfiker R. Sensorimotor tests, such as movement control and laterality judgment accuracy, in persons with recurrent neck pain and controls. A case-control study. *Man Ther*. 2014;19(6):555–561.
18. Grod JP, Diakow PR. Effect of neck pain on verticality perception: a cohort study. *Arch Phys Med Rehabil*. 2002;83(3):412–415.
19. Hansson EE, Mansson NO, Hakansson A. Balance performance and self-perceived handicap among dizzy patients in primary health care. *Scand J Prim Health Care*. 2005;23(4):215–220.
20. Juul-Kristensen B, Clausen B, Ris I, et al. Increased neck muscle activity and impaired balance among females with whiplash-related chronic neck pain: a cross-sectional study. *J Rehabil Med*. 2013;45(4):376–384.
21. Lange B, Murray M, Chreiteh SS, et al. Postural control and shoulder steadiness in f-16 pilots: a randomized controlled study. *Aviat Space Environ Med*. 2014;85(4):420–425.

22. Nagai T, Abt JP, Sell TC, et al. Neck proprioception, strength, flexibility, and posture in pilots with and without neck pain history. *Aviat Space Environ Med.* 2014;85(5):529–535.
23. Oddsdottir GL, Kristjansson E. Two different courses of impaired cervical kinaesthesia following a whiplash injury. A one-year prospective study. *Man Ther.* 2012;17(1):60–65.
24. Quek J, Brauer SG, Clark R, Treleaven J. New insights into neck-pain-related postural control using measures of signal frequency and complexity in older adults. *Gait Posture.* 2014;39(4):1069–1073.
25. Rix GD, Bagust J. Cervicocephalic kinesthetic sensibility in patients with chronic, nontraumatic cervical spine pain. *Arch Phys Med Rehabil.* 2001;82(7):911–919.
26. Röijezon U, Björklund M, Djupsjöbacka M. The slow and fast components of postural sway in chronic neck pain. *Man Ther.* 2011;16(3):273–278.
27. Sa S, Silva AG. Repositioning error, pressure pain threshold, catastrophizing and anxiety in adolescents with chronic idiopathic neck pain. *Musculoskelet Sci Pract.* 2017;30:18–24.
28. Sjolander P, Michaelson P, Jaric S, Djupsjöbacka M. Sensorimotor disturbances in chronic neck pain – range of motion, peak velocity, smoothness of movement, and repositioning acuity. *Man Ther.* 2008;13(2):122–131.
29. Woodhouse A, Vasseljen O. Altered motor control patterns in whiplash and chronic neck pain. *BMC Musculoskelet Disord.* 2008;9:90.
30. Michiels S, De Hertogh W, Truijzen S, November D, Wuyts F, Van de Heyning P. The assessment of cervical sensory motor control: a systematic review focusing on measuring methods and their clinimetric characteristics. *Gait Posture.* 2013;38(1):1–7.
31. Strimpakos N. The assessment of the cervical spine. Part 1: Range of motion and proprioception. *J Bodyw Mov Ther.* 2011;15(1):114–124.
32. Treleaven J, LowChoy N, Darnell R, Panizza B, Brown-Rothwell D, Jull G. Comparison of sensorimotor disturbance between subjects with persistent whiplash-associated disorder and subjects with vestibular pathology associated with acoustic neuroma. *Arch Phys Med Rehabil.* 2008;89(3):522–530.
33. Riemann BL, Lephart SM. The sensorimotor system. Part II: The role of proprioception in motor control and functional joint stability. *J Athl Train.* 2002;37(1):80–84.
34. Takasaki H, Treleaven J, Johnston V, Jull G. Minimum repetitions for stable measures of visual dependency using the dot version of the computer-based Rod–Frame test. *Man Ther.* 2012;17(5):466–469.
35. Kristjansson E, Hardardottir L, Asmundardottir M, Gudmundsson K. A new clinical test for cervicocephalic kinesthetic sensibility: “the fly”. *Arch Phys Med Rehabil.* 2004;85(3):490–495.
36. Kristjansson E, Treleaven J. Sensorimotor function and dizziness in neck pain: implications for assessment and management. *J Orthop Sports Phys Ther.* 2009;39(5):364–377.
37. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol.* 2008;61(4):344–349.
38. Osborne A, Costello J. Best practices in exploratory factor analysis: four recommendations for getting the most from your analysis. *Pract Assess Res Eval.* 2005;10(7).
39. O’Rourke N, Hatcher L. *A Step-by-Step Approach to Using SAS® for Factor Analysis and Structural Equation Modeling, Second Edition.* Cary, North Carolina, USA: SAS Institute Inc.; 2013.
40. Kamper SJ, Grootjans SJ, Michaleff ZA, Maher CG, McAuley JH, Sterling M. Measuring pain intensity in patients with neck pain: does it matter how you do it? *Pain Pract.* 2015;15(2):159–167.
41. Vernon H, Mior S. The neck disability index: a study of reliability and validity. *J Manipulative Physiol Ther.* 1991;14(7):409–415.
42. Godin G, Shephard RJ. A simple method to assess exercise behavior in the community. *Can J Appl Sport Sci.* 1985;10(3):141–146.
43. Audette I, Dumas JP, Cote JN, De Serres SJ. Validity and between-day reliability of the cervical range of motion (CROM) device. *J Orthop Sports Phys Ther.* 2010;40(5):318–323.
44. de Zoete RMJ, Osmotherly PG, Rivett DA, Farrell SF, Snodgrass SJ. Sensorimotor control in individuals with idiopathic neck pain and healthy individuals: a systematic review and meta-analysis. *Arch Phys Med Rehabil.* 2016.
45. Heide W, Koenig E, Trillenber P, Kömpf D, Zee DS. Electrooculography: technical standards and applications. *Electroencephalogr Clin Neurophysiol Suppl.* 1999;52:223–240.
46. Michaelson P, Michaelson M, Jaric S, Latash ML, Sjolander P, Djupsjöbacka M. Vertical posture and head stability in patients with chronic neck pain. *J Rehabil Med.* 2003;35(5):229–235.
47. Falla D. Unravelling the complexity of muscle impairment in chronic neck pain. *Man Ther.* 2004;9(3):125–133.
48. Falla DL, Jull GA, Hodges PW. Patients with neck pain demonstrate reduced electromyographic activity of the deep cervical flexor muscles during performance of the craniocervical flexion test. *Spine (Phila Pa 1976).* 2004;29(19):2108–2114.
49. Kristjansson E, Dall’Alba P, Jull G. A study of five cervicocephalic relocation tests in three different subject groups. *Clin Rehabil.* 2003;17:768–774.