Published in final edited form as:

Clin Lung Cancer. 2020 January ; 21(1): e25–e29. doi:10.1016/j.clcc.2019.10.002.

Complete Pathological Response to Crizotinib in a Patient with *ALK*-rearranged Lung Adenocarcinoma

Marissa S. Mattar, BS¹, Jason Chang, MD², Ryma Benayed, PhD², Darragh Halpenny, MD³, Astin Powers, MD⁴, David Kleiner, MD, PhD⁴, Alexander Drilon, MD⁵, Mark G. Kris, MD⁵

¹Antitumor Assessment Core Facility, Molecular Pharmacology Program, Memorial Sloan Kettering Cancer Center, New York, NY, USA.

²Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY, USA.

³Department of Radiology, Memorial Sloan Kettering Cancer Center, New York, NY, USA.

⁴Laboratory of Pathology, National Cancer Institute, National Institutes of Health, Bethesda, MD, USA.

⁵Department of Medicine, Division of Solid Tumor Oncology, Thoracic Oncology Service, Memorial Sloan Kettering Cancer Center, New York, NY, USA.

Keywords

EML4-ALK; EML4-ALK variants; Non-small cell lung cancer; Tyrosine kinase inhibitor; Tandem atypical propeller EML domain

Introduction

The identification of anaplastic lymphoma kinase (*ALK*) gene fusions as actionable oncogenic drivers in lung cancers led to the development and approval of multiple *ALK* tyrosine kinase inhibitors (TKIs); the first of these was crizotinib. Objective response rates with *ALK* TKIs are high,¹ and disease regression is achieved in most patients.² Here, we report the first complete pathological response to first-line crizotinib in a patient with *ALK*-rearranged lung cancer.

Case Presentation

A 75-year-old never-smoker male developed lethargy, chest pain, cough, and dyspnea. Computed chest tomography (CT) showed a 2 cm lesion in the right lower lobe, multiple pleural-based masses in the right hemithorax, and a right pleural effusion (Figure 1). The

Address for correspondence Alexander Drilon, MD, Memorial Sloan Kettering Cancer Center, 1275 York Avenue, New York, NY 10065, drilona@mskcc.org.

Conflict of interest disclosures:

Marissa S. Mattar, Jason Chang, Darragh Halpenny, Astin Powers, and David Kleiner have no conflicts of interest to disclose.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

patient underwent a bronchoscopy, cervical mediastinoscopy, and right-sided video-assisted thoracoscopy. Pleural biopsy revealed adenocarcinoma with signet ring cell features (Figure 2a). Genotyping of the tumor via Sequenom mass-spectrometry revealed no mutations in the following genes: AKT1, BRAF, EGFR, ERBB2, KRAS, MEK1 (MAP2K1), NRAS, and PIK3CA. Fluorescence in situ hybridization (FISH) revealed an *ALK* rearrangement (Figure 2b). First-line treatment with crizotinib (250 mg twice daily) was tolerated without undue toxicity. Imaging one month following treatment initiation showed reduction in the primary tumor and pleural metastases and resolution of the pleural effusion (Figure 1). Response to crizotinib was maintained, and the patient continued crizotinib for five years until death (Figure 1). An autopsy revealed death was due to pulmonary hemorrhage in the setting of cardiac arrhythmia. No cancer was found at autopsy (Figure 2c); however, complete resolution of disease was never radiologically observed. Discordance between the observed radiographic and pathological responses are not uncommon. As in our report, previous studies have found that the radiographic response rate underestimates the response rate found upon pathological evaluation.^{3, 4}

Discussion

The discovery of genomic aberrations that drive cellular proliferation and confer a fitness advantage in resulting tumors has led to a paradigm shift in the treatment of lung cancers. *ALK* rearrangements are present in 4% of patients with lung adenocarcinomas.^{5, 6} Crizotinib, a TKI of ALK, MET, and ROS, improves response rates and progression-free survival versus chemotherapy in patients with *ALK*-rearranged lung cancers.^{7, 8} To our knowledge, this is the first report of a complete pathological response to crizotinib confirmed by autopsy in a patient with metastatic *ALK*-rearranged lung adenocarcinoma. Similar complete responses were radiologically observed in 2 patients (3%) receiving first-line crizotinib in the frontline PROFILE 1014 clinical trial in which crizotinib was tested against standard chemotherapy.⁸ Additionally, the phase III clinical trial (PROFILE 1007) in which previously treated patients with *ALK*+ NSCLC were treated with crizotinib versus standard chemotherapy reported 1 case (1%) of complete radiological response.⁹ The incidence of complete responses in these studies in conjunction with our observed complete pathological response point to the likely existence of other dramatic, durable responses to TKI treatment.

It is likely the magnitude of response in *ALK*-rearranged cancers can vary with the *ALK* fusion partner. *EML4-ALK* rearrangements are inversions in chromosome 2p characterized by fusions of *ALK* exons 20 to 29 to varying regions of *EML4*. The coiled-coil domain in the N-terminus of *EML4* mediates kinase activation by serving as the trimerization domain and is thus required for the trimeric oligomerization of *EML4-ALK*.⁷ The fusion of *EML4* to the intracellular kinase domain of ALK causes ALK to localize within the cytoplasm instead of the plasma membrane.¹⁰ The intact ALK kinase domain found within the *EML4-ALK* fusions exhibits increased transforming activity and promotes oncogenesis via autophosphorylation and subsequent constitutive activation of the downstream pathways of ALK.¹¹⁻¹⁴ There is growing evidence to support the notion that tumors bearing *EML4-ALK* variant 1 (exon 13 of *EML4* fused to *ALK* exon 20) and variant 2 (exon 20 of *EML4* fused to *ALK* exon 20) experience greater sensitivity to ALK inhibitors than tumors with other

variants, namely variant 3 (exon 6a/b of *EML4* fused to *ALK* exon 20).¹¹ Two small, retrospective studies failed to note different levels of efficacy amongst the variant subgroups.^{15, 16} However, other studies, including a recent larger study with a longer follow-up period conducted by Christopoulos et al., found patients with *EML4-ALK* variant 3 rearrangements exhibited inferior outcomes compared to their variant 1 and variant 2 counterparts.^{14, 17, 18} Additionally, tumors with variant 3 fusions exhibited more aggressive behavior, including a later stage at initial presentation, a higher incidence of atypical metastatic sites, and an increased propensity for metastasis in general.^{18, 19}

Variants 1 and 2 are characterized by truncated tandem atypical propeller EML (TAPE) domains, whose resulting partial structure has been shown to induce protein instability. This instability is hypothesized to confer sensitivity to TKIs.^{5, 12, 15, 20} Given this patient's outstanding response to crizotinib, we suspected the patient had an *EML4-ALK* variant 1 or 2. To confirm our suspicions, we employed an Archer assay, an amplicon-based NGS method utilized to detect gene fusions.²¹ However, the Archer assay revealed a novel, in-frame fusion of *EML4* exon 17 to exon 20 of *ALK* with an 80 bp insertion from intron 19 (Figure 3). Fusions between *ALK* exon 20 and *EML4* exon 17 have been recently described as variant 8, which also exhibits a dysfunctional TAPE domain^{16, 22} and may explain the patient's lifelong response to crizotinib. This specific *ALK*-fusion variant appears to have imparted an exceptional sensitivity to the *ALK* TKI crizotinib that resulted in a complete pathological response.

Conclusion

To our knowledge, this is the first report of a complete pathological response to crizotinib in a patient with *ALK*-rearranged lung adenocarcinoma. The complete resolution of this patient's disease highlights the potential dominance of targeted therapy over standard-of-care cytotoxic chemotherapy. Recent studies have begun to illuminate the link between specific structural variants in *ALK* fusions and their sensitivities and eventual resistance to TKIs, including crizotinib. Lin et al. showed the *ALK* variant can serve as a predictive biomarker for mechanisms of resistance. They found that tumors bearing variant 3 *EML4-ALK* fusions were more likely to acquire secondary *ALK*-resistant mutations in response to treatments with TKIs. Specifically, variant 3 tumors were found to be significantly more likely to acquire a solvent-front mutation, *ALK* G1202R. The *ALK* G1202R mutation inhibits proper drug binding, thereby producing resistance to both first and second generation *ALK* inhibitors.²³ The acquired mutations are thought to decrease overall survival via accelerated metastasis and premature failure to multiple treatment modalities.¹⁸ Conversely, variant 1 tumors became resistant after a longer duration of TKI therapy and without the acquisition of an additional *ALK* resistance mutation such as *ALK* G1202R.^{23, 24} The influence of the *EML4-ALK* fusion variant on response to treatment with *ALK* inhibitors was recently explored in the Phase III ALEX trial, which compared the efficacy of crizotinib to alectinib, a second-generation *ALK* inhibitor. The study showed alectinib produced better clinical outcomes compared to crizotinib and found comparable response rates between patients with variants 1 and 3. The data from this exploratory study should be approached with caution, as the patient subgroups were small, post-progression samples have yet to be assessed, and the follow-up duration for overall survival is still ongoing.²⁵

Nevertheless, the gross structural differences among the isoforms of EML4-ALK fusions likely confer varying responses to targeted inhibition and thus, it is imperative to continue to explore the oncogenic potential of each variant. The resulting characterizations would allow patients to be stratified into well-defined risk groups and treated accordingly. To that end, identifying the TKI that is most effective for each *ALK* variant can greatly improve treatment outcomes for patients with *ALK* fusions.

Acknowledgments

Funding information: Funded in part through National Institutes of Health/National Cancer Institute Grants [P01 CA129243 and P30 CA008748]. The funder had no role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication.

Ryma Benayed has received a grant from ArcherDx

Mark G. Kris has received consulting fees from AstraZeneca, Pfizer, and Regeneron. He has received honoraria for participation in educational programs from WebMD, OncLive, Physicians Education Resources, Prime Oncology, Intellisphere, Creative Educational Concepts, Peerview, i3 Health, Paradigm Medical Communications, AXIS, Carvive Systems, AstraZeneca, and Research to Practice. Funds for travel and lodging, and food and beverage have been provided by AstraZeneca, Genentech, and Regeneron. Dr. Kris is an employee of Memorial Sloan Kettering. Memorial Sloan Kettering has received funding from The National Cancer Institute (USA), The Lung Cancer Research Foundation, Genentech/Roche, and PUMA Biotechnology for research conducted by Dr. Kris. Memorial Sloan Kettering has an institutional agreement with IBM for Watson for Oncology and receives royalties from IBM.

Alexander Drilon has received honoraria for participation in educational programs from Medscape, OncLive, PeerVoice, Physicians Education Resources, Targeted Oncology, and Research to Practice. He has received honoraria and served on advisory boards for participation in programs with Ignyta, Loxo, TP Therapeutics, AstraZeneca, Pfizer, Blueprint Medicines, Genentech/Roche, Takeda/Ariad/Millennium, Helsinn, Beigene, BergenBio, Hengrui Therapeutics, Exelixis, Bayer, Tyra Biosciences, Verastem, and MORE Health. Funds for travel and lodging, and food and beverage were received from Merck. Dr. Drilon is an employee of Memorial Sloan Kettering. Memorial Sloan Kettering has received research funding from Pfizer, Exelixis, GlaxoSmithKlein, Teva, and Taiho. He has also received research funding from Foundation Medicine. He receives royalties from Wolters Kluwer.

References

1. Watanabe S, Hayashi H, Okamoto K, et al. Progression-Free and Overall Survival of Patients With ALK Rearrangement-Positive Non-Small Cell Lung Cancer Treated Sequentially With Crizotinib and Alectinib. *Clin Lung Cancer*. 2016;17:528–534. [PubMed: 27318655]
2. Kwak EL, Bang YJ, Camidge DR, et al. Anaplastic lymphoma kinase inhibition in non-small-cell lung cancer. *N Engl J Med*. 2010;363:1693–1703. [PubMed: 20979469]
3. Schaake EE, Kappers I, Codrington HE, et al. Tumor Response and Toxicity of Neoadjuvant Erlotinib in Patients With Early-Stage Non-Small-Cell Lung Cancer. 2012;30:2731–2738.
4. Rotow JK, Woodard GA, Urisman A, et al. Pathologic Complete Response to Neoadjuvant Crizotinib in a Lung Adenocarcinoma Patient With a MET Exon 14 Skipping Mutation. *Clin Lung Cancer*. 2019;20:e137–e141. [PubMed: 30553716]
5. Jordan EJ, Kim HR, Arcila ME, et al. Prospective Comprehensive Molecular Characterization of Lung Adenocarcinomas for Efficient Patient Matching to Approved and Emerging Therapies. *Cancer Discov*. 2017;7:596–609. [PubMed: 28336552]
6. Schram AM, Chang MT, Jonsson P, Drilon A. Fusions in solid tumours: diagnostic strategies, targeted therapy, and acquired resistance. *Nat Rev Clin Oncol*. 2017;14:735–748. [PubMed: 28857077]
7. Shaw AT, Kim DW, Nakagawa K, et al. Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *N Engl J Med*. 2013;368:2385–2394. [PubMed: 23724913]
8. Solomon BJ, Mok T. First-line crizotinib in ALK-positive lung cancer. *N Engl J Med*. 2015;372:782. [PubMed: 25693021]

9. Blackhall F, Kim D-W, Besse B, et al. Patient-Reported Outcomes and Quality of Life in PROFILE 1007: A Randomized Trial of Crizotinib Compared with Chemotherapy in Previously Treated Patients with ALK-Positive Advanced Non-Small-Cell Lung Cancer. *Journal of Thoracic Oncology*. 2014;9:1625–1633. [PubMed: 25436797]
10. Richards MW, O'Regan L, Roth D, et al. Microtubule association of EML proteins and the EML4-ALK variant 3 oncoprotein require an N-terminal trimerization domain. *Biochem J*. 2015;467:529–536. [PubMed: 25740311]
11. Heuckmann JM, Balke-Want H, Malchers F, et al. Differential protein stability and ALK inhibitor sensitivity of EML4-ALK fusion variants. *Clin Cancer Res*. 2012;18:4682–4690. [PubMed: 22912387]
12. Sabir SR, Yeoh S, Jackson G, Bayliss R. EML4-ALK Variants: Biological and Molecular Properties, and the Implications for Patients. *Cancers (Basel)*. 2017;9.
13. Shaw AT, Engelman JA. ALK in lung cancer: past, present, and future. *J Clin Oncol*. 2013;31:1105–1111. [PubMed: 23401436]
14. Woo CG, Seo S, Kim SW, et al. Differential protein stability and clinical responses of EML4-ALK fusion variants to various ALK inhibitors in advanced ALK-rearranged non-small cell lung cancer. *Ann Oncol*. 2017;28:791–797. [PubMed: 28039177]
15. Cha YJ, Kim HR, Shim HS. Clinical outcomes in ALK-rearranged lung adenocarcinomas according to ALK fusion variants. *J Transl Med*. 2016;14:296. [PubMed: 27756333]
16. Lei YY, Yang JJ, Zhang XC, et al. Anaplastic Lymphoma Kinase Variants and the Percentage of ALK-Positive Tumor Cells and the Efficacy of Crizotinib in Advanced NSCLC. *Clin Lung Cancer*. 2016;17:223–231. [PubMed: 26454342]
17. Yoshida T, Oya Y, Tanaka K, et al. Differential Crizotinib Response Duration Among ALK Fusion Variants in ALK-Positive Non-Small-Cell Lung Cancer. *J Clin Oncol*. 2016;34:3383–3389. [PubMed: 27354483]
18. Christopoulos P, Endris V, Bozorgmehr F, et al. EML4-ALK fusion variant V3 is a high-risk feature conferring accelerated metastatic spread, early treatment failure and worse overall survival in ALK(+) non-small cell lung cancer. *Int J Cancer*. 2018;142:2589–2598. [PubMed: 29363116]
19. Noh K-W, Lee M-S, Lee SE, et al. Molecular breakdown: a comprehensive view of anaplastic lymphoma kinase (ALK)-rearranged non-small cell lung cancer. 2017;243:307–319.
20. Tu H-Y, Zhang Y-C, Wu Y-L. A story of ALK variants and the efficacy of ALK inhibitors: moving toward precision oncology. *Translational Cancer Research*; Vol 6, Supplement 2 (3 2017): *Translational Cancer Research*. 2017.
21. Hechtman JF, Benayed R, Hyman DM, et al. Pan-Trk Immunohistochemistry Is an Efficient and Reliable Screen for the Detection of NTRK Fusions. *Am J Surg Pathol*. 2017;41:1547–1551. [PubMed: 28719467]
22. Bayliss R, Choi J, Fennell DA, Fry AM, Richards MW. Molecular mechanisms that underpin EML4-ALK driven cancers and their response to targeted drugs. *Cell Mol Life Sci*. 2016;73:1209–1224. [PubMed: 26755435]
23. Lin JJ, Zhu VW, Yoda S, et al. Impact of EML4-ALK Variant on Resistance Mechanisms and Clinical Outcomes in ALK-Positive Lung Cancer. *J Clin Oncol*. 2018;36:1199–1206. [PubMed: 29373100]
24. Gainor JF, Dardaei L, Yoda S, et al. Molecular Mechanisms of Resistance to First- and Second-Generation ALK Inhibitors in *ALK*-Rearranged Lung Cancer. 2016;6:1118–1133.
25. Camidge DR, Dziadziuszko R, Peters S, et al. Updated Efficacy and Safety Data and Impact of the EML4-ALK Fusion Variant on the Efficacy of Alectinib in Untreated ALK-Positive Advanced Non-Small Cell Lung Cancer in the Global Phase III ALEX Study. *Journal of thoracic oncology: official publication of the International Association for the Study of Lung Cancer*. 2019;14:1233–1243.

Clinical Practice Points

- Gene fusions of anaplastic lymphoma kinase (*ALK*) are common drivers in non-small cell lung cancer that are effectively treated with crizotinib.
- Certain structural alterations may predict increased sensitivity to crizotinib in patients with non-small cell lung cancer.
- In some patients with lung cancer, crizotinib can induce complete pathological remission.

Figure 1.

Computed tomography scans of the chest demonstrating response over time. At baseline, there was a 2.0 x 1.7 cm right lower lobe tumor (arrow), a right pleural effusion, and compressive atelectasis. One month after commencing treatment with crizotinib, there was a reduction in the size of the tumor (1.9 x 0.8 cm; arrow) and the pleural effusion resolved. Fifty-seven months after starting crizotinib, regression noted from the baseline study is maintained.

a.

b.

C. **Figure 2.**

(A) Histologic sections from the pleural implant showed metastatic adenocarcinoma with signet ring cell features. Higher magnification (right) showed sheets of tumor cells with abundant intracytoplasmic mucin displacing nuclei to the periphery of the cells, forming signet ring cells. (B) Fluorescence in situ hybridization (FISH) with break apart probes detected an *ALK* rearrangement (2p23) in 52% of cells, including 22% of the cells with the standard pattern (bottom cell) and 30% of the cells with additional copies of the non-rearranged *ALK* gene (top two cells). (C) Histologic section from previous site of disease collected during autopsy showed an area of pulmonary consolidation characterized by hemorrhage and organizing pneumonia. No tumor is present.

a.

EML4 ex17 // ALK Intron19 ALK Exon20

CCTGTGGAACCTCAATGGAACACAGGCTGGAATGGACCAGGCTGGTAGATGAACCAGGACA
 CTGTGCAGATTTTCATCCAAGTGGCACAGTG GTG GCC ATA GGA ACG CAC TCA GGC
 AG//G TTT CCT GTA GGA AGT GGC CTG TGT AGT GCT TCA AAG GCC AGG CTG CCA
 GGC CAT GTT GCA GCT GAC CAC CCA CCT GCA GTG TAC CGC CGG AAG CAC CAG
 GAG CTG CAA GCC ATG CAG ATG GAG CTG

b.

Figure 3.

(A) Sequence of the E17; ins80A20 *EML4-ALK* in-frame fusion variant. Purple indicates *EML4* exon 17. Blue signifies the 80 bp insertion from intron 19. Green denotes *ALK* exon 20. (B) Schematic diagram of the in-frame fusion of exon 17 of *EML4* and exon 20 of *ALK* with an 80 bp insertion from intron 19.