Published in final edited form as:

J Am Chem Soc. 2018 November 28; 140(47): 16058–16061. doi:10.1021/jacs.8b10891.

Synthesis of Myrocin G, the Putative Active Form of the Myrocin Antitumor Antibiotics

Christos Economou[†], Martin Tomanik[†], Seth B. Herzon^{*,†,‡}

[†]Department of Chemistry, Yale University, New Haven, Connecticut 06520, United States

[‡]Department of Pharmacology, Yale School of Medicine, New Haven, Connecticut 06520, United States

Abstract

The antiproliferative antimicrobial fungal metabolites known as the myrocins have been proposed to cross-link DNA by double nucleotide addition. However, the nature of the DNA-reactive species is ambiguous, as myrocins have been isolated as functionally distinct 5-hydroxy- γ -lactone and diosphenol isomers. Based on literature precedent, we hypothesized that the diosphenol **7** (assigned here the trivial name myrocin G) is the biologically active form of the representative isolate (+)-myrocin C (**1**). To probe this, we developed a short enantioselective route to **7**. A powerful fragment-coupling reaction that forms the central ring of the target in 38% yield and in a single step was developed. In support of our hypothesis, **7** was efficiently transformed to the bis(sulfide) **6**, a product previously isolated from reactions of **1** with excess benzenethiol. This work provides the first direct access to the diosphenol **7**, sets the stage for elucidating the mode of interaction of the myrocins with DNA, and provides a foundation for the synthesis of other pimarane diterpenes.

Efforts to elucidate the mechanism of action of natural products are complicated when the metabolite can adopt two or more functionally distinct forms. This issue is exemplified by the antiproliferative antimicrobial metabolites myrocins C (**1**)¹ and B (**2**),^{2,3} fungal isolates that contain a sensitive 5-hydroxy- γ -lactone residue (Scheme 1A, blue in **1** and **2**). The literature indicates⁴ this substructure undergoes facile ring opening to the corresponding diosphenol under mildly acidic or basic conditions, raising uncertainty about its fidelity under biological conditions. Consistent with this, the diosphenol isomer of **2**, (–)-myrocin A (**3**), has been identified in fungal cultures.⁵

Following their landmark total synthesis of (±)-myrocin C (**1**),⁶ Chu-Moyer and Danishefsky disclosed that treatment of synthetic (±)-**1** with excess thiophenol and triethylamine generated the bis(sulfide) **6** (63%, Scheme 1B).⁷ The mechanism for formation of **6** was

*Corresponding Author: seth.herzon@yale.edu.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.8b10891.

X-ray structure determination for 007b-18079 (CIF) Detailed experimental procedures and characterization data for all new compounds (PDF)

The authors declare no competing financial interest.

proposed to comprise S_N2' substitution of the tertiary hydroxyl group (**1** $\rightarrow$ **4**), isomerization to the diosphenol **5**, and addition to the resulting activated cyclopropane. This reactivity led the authors to speculate that the myrocins cross-link DNA by sequential nucleotide addition reactions.⁸

The isolation of **3** suggests the existence of an analogous diosphenol isomer of **1**, “myrocin G (**7**)” (Scheme 1C). Ring opening of a lactone structurally related to **1** has been reported to occur under acidic or basic conditions.^{4b} The diosphenol isomer may be stabilized by a strong intramolecular hydrogen bond between the hydroxyl group and the adjacent carbonyl. Collectively, these data suggested to us an alternative order of events for **1** $\rightarrow$ **6** wherein ring opening of **1** to the diosphenol **7** precedes the initial alkylation.

Motivated by this analysis, we targeted **7** as the initial entry into this natural product family. In addition, we identified the diosphenol double bond as a strategic locus that could be converted retrosynthetically to the diketone **8** in a redox-neutral fashion (Scheme 1D). Further disconnection of the C9–C10 bond by a fragment-coupling reaction⁹ reveals the α,β -cyclopropylketone **9** and the unsaturated ketone **10** as two precursors of similar complexity. This synthetic strategy features the direct, stereocontrolled installation of the C9 alcohol, high modularity, and independent introduction of the peripheral C4 and C13 quaternary centers.

The coupling fragments **9** and **10** were prepared in three steps from known compounds (Scheme 2). Beginning with the Diels–Alder adduct **11**,¹⁰ Wittig olefination [potassium bis(trimethylsilyl)amide, methyl triphenylphosphonium bromide], tandem enoxysilane hydrolysis and β -carbamate elimination (aqueous hydrochloric acid), and α -dehydroiodination (iodine, pyridine)¹¹ provided the C-ring fragment **10** (22% over three steps, Scheme 2A). The A-ring fragment **9** was synthesized from the β -ketoester **12** (Scheme 2B).¹² Stereoselective Robinson annulation¹³ between **12** and acrolein diethylacetal provided the enone **13** (32%, 92% ee). α -Dehydroiodination¹¹ of **13** proceeded in 97% yield. Corey–Chaykovsky cyclopropanation¹⁴ (trimethylsulfoxonium iodide, sodium hydride) provided a 2.3:1 mixture of diastereomeric α,β -cyclopropylketones. The major (desired) diastereomer **9** was isolated in 64% yield after recrystallization.

The fragment-coupling product **14** was obtained by activation of the iodocyclopropane **9** with *iso*-propylmagnesium chloride–lithium chloride complex,¹⁵ followed by addition of the α -iodoenone **10** (92%, 8.2:1 dr, Scheme 3). The stereoselectivity in the coupling was anticipated based on the known stereoelectronic preferences for nucleophilic addition to cycloalkanones¹⁶ and consideration of nonbonded interactions in the transition state. Notably, retro-aldol reaction of **14** is prevented by the geometric constraints imposed by the cyclopropane ring.

The ring closure precursor **16** was prepared by a five-step sequence comprising Stille cross-coupling [tetrakis(triphenylphosphine)palladium(0), copper(I) iodide, cesium fluoride] with tributyl(1-ethoxyvinyl) tin, hydrolysis of the resulting vinyl ether (aqueous hydrochloric acid), tandem alcohol silylation–enoxysilane formation (trimethylsilyl trifluoromethanesulfonate, triethylamine), Rubottom oxidation (3-chloroperoxybenzoic

acid), and conversion of the primary alcohol to an allyl carbonate (allyl chloroformate, pyridine, 37% overall). The intermediate Rubottom oxidation product was desilylated and the structure of the desilylated product was confirmed by X-ray analysis.¹⁷

After much experimentation, we found that treatment of the allyl carbonate **16** with sodium *tert*-butoxide in tetrahydrofuran at 0 °C generated the diosphenol **19** (64%). Mechanistic studies suggest that **19** is formed via aldol addition (**16** → **17**), carbonate migration (**17** → **18**), and β -elimination. The silyl migration product **20** was isolated separately in 15% yield.

This cyclization cascade provides expedient access to a protected form of myrocin G (**7**). After some consideration, we recognized that the fragment-coupling–ring-closure cascade could potentially be carried out in one flask by embedding a latent enolate nucleophile in the C-ring electrophile. Toward this end, we prepared the enoxysilane **24** by the sequence shown in Scheme 4. Beginning with the α -iodoenone **10**, ketalization (ethylene glycol, triethylorthoformate, *p*-toluenesulfonic acid) followed by lithium–halogen exchange and addition of the Weinreb amide¹⁸ **21** provided the α,β -unsaturated ketone **22** (70%, two steps). Removal of the silyl ether (tetra-*n*-butylammonium fluoride), installation of the allyl carbonate (allyl chloroformate, pyridine), and removal of the acetal (aqueous hydrochloric acid) generated the β -diketone **23** (46%, three steps). Site-selective deprotonation of **23** [lithium bis(trimethylsilyl)amide] and trapping of the resulting enolate with chlorotrimethylsilane provided the target enoxysilane **24** (59%).

Attempts to effect the fragment coupling of the enoxysilane **24** with the organomagnesium reagent derived from **9** were unsuccessful. We found, however, that lithium–halogen exchange (*n*-butyllithium, –78 °C), followed by immediate addition of the enoxysilane **24** and warming to 0 °C provided the fully annulated product **19** in 38% yield (Scheme 5). In this coupling reaction, the ring closure is triggered by migration of the trimethylsilyl substituent in the addition product **26**. The modest yield of this transformation is offset to some extent by the rapid increase in molecular complexity achieved. Deprotection of **18** (tetra-*n*-butylammonium fluoride) then provided myrocin G (**7**, 64%). Subjecting synthetic **7** to the conditions disclosed by Chu-Moyer and Danishefsky⁷ generated the bis(sulfide) **6** (74%, Scheme 6). This result indicates that myrocin G (**7**) is a competent intermediate in the double nucleophilic addition of thiols and provides support for our hypothesis. We reasoned that (+)-myrocin C (**1**) itself might be accessible by temporarily disrupting the diosphenol hydrogen bond in **19** or **7**, thereby making lactonization thermodynamically favorable. However, exploratory experiments toward this end have thus far been unsuccessful.

In summary, we have developed a concise, enantioselective synthesis of myrocin G (**7**), the putative active form of the antiproliferative antimicrobial metabolite myrocin C (**1**). Key to the success of this approach was the development of a powerful annulation strategy that forges the central ring of the target in a single step from two synthetic precursors of similar complexity. With access to **7**, we are now in position to probe its antiproliferative activity and determine its biological target. The fragment-coupling reaction we have developed should be amenable to the synthesis of other pimarane diterpenes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

Financial support from the National Institutes of Health (Ruth L. Kirschstein National Research Service Award F31CA213964-03 to C.E.) and Yale University is gratefully acknowledged.

REFERENCES

- (1). (a)Hsu YH; Hirota A; Shima S; Nakagawa M; Nozaki H; Tada T; Nakayama M Structure of Myrocin C, A New Diterpene Antibiotic Produced by a Strain of *Myrothecium* sp. Agric. Biol. Chem 1987, 51, 3455.(b)Nakagawa M; Hsu YH; Hirota A; Shima S; Nakayama M; Myrocin C, A New Diterpene Antitumor Antibiotic from *Myrothecium Verrucaria*. I. Taxonomy of the Producing Strain, Fermentation, Isolation and Biological Properties. J. Antibiot 1989, 42, 218. [PubMed: 2494145] (c)Hsu YH; Hirota A; Shima S; Nakagawa M; Adachi T; Nozaki H; Nakayama M Myrocin C, A New Diterpene Antitumor Antibiotic from *Myrothecium verrucaria*. II. Physicochemical Properties and Structure Determination. J. Antibiot 1989, 42, 223. [PubMed: 2925513]
- (2). (a)Hsu YH; Nakagawa M; Hirota A; Shima S; Nakayama M Structure of Myrocin B, A New Diterpene Antibiotic Produced by *Myrothecium verrucaria*. Agric. Biol. Chem 1988, 52, 1305. (b)Lehr N-A; Meffert A; Antelo L; Sterner O; Anke H; Weber RWS Antiamoebins, myrocin B and the basis of antifungal antibiosis in the coprophilous fungus *Stilbella erythrocephala* (syn. *S. fimetaria*). FEMS Microbiol. Ecol 2006, 55, 105. [PubMed: 16420619]
- (3). For a review of pimarane diterpenes, see:Wang X; Yu H; Zhang Y; Lu X; Wang B; Liu X Bioactive Pimarane-type Diterpenes from Marine Organisms. Chem. Biodiversity 2018, 15, No. e1700276.
- (4). (a)Langschwager W; Hoffmann HMR Ring-Chain Tautomerism Provides a Route to 7a-Hydroxy-3a-methyl-2,7-dioxoperhydrobenzofuran. Synthesis of the Hydroxy γ -Lactone Substructure of Myrocin and Other Bioactive Natural Products. Liebigs Ann. 1995, 797. (b)Zander N; Langschwager W; Hoffmann HMR The Enol Lactone Approach to Protected Hydroxy γ -Lactones (5-Hydroxy-dihydro-furan-2-ones). Synth. Commun 1996, 26, 4577.
- (5). (a)Klemke C; Kehraus S; Wright AD; König GM New Secondary Metabolites from the Marine Endophytic Fungus *Apiospora montagnei*. J. Nat. Prod 2004, 67, 1058. [PubMed: 15217297] (b)Tsukada M; Fukai M; Miki K; Shiraishi T; Suzuki T; Nishio K; Sugita T; Ishino M; Kinoshita K; Takahashi K; Shiro M; Koyama K Chemical Constituents of a Marine Fungus, *Arthrinium sacchari*. J. Nat. Prod 2011, 74, 1645. [PubMed: 21718054]
- (6). (a)Chu-Moyer MY; Danishefsky SJ A Remarkable Cyclopropanation: The Total Synthesis of Myrocin C. J. Am. Chem. Soc 1992, 114, 8333.(b)Chu-Moyer MY; Danishefsky SJ; Schulte GK Total Synthesis of ($\pm$)-Myrocin C. J. Am. Chem. Soc 1994, 116, 11213.
- (7). Chu-Moyer MY; Danishefsky SJ The Mode of Action of Myrocin C: Evidence For a CC-1065 Connection. Tetrahedron Lett. 1993, 34, 3025.
- (8). For a review of natural products that damage DNA by nucleotide addition to an activated cyclopropane, see:Wolkenberg SE; Boger DL Mechanisms of in Situ Activation for DNA-Targeting Antitumor Agents. Chem. Rev 2002, 102, 2477. [PubMed: 12105933]
- (9). This disconnection drew inspiration from a related elegant fragment-coupling employed by Baran and co-workers en route to (–)-maoecrystal V. SeeCernijenko A; Risgaard R; Baran PS 11-Step Total Synthesis of (–)-Maoecrystal V. J. Am. Chem. Soc 2016, 138, 9425. [PubMed: 27457680]
- (10). Huang Y; Iwama T; Rawal VH Highly Enantioselective Diels-Alder Reactions of 1-Amino-3-siloxy-dienes Catalyzed by Cr(III)-Salen Complexes. J. Am. Chem. Soc 2000, 122, 7843.
- (11). Johnson CR; Adams JP; Braun MP; Senanayake CBW; Wovkulich PM; Uskokovic MR Direct α -Iodination of Cycloalkenones. Tetrahedron Lett. 1992, 33, 917.
- (12). (a)Ueda Y; Roberge G; Vinet V A simple method of preparing trimethylsilyl- and *tert*-butyldimethylsilyl-enol ethers of α -diazoacetoacetates and their use in the synthesis of a chiral

precursor to thienamycin analogs. *Can. J. Chem* 1984, 62, 2936.(b)Meng Z; Yu H; Li L; Tao W; Chen H; Wan M; Yang P; Edmonds DJ; Zhong J; Li A Total synthesis and antiviral activity of indolosesquiterpenoids from the xiamycin and oridamycin families. *Nat. Commun* 2015, 6, 6096. [PubMed: 25648883]

- (13). Xu C; Zhang L; Luo S Asymmetric Enamine Catalysis with β -Ketoesters by Chiral Primary Amine: Divergent Stereocontrol Modes. *J. Org. Chem* 2014, 79, 11517. [PubMed: 25370930]
- (14). Corey EJ; Chaykovsky M Dimethyloxosulfonium Methylide ((CH₃)₂SOCH₂) and Dimethylsulfonium Methylide ((CH₃)₂SCH₂). Formation and Application to Organic Synthesis. *J. Am. Chem. Soc* 1965, 87, 1353.
- (15). (a)Ren H; Krasovskiy A; Knochel P Preparation of cyclic alkenylmagnesium reagents *via* an iodine/magnesium exchange. *Chem. Commun* 2005, 543. For a review, see: (b) Klatt T; Markiewicz JT; Samann C; Knochel P Strategies to Prepare and Use Functionalized Organometallic Reagents. *J. Org. Chem* 2014, 79, 4253. [PubMed: 24697240]
- (16). Bürgi HB; Dunitz JD; Lehn JM; Wipff G Stereochemistry of reaction paths at carbonyl centres. *Tetrahedron* 1974, 30, 1563.
- (17). See the Supporting Information.
- (18). Nahm S; Weinreb SM *N*-Methoxy-*N*-Methylamides as Effective Acylating Agents. *Tetrahedron Lett.* 1981, 22, 3815.

Scheme 1.

(A) Structures of Myrocins A–C (1–3); (B) Structure of the Bis(sulfide) 6 and the Originally Proposed Mechanism for Its Formation; (C) Diosphenol 7, Which is Hypothesized to Be the Biologically-Active Form of 1; (D) Retrosynthetic Analysis of 7

Scheme 2. Synthesis of the Fragment Coupling Partners 9 and 10^a^aTMSE = 2-(trimethylsilyl)ethyl.

Scheme 3.
Synthesis of 19

Scheme 5.
One-Step Synthesis of 19 from 9 and 24

Scheme 6.
Synthesis of the Bis(sulfide) 6 from the Diosphenol 7