Published in final edited form as:

Theor Comput Sci. 2017 November 21; 701: 109–119. doi:10.1016/j.tcs.2017.05.019.

A Graph Isomorphism Condition and Equivalence of Reaction Systems*

Daniela Genova,

Department of Mathematics and Statistics, University of North Florida, Jacksonville, USA

Hendrik Jan Hoogeboom,

LIACS, Leiden University, the Netherlands

Nataša Jonoska

Department of Mathematics and Statistics, University of South Florida, Tampa, USA

Nataša Jonoska: jonoska@math.usf.edu

Abstract

We consider global dynamics of reaction systems as introduced by Ehrenfeucht and Rozenberg. The dynamics is represented by a directed graph, the so-called transition graph, and two reaction systems are considered equivalent if their corresponding transition graphs are isomorphic. We introduce the notion of a skeleton (a one-out graph) that uniquely determines a directed graph. We provide the necessary and sufficient conditions for two skeletons to define isomorphic graphs. This provides a necessary and sufficient condition for two reactions systems to be equivalent, as well as a characterization of the directed graphs that correspond to the global dynamics of reaction systems.

Keywords

directed graphs; graph isomorphism; graphs on posets; dynamics of reaction systems; equivalence of reaction systems

1. Introduction

Determining whether two graphs are isomorphic is one of the archetypical problems in graph theory and plays an important role in many applications and network analysis problems. Although there have been significant advances for this problem in the past year [2], the problem remains difficult. On the other side, often in network analysis, graphs are partitioned in so called ‘modules’ where each vertex in a module is adjacent to the same set of vertices outside the module [10]. Modules in directed graphs are defined as sets of vertices that have incoming and outgoing edges from, and to, the same vertices outside the module and it is shown that modular decompositions can be performed in linear time [12]. In this paper we study graph isomorphism through a variation of this notion, where we consider

*This work has been supported in part by the NSF grant CCF-1526485 and NIH grant R01 GM109459.

Correspondence to: Nataša Jonoska, jonoska@math.usf.edu.

vertices that have the “same” incoming edges, and we call such vertices “companions”. These vertices are precisely those that belong to the same region in the Venn diagram constructed out of the family of out-sets (an out-set for v is the set of vertices that have incoming edges starting at v). We further define a “skeleton” of a graph $G = (V, E)$ as a one-out graph over a set V such that the set of vertices that have non-zero in-degree are representatives of the family of out-sets. A skeleton defines uniquely a directed graph and we characterize skeletons of isomorphic graphs. Skeletons of isomorphic graphs are called “companion skeletons”. In particular, skeleton edges swapped at companion vertices produce companion skeletons. This observation allows characterizations of reaction systems (described below) that exhibit the same global dynamical behavior.

A formal description of biochemical interactions within a confined region bounded with a porous membrane that can interact with the environment has been introduced in [6], see [3] for an overview of the theory. This formal model, called “reaction systems”, is based on the idea that each reaction depends on the presence of a compound of enzymes, or facilitators, and the absence of any other control substance that inhibits the process. It is assumed further that the reaction is enabled only if the region contains all of the enabling ingredients and none of the inhibitors. In addition, if some ingredients are present in the system, the model allows their presence to be sufficient to enable all reactions where they participate. Formally, a reaction is modeled by a triple of sets (reactants, inhibitors, results) while the reaction system then represents a set of such triples. In each step, the system produces resulting elements according to the set of reactants that are enabled. It is further assumed that there is a universal set of elements that can enter the system from the outside environment and interact with the reactants at any given time. Several studies have addressed the question of the dynamics of the system (the step by step changes of the states of the system), such as reachability [5], convergence [9], fixed points and cycles [8, 7]. It has been observed that the complexity of deciding existence of certain dynamical properties falls within PSPACE (reachability) or NP-completeness (fixed points and fixed point attractors). In all of these studies, however, the changes in the dynamics through inclusion of new elements entering from the outside environment has not been considered. We call this condition of no outside involvement within the system as a 0-context reaction system. In this paper we study the relationship between the dynamics of the 0-context reaction systems and the global dynamics of the reaction system that depends on the environmental context. We observe that quite different dynamical properties of 0-context reaction systems produce equivalent global dynamics.

We represent the dynamics of a reaction system as a directed graph where each vertex is a state of the system represented as a set of elements present at the system at a given time. A directed edge from a vertex terminates at a vertex representing the new state of the system after all reactions enabled at the origin, with possible additions from the outside environment, are performed. In this way, the graph of the 0-context reaction system is a one-out graph (a skeleton) and is a subgraph of the graph of the full dynamics of the system. We characterize the graphs representing the global dynamics of reaction systems and show that two reaction systems are equivalent if their 0-context graphs are companion skeletons.

2. Subsets and Companions

We denote $[n] = \{0, 1, \dots, n-1\}$. The power set of a set A is denoted by 2^A . The number of elements of a finite set A is denoted by $|A|$ and is called the *size* of A . Given a function $f: X \rightarrow Y$, the natural equivalence on X defined by f is denoted with $\ker f$, i.e., $x \ker f y$ if and only if $f(x) = f(y)$. For $x \in X$ the equivalence class of $\ker f$ is denoted $[x]_f$. For a finite set V , let $\mathcal{O} \subseteq 2^V$ be a family of subsets of V . We say that $\mathcal{O}$ is a family of sets with domain V . The elements in V that appear in the same region of the Venn diagram for $\mathcal{O}$ are “companions” with respect to $\mathcal{O}$. Formally, let $\mathcal{N}_{\mathcal{O}}(x) = \{X \in \mathcal{O} \mid x \in X\}$ be the subfamily containing all sets that include x and $\mathcal{N}_{\mathcal{O}}^c(x)$ its complement in $\mathcal{O}$, the subfamily of those sets that do not contain x . We call $\mathcal{N}_{\mathcal{O}}(x)$ the *neighborhood* of x .

Definition 2.1. Let V be a finite set and $\mathcal{O} \subseteq 2^V$ be a family of subsets of V . Two elements $x, y \in V$ are companions with respect to $\mathcal{O}$ if $\mathcal{N}_{\mathcal{O}}(x) = \mathcal{N}_{\mathcal{O}}(y)$. We write $x \sim_{\mathcal{O}} y$ and denote the equivalence class of x by $C_{\mathcal{O}}(x)$. The set $C_{\mathcal{O}}(x)$ is called a companion set.

Thus, the equivalence class of every element $x \in V$, the set of companions of x relative to $\mathcal{O}$, is the intersection of all sets in $\mathcal{O}$ that include x minus the union of the remaining sets in $\mathcal{O}$, i.e. $C_{\mathcal{O}}(x) = \cap \mathcal{N}_{\mathcal{O}}(x) \setminus \cup \mathcal{N}_{\mathcal{O}}^c(x)$. The same equivalence based on neighborhoods of elements with respect to a family was also used in [4] where authors study activity regions for a set of neurons and the convexity of these regions was considered. A special case is when $\mathcal{N}_{\mathcal{O}}(x) = \emptyset$, i.e., when $x \notin \cup \mathcal{O}$, in which case it is in the outer region, $V \setminus (\cup \mathcal{O})$ denoted by $(\cup \mathcal{O})^c$, of the Venn diagram for $\mathcal{O}$. That is, by convention, $\cap \mathcal{N}_{\mathcal{O}}(x) = \cap \emptyset = V$ and $C_{\mathcal{O}}(x) = (\cup \mathcal{O})^c$.

The converse also holds. Any intersection of sets in $\mathcal{P} \subseteq \mathcal{O}$ minus the union of the remaining sets $\mathcal{P}^c = \mathcal{O} \setminus \mathcal{P}$ forms an equivalence class $C_{\mathcal{P}}$, provided it is non-empty. More precisely, any non-empty $C_{\mathcal{P}} = \cap \mathcal{P} \setminus \cup \mathcal{P}^c$ for some $\mathcal{P} \subseteq \mathcal{O}$ coincides with $C_{\mathcal{O}}(x)$ for some $x \in V$.

Assuming $x \in C_{\mathcal{P}}$, it implies $x \in X$ for every $X \in \mathcal{P}$ and $x \notin Y$ for every $Y \in \mathcal{P}^c$. Hence, $x \in \cap \mathcal{N}_{\mathcal{O}}(x) \setminus \cup \mathcal{N}_{\mathcal{O}}^c(x) = C_{\mathcal{O}}(x)$ and $C_{\mathcal{P}} \subseteq C_{\mathcal{O}}(x)$. Conversely if $y \in C_{\mathcal{O}}(x)$ then $y \in \cap \mathcal{N}_{\mathcal{O}}(x)$ which is precisely $\cap \mathcal{P}$ and $y \notin \cup \mathcal{N}_{\mathcal{O}}^c(x) = \cup \mathcal{P}^c$ for $\mathcal{P} = \mathcal{N}_{\mathcal{O}}(x)$ and hence $C_{\mathcal{O}}(x) \subseteq C_{\mathcal{P}}$.

Thus, $C_{\mathcal{P}} = C_{\mathcal{O}}(x)$.

Therefore, every equivalence class C of $\sim_{\mathcal{O}}$ is characterized by a subset $\mathcal{P} \subseteq \mathcal{O}$, its neighborhood, such that $C = \cap \mathcal{P} \setminus \cup \mathcal{P}^c$. In general, not every $\mathcal{P} \subseteq \mathcal{O}$ defines an equivalence class, i.e., $\cap \mathcal{P} \setminus \cup \mathcal{P}^c$ might be empty. This is the case when the corresponding region of the Venn diagram of $\mathcal{O}$ is empty.

For a family of sets $\mathcal{O}$ we denote with $\mathcal{O}^{\cap}$ the smallest family of sets that contains $\mathcal{O}$ and is closed under intersection. We say that $\mathcal{O}^{\cap}$ is the *intersection closure* of $\mathcal{O}$. If $\mathcal{O} = \mathcal{O}^{\cap}$ we say that $\mathcal{O}$ is *intersection closed*.

Example 2.1. Consider the finite set $V = \{1, 2, \dots, 8\}$ and the family of subsets $\mathcal{O} \subseteq 2^V$ given by $\mathcal{O} = \{\{1, 2, 3, 4\}, \{4, 5\}, \{5\}\}$. Then $\mathcal{O}^{\cap} = \mathcal{O} \cup \{\{4\}\}$. Note that $C_{\mathcal{O}}(1) = \cap \mathcal{N}_{\mathcal{O}}(1) \setminus \cup \mathcal{N}_{\mathcal{O}}^c(1) = \{1, 2, 3, 4\} \setminus (\{4, 5\} \cup \{5\}) = \{1, 2, 3\} = C_{\mathcal{O}}(2) = C_{\mathcal{O}}(3)$ and $C_{\mathcal{O}}(4) =$

{4}. Thus, the family $\mathcal{O}$ defines the following companion sets: {1, 2, 3}, {4}, {5}, and {6, 7, 8}, where each nonempty region in the corresponding Venn diagram is a companion set.

In the sections that follow we use a correspondence of families of sets, that have the same sizes of the sets as well as their intersections.

Definition 2.2. Let $\mathcal{O}_1$ and $\mathcal{O}_2$ be two families of sets over the domains V and W respectively, i.e. $\mathcal{O}_1 \subseteq 2^V$ and $\mathcal{O}_2 \subseteq 2^W$. A faithful correspondence between $\mathcal{O}_1$ and $\mathcal{O}_2$ is a bijection $\eta: \mathcal{O}_1^\cap \rightarrow \mathcal{O}_2^\cap$ that satisfies

- i. $|X| = |\eta(X)|$ for all $\mathcal{O}_1^\cap$, and
- ii. $\eta(X \cap Y) = \eta(X) \cap \eta(Y) \in \mathcal{O}_2^\cap$ for all $X, Y \in \mathcal{O}_1^\cap$

As shown below in Lemma 2.3, faithful correspondences preserve not only the size of the sets and their intersections, but also the sizes of the companion sets defined by these families. This is only true if the sizes of the domains are equal since otherwise the outer regions differ in size: $|(U\mathcal{O}_1)^c| \neq |(U\mathcal{O}_2)^c|$.

Lemma 2.3. Let η be a faithful correspondence between $\mathcal{O}_1 \subseteq 2^V$ and $\mathcal{O}_2 \subseteq 2^W$, with $|V| = |W|$. Then, there exists a bijection φ from V to W whose extension to 2^V , restricted to $\mathcal{O}_1$, coincides with η . Moreover, the extension bijectively maps companion sets of $\mathcal{O}_1$ into companion sets of $\mathcal{O}_2$, respecting size.

Proof. Observe that the equivalence classes of $\sim_{\mathcal{O}_1}$, which are companion sets, are precisely the *non-empty* sets $C\mathcal{P} = \cap \mathcal{P} \setminus \cup \mathcal{P}^c$ for some family $\mathcal{P} \subseteq \mathcal{O}_1$, where $\mathcal{P}^c = \mathcal{O}_1 \setminus \mathcal{P}$. Similarly, this holds for $\mathcal{O}_2$. We show that there is a bijection between the companion sets of $\mathcal{O}_1$ and $\mathcal{O}_2$ that preserves their sizes. Once this is established, the required bijection φ that respects the equivalences can naturally be constructed.

We find a correspondence between the companion sets of $\mathcal{O}_1$ and $\mathcal{O}_2$ and define $C_{\mathcal{N}(\mathcal{P})} = \cap \eta(\mathcal{P}) \setminus \cap \eta(\mathcal{P}^c)$, where $\eta(\mathcal{P}) = \{\eta(X) \mid X \in \mathcal{P}\} \subseteq \mathcal{O}_2$ and $\eta(\mathcal{P}^c) = \{\eta(X) \mid X \in \mathcal{P}^c\} = \eta(\mathcal{P})^c$, the latter equality holds because η is a bijection. When $\mathcal{P}$ ranges over the subsets of $\mathcal{O}_1$, then non-empty $C\mathcal{P}$ range over the equivalence classes of $\mathcal{O}_1$ while non-empty $C_{\eta(\mathcal{P})}$ range over the equivalence classes of $\mathcal{O}_2$.

We argue that $|C\mathcal{P}| = |C_{\eta(\mathcal{P})}|$ – this then takes care of the cases where $\mathcal{P}$ defines the empty set instead of an equivalence class, as the corresponding image under η is void too. First observe that $|\cap \mathcal{P} \setminus \cup \mathcal{P}^c| = |\cap \mathcal{P}| - |\cap \mathcal{P} \cap \cup \mathcal{P}^c|$.

The equality $|\cap \mathcal{P}| = |\cap \eta(\mathcal{P})|$ follows from the first and the second requirement in Definition 2.2, except for the special case when $\mathcal{P} = \emptyset$, but then $\cap \mathcal{P} = V$ and consequently, $\eta(\mathcal{P}) = \emptyset$ and $\cap \eta(\mathcal{P}) = W$. The inclusion-exclusion principle states that we can express the size of a union of sets as sums of sizes of intersections, $|\cup_{i=1}^m Y_i| = \sum_{\emptyset \subset I \subseteq [m]} (-1)^{|I|+1} |\cap_{i \in I} Y_i|$. We can apply this to the sets $\cap \mathcal{P} \cap Z$, $Z \in \mathcal{P}^c$, to obtain $|\cap \mathcal{P} \cap \cup \mathcal{P}^c| = |\cap \eta(\mathcal{P}) \cap \cup \eta(\mathcal{P}^c)|$. Now that the corresponding companion sets in $\mathcal{O}_1$ and $\mathcal{O}_2$ have the same number of elements, we can construct a bijection φ from V to W that respects companions.

3. Graphs and Companion Skeletons

In this paper all graphs are directed, in the sense that a *graph* G is a pair of sets (V, E) where V is a finite set whose elements are called *vertices* and $E \subseteq V \times V$ is the set of *edges*. We also write $V(G)$ (resp. $E(G)$) to denote the set of vertices (resp. edges) of G . For an edge $e = (v, v')$ we say that v is the *initial* vertex of e and v' is the *terminal* vertex of e . For a vertex $v \in V$ we define $\text{in}_G(v)$ to be the set of all vertices that are initial for edges whose terminal vertex is v , i.e., $\text{in}_G(v) = \{w \mid (w, v) \in E\}$. Similarly, the *out-set* $\text{out}_G(v)$ is the set of all vertices that are terminal to all edges whose initial vertex is v , that is, $\text{out}_G(v) = \{w \mid (v, w) \in E\}$. If $|\text{out}_G(v)| = 1$ for all vertices v , then we say that G is a *one-out* graph. The *out-family* is defined:

$$\mathcal{O}(G) = \{\text{out}_G(v) \mid v \in V\}$$

We drop the subscript G in $\text{in}_G(v)$ and $\text{out}_G(v)$ whenever the graph is understood from the context.

The following observation is the main motivation for considering the notion of companions.

Lemma 3.1. Let G be a graph and let $\mathcal{O} = \mathcal{O}(G)$. Then $\text{in}_G(x) = \text{in}_G(y)$ if and only if $x \sim_{\mathcal{O}} y$.

Proof. Observe that (w, x) is an edge in G if and only if $x \in \text{out}_G(w)$. Thus two nodes x and y have the same incoming edges if and only if they are in the same out-sets in $\mathcal{O}$, thus if and only if they are companions with respect to $\mathcal{O}$, i.e., $x \sim_{\mathcal{O}} y$.

In the case when $\mathcal{O}$ is $\mathcal{O}(G)$ for a graph G , the number of elements in $\mathcal{O}$ cannot exceed the number of vertices $V = V(G)$. Therefore, in this case we can always assume that $\mathcal{O}$ is indexed by a subset $R_{\mathcal{O}} \subseteq V$, i.e., $\mathcal{O} = \{O_z \mid z \in R_{\mathcal{O}}\}$. Given this representation, for $z \in R_{\mathcal{O}}$ the element z is called the *representative* of $O_z \in \mathcal{O}$ and $R_{\mathcal{O}}$ is the *set of representatives* for $\mathcal{O}$. We assume that representatives are unique, i.e., $O_z = O_{z'}$, if and only if $z = z'$.

Formally, representatives are fixed as a bijection $\rho: R_{\mathcal{O}} \rightarrow \mathcal{O}$. Clearly, the choice of a representative for a set O in $\mathcal{O}$ depends on the function ρ , but changing the map ρ is the same as renaming the sets in $\mathcal{O}$.

We use representatives later in the paper where they are used as convenient names for the out-sets in a structured context, see Section 4.

Convention. In order to ease our notation, if $\mathcal{O}$ is understood, we drop the subscript in $R_{\mathcal{O}}$ and simply denote the set of representatives for $\mathcal{O}$ by R . Also, in this context, we always assume that R is a subset of V and R is an index set for the family $\mathcal{O}$ such that each $z \in R$ uniquely determines a set O_z in $\mathcal{O}$.

Let $G = (V, E)$ be a graph without isolated vertices. Let $\mathcal{O} = \mathcal{O}(G) = \{O_z \mid z \in R\}$ for a set of representatives $R \subseteq V$. Define $f: V \rightarrow R$ such that $f(x) = z$ if and only if $\text{out}_G(x) = O_z$. Consider a one-out graph $G_f = (V, E_f)$ where $E_f = \{(x, f(x)) \mid x \in V\}$. Then by the choice of

$f(x)$ as the representative of $\text{out}_G(x)$ we have that $(x, z), (y, z) \in E_f$ if and only if $\text{out}_G(x) = \text{out}_G(y)$. This induces the following definition.

In the rest of this section we assume that V is fixed and finite.

Definition 3.2. A triple $\sigma = (\mathcal{O}, R, f)$ is called a skeleton over set V if $\mathcal{O} = \{O_z \mid z \in R\}$ is a family of subsets of V indexed by R with $|\mathcal{O}| = |V|$, and $f: V \rightarrow R$ is a surjection.

The graph defined by the skeleton σ is the graph $G_\sigma = (V, E)$ where $E = \{(x, w) \mid x \in V, w \in O_{f(x)}\}$.

The graph G_σ is uniquely determined by the skeleton σ . Directly from the definition we have that $\mathcal{O}(G_\sigma) = \mathcal{O}$ and $\text{out}_{G_\sigma}(x) = O_{f(x)}$. Moreover, for every graph G there is a skeleton σ such that $G = G_\sigma$. It is sufficient to take $\sigma = (\mathcal{O}(G), R, f)$ where R is a set of representatives of $\mathcal{O}(G)$ and $f(x) = r \in R$ if and only if $\text{out}_G(x) = O_r$.

Remark 3.3. Note that when every representative $z \in R$ is such that $z \in O_z$, i.e., the representative of every set in $\mathcal{O}$ is inside that set, then the one-out graph $G_f = (V, E_f)$ where $E_f = \{(x, f(x)) \mid x \in V\}$ defined by a skeleton $\sigma = (V, \mathcal{O}, f)$ is isomorphic to a subgraph of G_σ . Being an element of the set seems as a natural requirement for the representatives, but unfortunately this is not always possible. For instance, the sets $\{1, 2\}$, $\{2, 3\}$, $\{1, 3\}$ and $\{1\}$ have three elements altogether, so they cannot each contain their own representative. In general the possibility of choosing distinct representatives within a collection of sets is governed by Hall's Theorem [11]. In cases where the outsets of graph G do not allow all representatives to be chosen within the sets, the one-out graph defined by a skeleton for G cannot be a subgraph of G .

Example 3.1. Consider the collection $\{X = \{1, 2, 3\}, Y = \{2, 3, 4\}\}$ over $V = \{1, 2, \dots, 5\}$. Suppose G is a graph where $\mathcal{O}(G)$ is such that $\text{out}(1) = \text{out}(3) = X$ and $\text{out}(2) = \text{out}(4) = \text{out}(5) = Y$. We consider three skeletons $\nu = (\mathcal{O}, R_\nu, f)$, $\sigma = (\mathcal{O}', R_\sigma, g)$ and $\tau = (\mathcal{O}'', R_\tau, h)$ where $\mathcal{O} = \mathcal{O}' = \mathcal{O}''$ and the representatives are defined as follows.

$$\begin{aligned} R_\nu &= \{2, 3\} \text{ with } X = O_2 \text{ and } Y = O_3, \\ R_\sigma &= \{1, 4\} \text{ with } X = O'_1 \text{ and } Y = O'_4, \\ R_\tau &= \{4, 5\} \text{ with } X = O''_4 \text{ and } Y = O''_5. \end{aligned}$$

The three one-out graphs G_f , G_g and G_h are depicted in Fig. 1 (top). All three skeletons define the same graph G whose edges are $\{1, 3\} \times \{1, 2, 3\} \cup \{2, 4, 5\} \times \{2, 3, 4\}$, i.e., $G = G_\nu = G_\sigma = G_\tau$. Because both R_ν and R_σ have the property that the representative of X is in X and the representative of Y is in Y , the one-out graphs G_f and G_g are subgraphs of G (see Fig. 1). However, R_τ doesn't have that property, and G_h is not a subgraph of G (the vertex 5 is not in any out-set of G).

Given two skeletons $\sigma = (\mathcal{O}, R, f)$, and $\tau = (\mathcal{P}, Q, g)$, we are interested under which conditions their graphs G_σ and G_τ are isomorphic. We observe that, structurally, the one-out

graphs G_f and G_g may be quite different (as seen in Example 3.1) and yet, G_σ and G_τ may be isomorphic. We utilize the following definition.

Definition 3.4. Two skeletons $\sigma = (\mathcal{O}, R, f)$ over set V , and $\tau = (\mathcal{P}, Q, g)$ over set W are called skeleton companions, or just companions, if there is a bijection $\eta: V \rightarrow W$ that extends to a faithful correspondence $\eta: \mathcal{O}^\cap \rightarrow \mathcal{P}^\cap$ such that $\eta(O_{f(x)}) = P_{g(\eta(x))}$.

Note that because η is a bijection and extends to a faithful correspondence, the relationship “skeleton companions” is an equivalence relation on skeletons. First we see that any pair of skeletons for the same graph are companions.

Lemma 3.5. Any two skeletons $\sigma = (\mathcal{O}, R, f)$ and $\tau = (\mathcal{P}, Q, g)$ over V such that $G_\sigma = G_\tau$ are companions.

Proof. By definition of G_σ we have that $\text{out}_{G_\sigma}(x) = O_{f(x)}$, and $\mathcal{O}(G) = \mathcal{O} = \mathcal{P}$. Hence, $\text{out}_G(x) = O_{f(x)} = P_{g(x)}$. Thus the identity map on V extends to $\eta: \mathcal{O} \rightarrow \mathcal{P}$ mapping $O_{f(x)}$ to $P_{g(x)}$ for all $x \in V$ and it is a faithful correspondence satisfying $\eta(O_{f(x)}) = O_{f(x)} = P_{g(x)} = P_{g(\eta(x))}$.

The correspondence η between two companion skeletons in Definition 3.4 can take care of internal symmetry within the set V . A case of such a symmetry appears when η interchanges two companion vertices, which can be reflected as a swap of the outgoing edges in the corresponding skeletons. Let $\sigma = (\mathcal{O}, R, f)$ be a skeleton. Consider two companions x and y with respect to $\mathcal{O}$ in V along with the edges $(x, f(x))$ and $(y, f(y))$ in G_f . Consider the function $f_{x,y}: V \rightarrow V$ such that $f_{x,y}(x) = f(y)$, $f_{x,y}(y) = f(x)$ and $f_{x,y}(z) = f(z)$ for all $z \neq x, y$, that is, $f_{x,y}$ is equal to f , except that in $f_{x,y}$ the images of x and y are swapped. Then $\sigma_{x,y} = (\mathcal{O}, R, f_{x,y})$ is a skeleton companion to σ through the bijection $\eta: V \rightarrow V$ where $\eta(x) = y$, $\eta(y) = x$ and $\eta(z) = z$ for $z \neq x, y$. Note that since x and y are companions, by Lemma 3.1, these nodes have the same incoming edges in G_σ and hence also in $G_{\sigma_{x,y}}$. Swapping the out-edges of x and y means that the outgoing edges of x and y are interchanged in $G_{\sigma_{x,y}}$ compared to G_σ , and hence the two graphs are isomorphic (via the bijection that interchanges vertices x and y). We call $G_{\sigma_{x,y}}$ a result of G_σ through *companion edge swapping*.

Example 3.2. Consider the skeleton $\nu = (\mathcal{O}, R_\nu, f)$ from Example 3.1; its one-out graph is G_f depicted in Fig. 1 (top-left), and repeated in Fig. 2. The only pair of companions defined by the out-sets $O_2 = \{1, 2, 3\}$ and $O_3 = \{2, 3, 4\}$ is 2, 3. That means the only possibility for companion edge swapping is to exchange the outgoing edges of 2, 3 in f . As 2, 3 happen to be mapped to 3, 2 respectively, edge swapping yields the one-out graph $G_{f_{2,3}}$ shown in Fig. 2. The skeleton $\nu_{2,3} = (\mathcal{O}, R_\nu, f_{2,3})$ differs in the one-out graph, but has the same set of representatives for the out-sets. The graphs G_ν and $G_{\nu_{2,3}}$ (see Fig. 1 (bottom) and Fig. 2 (right)) are isomorphic.

Companion skeletons indeed capture isomorphic graphs. Companion edge swapping is a special case of the following general result.

Theorem 3.6. Two graphs $G = (V, E)$ and $G' = (V', E')$ are isomorphic if and only if there are companion skeletons $\sigma = (\mathcal{O}, R, f)$ over set V and $\tau = (\mathcal{P}, Q, g)$ over set V' such that $G = G_\sigma$ and $G' = G_\tau$.

Proof. Let $\Psi : V \rightarrow V'$ be a graph isomorphism. In a natural way we also have a faithful correspondence, $\Psi : \mathcal{O}(G)^\cap \rightarrow \mathcal{O}(G')^\cap$ that is an extension of Ψ , i.e., $\Psi(\text{out}_G(x)) = \text{out}_{G'}(\Psi(x))$. We write $\mathcal{O} = \mathcal{O}(G)$ and $\mathcal{P} = \mathcal{O}(G')$. Let $R \subseteq V$ be a set of representatives for $\mathcal{O} = \mathcal{O}(G)$ and define $f : V \rightarrow R$ such that $f(x) = z$ iff $\text{out}_G(x) = O_z$. Then $\sigma = (\mathcal{O}, R, f)$ is a skeleton and $G = G_\sigma$. Similarly, let $Q \subseteq V'$ be any set of representatives for $\mathcal{P}$ and $g : V' \rightarrow Q$ such that $g(x') = z'$ if and only if $\text{out}_{G'}(x') = P_{z'}$. Then $\tau = (\mathcal{P}, Q, g)$ is a skeleton and $G' = G_\tau$ and, since Ψ is an isomorphism, $\Psi(O_{f(x)}) = \Psi(\text{out}_G(x)) = \text{out}_{G'}(\Psi(x)) = P_{g(\Psi(x))}$, we have that τ and σ are companion skeletons.

Conversely, suppose $\sigma = (\mathcal{O}, R, f)$ and $\tau = (\mathcal{P}, Q, g)$ are companion skeletons and let $G = G_\sigma$ and $G' = G_\tau$. Then, by Definition 3.4 there is a bijection $\eta : V \rightarrow V'$ extending to a faithful correspondence $\eta : \mathcal{O}^\cap \rightarrow \mathcal{P}^\cap$ such that $\eta(O_{f(x)}) = P_{g(\eta(x))}$. We claim that η is an isomorphism from G to G' . Since $\eta : V \rightarrow V'$ is a bijection, we only need to show that $(x, y) \in E$ if and only if $(\eta(x), \eta(y)) \in E'$. Note that $(x, y) \in E$ if and only if $y \in \text{out}_G(x) = O_{f(x)}$ if and only if $\eta(y) \in \eta(\text{out}_G(x)) = \eta(O_{f(x)}) = P_{g(\eta(x))} = \text{out}_{G'}(\eta(x))$.

The proof above is somewhat high-level and does not provide insight into the construction of an isomorphism for a given pair of companion skeletons. In the following we show how different isomorphisms can be generated.

Let $\sigma = (\mathcal{O}, R, f)$ and $\tau = (\mathcal{P}, Q, g)$ be two companion skeletons over V , and $\eta : \mathcal{O}^\cap \rightarrow \mathcal{P}^\cap$ the corresponding faithful correspondence as in Definition 3.4. We show how η generates an isomorphism $\Psi : G \rightarrow G'$.

Let C be a set of companions corresponding to $\mathcal{O}$, i.e., C is an equivalence class of $\sim_{\mathcal{O}}$. Then by Lemma 2.3, $\eta(C)$ is a set of companions corresponding to $\mathcal{P}$. We have the following observation.

Lemma 3.7. For every $x \in V$, $|\llbracket x \rrbracket_f \cap C| = |\llbracket \eta(x) \rrbracket_g \cap \eta(C)|$

Proof. We show that $\eta(X) = Y$ for $X = \llbracket x \rrbracket_f \cap C$ and $Y = \llbracket \eta(x) \rrbracket_g \cap \eta(C)$. If $c \in X$ then $f(c) = f(x)$ and $c \in C$, so $O_{f(c)} = O_{f(x)}$ and therefore $\eta(O_{f(c)}) = O_{g(\eta(c))} = \eta(O_{f(x)}) = O_{g(\eta(x))}$ and so $g(\eta(c)) = g(\eta(x))$ implying $\eta(c) \in \llbracket \eta(x) \rrbracket_g$. So, $\eta(c) \in Y$. Due to the symmetry of the argument (working with η^{-1} instead of η), $\eta(X) = Y$, and because η is a bijection on V , $|X| = |Y|$.

Fig. 3 illustrates a case where $\mathcal{O} = \mathcal{P}$ and $R = Q$ but f and g are distinct. In this case η must permute the elements within a given companion set C with respect to the family $\mathcal{O}$, and could be the identity on the vertices outside of C . Indeed, by Lemma 3.7, for every companion set C , the number of elements in C that map with f to element y is the same with the number of elements in C that map to y with g . Fig. 3 shows an example of f and g on a companion set C . Because $O_{y1} = P_{y1}$, and $O_{f(x1)} = \eta(O_{f(x1)}) = P_{g(\eta(x1))}$, it must be that $\eta(x1) = x2$ or $\eta(x1) =$

x_3 . Say it is x_2 . Then $O_{y1} = \eta(O_{y1}) = \eta(O_{\eta(x1)}) = O_{g(\eta(x1))} = O_{g(x2)} = O_{y1}$. In this case G_σ and G_τ are isomorphic and G_g is obtained from G_f by multiple edge swapping.

We show a general construction of isomorphism Ψ .

Fix $x \in V$. By Lemma 3.7 for every companion set C with respect to θ we have that $|[x]_f \cap C| = |[\eta(x)]_g \cap \eta(C)|$. Let $C = C_\theta(x)$ be the companion set that contains x , and so $\eta(C)$ is a companion set with respect to $\mathcal{P}$ that contains $\eta(x)$. We denote with $X_{x,C} = [x]_f \cap C$ and $Y_{x,C} = [\eta(x)]_g \cap \eta(C)$. Therefore, there is a bijection $\Psi_{x,C}: X_{x,C} \rightarrow Y_{x,C}$. Observe that the companion classes with respect to θ (and similarly $\mathcal{P}$) form a partition of V (also V') and so the sets $X_{x,C}$ (resp. $Y_{x,C}$) form a partition on V (resp. V'). We can extend the bijections $\Psi_{x,C}$ to the whole set V : define $\Psi: V \rightarrow V'$ such that $\Psi|_{X_{x,C}} = \Psi_{x,C}$ for all $x \in V$ and companion sets C with respect to θ . Since Ψ is an extension of bijections of a partition of V , Ψ is well defined and a bijection itself. Observe that by definition of Ψ , because $Y_{x,C} \subseteq \eta(C)$, for each $x \in V$, $\Psi(x)$ and $\eta(x)$ belong to the same companion set with respect to $\mathcal{P}$.

Isomorphic graphs have companion skeletons, regardless which skeleton we choose.

Corollary 3.8. Let $G = (V, E)$ and $G' = (V', E')$ be isomorphic graphs. If $\sigma = (\theta, R, f)$ and $\tau = (\mathcal{P}, Q, g)$ are skeletons over set V and V' , respectively, such that $G = G_\sigma$ and $G' = G_\tau$ then σ and τ are companions.

Proof. As G and G' are isomorphic by Theorem 3.6 there exist companion skeletons for G and G' . By Lemma 3.5 these are companions to σ and τ respectively. The result follows by transitivity of companionship of skeletons.

4. Partially ordered sets

In this section we consider partially ordered sets in preparation for our discussion on reaction systems in the next section. The skeletons and their corresponding graphs have a special structure. The nodes are elements from a partially ordered set. Additionally we assume that out-sets of the graphs are cones in that partial order. This has a convenient benefit that the minimal element of a cone can be taken as a natural representative of that cone.

Let $(P, \leq)$ be a partially ordered finite set (poset). For $x \in P$ we define $\text{Up}(x) = \{y \in P \mid x \leq y\}$ to be the *upper cone* of x or simply just the *cone* of x . Let $R \subseteq P$. We define a set of cones *based at* R to be $\text{Up}[R] = \{\text{Up}(x) \mid x \in R\}$. The poset $\mathcal{P}$ is called an *upper semi-lattice* if every subset of P has a least upper bound (which must be unique). Observe that in the case when P is an upper semi-lattice, an intersection of two cones of elements x and y in P is a cone of the least upper bound z of $\{x, y\}$ i.e., the smallest z such that $x \leq z$ and $y \leq z$. Therefore, the set of cones of an upper semi-lattice is closed under intersection.

Convention. All posets P considered here are upper semi-lattices.

Consider $\theta_R = \text{Up}[R]$ for some $R \subseteq P$, a family of sets consisting of cones based at R . The set of companions of $x \subseteq R$ with respect to θ_R becomes

$$C_{\mathcal{O}_R}(x) = C_R(x) = \text{Up}(x) \setminus \left(\bigcup_{\substack{y \in R \\ y \not\leq x}} \text{Up}(y) \right). \text{ If } x \in R, x \text{ itself is the smallest element of its}$$

companion equivalence class, and it is called the *main representative* of $C_{\mathcal{O}_R}(x)$. Observe that for each $x \in R$, $C_{\mathcal{O}_R}(x) \cap R = \{x\}$, i.e., no two elements of R are companions, and R is a set of (main) representatives for $\text{Up}[R]$.

In particular, when P is the poset $(2^S, \subseteq)$ for some finite set S and $X \in R \subseteq 2^S$, the set of

$$\text{companions of } X \text{ with respect to } \mathcal{O}_R = \text{Up}[R] \text{ equals } C_{\mathcal{O}_R}(X) = \text{Up}(X) \setminus \left(\bigcup_{\substack{Y \in R \\ Y \not\subseteq X}} \text{Up}(Y) \right).$$

Example 4.1. We consider the poset $(2^S, \subseteq)$ for $S = \{a, b, c\}$. Let $R = \{\{a\}, \{b, c\}\}$, and let $\mathcal{O}_R = \text{Up}[R]$. Note $\mathcal{O}_R^\cap = \mathcal{O}_R \cup \{\{a, b, c\}\}$. Let $X = \{a\}$. Then $C_R(X) = \{\{a\}, \{a, b\}, \{a, c\}\}$ as illustrated in Fig. 4 (left). Note that $\{a, b, c\}$ is not a companion for X because it contains a subset $\{b, c\}$ which is in R . The element $\{a, b, c\}$ is in a companion class by itself. Note that also $\{b, c\}$ has no companions except itself, and $\emptyset$ is a companion to both $\{b\}$ and $\{c\}$.

In the remainder of the paper we consider graphs where the vertices are the elements of a poset. Let $(P, \leq)$ be a poset and $f: P \rightarrow P$ a function on P . Then $G_f = (P, f)$ is the usual one-out graph associated with f . As we have observed, cones in the partial order have a natural representative, so a family of cones has a natural set of representatives. The *main skeleton* associated with G_f equals $\sigma = (\text{Up}[R], R, f)$, with $f(P) = R$. It defines the graph G_σ with edges $\{(x, y) \mid x, y \in P, f(x) \leq y\}$. Note that the element x is a member of the set $\text{Up}(x)$ it represents, and hence G_f is a subgraph of G_σ .

Conversely, if G is a graph with nodes P , such that its out-sets are cones in poset $(P, \leq)$, then G has a unique main skeleton σ such that $G_\sigma = G$.

Thus in the case when out-sets are cones, the main skeleton is fixed by the function $f: P \rightarrow P$. This has a consequence for the notion of companions. We say that two functions $f, g: P \rightarrow P$ are *companions* if the main skeletons they define are companions, i.e., if there is a bijection η on P such that $\eta(\text{Up}(f(z))) = \text{Up}(g(\eta(z)))$ for $z \in P$. Identifying a function f with its one-out graph G_f we call G_f and G_g *companions* when f and g are.

Summarizing, graphs that have their nodes from a partial order, such that the out-sets all are cones in the partial order have an efficient ‘summary’ where each node is mapped to the minimal element of its out-set. Such a skeleton is unique, given the graph.

We can characterize graphs that are isomorphic to graphs that are defined by (main) skeletons G_α . It suffices to consider the structure of the family of out-sets.

Lemma 4.1. Let P be a poset. A graph G is isomorphic to a graph G_α , where α is a main skeleton over P if and only if $|V(G)| = |P|$ and there exists a faithful correspondence between $\mathcal{O}(G)$ and $\text{Up}[R]$ for some $R \subseteq P$.

Proof. Obviously the out-sets of the graph G_α consist of upper cones of P , hence the forward implication. Conversely, assume G a graph over $V = V(G)$ with $|P|$ nodes and such that there exists a faithful correspondence η between $\mathcal{O}(G)$ and $\text{Up}[R]$ for some $R \subseteq P$. By Lemma 2.3 we can find a bijection φ between $V(G)$ and P whose extension coincides with η .

Let $\sigma = (\mathcal{O}(G), Q, f)$ be a skeleton for G . We define a main skeleton $\alpha = (\text{Up}[R], R, g)$ over P which is a companion to σ ; from this the result follows by the characterization in Theorem 3.6.

For each $x \in V$ the out-set $\text{out}_G(x) = \mathcal{O}_{f(x)}$ of x in G corresponds via η , and φ , to a cone $\text{Up}(z)$ for some $z \in P$. Now define $g(\varphi(x))$ to be z . Then σ and α are companions by construction.

We illustrate the notions in the proof above by an example.

Example 4.2. Let graph G with node set $V = \{1, 2, \dots, 8\}$ be given by the following adjacency matrix.

	1	2	3	4	5	6	7	8
1				X	X			
2	X	X	X	X				
3	X	X	X	X	X	X	X	X
4				X	X			
5	X	X	X	X				
6	X	X	X	X	X	X	X	X
7				X	X			
8	X	X	X	X				

Then $\mathcal{O}(G) = \{V, \{1, 2, 3, 4\}, \{4, 5\}\}$, and $\mathcal{O}^\cap(G) = \mathcal{O}(G) \cup \{\{4\}\}$ (see Fig. 4 (right)). Now $\mathcal{O}(G)$ defines the following companion sets: $\{1, 2, 3\}$, $\{4\}$, $\{5\}$, and $\{6, 7, 8\}$.

One can obtain a faithful correspondence η between $\mathcal{O}(G)$ and a family of upper cones R within $P = (2^{\{a,b,c\}}, \subseteq)$. Take $R = \{\emptyset, \{a\}, \{b, c\}\}$. Then the intersection closure of its upsets $\text{Up}[R]^\cap = \{\text{Up}(\emptyset), \text{Up}(\{a\}), \text{Up}(\{b, c\}), \text{Up}(\{a, b, c\})\}$ contains sets of size 8, 4, 2 and 1, respectively matching those in $\mathcal{O}^\cap(G)$.

Sets in $\mathcal{O}(G)$ and $\text{Up}[R]$ are illustrated in Fig. 4. Let $\varphi: V \rightarrow 2^{\{a,b,c\}}$ be the bijection as shown in the table below, extended to sets it matches η . The notation follows the one from the proof of Lemma 4.1; g is the function of the main skeleton.

x	1	2	3	4	5	6	7	8
$\varphi(x)$	$\{a\}$	$\{a, b\}$	$\{a, c\}$	$\{a, b, c\}$	$\{b, c\}$	$\emptyset$	$\{b\}$	$\{c\}$
$g(\varphi(x)) = z$	$\{b, c\}$	$\{a\}$	$\emptyset$	$\{b, c\}$	$\{a\}$	$\emptyset$	$\{b, c\}$	$\{a\}$

The bottom rows of the above table define the (one-out function g of the) main skeleton on $P = (2^{\{a,b,c\}}, \subseteq)$ that defines a graph isomorphic to G . In this table we can swap the elements

$\varphi(x)$ within each companion set, and obtain the main skeleton for a different but an isomorphic graph.

5. Reaction Systems

We start by recalling some basic notions of reaction systems [6]. A reaction is formalized as a triple that represents the *reactant*, *inhibitor* and *product* of the reaction. Whenever all reactants and none of the inhibitors are present, the reaction will yield the product. The effect of separate reactions is cumulative, the union of the products for applicable reactions. More precisely, we have the following.

Definition 5.1. A reaction system (RS) is a pair $\mathcal{A} = (S, A)$ where S is a finite set, the background set, and $A \subseteq (2^S \setminus \{\emptyset\}) \times (2^S \setminus \{\emptyset\}) \times 2^S$ is a set of reactions in S .

Let $X \subseteq S$. For a reaction $a = (R, I, P)$ we say that a is enabled in X if $R \subseteq X$ and $I \cap X = \emptyset$. The result of a on X , denoted $\text{res}_a(X)$, equals P if a is enabled in X , and $\emptyset$ otherwise. The result of X in $\mathcal{A}$ equals $\text{res}_{\mathcal{A}}(X) = \bigcup_{a \in A} \text{res}_a(X)$.

Note that it is required that the reactant and the inhibitor sets are non-empty. This technical assumption has the consequence that no reaction is enabled in either $\emptyset$ or S , thus $\text{res}_{\mathcal{A}}(\emptyset) = \text{res}_{\mathcal{A}}(S) = \emptyset$.

Given a reaction system $\mathcal{A} = (S, A)$ define $\text{RES}_{\mathcal{A}} \subseteq 2^S$ to be the set of all $Y \subseteq S$ such that there is an X with $\text{res}_{\mathcal{A}}(X) = Y$. Thus $\text{res}_{\mathcal{A}} : 2^S \rightarrow \text{RES}_{\mathcal{A}}$ is a surjection. Note that $\emptyset \in \text{RES}_{\mathcal{A}}$.

Example 5.1. We use the reaction system $\mathcal{A} = (S, A)$ from [3, Example 7]. It has background set $S = \{1, 2, 3, 4\}$ and the following six reactions belong to A :

$a_1 = (\{1\}, \{3\}, \{2\})$, $a_2 = (\{2\}, \{1\}, \{1\})$, $a_3 = (\{2\}, \{3\}, \{3\})$, $a_4 = (\{3\}, \{1, 2\}, \{1, 2, 4\})$, $a_5 = (\{4\}, \{3\}, \{1, 2\})$, and $a_6 = (\{1, 3\}, \{2, 4\}, \{2, 3\})$.

In $\{2, 3, 4\}$ only a_2 is enabled, so we have $\text{res}_{\mathcal{A}}(\{2, 3, 4\}) = \{1\}$. In $\{1, 2, 3\}$ no reactions are enabled, so $\text{res}_{\mathcal{A}}(\{1, 2, 3\}) = \emptyset$. In $\{1, 2, 4\}$ all three a_1 , a_3 and a_5 are enabled, so $\text{res}_{\mathcal{A}}(\{1, 2, 4\}) = \{1, 2, 3\}$.

The dynamic behaviour of a reaction system is given by the notion of state sequence of an interactive process. Let S be the background set. Then a *state sequence* is of the form $W_0, W_1, \dots, W_n$ where $W_i \subseteq S$, is such that $\text{res}(W_i) \subseteq W_{i+1}$ for $0 \leq i < n$. The intuition behind this computational process is as follows. In each step the new products are generated by the enabled reactions. Elements that are not produced by the reactions vanish, the so-called principle of *non-permanency*. On the other hand, in each step the context, the environment in which the reactions take place, may add new elements in the state of the system. Hence the new state of the system is a step from W_i to any superset of $\text{res}(W_i)$.

From this perspective we introduce two graphs to represent the stepwise behavior of reaction systems, without and with context.

Definition 5.2. For a RS $\mathcal{A}$ the 0-context graph of $\mathcal{A}$ is the one-out graph $G_{\mathcal{A}}^0 = (2^S, E)$ with edge set $E = \{(\emptyset, \text{res}_{\mathcal{A}}(\emptyset)) \mid \emptyset \in 2^S\}$.

For a RS $\mathcal{A}$ the transition graph of $\mathcal{A}$ is the graph $G_{\mathcal{A}} = (2^S, E)$ with edge set $E = \{(\emptyset, w) \mid \emptyset \in 2^S, \text{res}_{\mathcal{A}}(\emptyset) \subseteq w\}$.

When $\mathcal{A}$ is understood, we allow to drop the subscript in res.

By definition the 0-context graph $G_{\mathcal{A}}^0$ is a subgraph of the transition graph $G_{\mathcal{A}}$ of the same reaction system.

Remark 5.3. The link from these two graphs for a reaction system to one-out graphs G_f and graphs G_a defined by the main skeleton α in a poset as defined in Section 4 is obtained as follows. Defined on node set 2^S , the 0-context graph $G_{\mathcal{A}}^0$ equals the one-out graph $G_{\text{res}_{\mathcal{A}}}$ defined by $\text{res}_{\mathcal{A}}$, while the transition graph $G_{\mathcal{A}}$ is the graph defined by the main skeleton fixed by $\text{res}_{\mathcal{A}}$ in the partial order $(2^S, \subseteq)$.

Example 5.2. For the RS $\mathcal{A}$ from Example 5.1 the 0-context graph $G_{\mathcal{A}}^0$ is given in Figure 5. The family $\text{RES}_{\mathcal{A}}$ matches the family $\mathcal{R}$ of sets that have a nonempty set of incoming edges.

As $G_{\mathcal{A}}^0$ is a one-out graph, it consists of one or more components. Each of these components is tree-like, ‘ending in’ a single cycle. By definition of reaction systems one component must have a loop at $\emptyset$. It turns out that virtually any graph with domain 2^S is a 0-context graph of a reaction system. We only have to respect the special position of the minimal and maximal set $\emptyset$ and S . This follows from a common construction in reaction systems, see, e.g., the implementation of a transition system in Section 4.2 of [3].

Proposition 5.4. A one-out graph G with vertex set 2^S is a 0-context graph of a RS if and only if $(\emptyset, \emptyset)$ and $(S, \emptyset)$ are edges of G .

Proof. For every RS we have $\text{res}(\emptyset) = \text{res}(S) = \emptyset$ as no reactions are enabled in the empty set, and all reactions are inhibited in the full set S due to the requirement that the reactant and the inhibitor are non-empty. Therefore, every 0-context graph of a RS contains the edges $(\emptyset, \emptyset), (S, \emptyset)$.

Consider a one-out graph with vertices 2^S containing edges $(\emptyset, \emptyset), (S, \emptyset)$. Then we define a set of reactions matching the rest of the edges in the graph: $A = \{(X, S \setminus X, Y) \mid (X, Y) \in E, X \not\subseteq \emptyset, S\}$. The complementarity of the first and second component of the reactions ensures that each reaction is enabled only at a single set, and that $(X, Y) \in E$ if and only if $Y = \text{res}(X)$.

As a consequence of Lemma 4.1 we can characterize graphs that are isomorphic to transition graphs of reaction systems. Recall that a faithful correspondence maps companion sets between two families of sets, respecting their sizes. As observed in Lemma 4.1, the out-sets of transition graphs must faithfully correspond to the structure of the upper cones in $(2^S, \subseteq)$.

Additionally, by Proposition 5.4, each transition graph must have the edges $(\emptyset, \emptyset)$ and $(S, \emptyset)$. Similar edges must be present in any graph isomorphic to a transition graph, with one vertex in the intersection of all out-sets and the other vertex in none of the out-sets (except the out-set that consists of all vertices). This characterization is given formally in the following theorem.

Theorem 5.5. A graph $G = (V, E)$ is isomorphic to a transition graph of a reaction system if and only if

1. there is a faithful correspondence η between $\mathcal{O}(G)$ and a family of upper cones of $(2^{[n]}, \subseteq)$, for some $n \geq 1$,
2. there is a vertex $v_{\perp} \in V \setminus \bigcup_{X \in \mathcal{O}(G), X \neq V} X$, and a vertex $v_{\top} \in \bigcap \mathcal{O}(G)$ such that $(v_{\perp}, v_{\perp}), (v_{\top}, v_{\perp})$ are edges in G .

Proof If $\mathcal{A}$ is a RS with background set S , and taking $v_{\perp} = \emptyset$ and $v_{\top} = S$, by construction requirements (1) and (2) hold for the transition graph $G_{\mathcal{A}}$, so they must hold for any graph isomorphic to it.

Assume G is a graph satisfying the two conditions of the theorem. We show it is isomorphic to a transition graph of a RS.

Note that the incoming edge $(v_{\top}, v_{\perp})$ to $v_{\perp}$ means that $v_{\perp}$ is an element of one of the out-sets. As we have chosen $v_{\perp}$ to be in none of the out-sets, except (possibly) V , we confirm that V is an out-set. As all out-sets are subsets of V , and the faithful correspondence matches V to the largest upper cone in the family, we may assume that $|V| = 2^n$. We set $S = [n]$.

By (1) and Lemma 4.1, G is isomorphic to a graph G_{α} over the poset $(2^S, \subseteq)$, where $\alpha = (\text{Up}[R], R, g)$ is a main skeleton. In order for G_{α} to be a transition graph, the function $g: 2^S \rightarrow 2^S$ must additionally satisfy $g(\emptyset) = \emptyset$ and $g(S) = \emptyset$, cf. Proposition 5.4.

Note that the skeleton α as constructed in the proof of Lemma 4.1 is based on the node-to-node bijection ϕ between V and 2^S that extends the bijection η between $\mathcal{O}(G)$ and $\text{Up}[R]$, see Lemma 2.3. There is some freedom in constructing this bijection ϕ , as long as we respect companion sets.

By (2) vertex $v_{\perp}$ belongs to the companion set that is within V and outside the other sets from $\mathcal{O}(G)$. By the faithful correspondence η , that companion set must match the corresponding ‘outer’ companion set $C_{\text{Up}[R]}(\emptyset)$. That means we can take $\phi(v_{\perp}) = \emptyset$. As the only out-set that contains $v_{\perp}$ must be V , we conclude that in G both $v_{\perp}$ and $v_{\top}$ have out-set V .

Similarly, the intersection of all out-sets in $\mathcal{O}(G)$ is a companion set which must correspond to the ‘inner’ companion set $\bigcap \text{Up}[R]$, which contains S , and we may assume that $\phi(v_{\top}) = S$.

To conclude, we follow the proof of Lemma 4.1. For each $x \in V$ with out-set X in that proof we set $g(\phi(x)) = z$ where $\text{Up}(z) = \phi(X)$ is the set in $\text{Up}[R]$ that corresponds to X . If we apply

this to $v_{\perp}$ we set $g(\varphi(v_{\perp})) = g(\emptyset) = \emptyset$ as $v_{\perp}$ has out-set V which must correspond to $\text{Up}(\emptyset)$. Also $v_{\top}$ has out-set V , thus $g(S) = \emptyset$, as required.

We now state a characterization when reaction systems have isomorphic transition graphs. This is an immediate application of Theorem 3.6, where it is shown that two graphs are isomorphic iff they have companion skeletons. As observed in Section 4, in the case when the out-sets of graph G are cones in a poset, then the main skeleton of G is fixed by a function f , or equivalently a one-out graph G_f . This can be further specialised to reaction systems, see Remark 5.3, where the transition graph $G_{\mathcal{A}}$ has its main skeleton associated to function $\text{res}_{\mathcal{A}}$ and one-out graph $G_{\mathcal{A}}^0$.

Theorem 5.6. For reaction systems $\mathcal{A}$ and $\mathcal{A}'$, their transition graphs $G_{\mathcal{A}}$ and $G_{\mathcal{A}'}$ are isomorphic if and only if the 0-context graphs $G_{\mathcal{A}}^0$ and $G_{\mathcal{A}'}^0$ are companions.

We see that there is no obvious structural relationship between two RS's $\mathcal{A}$ and $\mathcal{A}'$ such that $G_{\mathcal{A}}$ and $G_{\mathcal{A}'}$ are isomorphic. A basic operation on 0-context graphs that yields a transition graph isomorphic to that of the original 0-context graph is based on the companion edge swapping on the main skeletons.

Let $\mathcal{A}$ be a RS, and let $x, y \in 2^S$ be two elements such that $x \sim_{\text{RES}_{\mathcal{A}}} y$. Consider the pair of edges $(x, \text{res}_{\mathcal{A}}(x))$ and $(y, \text{res}_{\mathcal{A}}(y))$ in $G_{\mathcal{A}}^0$ (with $\text{res}_{\mathcal{A}}(x) \leq \text{res}_{\mathcal{A}}(y)$). The graph $G_{x,y}$ that is obtained from $G_{\mathcal{A}}^0$ by swapping the targets of these edges, introducing the new pair $(x, \text{res}_{\mathcal{A}}(y))$ and $(y, \text{res}_{\mathcal{A}}(x))$, is again a one-out graph and hence the 0-context graph $G_{\mathcal{A}'}^0$, of a RS $\mathcal{A}'$ (provided x, y are unequal to either $\emptyset$ or S , see Proposition 5.4).

By Lemma 3.1 the sources of the incoming edges of x and y in $G_{\mathcal{A}}$ are equal. Switching outgoing edges of x and y in $G_{\mathcal{A}}^0$ to obtain $G_{\mathcal{A}'}^0$, swaps all outgoing edges of x and y in $G_{\mathcal{A}}$ to obtain $G_{\mathcal{A}'}$. All other edges from $G_{\mathcal{A}}$ remain unchanged in $G_{\mathcal{A}'}$, so we can conclude that $G_{\mathcal{A}}$ and $G_{\mathcal{A}'}$, are isomorphic.

Example 5.3. Reconsider the RS $\mathcal{A}$ from Example 5.1, see Figure 5 for its 0-context graph $G_{\mathcal{A}}^0$. The elements $\{1, 3\}$ and $\{1, 3, 4\}$ are companions with respect to $\text{RES}_{\mathcal{A}}$. After companion edges swapping we obtain the graph from Figure 6 that can be considered a 0-context graph $G_{\mathcal{A}'}^0$, for RS $\mathcal{A}'$ where the reactions can be constructed as in the ‘normal form’ construction of Proposition 5.4. This graph has a single component; it has no cycles except for the (unavoidable) loop at $\emptyset$, where $G_{\mathcal{A}}^0$ from Figure 5 has two components.

This change in the 0-context graphs results in the following changes in the transition graphs. The targets of the edges originating in $\{1, 3\}$ in $G_{\mathcal{A}}$ are swapped with the targets of the edges starting at $\{1, 3, 4\}$ to obtain $G_{\mathcal{A}'}$. In particular, the loop $(\{1, 3, 4\}, \{1, 3, 4\})$ and the edge $(\{1, 3, 4\}, \{1, 3\})$ from $G_{\mathcal{A}}$ are replaced by the edge $(\{1, 3\}, \{1, 3, 4\})$ and the loop $(\{1, 3\}, \{1, 3\})$ in $G_{\mathcal{A}'}$. The bijection on the vertices of $G_{\mathcal{A}}$ which interchanges vertices

$\{1, 3\}$ and $\{1, 3, 4\}$, while mapping all other vertices to themselves is an isomorphism from the transition graph $G_{\mathcal{A}}$ to the transition graph $G_{\mathcal{A}'}$.

Acknowledgments

The authors thank the anonymous referees for their valuable comments. This research was initiated while the first and third author spent part of their sabbatical at Leiden University.

References

1. Baixeries J, Szathmary L, Valtchev P, Godin R. Yet a faster Algorithm for Building the Hasse Diagram of a Concept Lattice. Formal Concept Analysis (ICFCA 2009), Lecture Notes in Computer Science. 55482009; :162–177. DOI: 10.1007/978-3-642-01815-2_13
2. Babai, L. Proceedings of the 48th Annual ACM SIGACT Symposium on Theory of Computing (STOC 2016). ACM; New York, NY, USA: 2016. Graph isomorphism in quasipolynomial time; 684–697. extended abstract
3. Brijder R, Ehrenfeucht A, Main M, Rozenberg G. A Tour of Reaction Systems. International Journal of Foundations of Computer Science. 222011; :1499–1517. DOI: 10.1142/S0129054111008842
4. Cruz J, Giusti C, Itskov V, Kronholm B. On open and closed convex codes. arXiv:1609.03502.
5. Dennunzio A, Formenti E, Manzoni L, Porreca AE. Reachability in Resource-Bounded Reaction Systems, Language and Automata Theory and Applications (LATA 2016). Lecture Notes in Computer Science. 96182016; :592–602. DOI: 10.1007/978-3-319-30000-9_45
6. Ehrenfeucht A, Rozenberg G. Reaction systems. Fundamenta Informaticae. 752007; :263–280.
7. Formenti E, Manzoni L, Porreca AE. Cycles and Global Attractors of Reaction Systems, Descriptive Complexity of Formal Systems (DCFS 2014). Lecture Notes in Computer Science. 86142014; :114–125. DOI: 10.1007/978-3-319-09704-6_11
8. Formenti E, Manzoni L, Porreca AE. Fixed Points and Attractors of Reaction Systems Computability in Europe (CiE 2014). Lecture Notes in Computer Science. 84932014; :194–203. DOI: 10.1007/978-3-319-08019-2_20
9. Formenti E, Manzoni L, Porreca AE. On the complexity of occurrence and convergence problems in reaction systems. Natural Computing. 142015; :185–191. DOI: 10.1007/s11047-014-9456-3
10. Gallai T. Transitiv orientierbare Graphen. Acta Mathematica Academiae Scientiarum Hungaricae. 181967; :25–66. DOI: 10.1007/BF02020961
11. Hall P. On Representatives of Subsets. Journal of the London Mathematical Society. 101935; :26–30. DOI: 10.1112/jlms/s1-10.37.26
12. McConnell RM, de Montgolfier F. Linear-time modular decomposition of directed graphs. Discrete Applied Mathematics. 1452005; :198–209. DOI: 10.1016/j.dam.2004.02.017

Figure 1.

From Example 3.1: Top: the one-out graphs G_f , G_g and G_h corresponding to the skeletons ν , σ , and τ . In each graph, the representative of X is indicated in red and the representative of Y is in blue. Bottom: the graph $G = G_\nu = G_\sigma = G_\tau$. Graphs G_f and G_g are subgraphs of G , but G_h is not.

Figure 2. Companion skeletons related by edge swapping. The graph $G_{\nu_{2,3}}$ is isomorphic to $G = G_{\nu}$ in Fig. 1.

Figure 3.

Functions f and g of companion skeletons, cf. Lemma 3.7, a case where $\mathcal{O} = \mathcal{P}$ and $R = Q$ but f and g are distinct

Figure 4. Venn diagram of the out-sets of a graph, and collection of cones which is in faithful correspondence, see Example 4.2

Figure 5.
Zero context graph G_S^0 for the RS from Example 5.1.

Figure 6.

The 0-context graph for a RS equivalent to the one in Example 5.1 after swapping companion edges. Dashed edges are the replacements of the two original edges (compare to Fig. 5).