Published in final edited form as:

J Int Neuropsychol Soc. 2020 February ; 26(2): 197–209. doi:10.1017/S1355617719001140.

Profiles of Normal Cognition in Essential Tremor

Tess E.K. Cersonsky¹, Sarah Kellner¹, Silvia Chapman², Edward D. Huey^{2,3,4}, Elan D. Louis^{1,5,6}, Stephanie Cosentino^{2,3}

¹Division of Movement Disorders, Department of Neurology, Yale School of Medicine, Yale University, New Haven, CT, USA

²Taub Institute for Research on Alzheimer's Disease and the Aging Brain, College of Physicians and Surgeons, Columbia University, New York, NY, USA

³Department of Psychiatry, College of Physicians and Surgeons, Columbia University, New York, NY, USA

⁴Department of Neurology, College of Physicians and Surgeons, Columbia University, New York, NY, USA

⁵Department of Chronic Disease Epidemiology, Yale School of Public Health, Yale University, New Haven, CT, USA

⁶Center for Neuroepidemiology and Clinical Neurological Research, Yale School of Medicine, Yale University, New Haven, CT, USA

Abstract

Objectives: Patients with essential tremor exhibit heterogeneous cognitive functioning. Though the majority of patients fall under the broad classification of cognitively “normal,” essential tremor is associated with increased risk for Mild Cognitive Impairment and dementia. It is possible that patterns of cognitive performance within the wide range of normal functioning have predictive utility for Mild Cognitive Impairment or dementia. These cross-sectional analyses sought to determine whether cognitive patterns, or “clusters,” could be identified among individuals with essential tremor diagnosed as cognitively normal. We also determined whether such clusters, if identified, were associated with demographic or clinical characteristics of patients.

Methods: Elderly subjects with essential tremor (age >55 years) underwent comprehensive neuropsychological testing. Domain means (memory, executive function, attention, visuospatial abilities, language) from 148 individuals diagnosed as cognitively normal were partitioned using k-means cluster analysis. Individuals in each cluster were compared according to cognitive functioning (domain means and test scores), demographic factors, and clinical variables.

Results: There were three clusters. Cluster 1 (n=64) was characterized by comparatively low memory scores ($p<0.001$), Cluster 2 (n=39) had relatively low attention and visuospatial scores ($p<0.001$), and Cluster 3 (n=45) exhibited consistently high performance across all domains.

Cluster 1 had lower Montreal Cognitive Assessment scores and reported more prescription medication use and lower balance confidence.

Conclusions: Three patterns of cognitive functioning within the normal range were evident and tracked with certain clinical features. Future work will examine the extent to which such patterns predict conversion to Mild Cognitive Impairment and/or dementia.

Keywords

Cognitive Aging; Cerebellar diseases; Movement Disorders; Attention; Memory; Postural Balance

Introduction

Diagnosed on the basis of kinetic tremor (Benito-León & Louis, 2007), essential tremor (ET), a common movement disorder with a prevalence of 4% in adults age 40 (Dogu et al., 2003; Louis & Ferreira, 2010), is associated with heterogeneous cognitive functioning ranging from normal cognition to dementia (Benito-León, Louis, Bermejo-Pareja, & Neurological Disorders in Central Spain (NEDICES) Study Group, 2006a; Benito-León, Louis, Mitchell, & Bermejo-Pareja, 2011; Bermejo-Pareja, Louis, Benito-León, & Neurological Disorders in Central Spain (NEDICES) Study Group, 2007; Louis, Benito-León, Vega-Quiroga, Bermejo-Pareja, & Neurological Disorders in Central Spain (NEDICES) Study Group, 2010a; Louis, Rao, & Gerbin, 2012; Sengul et al., 2015; Thawani, Schupf, & Louis, 2009). While the majority of individuals with ET perform within the spectrum of normal cognition, there is accumulating evidence that they progress to Mild Cognitive Impairment (MCI) and dementia at higher rates than age matched controls (Benito-León et al., 2011; Bermejo-Pareja et al., 2007; Thawani et al., 2009). Identifying the earliest manifestations of cognitive compromise and improving knowledge of cognitive progression in ET is important for diagnosis, prognosis, and treatment.

Toward these ends, examining the variability of performance in normal cognitive function may help identify early patterns of cognitive performance that predict MCI, dementia, or other disease outcomes in ET. In general, cognitive performance is considered to be within normal limits when no lower than 1.0 to 2.0 standard deviations (SD) below the mean; normal performance is thus frequently defined as performance above a cut-off of -1.5 SD, corresponding to approximately the 7th percentile and extending up through the 99th percentile. Within this very broad spectrum, individuals can exhibit certain performance patterns or relative strengths and weaknesses across the domains of memory, attention, language, executive functioning, and visuospatial abilities (Chamorro-Premuzic, Von Stumm, & Furnham 2011; Salthouse, 2004). Whether there are identifiable different patterns of performance among individuals with ET considered to be cognitively normal, and whether such patterns have relevance for heterogeneity in cognitive progression or the substrates of future cognitive decline, is not known.

With regard to clinically significant cognitive impairment in ET, executive dysfunction is most often noted (Frisina, Tse, Hälbig, & Libow, 2009; Gasparini et al., 2001; Higginson et al., 2008; Kim et al., 2009; Lombardi, Woolston, Roberts, & Gross, 2001; Passamonti et al., 2011) and is thought to reflect alterations in the cerebello-thalamo-cortical loop (Deuschl,

Wenzelburger, Löffler, Raethjen, & Stolze, 2000; Middleton & Strick, 2000a, 2000b, 2001; Montgomery, Baker, Lyons, & Koller, 2000). However, cognitive impairment in ET is not homogeneous, and studies have documented deficits in memory, attention, and visuospatial abilities as well (Benito-León, Louis, Sánchez-Ferro, & Bermejo-Pareja, 2013; Gasparini et al., 2001; Janicki, Cosentino, & Louis, 2013; Kim et al., 2009; Lombardi et al., 2001; Louis, 2010; Louis & Rao, 2014; Mameli et al., 2013; Sinoff & Badarny, 2014; Tröster et al., 2002). In fact, in a recent study of ET-MCI, more patients had amnesic (deficient in memory) rather than non-amnesic (deficient in other cognitive domains) presentations of MCI (Cersonsky et al., 2018; Collins et al., 2017). The presence of amnesic deficits in a subset of individuals with ET aligns with epidemiologic and pathological studies suggesting that neurodegenerative disorders such as Alzheimer's disease (AD) or other tauopathies may be more common in ET (Benito-León, et al., 2006a, 2006b; Benito-León et al., 2011; Farrell et al., 2019; LaRoia & Louis, 2011; Louis, Benito-León, et al., 2010; Louis, Benito-León, Vega-Quiroga, Bermejo-Pareja, & Neurological Disorders in Central Spain (NEDICES) Study Group, 2010b; Louis, Babij, Ma, Cortés, & Vonsattel, 2013; Pan et al., 2014; Thawani et al., 2009). The nature and bases of cognitive symptoms in ET, as well as their longitudinal course, are thus seemingly heterogeneous.

This cross-sectional study utilized a cluster-based approach to characterize patterns of cognitive functioning among cognitively normal individuals with ET. Although we do not examine the specificity of cognitive patterns in ET, identifying clusters within an ET population may be useful as it may reveal sub-clinical patterns of performance that foreshadow the emergence of different types of cognitive impairment and different cognitive trajectories among individuals with ET. Identification of cognitive profiles was accomplished using k-means cluster analysis to group individuals by domain scores (memory, executive function, attention, visuospatial abilities, and language). K-means cluster analysis partitions subjects according to the variables under investigation (i.e., cognitive domain scores) into a given number of clusters such that each subject belongs to the cluster with the nearest mean. A secondary goal was to assess whether these patterns were associated with demographic variables or clinical features such as tremor severity, gait/balance difficulty, and/or depressive symptoms. The current analyses are a necessary first step in determining whether certain cognitive profiles have implications for progression to MCI, dementia or other disease outcomes. Moreover, they may improve understanding of the bases of cognitive deficits in those with progressive cognitive decline versus those whose cognitive deficits remain relatively static.

Methods

Subjects

Subjects were assessed using a cognitive test battery designed by a neuropsychologist (SC) as part of COGNET (Clinical-Pathological Study of Cognitive Impairment in Essential Tremor, NINDS R01NS086736), a prospective, longitudinal study of cognitive function in ET. Since July 2014, subjects have been clinically characterized over three assessments (baseline, 18 months, 36 months); baseline data utilized for these analyses were collected from July 2014 through July 2016. Signed, written, informed consent was obtained upon

enrollment from eligible subjects recruited online through study and International Essential Tremor Foundation websites. Subjects met each of the following criteria: 1. diagnosed with ET; 2. age >55 years; 3. no history of deep brain stimulation surgery; 4. concurrent enrollment as a brain donor and agreement to complete study measures. Yale University and Columbia University Internal Review Boards approved study procedures.

Subjects were excluded from analyses if they fulfilled any of the following criteria: 1. Diagnosis of dementia (n = 21), MCI (of any type, n = 35), or cognitive impairment related to substance use, stroke, or other injury (n = 3); 2. Diagnosis of PD, dystonia, or other non-ET cause of tremor (n = 32), for a final sample size of 148.

In-person assessments

Subjects were assessed during in-person study visits consisting of a clinical questionnaire, 19 neuropsychological tests, neuropsychiatric measures, and a videotaped neurological examination. The clinical questionnaire included demographic and clinical data on age, gender, ethnicity, and education as well as questions regarding medication usage, including those with cognition-enhancing, cognition-decreasing, and mood-modulating effects (determined by EDL). Additional questionnaires evaluated mood (Geriatric Depression Scale (GDS; range 0-30, higher scores indicate more depressive symptoms (Yesavage et al., 1982)), balance (number of falls in past year and short Activities-specific Balance Confidence scale (ABC-6; range 0-100, lower scores indicate less balance confidence (Peret, Herman, Hausdorff, & Giladi, 2006)), and tremor experience (tremor duration and Tremor Disability Scale [range 0-100, higher scores indicate more disability (Louis et al., 1999)]).

Diagnosis of ET was confirmed by a movement disorders neurologist (EDL) from the videotaped neurological examination using the Washington Heights-Inwood Genetic Study of ET (WHIGET) diagnostic criteria (Louis et al., 1997), which require moderate or greater amplitude kinetic tremor during three or more tests or head tremor in the absence of Parkinson's Disease (PD), dystonia, or other cause. These criteria have been shown to be reliable (Louis, Ford, & Bismuth, 1998) and valid (Louis et al., 1999). Kinetic or postural tremor were rated (0-3) on 12 items, resulting in a total tremor score (range 0-36, higher scores indicate more severe tremor). For tandem gait assessment in the neurological videotape, individuals were asked to walk heel-to-toe in a straight line for at least 10 feet; the number of steps taken off a straight line was reported.

The cognitive test battery was designed to minimize disadvantage towards ET subjects with moderate or severe tremors by incorporating only tests whose scores had little to no reliance on motor functioning and discounting tremor-related difficulty on those tests requiring motor utilization (i.e. in the Token test, not considering an answer incorrect if a participant touches other tokens in addition to the target token because of their tremor). In addition to assessments of global functioning (Mini-Mental State Examination [MMSE (Folstein, Folstein, & McHugh, 1975)] and Montreal Cognitive Assessment [MoCA (Nasreddine et al., 2005)]), participants completed detailed tests across five domains including: 1. **memory** (California Verbal Learning Test II (Delis, Kramer, Kaplan, & Ober, 2000), Wechsler Memory Scale Revised: Logical Memory (Wechsler, 1987), Wechsler Memory Scale IV: Verbal Paired Associates (Wechsler, 2008)), 2. **executive function** (Wechsler Adult

Intelligence Scale IV [WAIS-IV] Digit Span Backward (Wechsler, 1997), Delis-Kaplan Executive Function System: Verbal Fluency Test, Color-Word Interference, Sorting, 20-Questions (Delis, Kaplan, & Kramer, 2001)), 3. **attention** (Oral Symbol-Digit Modalities Test (Smith, 1982), WAIS-IV Digit Span Forward (Wechsler, 1997)), 4. **visuospatial abilities** (Benton Judgment of Line Orientation (Benton, Sivan, des Hamsher, Varney, & Spreen, 1994), Benton Facial Recognition Test (Benton & Van Allen, 1968), WAIS-IV Visual Puzzles (Wechsler, 1997)), and 5. **language** (Multilingual Aphasia Examination: Token Test (Benton, des Hamsher, Rey, & Sivan, 1994), Boston Naming Test (Kaplan, Goodglass, & Weintraub, 1983)). Raw scores of these assessments were converted to Z-scores using published normative data according to age, gender, and/or education, which offer objective references against which we can compare individual performance, rather than relying on the heterogeneity of the specific sample to provide normative data (Busch, Chelune, & Suchy, 2006). We also chose our approach to enable classification of cognitive performance in relation to the expected level of performance among the “normal” population rather than in relation to sample specific performance which may or may not be within normal limits. A disadvantage of this approach is that normative scores are thus based on different sample for different tests rather than from a single sample. Nonetheless, this is the standard approach to interpreting neuropsychological performance in clinical settings. Z-scores were considered to be “low average” when between -1.49 and -0.51 , “average” when between -0.50 and 0.50 , and “high average” when between 0.51 and 1.49 . Domain aggregates were calculated using a selection of tests per domain meant to capture heterogeneous cognitive profiles by including tests that measured different sub-domain attributes (i.e. incorporating immediate and delayed recall sub-scores to include both an index of verbal learning ability and a measure of delayed retention and forgetting rates). In the California Verbal Learning Test, both Total Recall and Long-Delay Free Recall were chosen to capture both immediate and delayed recall, but individual trials, short-delay recall, and recognition hits were not included; additionally, Number Span length of the longest series were excluded for both forward and backward assessments.

Cognitive diagnosis assignment

Informants designated by participants were asked to assess functioning across the six Clinical Dementia Rating (CDR (Morris, 1993, 1997)) dimensions (Memory, Orientation, Judgment and Problem Solving, Community Affairs, Home and Hobbies, and Personal Care; range 0-3, higher scores indicate more cognitive impairment) and completed the Lawton Instrumental Activities of Daily Living Scale (IADL; range 0-8, lower scores indicate less functional ability (Lawton & Brody, 1969)). Aggregate CDR score was calculated according to informant and examiner impressions. Tremor-related disability was not included in scoring. If an informant was not available, CDR score was calculated according to examiner impression and participant self-report ($n = 42$, 28.4%).

Cognitive diagnoses (normal, MCI, and dementia) were assigned during a diagnostic case conference with trained experts (EDH, SC) using the CDR (Morris, 1993), neuropsychological tests selected *a priori* for diagnosis of MCI, and clinical judgment, as described previously (Cersonsky et al., 2018; Collins et al., 2017). Within the group of individuals diagnosed as cognitively normal, subjects were assigned subdiagnoses to capture

apparent differences among individuals at a global level, including: fully normal cognition (CDR = 0, $z > -1.5$ for all scores), test impairment of unlikely clinical significance, test impairment of possible clinical significance, and questionable/isolated functional impairment (CDR = 0.5, $z > -1.5$ for all scores).

Statistical analysis

Domain aggregates (mean Z-score per each of the cognitive domains) were calculated for each of the study subjects. Domains were first compared across subdiagnoses of this cognitively normal ET sample including those diagnosed with fully normal cognition (group 1, fully normal cognition) and those labeled as having test impairment of unlikely clinical significance, test impairment of possible clinical significance, and questionable/isolated functional impairment (group 2, not-fully normal cognition) using 2-sample t-tests. K-means cluster analysis was then performed (Chiu, Douglas, & Li, 2009; Souza, Oliveira, Foss, & Tumas, 2016; Szeto et al., 2015) to identify the extent to which a set of specific cognitive profiles best described the overall sample, and bar graphs were generated, using SPSS 24. Analysis was performed at a range of cluster numbers. Mean domain scores were compared across clusters using analyses of variance (ANOVAs) with Bonferroni corrections for post-hoc analyses. For each cluster solution (i.e., 2 through 5 clusters), we calculated the number of significant differences in cognitive domain scores across clusters, correcting for multiple comparisons. The ideal cluster solution was defined as that which generated the highest number of significant differences in domain scores across clusters with the largest number of individuals per cluster (Bruehl, Lofland, Semenchuk, Rokicki, & Penzien, 1999; Hauser & Rybakowski, 1997).

After determining the optimal cluster solution, individual neuropsychological test scores were compared within clusters using Kruskal-Wallis H-tests and between clusters using post-hoc Mann-Whitney U-tests (non-normal distributions according to Kolmogorov-Smirnov test).

Demographic and clinical analyses were completed using Kruskal-Wallis H-tests (continuous variables) or Pearson Chi-Square Tests (categorical variables) and post-hoc analyses were performed using Mann-Whitney U-tests or Pearson Chi-Square tests, respectively. Frequency of normal cognition sub-diagnosis (fully normal cognition, test impairment of unlikely clinical significance, test impairment of possible clinical significance, and questionable/isolated functional impairment) was compared across cluster membership using a Fisher's Exact Test. Values were considered significant if $p < 0.05$, unless otherwise stated. All analyses were performed using SPSS 24.

Results

Comparison of "fully normal cognition" to "not-fully normal cognition" (group 1 versus group 2) showed that memory, executive function, and attention domains significantly differed between these cognitively normal subgroups (Table 1).

The 3-cluster solution yielded the highest number of significant differences in cognitive domain scores (75%) across clusters in post-hoc analyses (Table 2, Figure 1). Attention and

visuospatial abilities were significantly different across all clusters, while memory and executive function differed only across two. Only language was comparable across all clusters. Cluster 1 (n=64) was characterized by average performance in all domains, though memory was significantly lower than Clusters 2 and 3; Cluster 2 (n=39) was characterized by high average memory, low average attention, and average executive function, visuospatial abilities, and language; and Cluster 3 (n=45) was characterized by high average memory, executive function, attention, and visuospatial abilities, and average language. In comparing individual neuropsychological test scores across clusters, several tests across all domains emerged as significantly different ($p < 0.0083$, Table 3).

The clusters did not differ according to demographic (age, gender, education, ethnicity) or tremor (total tremor score, tremor disability scale, tremor duration, presence of head tremor, presence of voice tremor) variables, nor did they differ with respect to the usage of cognition-decreasing or mood-modulating medications or levels of depression (Table 4).

However, number of medications, ABC-6, and MoCA were significantly different among clusters, with particular difference found between Clusters 1 and 3 (cluster 1 indicating more medications, lower balance confidence, and lower MoCA; post-hoc $p < 0.0167$) in those variables. Additionally, the frequencies of cognitively normal subdiagnoses (fully normal cognition, test impairment of unlikely clinical significance, test impairment of possible clinical significance, and questionable/isolated functional impairment) were also significantly different across clusters ($p > 0.001$, Table 5). Specifically, there were more individuals with fully normal cognition in Cluster 3 (n=30) than in the other clusters, and more individuals showing impairment of possible clinical significance in Cluster 1 (n=24) than in the other clusters.

Discussion

Cognitive patterns

This study aimed to identify patterns of cognitive functioning among individuals with ET who underwent comprehensive cognitive testing and who were assigned diagnoses of normal cognition during clinical diagnostic consensus conference. Subtle cognitive differences in the same domains that appeared to drive cluster formation (i.e. attention, executive function, and memory) were observed even between those classified as “fully normal cognition” and those with test impairment of unlikely clinical significance, test impairment of possible clinical significance, and questionable/isolated functional impairment, lending support to the cluster identities. K-means cluster analysis within this group of individuals indicated that a three-cluster solution best captured differences in patterns of performance across measures of memory, executive function, attention, visuospatial abilities, and language. Between-cluster comparisons of cognitive functions revealed sparing and weaknesses of specific abilities relative to the other clusters. Cluster 1 was average in all domains with a primary weakness in memory (worse than other two clusters), and relative weaknesses in attention (including weakness in some tests with attentional components, such as Immediate Recall memory tests and Color Naming and Reading), executive and visuospatial abilities compared to Cluster 3. Cluster 2 was largely average as well, but had high average memory and primary weakness in the attention

domain (worse than other two clusters) and a relative weakness in executive functioning compared to Cluster 3. Finally, Cluster 3 was high average across all domains except language, which was average. Several sub-scores, such as Logical Memory Delayed, Digit Span Forward, and Benton Facial Recognition Test, were significantly different within and between clusters.

Demographic and clinical correlates of clusters

The clusters were formed using cognitive scores that were standardized and generally adjusted for age, and gender and education when possible. This approach reduces the likelihood that such demographic variables would differ across the clusters, although it is certainly possible that differences could still have arisen. In our current study, the comparability of age, gender distribution, and educational attainment across the three clusters reinforces the idea that the clusters are capturing distinct cognitive profiles rather than differences in overall cognitive functioning secondary to the demographic characteristics of the participants. In particular, it is worth noting that the similarity in years of formal education across groups reduces the possibility that the cognitive advantage in Cluster 3 reflects premorbid cognitive characteristics. Similarly, the clusters did not differ with respect to the use of cognition-decreasing medication. With regard to indicators of overall cognitive performance, Cluster 1 contained the most individuals with sub-test impairment of possible clinical significance while Cluster 3 contained the most fully cognitively normal individuals compared to other clusters. Consistent with this finding, Cluster 1 evidenced lower performance on the MoCA, a global cognitive screening measure, than Cluster 3.

The current study also examined the extent to which cognitive clusters co-varied with clinical features of disease. Tremor characteristics including total tremor score, duration, and disability, and presence of head or voice tremor, were not different across clusters, consistent with previous work suggesting that tremor severity does not dictate cognitive performance (Passamonti et al., 2011). Moreover, objective measures of balance, such as number of falls in past year or number of steps off the line in tandem gait testing, did not differ across clusters. Interestingly, however, subjective ratings of balance confidence did differ, with Cluster 3 reporting better balance than Cluster 1. It is possible that the ABC-6, which has been shown to correlate well with falls and balance-impairment (Cho, Scarpance, & Alexander, 2004; Cumming, Salkeld, Thomas, & Szonyi, 2000), may capture subtle subjective changes in balance confidence. Taken together, it appears that measurable differences in cognition, specifically in memory, already exist among cognitively normal ET subjects and co-vary with subjective balance confidence, consistent with the idea that balance impairment may be a better marker of cognitive decline than tremor severity (Louis et al., 2017; Louis & Rao, 2014).

Number of medications also differed across clusters, specifically between Cluster 1 and Cluster 3, with Cluster 1 reporting more medication use, though cognition-enhancing and cognition-decreasing medications did not differ across or between clusters. Future analyses may reveal that cognition is more likely to decline among subjects in Cluster 1, as it is characterized by lower global cognitive scores (MoCA), less balance confidence (ABC-6)

and use of more medications, measures which are related to cognitive decline in ET (Louis et al., 2017).

Potential implications for the substrates and course of cognitive weaknesses

It is possible that Clusters 1 and 2 each reflect very mild and/or early compromise of specific neuroanatomic networks that may be vulnerable in ET. In contrast, Cluster 3 may reflect a subset of cases in which cognitive functioning and their underlying substrates are relatively robust. Compared to Cluster 3, individuals in Clusters 1 and 2 are largely similar apart from the primary memory weakness in Cluster 1. If indeed the variability in cognitive abilities reflects incipient pathological changes in the neuroanatomic networks that support these functions, several interpretations could be extrapolated. For example, Cluster 2 may be reflective of a fairly “pure” ET cognitive profile (“cortico-cerebellar”, or “fronto-thalamic-cerebellar”) as ET-associated cerebellar dysfunction and defects in the cerebellar-cortical loop have been shown to contribute to deficits in attention (Bareš, Husárová, & Lungu, 2012; Deuschl et al., 2000; Hanajima et al., 2016; Louis, 2016; Louis et al., 2006). Moreover, this pathology does not appear to be directly linked to deficits in the memory domain (Schmahmann & Caplan, 2006). If this Cluster were to represent a “pure” ET group, the relatively intact executive function in comparison to Cluster 1 may be somewhat unexpected. It is possible that restriction of low performance to the attention domain is a product of the specific clusters formed in this sample; it is also possible that the primary profile of “pure” ET is not characterized by significant executive dysfunction. Future longitudinal and clinical-pathological analyses will better inform these questions.

In contrast, Cluster 1 shows relative weaknesses of attention, in addition to memory and executive function. This group may have pathological changes characteristic of ET-specific generation in conjunction with another neurodegenerative disorder (e.g. AD, progressive supranuclear palsy, frontotemporal dementia, etc.). Finally, Cluster 3 appears to be an intact, high functioning group. These individuals may have an inherent resilience to the cognitive change observed in others with ET. This resilience could be reflective of some intrinsic, unmeasured characteristic of this group such as quality rather than quantity of their educational experiences for example (Manly, Touradji, Tang, & Stern, 2003), or it may reflect the particular manner in which ET expresses itself.

Limitations

There are several limitations inherent to the current study. Following the criteria for diagnosing PD-MCI from the Movement Disorders Society (Goldman et al., 2015), we selected various tests per domain; unfortunately, some domains had fewer tests than others, which could have affected the domain measurements as some may be more reliable than others. We took an agnostic approach to the cognitive profiles that might emerge in the cluster analysis, grouping neuropsychological tests according to their primary domain. While the domain groupings are thus psychometrically and conceptually meaningful, they may obscure relevant heterogeneity in the cognitive processes measured by tests in each domain. However, as the first study to examine cognitive profiles among cognitively normal individuals with ET, there was not sufficient evidence to group cognitive performance according to particular pre-clinical disease profiles (e.g., AD profile). Without a non-ET

group or another clinical population, we are unable to determine the extent to which these clusters are specific to ET and different than what is seen among neurologically healthy elders. However, the potential utility of these clusters is not dependent on the extent to which they are specific to ET. Second, while the neuropsychological test battery was designed to reduce any effect of kinetic or voice tremor on performance, it is possible that certain tests with rapid verbal or manual responses could be influenced by tremor characteristics. However, this does not appear to be the case, as tremor characteristics were not different among clusters.

Though there was no significant difference in the number of individuals using cognition-enhancing or cognition-decreasing medications across or between clusters, differences in the number of medications or dosage effects could have contributed to the observed differences in cognition. We also did not include a measure of medical comorbidity other than number of medications in our analyses; it is therefore possible that individual cognitive profiles were affected by unmeasured medical co-morbidity. However, scores on the Cumulative Illness Rating Scale (CIRS; range 0-42, higher scores indicate more medical co-morbidity), measured at follow up only, did not differ between clusters after 18 months (Mann-Whitney U-test p -Value = 0.65). It will be vital to assess medical co-morbidity in future analyses on this cohort to ensure that other conditions besides ET are not significantly contributing to cognitive decline. Another consideration is that subjects chose to participate in this study on their own volition, thus introducing potential selection bias to the study by enrolling individuals who may have had more subjective tremor- or cognition-related complaints.

Finally, though k-means cluster analysis offers a systematic approach for grouping individuals into clusters according to several characteristic measures (in our case, domain means), this method of unsupervised machine learning can be limited in its utility without a clinical gold-standard for validation. Therefore, future analyses should include pathological or cognitive outcome variables that can guide the modeling and interpretation of clusters with more conclusive clinical outcomes or pathological correlates.

Conclusions and future directions

In this paper, we were able to identify three patterns of cognitive performance in ET subjects diagnosed as cognitively normal. The three clusters differed to some extent from one another with regard to performance in memory, attention, visuospatial abilities, and executive function. Cluster 1 displayed relatively low memory compared to the other two clusters, Cluster 2 displayed relatively low attention, and Cluster 3 performed relatively high across all areas. We were also able to identify clinical correlates of these clusters, in particular that Cluster 1 had lower overall global cognition, less balance confidence, and more medications, suggesting that this cluster may be at particular risk for cognitive progression or functional decline.

With these cognitive profiles now identified, future work can assess their predictive utility for conversion to MCI, dementia, or other disease outcomes. This would not only provide information regarding the progress of cognitive decline in ET, but would also provide pre-impairment identifiers for clinicians and patients. While speculative, the patterns of performance may map onto different neuroanatomic vulnerabilities in ET. One might

hypothesize that individuals in the high-performing group will not develop cognitive impairment, whereas individuals in Cluster 2 may convert only to ET-MCI, but individuals in Cluster 1 may ultimately convert to dementia.

Acknowledgements

No conflicts of interest or competing financial interests are held by any of the authors. The National Institutes of Health supported this work (grant number NINDSR01NS086736) but played no role in the design of the study, the collection, analysis, and interpretation of data, or the writing of the manuscript.

References

- Bareš M, Husárová I, & Lungu OV (2012). Essential tremor, the cerebellum, and motor timing: towards integrating them into one complex entity Tremor and Other Hyperkinetic Movements (New York, N.Y.), 2.
- Benito-León J, & Louis ED (2007). Clinical update: diagnosis and treatment of essential tremor. *The Lancet*, 369(9568), 1152–1154. doi:10.1016/S0140-6736(07)60544-3
- Benito-León J, Louis ED, Bermejo-Pareja F, & Neurological Disorders in Central Spain (NEDICES) Study Group. (2006a). Elderly-onset essential tremor is associated with dementia. *Neurology*, 66(10), 1500–1505. doi:10.1212/01.wnl.0000216134.88617.de [PubMed: 16717208]
- Benito-León J, Louis ED, Bermejo-Pareja F, & Neurological Disorders in Central Spain (NEDICES) Study Group. (2006b). Population-based case-control study of cognitive function in essential tremor. *Neurology*, 66(1), 69–74. doi:10.1212/01.wnl.0000192393.05850.ec [PubMed: 16401849]
- Benito-León J, Louis ED, Mitchell AJ, & Bermejo-Pareja F (2011). Elderly-Onset Essential Tremor and Mild Cognitive Impairment: A Population-Based Study. *Journal of Alzheimer's Disease*, 23(4), 727–735.
- Benito-León J, Louis ED, Sánchez-Ferro Á, & Bermejo-Pareja F (2013). Rate of cognitive decline during the premotor phase of essential tremor: a prospective study. *Neurology*, 81(1), 60–66. [PubMed: 23700331]
- Benton A, des Hamsher K, Rey G, & Sivan A (1994). *Multilingual Aphasia Examination* (3rd ed.). Iowa City, IA: AJA Associates.
- Benton AL, & Van Allen MW (1968). Impairment in facial recognition in patients with cerebral disease. *Transactions of the American Neurological Association*, 93, 38–42. [PubMed: 5711050]
- Benton A, Sivan A, des Hamsher K, Varney N, & Spreen O (1994). *Contributions to neuropsychological assessment* (2nd ed.). New York, NY: Oxford University Press.
- Bermejo-Pareja F, Louis ED, Benito-León J, & Neurological Disorders in Central Spain (NEDICES) Study Group. (2007). Risk of incident dementia in essential tremor: A population-based study. *Movement Disorders*, 22(11), 1573–1580. doi:10.1002/mds.21553 [PubMed: 17516478]
- Bruehl S, Lofland KR, Semenchuk EM, Rokicki LA, & Penzien DB (1999). Use of Cluster Analysis to Validate IHS Diagnostic Criteria for Migraine and Tension-Type Headache. *Headache*, 39, 181–189. [PubMed: 15613212]
- Busch RM, Chelune GJ, & Suchy Y (2006). Using Norms in Neuropsychological Assessment of the Elderly In Attix DK & Welsh-Bohmer KA (Eds.), *Geriatric Neuropsychology: Assessment and Intervention* (pp. 133–157).
- Cersonsky TEK, Morgan S, Kellner S, Robakis D, Liu X, Huey ED, ... Cosentino S (2018). Evaluating Mild Cognitive Impairment in Essential Tremor: How Many and Which Neuropsychological Tests? *Journal of the International Neuropsychological Society*, 24(10), 1084–1098. [PubMed: 30303051]
- Chamorro-Premuzic T, Von Stumm S, & Furnham A (2011). To What Extent Do Individual Differences Exist in Aging-Related Cognitive Changes? In *The Wiley-Blackwell handbook of individual differences* (p. 820). Wiley-Blackwell.
- Chiu C-Y, Douglas JA, & Li X (2009). Cluster Analysis for Cognitive Diagnosis: Theory and Applications. *PSYCHOMETRIKA*, 74(4), 633–665. doi:10.1007/S11336-009-9125-0

- Cho B, Scarpace D, & Alexander NB (2004). Tests of Stepping as Indicators of Mobility, Balance, and Fall Risk in Balance-Impaired Older Adults. *Journal of the American Geriatrics Society*, 52(7), 1168–1173. doi:10.1111/j.1532-5415.2004.52317.x [PubMed: 15209657]
- Collins K, Rohl B, Morgan S, Huey ED, Louis ED, & Cosentino S (2017). Mild Cognitive Impairment Subtypes in a Cohort of Elderly Essential Tremor Cases. *Journal of the International Neuropsychological Society*, 23(05), 390–399. doi:10.1017/S1355617717000170 [PubMed: 28367776]
- Cumming RG, Salkeld G, Thomas M, & Szonyi G (2000). Prospective study of the impact of fear of falling on activities of daily living, SF-36 scores, and nursing home admission. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*, 55(5), M299–305.
- Delis D, Kaplan E, & Kramer J (2001). *Delis Kaplan Executive Function System (DKEFS) Examiner's Manual*. San Antonio, TX: Pearson.
- Delis D, Kramer K, Kaplan E, & Ober B (2000). *California Verbal Learning Test - second edition (CVLT-II)*. San Antonio, TX: Psychological Corporation.
- Deuschl G, Wenzelburger R, Löffler K, Raethjen J, & Stolze H (2000). Essential tremor and cerebellar dysfunction Clinical and kinematic analysis of intention tremor. *Brain*, 123(8), 1568–1580. doi: 10.1093/brain/123.8.1568 [PubMed: 10908187]
- Dogu O, Sevim S, Camdeviren H, Sasmaz T, Bugdayci R, Aral M, ... Louis ED (2003). Prevalence of essential tremor: door-to-door neurologic exams in Mersin Province, Turkey. *Neurology*, 61(12), 1804–1806. [PubMed: 14694055]
- Farrell K, Cosentino S, Iida MA, Chapman S, Bennett DA, Faust PL, ... Crary JF (2019). Quantitative Assessment of Pathological Tau Burden in Essential Tremor: A Postmortem Study. *Journal of Neuropathology & Experimental Neurology*, 78(1), 31–37. doi:10.1093/jnen/nly104 [PubMed: 30476290]
- Folstein MF, Folstein SE, & McHugh PR (1975). "Mini-mental state": A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*, 12(3), 189–198. [PubMed: 1202204]
- Frisina PG, Tse W, Hälbig TD, & Libow LS (2009). The Pattern of Cognitive-Functional Decline in Elderly Essential Tremor Patients: An Exploratory-Comparative Study With Parkinson's and Alzheimer's Disease Patients. *Journal of the American Medical Directors Association*, 10(4), 238–242. doi: 10.1016/j.jamda.2008.10.013 [PubMed: 19426939]
- Gasparini M, Bonifati V, Fabrizio E, Fabbrini G, Brusa L, Lenzi GL, & Meco G (2001). Frontal lobe dysfunction in essential tremor: a preliminary study. *Journal of Neurology*, 248(5), 399–402. [PubMed: 11437162]
- Goldman JG, Holden S, Ouyang B, Bernard B, Goetz CG, & Stebbins GT (2015). Diagnosing PD-MCI by MDS Task Force criteria: how many and which neuropsychological tests? *Movement Disorders: Official Journal of the Movement Disorder Society*, 30(3), 402–406. doi: 10.1002/mds.26084 [PubMed: 25449653]
- Hanajima R, Tsutsumi R, Shirota Y, Shimizu T, Tanaka N, & Ugawa Y (2016). Cerebellar dysfunction in essential tremor. *Movement Disorders*, 31(8), 1230–1234. doi: 10.1002/mds.26629 [PubMed: 27062434]
- Hauser J, & Rybakowski J (1997). Three clusters of male alcoholics. *Drug and Alcohol Dependence*, 48(3), 243–250. doi: 10.1016/S0376-8716(97)00130-0 [PubMed: 9449024]
- Higginson CI, Wheelock VL, Levine D, King DS, Pappas CTE, & Sigvardt KA (2008). Cognitive deficits in essential tremor consistent with frontosubcortical dysfunction. *Journal of Clinical and Experimental Neuropsychology*, 30(7), 760–765. doi: 10.1080/13803390701754738 [PubMed: 18608666]
- Janicki SC, Cosentino S, & Louis ED (2013). The cognitive side of essential tremor: what are the therapeutic implications? *Therapeutic Advances in Neurological Disorders*, 6(6), 353–368. doi: 10.1177/1756285613489591 [PubMed: 24228071]
- Kaplan E, Goodglass H, & Weintraub S (1983). *Boston Naming Test*. Philadelphia, PA: Lea & Febiger.
- Kim J-S, Song I-U, Shim Y-S, Park J-W, Yoo J-Y, Kim Y-I, & Lee K-S (2009). Cognitive Impairment in Essential Tremor without Dementia. *Journal of Clinical Neurology*, 5(2), 81. doi:10.3988/jcn.2009.5.2.81 [PubMed: 19587814]

- LaRoia H, & Louis ED (2011). Association between essential tremor and other neurodegenerative diseases: what is the epidemiological evidence? *Neuroepidemiology*, 37(1), 1–10. doi: 10.1159/000328866 [PubMed: 21757958]
- Lawton MP, & Brody EM (1969). Assessment of Older People: Self-Maintaining and Instrumental Activities of Daily Living. *The Gerontologist*, 9(3), 179–186. doi:10.1093/geront/9.3_Part_1.179 [PubMed: 5349366]
- Lombardi WJ, Woolston DJ, Roberts JW, & Gross RE (2001). Cognitive deficits in patients with essential tremor. *Neurology*, 57(5), 785–790. doi:10.1212/WNL.57.5.785 [PubMed: 11552004]
- Louis ED (2010). Functional correlates of lower cognitive test scores in essential tremor. *Movement Disorders*, 25(4), 481–485. doi:10.1002/mds.22920 [PubMed: 20108374]
- Louis ED (2016). Linking Essential Tremor to the Cerebellum: Neuropathological Evidence. *The Cerebellum*, 15(3), 235–242. doi:10.1007/s12311-015-0692-6 [PubMed: 26129713]
- Louis ED, Babij R, Ma K, Cortés E, & Vonsattel J-PG (2013). Essential tremor followed by progressive supranuclear palsy: postmortem reports of 11 patients. *Journal of Neuropathology and Experimental Neurology*, 72(1), 8–17. doi:10.1097/NEN.0b013e31827ae56e [PubMed: 23242279]
- Louis ED, Benito-León J, Vega-Quiroga S, Bermejo-Pareja F, & Neurological Disorders in Central Spain (NEDICES) Study Group. (2010a). Cognitive and motor functional activity in non-demented community-dwelling essential tremor cases. *Journal of Neurology, Neurosurgery & Psychiatry*, 81(9), 997–1001. doi:10.1136/jnnp.2009.202838
- Louis ED, Benito-León J, Vega-Quiroga S, Bermejo-Pareja F, & Neurological Disorders in Central Spain (NEDICES) Study Group. (2010b). Faster rate of cognitive decline in essential tremor cases than controls: a prospective study. *European Journal of Neurology*, 17(10), 1291–1297. doi: 10.1111/j.1468-1331.2010.03122.x [PubMed: 20561042]
- Louis ED, & Ferreira JJ (2010). How common is the most common adult movement disorder? Update on the worldwide prevalence of essential tremor. *Movement Disorders*, 25(5), 534–541. doi: 10.1002/mds.22838 [PubMed: 20175185]
- Louis ED, Ford B, & Bismuth B (1998). Reliability between two observers using a protocol for diagnosing essential tremor. *Movement Disorders*, 13(2), 287–293. doi:10.1002/mds.870130215 [PubMed: 9539343]
- Louis ED, Kellner S, Morgan S, Collins K, Rohl B, Huey ED, & Cosentino S (2017). Cognitive Dysfunction Is Associated with Greater Imbalance and Falls in Essential Tremor. *Frontiers in Neurology*, 8, 154. doi:10.3389/fneur.2017.00154 [PubMed: 28469597]
- Louis ED, Ottman R, Ford B, Pullman S, Martinez M, Fahn S, & Hauser WA (1997). The Washington Heights-Inwood Genetic Study of Essential Tremor: Methodologic Issues in Essential-Tremor Research. *Neuroepidemiology*, 16(3), 124–133. doi:10.1159/000109681 [PubMed: 9159767]
- Louis ED, & Rao AK (2014). Tandem gait performance in essential tremor patients correlates with cognitive function. *Cerebellum & Ataxias*, 1, 19. doi:10.1186/s40673-014-0019-2 [PubMed: 26331043]
- Louis ED, Rao AK, & Gerbin M (2012). Functional correlates of gait and balance difficulty in essential tremor: balance confidence, near misses and falls. *Gait & Posture*, 35(1), 43–47. doi: 10.1016/j.gaitpost.2011.08.002 [PubMed: 21930384]
- Louis ED, Vonsattel JPG, Honig LS, Lawton A, Moskowitz C, Ford B, & Frucht S (2006). Essential Tremor Associated With Pathologic Changes in the Cerebellum. *Archives of Neurology*, 63(8), 1189. doi:10.1001/archneur.63.8.1189 [PubMed: 16908751]
- Louis ED, Wendt KJ, Albert SM, Pullman SL, Yu Q, & Andrews H (1999). Validity of a performance-based test of function in essential tremor. *Archives of Neurology*, 56(7), 841–846. [PubMed: 10404986]
- Mameli F, Tomasini E, Scelzo E, Fumagalli M, Ferrucci R, Bertolasi L, & Priori A (2013). Lies tell the truth about cognitive dysfunction in essential tremor: an experimental deception study with the guilty knowledge task. *Journal of Neurology, Neurosurgery, and Psychiatry*, 84(9), 1008–1013. doi:10.1136/jnnp-2012-304674
- Manly JJ, Touradji P, Tang M-X, & Stern Y (2003). Literacy and Memory Decline Among Ethnically Diverse Elders. *Journal of Clinical and Experimental Neuropsychology*, 25(5), 680–690. doi: 10.1076/jcen.25.5.680.14579 [PubMed: 12815505]

- Middleton FA, & Strick PL (2000a). Basal ganglia and cerebellar loops: motor and cognitive circuits. *Brain Research. Brain Research Reviews*, 31(2–3), 236–250. [PubMed: 10719151]
- Middleton FA, & Strick PL (2000b). Basal Ganglia Output and Cognition: Evidence from Anatomical, Behavioral, and Clinical Studies. *Brain and Cognition*, 42(2), 183–200. doi: 10.1006/brcg.1999.1099 [PubMed: 10744919]
- Middleton FA, & Strick PL (2001). Cerebellar projections to the prefrontal cortex of the primate. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 21(2), 700–712. [PubMed: 11160449]
- Montgomery EB, Baker KB, Lyons K, & Koller WC (2000). Motor initiation and execution in essential tremor and Parkinson's disease. *Movement Disorders*, 15(3), 511–515. doi: 10.1002/1531-8257(200005)15:3<511::AID-MDS1014>3.0.CO;2-R [PubMed: 10830417]
- Morris JC (1993). The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology*, 43(11), 2412–2414.
- Morris JC (1997). Clinical dementia rating: a reliable and valid diagnostic and staging measure for dementia of the Alzheimer type. *International Psychogeriatrics*, 9 Suppl 1, 173–6; discussion 177–8. [PubMed: 9447441]
- Nasreddine ZS, Phillips NA, Bédirian V, Charbonneau S, Whitehead V, Collin I, ... Chertkow H (2005). The Montreal Cognitive Assessment, MoCA: A Brief Screening Tool For Mild Cognitive Impairment. *Journal of the American Geriatrics Society*, 53(4), 695–699. doi:10.1111/j.1532-5415.2005.53221.x [PubMed: 15817019]
- Pan JJ, Lee M, Honig LS, Vonsattel J-PG, Faust PL, & Louis ED (2014). Alzheimer's-related changes in non-demented essential tremor patients vs. controls: Links between tau and tremor? *Parkinsonism & Related Disorders*, 20(6), 655–658. doi:10.1016/j.parkreldis.2014.03.003 [PubMed: 24679899]
- Passamonti L, Novellino F, Cerasa A, Chiriac C, Rocca F, Matina MS, ... Quattrone A (2011). Altered cortical-cerebellar circuits during verbal working memory in essential tremor. *Brain*, 134(8), 2274–2286. doi:10.1093/brain/awr164 [PubMed: 21747127]
- Peretz C, Herman T, Hausdorff JM, & Giladi N (2006). Assessing fear of falling: Can a short version of the Activities-specific Balance Confidence scale be useful? *Movement Disorders*, 21(12), 2101–2105. doi:10.1002/mds.21113 [PubMed: 16991140]
- Salthouse TA (2004). What and When of Cognitive Aging. *Current Directions in Psychological Science*, 13(4), 140–144. doi:10.1111/j.0963-7214.2004.00293.x
- Schmahmann JD, & Caplan D (2006). Cognition, emotion and the cerebellum. *Brain*, 129(2), 290–292. doi:10.1093/brain/awh729 [PubMed: 16434422]
- Sengul Y, Sengul HS, Yucekaya SK, Yucel S, Bakim B, Pazarci NK, & Özdemir G (2015). Cognitive functions, fatigue, depression, anxiety, and sleep disturbances: assessment of nonmotor features in young patients with essential tremor. *Acta Neurologica Belgica*, 115(3), 281–287. doi: 10.1007/s13760-014-0396-6 [PubMed: 25471376]
- Sinoff G, & Badarny S (2014). Mild cognitive impairment, dementia, and affective disorders in essential tremor: a prospective study. *Tremor and Other Hyperkinetic Movements (New York, N.Y.)*, 4, 227. doi:10.7916/D85B00KN
- Smith A (1982). Symbol Digit Modalities Test. Los Angeles, CA: Western Psychological Services.
- Souza CP, Oliveira GN, Foss MP, & Tumas V (2016). Cluster analysis of cognitive performance in a sample of patients with Parkinson's disease. *Dementia & Neuropsychologia*, 10(4), 315–319. doi: 10.1590/s1980-5764-2016dn1004010 [PubMed: 29213475]
- Szeto JY, O'Callaghan C, Shine JM, Walton CC, Mowszowski L, Naismith SL, ... Lewis SJ (2015). The relationships between mild cognitive impairment and phenotype in Parkinson's disease. *Npj Parkinson's Disease*, 1(1), 15015. doi:10.1038/npjparkd.2015.15
- Thawani SP, Schupf N, & Louis ED (2009). Essential tremor is associated with dementia: prospective population-based study in New York. *Neurology*, 73(8), 621–625. doi:10.1212/WNL.0b013e3181b389f1 [PubMed: 19704081]
- Tröster AI, Woods SP, Fields JA, Lyons KE, Pahwa R, Higginson CI, & Koller WC (2002). Neuropsychological deficits in essential tremor: an expression of cerebello-thalamo-cortical pathophysiology? *European Journal of Neurology*, 9(2), 143–151. [PubMed: 11882055]

- Wechsler D (1987). Wechsler Memory Scale - Revised. San Antonio, TX: Psychological Corporation.
- Wechsler D (1997). Manual for the Wechsler Adult Intelligence Scale (3rd *ed.*). San Antonio, TX: Psychological Corporation.
- Wechsler D (2008). Wechsler Memory Scale (4th *ed.*). San Antonio, TX: Pearson.
- Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey M, & Leirer VO (1982). Development and validation of a geriatric depression screening scale: A preliminary report. *Journal of Psychiatric Research*, 17(1), 37–49. doi:10.1016/0022-3956(82)90033-4 [PubMed: 7183759]

Figure 1: Comparisons with three clusters

Cluster domain means; □ = Memory Aggregate; ▨ = Executive Function Aggregate; ▩ = Attention Aggregate; ▤ = Visuospatial Abilities Aggregate; ▦ = Language Aggregate.

Table 1:

Cognitive Domain Scores according to Cognitive Diagnosis

	Group 1 (Fully Normal Cognition, n = 61)	Group 2 (Not-fully Normal Cognition, n = 87)	p-Value ¹
Memory	0.56 ± 0.61	0.02 ± 0.66	<0.001
Executive Function	0.49 ± 0.38	0.06 ± 0.50	<0.001
Attention	0.23 ± 0.64	-0.32 ± 0.66	<0.001
Visuospatial Abilities	0.70 ± 0.62	0.53 ± 0.64	0.12
Language	0.27 ± 0.38	0.60 ± 0.63	0.02

Values are reported as mean ± standard deviation (SD).

¹ 2-sample t-test. Threshold corrected for multiple comparisons using Bonferroni correction; p < 0.01 is considered to be significant.

Table 2:

Comparisons with three clusters

Domain	Cluster ^J			ANOVA p-Value	Post-Hoc p-Value (Bonferroni)		
	1	2	3				
	N = 64	N = 39	N = 45				
Memory	-0.31 (-1.31 – 0.41) Average	0.78 (0.03 – 2.27) High average	0.56 (-0.71 – 1.77) High average	<0.001	1 to 2 1 to 3 2 to 3	<0.001 <0.001 0.13	
Executive Function	0.08 (-1.13 – 0.85) Average	0.06 (-1.10 – 0.95) Average	0.60 (-0.26 – 1.29) High average	<0.001	1 to 2 1 to 3 2 to 3	1.00 <0.001 <0.001	
Attention	-0.29 (-1.45 – 0.80) Average	-0.57 (-1.73 – 0.67) Low average	0.60 (-0.49 – 2.00) High average	<0.001	1 to 2 1 to 3 2 to 3	0.027 <0.001 <0.001	
Visuospatial Abilities	0.48 (-1.04 – 1.95) Average	0.20 (-0.75 – 0.97) Average	1.12 (-0.16 – 2.24) High average	<0.001	1 to 2 1 to 3 2 to 3	0.029 <0.001 <0.001	
Language	0.16 (-1.01 – 1.26) Average	0.04 (-1.49 – 0.82) Average	0.27 (-0.80 – 1.08) Average	0.09	1 to 2 1 to 3 2 to 3	0.65 0.73 0.09	
Significant p-Values (indicated in bold)				15 (75.0%)			

^JValues are reported as mean domain score (range) per cluster.

Table 3:

Cognitive profiles by neuropsychological test scores

Domain	Test	Sub-score	All Cases (n = 148)	Cluster			Kruskal-Wallis H-Test p-Value	Post-hoc Mann-Whitney U-Test p-Value		
				1 (n = 64)	2 (n = 39)	3 (n = 45)		1 to 2	1 to 3	2 to 3
Memory	California Verbal Learning Test	Total Recall	0.09 ± 1.24	-0.48 ± 1.05	0.51 ± 1.07	0.52 ± 1.30	<0.001	<0.001	<0.001	<0.001
		Long-Delay Free Recall	0.01 ± 1.03	-0.58 ± 0.84	0.45 ± 0.81	0.46 ± 1.05	<0.001	<0.001	<0.001	0.84
		Immediate	0.67 ± 0.92	0.17 ± 0.82	1.12 ± 0.85	0.99 ± 0.77	<0.001	<0.001	<0.001	0.53
	Verbal Paired Associates	Delayed	0.54 ± 0.99	0.02 ± 0.97	1.08 ± 0.82	0.81 ± 0.81	<0.001	<0.001	<0.001	0.56
		Immediate	0.13 ± 0.95	-0.34 ± 0.83	0.68 ± 0.92	0.30 ± 0.83	<0.001	<0.001	<0.001	0.10
		Delayed	0.16 ± 0.91	-0.30 ± 0.69	0.80 ± 0.81	0.28 ± 0.91	<0.001	<0.001	<0.001	0.07
	Verbal Fluency Test	Letter	0.30 ± 1.08	0.15 ± 1.07	0.14 ± 0.98	0.64 ± 1.12	0.03	1 to 2	1 to 3	0.71
		Category	0.34 ± 1.04	0.10 ± 0.85	0.29 ± 1.16	0.72 ± 1.09	0.01	1 to 2	1 to 3	0.01
		Switching Fluency	0.27 ± 1.13	0.07 ± 0.97	0.04 ± 1.35	0.76 ± 1.02	0.005	1 to 2	1 to 3	0.043
	Executive Function	Letter	0.30 ± 1.08	0.15 ± 1.07	0.14 ± 0.98	0.64 ± 1.12	0.03	1 to 2	1 to 3	0.71
		Category	0.34 ± 1.04	0.10 ± 0.85	0.29 ± 1.16	0.72 ± 1.09	0.01	1 to 2	1 to 3	0.01
		Switching Fluency	0.27 ± 1.13	0.07 ± 0.97	0.04 ± 1.35	0.76 ± 1.02	0.005	1 to 2	1 to 3	0.043

Domain	Test	Sub-score	All Cases (n = 148)	Cluster			Kruskal-Wallis H-Test p-Value	Post-hoc Mann-Whitney U-Test p-Value	
				1 (n = 64)	2 (n = 39)	3 (n = 45)		1 to 3	2 to 3
		Switching Accuracy	0.31 ± 1.10	0.19 ± 0.96	0.03 ± 1.28	0.72 ± 1.03	0.01	1 to 2	0.71
								1 to 3	0.005
								2 to 3	0.013
	Color-Word Inhibition	Naming	-0.29 ± 1.02	-0.43 ± 0.89	-0.59 ± 1.18	0.17 ± 0.91	0.001	1 to 2	0.66
								1 to 3	<0.001
		Reading	-0.16 ± 0.95	-0.26 ± 1.05	-0.31 ± 0.92	0.13 ± 0.74	0.07	2 to 3	0.003
								1 to 2	0.63
								1 to 3	0.07
		Inhibition	0.25 ± 0.89	0.21 ± 0.77	-0.17 ± 1.14	0.65 ± 0.60	0.001	2 to 3	0.030
								1 to 2	0.20
	Sorting	Inhibition/Switching	0.27 ± 0.95	0.15 ± 1.05	0.07 ± 0.93	0.63 ± 0.73	0.02	1 to 3	0.003
								2 to 3	0.001
		Confirmed Correct Sorts	0.43 ± 0.75	0.27 ± 0.66	0.39 ± 0.76	0.71 ± 0.79	0.02	1 to 2	0.55
								1 to 3	0.022
								2 to 3	0.008
		Free Sort Description	0.25 ± 0.84	0.09 ± 0.77	0.15 ± 0.82	0.57 ± 0.87	0.02	1 to 2	0.81
								1 to 3	0.007
	Recognition Description	0.06 ± 0.95	-0.09 ± 0.94	-0.13 ± 0.81	0.45 ± 0.98	0.01	0.01	2 to 3	0.034
								1 to 2	0.85
								1 to 3	0.007
	20-Questions	Initial Abstraction Score	0.41 ± 1.03	0.28 ± 1.05	0.38 ± 0.90	0.62 ± 1.08	0.22	2 to 3	0.009
								1 to 2	0.65

Domain	Test	Sub-score	All Cases (n = 148)	Cluster			Kruskal-Hallis H-Test p-Value ¹	Post-hoc Mann-Whitney U-Test p-Value	
				1 (n = 64)	2 (n = 39)	3 (n = 45)		1 to 3	1 to 2
Attention								1 to 3	0.10
								2 to 3	0.22
		Total Questions Asked	0.55 ± 0.94	0.38 ± 0.98	0.55 ± 1.00	0.79 ± 0.77	0.02	1 to 2	0.14
		Weighted Achievement Score	0.57 ± 1.07	0.30 ± 1.17	0.47 ± 1.05	1.01 ± 0.80	0.001	1 to 2	0.59
	Digit Span Backwards		0.07 ± 1.00	-0.06 ± 0.88	-0.30 ± 1.10	0.57 ± 0.90	<0.001	1 to 2	0.14
								1 to 3	0.001
								2 to 3	<0.001
		Oral Symbol Digit Modalities Test	0.28 ± 0.85	0.05 ± 0.77	0.02 ± 0.86	0.83 ± 0.70	<0.001	1 to 2	0.52
								1 to 3	<0.001
								2 to 3	<0.001
Digit Span Forwards		-0.47 ± 1.10	-0.63 ± 0.93	-1.15 ± 0.82	0.37 ± 1.06	<0.001	1 to 2	0.006	
							1 to 3	<0.001	
							2 to 3	<0.001	
Visuospatial Abilities	Benton Judgment of Line Orientation		0.66 ± 0.90	0.49 ± 0.92	0.34 ± 0.85	1.18 ± 0.67	<0.001	1 to 2	0.36
								1 to 3	<0.001
	Benton Facial Recognition Test		0.92 ± 1.06	0.81 ± 0.97	0.33 ± 1.07	1.59 ± 0.76	<0.001	1 to 2	0.023
								1 to 3	<0.001
Visual Puzzles			0.24 ± 0.89	0.15 ± 0.89	-0.10 ± 0.63	0.61 ± 0.95	0.001	1 to 2	0.24
								1 to 3	<0.001
	Multilingual-Aphasia Examination Token Test		0.60 ± 0.42	0.62 ± 0.40	0.38 ± 0.45	0.76 ± 0.34	<0.001	2 to 3	<0.001
								1 to 2	0.011

Domain	Test	Sub-score	All Cases (n = 148)	Cluster			Kruskal-Wallis H-Test p-Value <i>I</i>	Post-hoc Mann-Whitney U-Test p-Value	
				1 (n = 64)	2 (n = 39)	3 (n = 45)			
	Boston Naming Test		-0.27 ± 0.78	-0.30 ± 0.85	-0.29 ± 0.78	-0.23 ± 0.68	0.96	1 to 3	0.08
								2 to 3	<0.001
								1 to 2	0.92
								1 to 3	0.80
								2 to 3	0.84

Values are reported as Mean ± SD.

Threshold corrected for multiple comparisons using Bonferroni correction; p < 0.0083 is considered to be significant.

Table 4:

Demographic and clinical characteristics

	All Cases (n = 148)	Cluster			Among group p- Value	Between group p-Value ²	
		1 (n = 64)	2 (n = 39)	3 (n = 45)			
Demographics	Age	78.2 ± 9.5	79.7 ± 9.6	77.2 ± 9.1	0.30 ^a	1 to 2	0.17 ^c
						1 to 3	0.24 ^c
						2 to 3	0.72 ^c
Gender (female)	95 (64.1)	38 (59.4)	31 (79.5)	26 (57.8)	0.08 ^b	1 to 2	0.04 ^b
						1 to 3	0.87 ^b
						2 to 3	0.03 ^b
Education	15.9 ± 2.5	15.8 ± 2.8	16.0 ± 2.4	16.2 ± 2.3	0.77 ^a	1 to 2	0.72 ^c
						1 to 3	0.50 ^c
						2 to 3	0.67 ^c
Hispanic	2 (1.4)	1 (1.6)	0 (0.0)	1 (2.2)	0.47 ^b	1 to 2	0.43 ^b
						1 to 3	0.80 ^b
						2 to 3	0.35 ^b
Total tremor Score	19.9 ± 4.9	19.6 ± 4.5	20.6 ± 5.9	19.7 ± 4.5	0.67 ^a	1 to 2	0.44 ^c
						1 to 3	0.76 ^c
						2 to 3	0.44 ^c
Tremor Disability Scale	67.5 ± 26.1	66.5 ± 28.7	70.1 ± 22.2	66.8 ± 25.6	0.89 ^a	1 to 2	0.74 ^c
						1 to 3	0.86 ^c
						2 to 3	0.63 ^c
Tremor Duration	37.8 ± 21.9	40.6 ± 23.6	38.5 ± 22.1	33.3 ± 18.7	0.30 ^a	1 to 2	0.67 ^c

	All Cases (n = 148)	Cluster			Among group p- Value ^f	Between group p-Value ²	
		1 (n = 64)	2 (n = 39)	3 (n = 45)			
Functional Characteristics						1 to 3 0.11 ^c	
						2 to 3 0.38 ^c	
	Head Tremor (present on examination)	93 (62.8)	43 (71.9)	28 (71.8)	22 (48.9)	0.06 ^b	1 to 2 0.62 ^b
						1 to 3 0.06 ^b	
						2 to 3 0.03 ^b	
	Voice Tremor (present on examination)	73 (49.3)	30 (46.9)	21 (53.8)	22 (48.9)	0.82 ^b	1 to 2 0.49 ^b
						1 to 3 0.84 ^b	
						2 to 3 0.65 ^b	
	Number of Medications	5.3 ± 3.9	6.4 ± 4.7	4.5 ± 2.9	4.2 ± 3.2	0.015 ^a	1 to 2 0.04 ^c
						1 to 3 0.008 ^c	
Balance						2 to 3 0.49 ^c	
	Taking Cognition-Enhancing Medications	1 (0.7)	1 (1.6)	0 (0.0)	0 (0.0)	0.52 ^b	1 to 2 0.43 ^b
						1 to 3 0.40 ^b	
						2 to 3 1.00 ^b	
	Taking Cognition-Decreasing Medications	72 (48.6)	32 (50.0)	21 (53.8)	19 (42.2)	0.47 ^b	1 to 2 0.71 ^b
						1 to 3 0.42 ^b	
						2 to 3 0.29 ^b	
	Lawton Instrumental Activities of Daily Living (IADL)	7.5 ± 1.1	7.6 ± 0.8	7.5 ± 1.3	7.4 ± 1.3	0.91 ^a	1 to 2 0.67 ^c
						1 to 3 0.91 ^c	
						2 to 3 0.68 ^c	
Number of steps off line in Tandem Gait	4.5 ± 4.0	5.2 ± 4.0	4.0 ± 4.1	3.9 ± 4.0	0.14 ^a	1 to 2 0.11 ^c	

	All Cases (n = 148)	Cluster			Among group p- Value ^c	Between group p-Value ²
		1 (n = 64)	2 (n = 39)	3 (n = 45)		
Cognitive Characteristics						1 to 3 0.09 ^c
						2 to 3 0.84 ^c
	Activities-specific Balance Confidence Scale (ABC-6)	56.3 ± 28.0	54.7 ± 29.6	65.2 ± 26.1	0.028^a	1 to 2 0.50 ^c
						1 to 3 0.006^c
						2 to 3 0.12 ^c
	Falls in Past Year	1.2 ± 3.5	1.0 ± 2.0	0.6 ± 1.2	0.57 ^a	1 to 2 0.75 ^c
						1 to 3 0.29 ^c
						2 to 3 0.52 ^c
	Mini-Mental State Examination (MMSE)	29.0 ± 1.3	29.0 ± 1.5	29.1 ± 1.2	0.81 ^a	1 to 2 0.56 ^c
						1 to 3 0.65 ^c
Psychological Characteristics						2 to 3 0.81 ^c
	Montreal Cognitive Assessment (MoCA)	25.7 ± 2.6	25.9 ± 2.6	26.9 ± 2.1	<0.001^a	1 to 2 0.02 ^c
						1 to 3 <0.001^c
						2 to 3 0.11 ^c
	Clinical Dementia Rating (CDR, score = 0.5)	14 (9.5)	3 (7.7)	4 (8.9)	0.69 ^b	1 to 2 0.59 ^b
						1 to 3 0.73 ^b
						2 to 3 0.84 ^b
	Geriatric Depression Scale (GDS)	6.0 ± 4.8	5.9 ± 5.0	5.7 ± 4.7	0.77 ^a	1 to 2 0.55 ^c
						1 to 3 0.54 ^c
						2 to 3 0.97 ^c
	Taking Mood-Modulating Medications	30 (20.3)	10 (25.6)	8 (17.8)	0.96 ^b	1 to 2 0.69 ^b

	All Cases (n = 148)	Cluster			Among group p- Value ^c	Between group p-Value ²	
		1 (n = 64)	2 (n = 39)	3 (n = 45)		1 to 3	2 to 3
						1 to 3	0.017 ^b
						2 to 3	0.77 ^b

All continuous values are given as mean ± standard deviation (SD); all categorical values are given as number (percentage).

^aKruskall-Wallis H-test

^bPearson Chi-Square Test

^cMann-Whitney U-Test

¹Significant if p < 0.05

²Significant if p < 0.0167.

Table 5:

Cognitively normal sub-diagnoses across clusters

Diagnostic Category	All Cases	Cluster ¹		
		1	2	3
Fully Normal Cognition	61	17 (27.9)	14 (22.9)	30 (49.2)
Test impairment with unlikely clinical significance	45	23 (51.1)	11 (24.4)	11 (24.4)
Test impairment with possible clinical significance	39	24 (61.5)	13 (33.3)	2 (5.1)
Questionable or isolated functional impairment	3	0 (0.0)	1 (33.3)	2 (66.7)
Fisher Exact Test p-Value: <0.001				

Values are given as number (percentage of diagnostic category).