Published in final edited form as:

Drug Alcohol Depend. 2019 January 01; 194: 45–50. doi:10.1016/j.drugalcdep.2018.09.012.

Prevalence, incidence, recovery, and recurrence of alcohol use disorders from childhood to age 30

John R. Seeley^{a,b}, Richard F. Farmer^b, Derek B. Kosty^b, Jeff M. Gau^b

^aUniversity of Oregon, College of Education, 1215 University St., Eugene, OR, 97403, USA

^bOregon Research Institute, 1776 Millrace Drive, Eugene, OR, 97403, USA

Abstract

Background: Little is known about the course of alcohol use disorders (AUDs) in representative samples during high-risk periods of adolescence and early adulthood. The primary objective of this research is to describe the prevalence and course of initial AUD episodes experienced between childhood and age 30 in a regionally representative cohort sample.

Methods: Study data are from an epidemiological study of 816 youth. Participants were initially selected at random from nine high schools in western Oregon, USA. Four waves of data collection were conducted between ages 16 and 30. AUD course milestones are referenced to participants' age.

Results: Results indicated that male participants (43%) were significantly more likely to be diagnosed with a lifetime AUD than female participants (28%), *OR* [*CI*₉₅] = 1.97 [1.47–2.65], and rate of first incidence was especially high between ages 18 and 24.9, a developmental period that also corresponded to the peak interval in prevalence rates. The rate of first AUD incidence substantially diminished beginning around age 25. Among those with an initial AUD episode, 87% recovered by age 30 and, of these, the average episode length was 23 months. Among recovered cases, 33% went on to experience a second AUD episode (i.e., a recurrence) after a minimum 12-month asymptomatic recovery period. Risk for recurrence remained relatively high within the 5 years following initial AUD offset.

Conclusions: AUDs are common lifetime conditions in representative samples, whereby most affected individuals by age 30 experience a time-limited course rather than a recurring or persistent course.

Correspondence: John R. Seeley, College of Education, University of Oregon, 1215 University St., Eugene, OR 97403-1215, jseeley@uoregon.edu.

Contributors

JRS had the initial study idea and was involved in the original study protocol and data collection. RFF conducted background literature searches. DBK and JMG programmed and conducted statistical analyses. JRS, RFF and DBK wrote the first draft of the manuscript. JRS, RFF, DBK and JMG contributed to interpreting the findings and writing further drafts of the manuscript. All authors reviewed and have approved the final manuscript.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest
No conflict declared.

Keywords

Alcohol Use Disorders; Natural Course; Onset; Recovery; Recurrence; Sex Differences

1. Introduction

Alcohol use disorders (AUDs) are prevalent globally, especially in western countries. Within the United States (U.S.), the cross-sectional 2001–2002 National Epidemiologic Survey on Alcohol and Related Conditions (NESARC-I) estimated the lifetime rate of AUD based on DSM-IV criteria to be 30% (Hasin et al., 2007). NESARC-III, a cross-sectional survey conducted during 2012–2013 and based on DSM-5 criteria, produced a 29% lifetime AUD rate estimate (Grant et al., 2015). Lifetime prevalence rates derived from multiple assessments with prospective longitudinal cohort samples also indicate that AUDs are common. In the Dunedin Multidisciplinary Health and Development Study, the lifetime rate of alcohol dependence based on DSM-III-R and DSM-IV criteria was 32% between ages 18 and 32 (Meier et al., 2013; Moffitt et al., 2010). In the Christchurch Health and Development Study (Fergusson and Horwood, 2001), lifetime rates of AUD from ages 15 to 35 based on DSM-III-R and DSM-IV criteria were 56% and 40% for male and female cohorts, respectively, or about 48% overall (Joseph M. Boden, personal communication, 2 July 2017). In the Zurich Cohort Study of Young Adults, which spanned ages 18 to 50, the estimated lifetime rate of AUD based on DSM-III, DSM-III-R, and DSM-IV was 29% (Angst et al., 2016). We (Farmer et al., 2016) earlier reported a 34% lifetime rate of DSM-IV AUD from childhood to age 30 for the Oregon Adolescent Depression Project (OADP) cohort. Despite the regional and methodological differences that characterize these studies, the overarching conclusion remains that AUDs are common in communities within Australasia, Europe, and North America.

AUDs, as defined in DSM-IV or earlier editions, typically first emerge and increase in prevalence during adolescence, peak during late adolescence and early adulthood, and gradually decline thereafter (Sher et al., 2005). Factors associated with the rapid escalation of AUD prevalence within this span include increased alcohol availability (Komro et al., 2007), reduced parental monitoring (Guilamo-Ramos et al., 2005), and peer influences (Patrick et al., 2016). Reductions in prevalence after early adulthood are often attributed to “maturing out” and the assumption of adult life roles incompatible with excessive alcohol consumption (Sher et al., 2005). A better understanding of AUD course in representative samples, particularly during the high-risk periods spanning childhood through young adulthood, is especially important given the adverse health, social, economic, and legal consequences that often accompany AUDs (Brown et al., 2000; Parry et al., 2011; Rehm et al., 2009; Whiteford et al., 2013) which may be difficult to overcome later in life.

Views on the course of AUD have largely been shaped by studies of treated or high-risk samples, leading some to conclude that AUDs are best characterized as a chronic medical illness punctuated by alternating periods of remission and relapse (McLellan et al., 2000). A different view on the course of AUDs, however, is emerging from representative and population samples. Remission and recovery concepts in these studies have been defined

variously as (a) the point at which an individual no longer meets diagnostic criteria for AUD, (b) consumption of alcohol at an amount below some threshold level of risk, or (c) complete abstinence from alcohol. Overall, these studies suggest that AUDs typically have a relatively brief course, with a large majority of baseline positive cases in remission or recovery within three years (Boschloo et al., 2012; Dawson et al., 2012; de Bruijn et al., 2006; Hasin et al., 2011; Tuithof et al., 2013). Although a large proportion of persons diagnosed with an AUD recover without participation in a treatment program (Dawson et al., 2005; de Bruijn et al., 2006; Sobell et al., 1996), those who report more severe alcohol-related problems are less likely to recover (Cunningham, 1999; de Bruijn et al., 2006).

Relapse or recurrence following a period of sustained remission represents a significant challenge to lasting change. Research on the recurrence of AUD following remission or recovery in representative samples is rare (Tuithof et al., 2014). A small number of such studies suggests that relapse or recurrence rates for remitted or recovered cases may be less than 15% within a 3 to 20 years follow-up interval (Dawson et al., 2007; de Bruijn et al., 2006; Tuithof et al., 2014). Altogether, the relatively few studies with representative prospective samples suggest that AUDs are common, generally of brief duration, and rarely recurrent.

Most research on the course of AUDs is based on treated samples, within which disorder severity and comorbid psychopathology are often substantially greater than in representative samples (Low et al., 2008). This investigation into the natural course of AUDs is based on data collected as part of the OADP, a longitudinal study of a regionally representative community sample. Previous OADP publications have presented AUD rates and sex distributions at ages 16 and 17 (Rohde et al., 1996), at age 24 (Rohde et al., 2001), and at age 30 (Farmer et al., 2016). The present research extends our earlier work by presenting first incidence and period prevalence data from childhood through age 30. We also examine several course indicators that signify milestones in change processes not previously reported in earlier publications, including time to onset, recovery, and recurrence when referenced to the initial (first) AUD episode. Additionally, limited indications suggest that females experience a course of AUD that is distinct from that of males, although relevant research with representative samples is sparse (e.g., Edens et al., 2008; Merikangas and McClair, 2012). We consequently conduct sex distribution comparisons of AUD for each time period or interval evaluated and in the cumulative hazard functions that describe various course indicators over time.

2. Materials and methods

2.1 Participants

OADP participants were assessed on four occasions between ages 16 and 30 (T_1 - T_4). The T_1 sample consisted of 1,709 adolescents randomly selected from 9 high schools representative of urban and rural districts in western Oregon (Lewinsohn et al., 1993). About one year later (T_2), 1,507 (88%) were again evaluated. At T_3 (7 years after T_2), a sampling stratification procedure was introduced whereby eligible participants included all persons with a history of a psychiatric diagnosis by T_2 ($n = 644$) and a randomly selected subset of participants with no mental disorder history (NMD) by T_2 ($n = 457$ of 863 persons). Comparisons

between the T₃ NMD randomly selected participants with unselected NMD participants revealed no significant differences with respect to T₂ data. Of these 1,101 eligible persons, 941 (85%) completed T₃. Among those who completed T₃, 816 (87%) completed the T₄ assessment at age 30 (59% female; 89% white; 53% married; 43% at least a bachelor's degree).

An analysis of attrition included comparisons between T₄ participants to those who dropped out after T₁ with respect to a lifetime diagnosis of a DSM-defined disorder and the summated number of distinct lifetime psychiatric disorders by T₁ (Farmer et al., 2013). The T₄ panel was not statistically different from the attrition group with respect to any psychiatric disorder history ($p = .96$) or the sum of all lifetime disorders ($p = .23$) at T₁. An attrition analysis conducted for this report found that those in the attrition group, when compared to the T₄ panel, did not have significantly higher rates of AUDs at T₁ (7% vs. 5%, respectively). Across-wave analyses, however, revealed one significant difference whereby discontinuation after T₃ was more common among those with an AUD history by T₃ (lifetime AUD rates: 41% for discontinuers, 31% for T₃ participants; Pearson $\chi^2 [1, n = 941] = 4.88, p = .027$).

Other differences across waves were noted between study continuers and dropouts. When compared to the T₂ sample, discontinuers after T₁ were more likely male, from smaller sized households, socioeconomically challenged, tobacco users, and more likely to have histories of disruptive behavior disorders and substance use disorders (SUDs) undifferentiated with respect to substance type (Lewinsohn et al., 1993). From T₂ to T₃ and T₃ to T₄, women were more likely to continue participation than men (Lewinsohn et al., 1999; Rohde et al., 2007). From T₃ to T₄, discontinuers had a higher rate of SUD undifferentiated with respect to substance. Because of the sample stratification procedures implemented at T₃ and the relatively modest attrition across waves, results presented below are based on the T₄ panel ($N = 816$).

2.2 AUD assessments

From T₁ through T₃, participants were interviewed with the Schedule for Affective Disorders and Schizophrenia for School-Age Children (K-SADS) that combined features of the Epidemiologic and Present Episode versions (Chambers et al., 1985; Orvaschel et al., 1982). T₂ and T₃ assessments also involved the administration of the Longitudinal Interval Follow-Up Evaluation (LIFE; Keller et al., 1987) that, in conjunction with the K-SADS, provided detailed information concerning the presence and course of disorders since the previous interview. The T₄ assessment included an administration of the LIFE and the Structured Clinical Interview for Axis I DSM-IV Disorders– Non-Patient Edition (First et al., 1994). Sufficient additional symptom information was collected during T₁ and T₂ to permit DSM-IV-based evaluations of substance-specific SUD categories (Rohde et al., 2007). Consequently, AUD diagnoses are based on DSM-IV diagnostic criteria. Information was collected at each assessment regarding treatment utilization for AUD, which was defined as outpatient treatment (individual, family, or alcohol support groups), inpatient treatment, hospitalizations, and medications. Agreement among raters for AUD diagnoses since the previous interview, as indexed by kappa (κ), was good to excellent (κ s: T₁ = .76,

$T_2 = .89$, $T_3 = .69$, $T_4 = .79$). When both raters agreed on the occurrence of an initial AUD episode, the intraclass correlation coefficient (ICC) associated with duration judgments (in months) was .89. ICCs corresponding to rater agreement for age (in months) of disorder onsets and offsets in instances where both raters agreed on an initial AUD episode occurrence were similarly high (ICCs = .92 and .87 for disorder onsets and offsets, respectively).

2.3 AUD recovery and recurrence defined

Definitions of recovery and recurrence used in this study are informed by earlier descriptions of these concepts (Chung and Maisto, 2006; Frank et al., 1991), LIFE interview naming conventions (Keller et al., 1987), and DSM-IV guidelines. In the present research, *remission* refers to offset of an initial AUD episode lasting at least 1 full month but < 12 months during which the participant did not endorse any diagnostic criteria for AUD. During this remission period, however, participants may have used alcohol in the absence of alcohol-related problems as defined in AUD diagnostic criteria. The re-emergence of any AUD symptomatology during the remission period was regarded as a continuation of the initial episode (i.e., a *relapse*). Initial episode resolution, defined by a period of uninterrupted remission of at least 12 months, is considered a *recovery* from the initial episode. A *recurrence* is regarded as the emergence of a new AUD episode after a period of recovery.

2.4 Analyses

Estimates of prevalence rates, incidence rates, and odds ratios (*ORs*) accounted for the unequal stratified sampling procedure implemented at T_3 with a case weighting procedure. Age ranges used when reporting first incidence and period prevalence rates were informed by Arnett's (2007) framework: childhood through emerging adolescence (childhood to age 13.9), adolescence (ages 14.0–17.9), adolescence transitioning to emerging adulthood (ages 18.0–24.9), and emerging adulthood transitioning into young adulthood (ages 25.0–30.0).

Time-to-event analyses were implemented with Mplus statistical software (Muthén and Muthén, 1998–2012). Cumulative hazard functions were used to describe AUD onset, recovery, and recurrence functions in the presence of censorship (i.e., participants who do not experience an AUD onset, recovery, or recurrence between childhood and age 30). Time-to-event was measured in months. When cumulative hazard functions exceeded 0.5, the median survival time was reported to facilitate data interpretation.

Sex moderation of the time-to-event functions was evaluated with Cox proportional hazard (PH) models. These models assume the absence of a significant time-by-predictor interaction. Preliminary analyses involved the inclusion of the interaction term in the model as recommended by Singer and Willett (1991). On no occasion was the interaction term statistically significant (all $ps > .05$), indicating that this model assumption was upheld. All models were subsequently rerun with the interaction term removed. Departures in onset curves as a function of sex were derived from hazard ratio (*HR*) estimates and accompanying 95% confidence intervals (CI_{95}).

3. Results

3.1 Prevalence rates, incidence rates, and age of onset for initial AUD episodes

3.1.1 Prevalence and incidence rates.—The weighted lifetime prevalence of AUD from childhood to age 30 was 34.3%. By T₄, men (42.9%) were more likely than women (27.6%) to be diagnosed with a lifetime AUD ($OR [CI_{95}] = 1.97 [1.47–2.65]$). Only 15.3% of those with a lifetime AUD reported receiving treatment for AUD.

Table 1 presents weighted first incidence and period prevalence rate information. AUD first incidence rates were especially high between ages 18.0 to 24.9 and substantially diminished after age 25. Ages 18.0 to 24.9 also corresponded to the peak AUD prevalence period for both men and women. *First incidence rates* significantly differed by sex within the 18.0 to 24.9 period only (29.4% of males, 16.3% of females; $OR [CI_{95}] = 2.14 [1.53–3.00]$). *Period prevalence rates* significantly differed by sex within the 18.0 to 24.9 age range (36.7% of males, 22.6% of females; $OR [CI_{95}] = 1.99 [1.46–2.70]$) and the 25.0 to 30.0 range (21.9% of males, 12.3% of females; $OR [CI_{95}] = 2.00 [1.37–2.91]$).

3.1.2 Time to AUD onset.—Among participants with a lifetime AUD, the mean onset age for the initial episode was 20.3 years ($SD = 3.8$). Mean onset ages did not significantly differ by sex (observed Ms : males = 20.6, $SD = 3.4$; females = 19.9). Cumulative hazard functions for AUD onset for the entire sample and separately by sex are presented in Figure 1. The cumulative hazard functions significantly differed by sex ($HR [CI_{95}] = 1.63, [1.30–2.03]$, $p < .001$), with males compared to females demonstrating a steeper rise and consistently greater cumulative rates beginning around age 18.

3.2 Recovery and non-recovery from the initial AUD episode

3.2.1 Rates of recovery.—For those with a lifetime AUD, 87.1% experienced recovery from the initial episode by age 30. Recovery rates did not significantly differ by sex (84.9% for males, 89.8% for females).

3.2.2 Episode duration and time to recovery.—The mean duration of the initial AUD episode for those who recovered was 23.1 months ($SD = 23.3$). A statistically significant sex difference in episode duration means was observed (males = 28.2 months, $SD = 27.2$; females = 17.3 months, $SD = 16.1$; Welch's $t[215] = -3.87$, $p < .001$, $\eta^2 = .055$).

Time-to-recovery estimates were based on the duration of the initial AUD episode plus a 12-month period of sustained remission without relapse following episode offset. Participants were right censored from the analysis if a 12-month period of sustained remission following offset of the initial AUD episode was not achieved before age 30. Cumulative hazard functions for recovery for the full subsample with AUD histories and separately by sex within this subsample are presented in Figure 2. For those who recovered, the mean time to recovery from the initial episode was 34.9 months ($SD = 23.3$). A statistically significant sex difference in mean time to recovery was observed (males = 40.0 months, $SD = 27.2$; females = 29.1 months, $SD = 16.1$; Welch's $t[215] = -3.89$, $p < .001$, $\eta^2 = .055$). Treatment utilization was not significantly related to time to recovery ($p = .598$).

3.2.3 Episode duration for those who did not recover from the initial episode.

—In the weighted sample, 18% of those with one AUD episode remained in episode at age 30. The mean duration of that episode for these individuals through age 30 was 59.2 months ($SD = 42.2$), which was significantly longer ($t = 4.87$, $p < .001$) than the episode length for those with a single AUD episode and who achieved recovery ($M = 23.9$ months, $SD = 22.2$).

3.3 Recurrence after the initial AUD episode

3.3.1 Recurrence rates following recovery.—Among participants who recovered from the initial AUD episode, 32.8% subsequently developed another AUD episode prior to age 30. Rates of recurrence were not significantly different between males (33.8%) and females (31.7%).

3.3.2 Time to recurrence.—For those who experienced a second AUD episode, the mean time to recurrence was 49.2 months ($SD = 28.0$). Time to recurrence did not significantly differ between males (46.0 months) and females (53.0 months). Cumulative hazard functions are presented in Figure 3 for time-to-recurrence for the full subsample as well as separately by sex for those who experienced a recovery from the initial AUD episode by age 30. Cumulative hazard functions that assessed time to recurrence did not differ by sex or AUD treatment received during the index episode. As depicted in Figure 3, recurrence rates began to level off about 60 months after recovery, with recurrence becoming relatively rare after the 75th consecutive month after offset of the initial episode. Overall, findings reported in Figure 3 suggest that AUD recurrence before age 30 takes place among a minority of those with a previous episode. An asymptomatic 12-month period following initial AUD offset, however, appears not to be an optimal threshold for denoting full recovery from AUDs. Rather, recurrence risk continues to be relatively high after 5 years of remission and recovery.

4. Discussion

Data on the natural course of AUDs can inform judgments about the expected prognosis of AUDs over time, influence theoretical conceptualizations of the nature of AUDs, inform the timing and design of preventive, treatment, and relapse reduction efforts, and serve as a benchmark for evaluating therapy effectiveness (Maisto et al., 2014). Our research suggests that in addition to being common among young community residents, AUDs are heterogeneous with respect to course among affected persons. AUDs are time-limited and non-recurrent for some, whereas for others AUDs constitute a persistent or recurring condition.

Sex differences in AUD rates across the age span, with the possible exception of early to mid-adolescence when rates are relatively low, have been previously noted (Wells et al., 2006). International research involving representative cross-sectional (Cheng et al., 2015; Cho et al., 2015; Grant et al., 2015; Hasin et al., 2007; Kessler et al., 2005; Subramaniam et al., 2012; Teesson et al., 2010) or community-based prospective samples (Angst et al., 2016; Meier et al., 2012; Wells et al., 2006) consistently indicate greater rates of AUDs among male versus female participants. We similarly found an absence of significant sex differences

in AUD rates prior to age 18 but generally small to moderate sex differences afterwards with men demonstrating consistently higher rates than women.

For both sexes, the peak incidence and prevalence rates for AUD occurred between ages 18 and 24.9. These findings are consistent with those reported in other studies with representative samples, where the typical age of onset for the first AUD episode was before age 30 (Grant et al., 2015; Hasin et al., 2007; Kessler et al., 2005; Subramaniam et al., 2012; Teesson et al., 2010). Cumulative hazard functions characterizing AUD onset functions in the present research differed significantly between male and female participants: males demonstrated a steeper rise beginning around age 18. For both sexes, risk for first AUD onsets substantially diminished beginning at age 25, although period prevalence rates for AUDs remained high between ages 25 and 30. Greater AUD persistence was noted for men, a finding consistent with earlier reports (Boschloo et al., 2012; Walitzer and Dearing, 2006). Overall, first incidence and period prevalence rate data in this study have strong resemblances to the “developmentally limited” patterns that have been reported earlier in cross-sectional and longitudinal studies with representative samples, whereby heavy alcohol consumption or AUDs increase during adolescence and early adulthood and then decline during the mid to late 20s (Meier et al., 2013; Sher et al., 2005; Teesson et al., 2010; Wells et al., 2006).

Among those with AUD histories, 87% recovered from the initial episode by age 30. No significant differences in recovery rates were observed between men and women. Cumulative hazard functions significantly differed by sex, however, whereby women recovered more quickly and at a faster rate than men. For those who recovered, the average length of the initial AUD episode was relatively brief (23 months) when referenced to AUD episode durations commonly reported within treatment samples. Of those who recovered, however, 33% experienced an episode recurrence. This rate is higher than those earlier reported for representative samples with follow-up intervals of 3 years duration (Dawson et al., 2007; de Bruijn et al., 2006). This study’s recurrence rate is also higher than that reported by Tuithof at a 20-year follow-up (Tuithof et al., 2014). This latter investigation, however, was primarily based on retrospective reports, which are known to generally underestimate lifetime disorder occurrences (Haeny et al., 2014; Moffitt et al., 2010). The absence of sex differences in recurrence rates found in the present study is generally consistent with earlier findings summarized elsewhere based on treatment samples (Walitzer and Dearing, 2006). Additional findings indicated that risk for episode recurrence remained considerable within the first five years following remission and recovery from the first episode. These observations are similar to those reported in the natural recovery literature, which generally suggests that recovery is more likely to be maintained after five years of non-recurrence (Sobell et al., 2000).

Perspective on the course of AUDs, based largely on treatment samples, has shifted in recent years from one where AUDs were regarded primarily as chronic medical illness punctuated by periods of remission and relapse (McLellan et al., 2000) to a perspective where the course of AUDs is relatively brief and largely time-limited (Boschloo et al., 2012; Dawson et al., 2012; de Bruijn et al., 2006; Hasin et al., 2011; Tuithof et al., 2013). We found evidence supportive of both of these perspectives. A third of those who recovered from the initial

AUD episode experienced a recurrence, and 18% of those with a single AUD episode did not recover from that episode by age 30. For this latter subgroup, the mean AUD episode duration by age 30 was about 5 years. A majority of the sample with AUD (71%), however, recovered from the initial episode and did not experience a recurrence by age 30. Since only 15% of the OADP sample with AUD received some form of treatment during the initial episode, a large majority of cases were unaided professionally in their transition from AUD to recovery. Because a small proportion of those with AUD received treatment services, it is unlikely that most of the OADP sample with AUD matched the degree severity and psychiatric comorbidity found in treatment samples (Low et al., 2008). In other studies with representative samples, AUD recurrence or persistence has been predicted by the severity of alcohol-related problems and psychiatric comorbidity (Boschloo et al., 2012; Dawson et al., 2007; Hasin et al., 2011; Tuithof et al., 2014).

Although there are several study strengths, there are also limitations. First, the ethnic diversity of the sample, although representative of the ethnic distribution of western Oregon, is restricted. Within U.S. population samples, AUD prevalence rates and some course features demonstrate differences depending on race or ethnicity (Dawson et al., 2005; Grant et al., 2012; Grant et al., 2015; Hasin et al., 2007). AUD rates and course features also vary across geographic regions and cultures (Cheng et al., 2015; Cho et al., 2015; Subramaniam et al., 2012; Zavos et al., 2015). Consequently, there are some limits on the generalizability of study findings reported here. Second, study discontinuation after T₃ was more common among those with a history of an AUD, and there were some other statistically significant differences between those who discontinued or remained in the study over time. The full impact of non-random discontinuation on this study's findings is unclear. Third, the sample was followed only through 30 years of age, which limits the generalization of the study findings beyond this period of life.

Directions for future research are several. Because AUDs include a heterogeneous set of conditions with problematic alcohol use in common (Sher et al., 2005), future natural course research should better accommodate such heterogeneity into the modeling process. In DSM-5, for example, AUDs are characterized as mild, moderate, or severe based on the number of symptom features present. Greater alcohol involvement and alcohol-related problems have been found to predict a poorer AUD course (Connor et al., 2016; Dawson et al., 2005; Moyer et al., 2002); consequently, future research might evaluate whether different levels of AUD severity arise through distinct sets of processes. A greater understanding of processes that underlie transitions in the course of AUD would also have significant value in informing theory and intervention development.

Role of the Funding Source

National Institutes of Health grants R01MH40501, R01MH50522, and R01DA12951 to Peter Lewinsohn and R01AA020968 to Richard Farmer and John Seeley supported this research. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The authors thank Peter Lewinsohn for the Oregon Adolescent Depression Project dataset used in this research.

References

- Angst J, Paksarian D, Cui L, Merikangas KR, Hengartner MP, Ajdacic-Gross V, Rössler W, 2016 The epidemiology of common mental disorders from age 20 to 50: Results from the Prospective Zurich Cohort Study. *Epidemiol. Psychiatr. Sci* 25, 24–32. [PubMed: 25802979]
- Arnett JJ, 2007 Emerging adulthood: What is it, and what is it good for? *Child Dev. Perspect* 1, 68–73.
- Boschloo L, Vogelzangs N, van den Brink W, Smit JH, Beekman AT, Penninx BW, 2012 Predictors of the 2-year recurrence and persistence of alcohol dependence. *Addiction* 107, 1639–1640.
- Brown SA, Tapert SF, Granholm E, Delis DC, 2000 Neurocognitive functioning of adolescents: Effects of protracted alcohol use. *Alcohol. Clin. Exp. Res* 24, 164–171. [PubMed: 10698367]
- Chambers WJ, Puig-Antich J, Hirsch M, Paez P, Ambrosini PJ, Tabrizi MA, Davies M, 1985 The assessment of affective disorders in children and adolescents by semistructured interview: Test-retest reliability of the Schedule for Affective Disorders and Schizophrenia for School-Age Children, present episode version. *Arch. Gen. Psychiatry* 4, 696–702.
- Cheng HG, Deng F, Xiong W, Phillips MR, 2015 Prevalence of alcohol use disorders in mainland China: A systematic review. *Addiction* 110, 761–74. [PubMed: 25678403]
- Cho MJ, Seong SJ, Park JE, Chung IW, Lee YM, Bae A, Ahn JH, Lee DW, Bae JN, Cho SJ, Park JI, Son J, Chang SM, Hahm BJ, Lee JY, Sohn JH, Kim JS, Hong JP, 2015 Prevalence and correlates of DSM-IV mental disorders in South Korean adults: The Korean epidemiologic catchment area study 2011. *Psychiatry Investig.* 12, 164–70.
- Chung T, Maisto SA, 2006 Relapse to alcohol and other drug use in treated adolescents: Review and reconsideration of relapse as a change point in clinical course. *Clin. Psychol. Rev* 26, 149–161. [PubMed: 16364524]
- Connor JP, Haber PS, Hall WD, 2016 Alcohol use disorders. *Lancet* 387, 988–998. [PubMed: 26343838]
- Cunningham JA, 1999 Resolving alcohol-related problems with and without treatment: The effects of different problem criteria. *J. Stud. Alcohol* 60, 463–466. [PubMed: 10463801]
- Dawson DA, Goldstein RB, Grant BF, 2007 Rates and correlates of relapse among individuals in remission from DSM-IV alcohol dependence: A 3-year follow-up. *Alcohol. Clin. Exp. Res* 31, 2036–2045. [PubMed: 18034696]
- Dawson DA, Goldstein RB, Ruan WJ, Grant BF, 2012 Correlates of recovery from alcohol dependence: A prospective study over a 3-year follow-up interval. *Alcohol. Clin. Exp. Res* 36, 1268–1277. [PubMed: 22309217]
- Dawson DA, Grant BF, Stinson FS, Chou PS, Huang B, Ruan W, 2005 Recovery from DSM-IV alcohol dependence: United States, 2001–2002. *Addiction* 100, 281–292. [PubMed: 15733237]
- De Bruijn C, Van Den Brink W, De Graaf R, Vollebergh WA, 2006 The three year course of alcohol use disorders in the general population: DSM-IV, ICD-10 and the Craving Withdrawal Model. *Addiction* 101, 385–392. [PubMed: 16499511]
- Edens EL, Glowinski AL, Grazier KL, Bucholz KK, 2008 The 14-year course of alcoholism in a community sample: Do men and women differ? *Drug Alcohol Depend.* 93, 1–11. [PubMed: 17935914]
- Farmer RF, Gau JM, Seeley JR, Kosty DB, Sher KJ, Lewinsohn PM, 2016 Internalizing and externalizing disorders as predictors of alcohol use disorder onset during three developmental periods. *Drug Alcohol Depend.* 164, 38–46. [PubMed: 27141839]
- Farmer RF, Kosty DB, Seeley JR, Olino TM, Lewinsohn PM, 2013 Aggregation of lifetime axis I psychiatric disorders through age 30: Incidence, predictors, and associated psychosocial outcomes. *J. Abnorm. Psychol* 122, 573–586. [PubMed: 23421525]
- Fergusson DM, Horwood LJ, 2001 The Christchurch Health and Development Study: Review of findings on child and adolescent mental health. *Aust. N. Z. J. Psychiatry* 35, 287–296. [PubMed: 11437801]
- First MB, Spitzer RL, Gibbon M, Williams JBW, 1994 Structured Clinical Interview for Axis I DSM-IV Disorders– Non-patient edition. Biometrics Research Department, New York.
- Frank E, Prien RF, Jarrett RB, Keller MB, Kupfer DJ, Lavori PW, Rush AJ, Weissman MM, 1991 Conceptualization and rationale for consensus definitions of terms in major depressive disorder:

- Remission, recovery, relapse, and recurrence. *Arch. Gen. Psychiatry* 48, 851–855. [PubMed: 1929776]
- Grant BF, Goldstein RB, Saha TD, Chou SP, Jung J, Zhang H, Pickering RP, Ruan WJ, Smith SM, Huang B, Hasin DS, 2015 Epidemiology of DSM-5 alcohol use disorder: Results from the National Epidemiologic Survey on Alcohol and Related Conditions III. *JAMA Psychiatry* 72, 757–766. [PubMed: 26039070]
- Grant JD, Vergés A, Jackson KM, Trull TJ, Sher KJ, Bucholz KK, 2012 Age and ethnic differences in the onset, persistence and recurrence of alcohol use disorder. *Addiction* 107, 756–765. [PubMed: 22085024]
- Guilamo-Ramos V, Jaccard J, Turrisi R, Johansson M, 2005 Parental and school correlates of binge drinking among middle school students. *Am. J. Public Health* 95, 894–899. [PubMed: 15855471]
- Haeny AM, Littlefield AK, Sher KJ, 2014 Repeated diagnoses of lifetime alcohol use disorders in a prospective study: Insights into the extent and nature of the reliability and validity problem. *Alcohol. Clin. Exp. Res* 38, 489–500. [PubMed: 24033811]
- Hasin DS, Stinson FS, Ogburn E, Grant BF, 2007 Prevalence, correlates, disability, and comorbidity of DSM-IV alcohol abuse and dependence in the United States: Results from the National Epidemiologic Survey on Alcohol and Related Conditions. *Arch. Gen. Psychiatry* 64, 830–842. [PubMed: 17606817]
- Hasin D, Fenton MC, Skodol A, Krueger R, Keyes K, Geier T, Greenstein E, Blanco C, Grant B, 2011 Personality disorders and the 3-year course of alcohol, drug, and nicotine use disorders. *Arch. Gen. Psychiatry* 68, 1158–1167. [PubMed: 22065531]
- Keller MB, Lavori PW, Friedman B, Nielsen E, Endicott J, McDonald-Scott PA, 1987 The longitudinal interval follow-up evaluation: A comprehensive method for assessing outcome in prospective longitudinal studies. *Arch. Gen. Psychiatry* 44, 540–548. [PubMed: 3579500]
- Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE, 2005 Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch. Gen. Psychiatry* 62, 593–602. [PubMed: 15939837]
- Komro KA, Maldonado-Molina MM, Tobler AL, Bonds JR, Muller KE, 2007 Effects of home access and availability of alcohol on young adolescents' alcohol use. *Addiction* 102, 1597–1608. [PubMed: 17854336]
- Lewinsohn PM, Hops H, Roberts RE, Seeley JR, Andrews JA, 1993 Adolescent psychopathology: I. Prevalence and incidence of depression and other DSM-III-R disorders in high school students. *J. Abnorm. Psychol* 102, 133–144. [PubMed: 8436689]
- Lewinsohn PM, Rohde P, Klein DN, Seeley JR, 1999 Natural course of adolescent major depressive disorder: I. Continuity into young adulthood. *J. Amer. Acad. Child Adoles. Psychiatry* 38, 56–63.
- Low NC, Cui L, Merikangas KR, 2008 Community versus clinic sampling: Effect on the familial aggregation of anxiety disorders. *Biol. Psychiatry* 63, 884–890. [PubMed: 17945198]
- Maisto SA, Kirouac M, Witkiewitz K, 2014 Alcohol use disorder clinical course research: Informing clinicians' treatment planning now and in the future. *J. Stud. Alcohol Drugs* 75, 799–807. [PubMed: 25208198]
- McLellan AT, Lewis DC, O'Brien CP, Kleber HD, 2000 Drug dependence, a chronic medical illness: Implications for treatment, insurance, and outcomes evaluation. *JAMA* 284, 1689–1695. [PubMed: 11015800]
- Meier MH, Caspi A, Houts R, Slutske WS, Harrington H, Jackson KM, Belsky DW, Poulton R, Moffitt TE, 2013 Prospective developmental subtypes of alcohol dependence from age 18 to 32 years: Implications for nosology, etiology, and intervention. *Dev. Psychopathol* 25, 785–800. [PubMed: 23880392]
- Merikangas KR, McClair VL, 2012 Epidemiology of substance use disorders. *Hum. Genet* 131, 779–789. [PubMed: 22543841]
- Moffitt TE, Caspi A, Kokaua J, Milne BJ, Polanczyk G, Poulton R, 2010 How common are common mental disorders? Evidence that lifetime prevalence rates are doubled by prospective versus retrospective ascertainment. *Psychol. Med* 40, 899–909. [PubMed: 19719899]

- Moyer A, Finney JW, Swearingen CE, Vergun P, 2002 Brief interventions for alcohol problems: A meta-analytic review of controlled investigations in treatment-seeking and non-treatment-seeking populations. *Addiction* 97, 279–292. [PubMed: 11964101]
- Muthén B, Muthén LK, 1998–2012 Mplus User's Guide (7th ed.). Muthén and Muthén, Los Angeles.
- Orvaschel H, Puig-Antich J, Chambers WJ, Tabrizi MA, Johnson R, 1982 Retrospective assessment of prepubertal major depression with the Kiddie-SADS-E. *J. Am. Acad. Child Psychiatry* 21, 392–397. [PubMed: 7119313]
- Parry CD, Patra J, Rehm J, 2011 Alcohol consumption and non-communicable diseases: Epidemiology and policy implications. *Addiction* 106, 1718–1724. [PubMed: 21819471]
- Patrick ME, Schulenberg JE, Maggs JL, Maslowsky J, 2016 Substance use and peers during adolescence and the transition into adulthood: Selection, socialization, and development In: Sher KJ (Ed.). *The Oxford handbook of substance use and substance use disorders* Vol. 1 Oxford University Press, New York pp. 526–548.
- Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J, 2009 Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *Lancet* 373, 2223–2233. [PubMed: 19560604]
- Rohde P, Lewinsohn PM, Kahler CW, Seeley JR, Brown RA, 2001 Natural course of alcohol use disorders from adolescence to young adulthood. *J. Am. Acad. Child Adolesc. Psychiatry* 40, 83–90. [PubMed: 11195569]
- Rohde P, Lewinsohn PM, Seeley JR, 1996 Psychiatric comorbidity with problematic alcohol use in high school students. *J. Am. Acad. Child Adolesc. Psychiatry* 35, 101–109. [PubMed: 8567601]
- Rohde P, Lewinsohn PM, Seeley JR, Klein DN, Andrews JA, Small JW, 2007 Psychosocial functioning of adults who experienced substance use disorders as adolescents. *Psychol. Addict. Behav* 21, 155–164. [PubMed: 17563135]
- Sher KJ, Grekin ER, Williams NA, 2005 The development of alcohol use disorders. *Annu. Rev. Clin. Psychol* 1, 493–523. [PubMed: 17716097]
- Singer JD, Willett JB, 1991 Modeling the days of our lives: Using survival analysis when designing and analyzing longitudinal studies of duration and the timing of events. *Psychol. Bull* 110, 268–290.
- Sobell LC, Cunningham JA, Sobell MB, 1996 Recovery from alcohol problems with and without treatment: Prevalence in two population surveys. *Am. J. Public Health* 86, 966–972. [PubMed: 8669520]
- Sobell LC, Ellingstad TP, Sobell MB, 2000 Natural recovery from alcohol and drug problems: Methodological review of the research with suggestions for future directions. *Addiction* 95, 749–764. [PubMed: 10885050]
- Subramaniam M, Abidin E, Vaingankar J, Phua AM, Tee J, Chong SA, 2012 Prevalence and correlates of alcohol use disorders in the Singapore Mental Health Survey. *Addiction*. 107, 1443–1452. [PubMed: 22296228]
- Teesson M, Hall W, Slade T, Mills K, Grove R, Mewton L, Baillie A, Haber P, 2010 Prevalence and correlates of DSM-IV alcohol abuse and dependence in Australia: Findings of the 2007 National Survey of Mental Health and Wellbeing. *Addiction* 105, 2085–2094. [PubMed: 20707771]
- Tuithof M, ten Have M, van den Brink W, Vollebergh W, de Graaf R, 2013 Predicting persistency of DSM-5 alcohol use disorder and examining drinking patterns of recently remitted individuals: A prospective general population study. *Addiction* 108, 2091–2099. [PubMed: 23889861]
- Tuithof M, ten Have M, van den Brink W, Vollebergh W, de Graaf R, 2014 Alcohol consumption and symptoms as predictors for relapse of DSM-5 alcohol use disorder. *Drug Alcohol Depend.* 140, 85–91. [PubMed: 24793368]
- Walitzer KS, Dearing RL, 2006 Gender differences in alcohol and substance use relapse. *Clin. Psychol. Rev* 26, 128–148. [PubMed: 16412541]
- Wells JE, Horwood LJ, Fergusson DM, 2006 Stability and instability in alcohol diagnosis from ages 18 to 21 and ages 21 to 25 years. *Drug Alcohol Depend.* 81, 157–165. [PubMed: 16046080]
- Whiteford HA, Degenhardt L, Rehm J, Baxter AJ, Ferrari AJ, Erskine HE, Charlson FJ, Norman RE, Flaxman AD, Johns N, Burstein R, Murray CJ, Vos T, 2013 Global burden of disease attributable

to mental and substance use disorders: Findings from the Global Burden of Disease Study 2010. *Lancet* 382, 1575–1586. [PubMed: 23993280]

Zavos HMS, Siribaddana S, Ball HA, Lynskey MT, Sumathipala A, Rijsdijk FV, Hotopf M, 2015 The prevalence and correlates of alcohol use and alcohol use disorders: A population based study in Colombo, Sri Lanka. *BMC Psychiatry* 15, 158. [PubMed: 26169683]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.
Cumulative hazard functions for the onset of alcohol use disorders (AUD).

Figure 2.
Cumulative hazard functions for the recovery from alcohol use disorders (AUDs).

Figure 3.
Cumulative hazard functions for the recurrence of alcohol use disorders (AUD).

Table 1.

Prevalence, incidence, recovery, and recurrence rates of alcohol use disorders from childhood to age 30 by sex.

	Female Participants % [CI ₉₅]	Male Participants % [CI ₉₅]	OR [CI ₉₅]
Lifetime prevalence	27.6 [23.5–31.7]	42.9 [37.8–48.0]	1.97 [1.47–2.65]
First incidence			
0.0 to 13.9 years	1.7 [0.5–2.9]	0.6 [0.0–1.4]	0.35 [0.08–1.57]
14.0 to 17.9 years	6.4 [4.2–8.6]	8.1 [5.4–10.8]	1.27 [0.75–2.17]
18.0 to 24.9 years	16.3 [13.0–19.6]	29.4 [24.7–34.1]	2.14 [1.53–3.00]
25.0 to 30.0 years	3.2 [1.6–4.8]	4.9 [2.7–7.1]	1.57 [0.77–3.21]
Period prevalence			
14.0 to 17.9 years	7.2 [4.8–9.6]	8.7 [5.8–11.6]	1.22 [0.73–2.03]
18.0 to 24.9 years	22.6 [18.7–26.5]	36.7 [31.6–41.8]	1.99 [1.46–2.70]
25.0 to 30.0 years	12.3 [9.4–15.2]	21.9 [17.6–26.2]	2.00 [1.37–2.91]
Recovery rates	89.8 [84.5–95.1]	84.9 [79.2–90.6]	0.64 [0.31–1.32]
Recurrence rates for those who recovered	31.7 [23.1–40.3]	33.8 [25.6–42.0]	1.10 [0.64–1.88]

Note. CI₉₅ = 95% confidence interval. OR = Odds ratio. Bolded ORs are statistically significant at $p < .05$. All summary statistics account for sample weighting.