Published in final edited form as:

Cardiol Young. 2020 January ; 30(1): 66–73. doi:10.1017/S1047951119002786.

Association between preoperative respiratory support and outcomes in pediatric cardiac surgery

Elizabeth C. Ciociola¹, Karan R. Kumar^{1,2}, Kanecia O. Zimmerman^{1,2}, Elizabeth J. Thompson^{1,2}, Melissa Harward², Laura N. Sullivan³, Joseph W. Turek³, Christoph P. Hornik^{1,2}

¹Duke Clinical Research Institute, Duke University School of Medicine, Durham, NC, USA

²Department of Pediatrics, Duke University Medical Center, Durham, NC, USA

³Department of Surgery, Duke University Medical Center, Durham, NC, USA

Abstract

Background—Preoperative mechanical ventilation is associated with morbidity and mortality following CHD surgery, but prior studies lack a comprehensive analysis of how preoperative respiratory support mode and timing affects outcomes.

Methods—We retrospectively collected data on children <18 years of age undergoing cardiac surgery at an academic tertiary care medical center. Using multivariable regression, we examined the association between modes of preoperative respiratory support (nasal cannula, high flow nasal cannula/noninvasive ventilation, or invasive mechanical ventilation), escalation of preoperative respiratory support, and invasive mechanical ventilation on the day of surgery for three outcomes: operative mortality, postoperative length of stay, and postoperative complications. We repeated our analysis in a subcohort of neonates.

Results—A total of 701 children underwent 800 surgical procedures, and 40% received preoperative respiratory support. Among neonates, 243 patients underwent 253 surgical procedures, and 79% received preoperative respiratory support. In multivariable analysis, all modes of preoperative respiratory support, escalation in preoperative respiratory support, and invasive mechanical ventilation on the day of surgery were associated with increased odds of prolonged length of stay in children and neonates. Children (odds ratio=3.69, 95% CI 1.2–11.4) and neonates (odds ratio=8.97, 95% CI 1.31–61.14) on high flow nasal cannula/noninvasive ventilation had increased odds of operative mortality compared to those on room air.

Conclusion—Preoperative respiratory support is associated with prolonged length of stay and mortality following CHD surgery. Knowing how preoperative respiratory support affects outcomes may help guide surgical timing, inform prognostic conversations, and improve risk stratification models.

Corresponding Author: Christoph P. Hornik, MD, MPH, Duke Clinical Research Institute, Durham, NC 27705; Phone: (919) 668-8935; Fax: (919) 668-7032; christoph.hornik@duke.edu.

Ethical Standards. The institutional review board at Duke University approved this study (IRB Pro00100843) and waived the need for informed consent.

Keywords

CHD; preoperative ventilation; pediatric cardiac surgery; respiratory support; noninvasive ventilation; postoperative length of stay

Nearly 40,000 children in North America underwent cardiac surgery between January 1, 2014, and December 31, 2017.¹ Although outcomes have improved over time, mortality continues to range from 0.4% to 15% depending on diagnosis, procedure, and site.² Among survivors, up to 30% suffer a major complication.³ In addition, postoperative length of stay can be prolonged, particularly for children with the most severe defects.² Given the prevalence and variability of these adverse outcomes, identifying preoperative risk factors in this population remains critical.

Preoperative mechanical ventilation is associated with morbidity and mortality following pediatric cardiac surgery. An analysis of 25,476 pediatric cardiac operations in the Society of Thoracic Surgeons Congenital Heart Surgery Database found that preoperative mechanical ventilation was associated with discharge mortality.⁴ A multicenter retrospective analysis of 2,557 Norwood patients similarly found that preoperative mechanical ventilation was associated with 30% increased odds of experiencing postoperative complications.⁵ Smaller single-center studies have shown that preoperative mechanical ventilation is associated with prolonged postoperative hospital stay.^{6,7} However, the impact of duration and timing of preoperative invasive mechanical ventilation, and other modes of preoperative respiratory support including noninvasive ventilation, on outcomes in children after cardiac surgery remains unknown.

In this study, we examined the association between mode, escalation, and timing of preoperative respiratory support and postoperative mortality, hospital length of stay, and complications in children undergoing cardiac surgery. An understanding of how ventilation practices affect postoperative outcomes has important implications for preoperative management strategies in this patient population.

Materials and Methods

Data source and study population

We conducted a retrospective observational cohort study of all children who underwent cardiac surgery at Duke University Medical Center between January 1, 2014, and December 31, 2017. The Duke congenital cardiothoracic surgical database and the Duke Enterprise Data Unified Content Explorer, a web-based electronic-health record query system, were used to identify the cohort and obtain data. We included all children younger than 18 years who underwent cardiac surgery, including those not admitted prior to surgery. We excluded children who received mechanical circulatory support prior to surgery, dropping 168 observations. In addition, we excluded those with missing data on key study variables including hospital length of stay, postoperative complications, mortality, preoperative respiratory support, and STAT mortality category. There were no patients with missing data on hospital length of stay, postoperative complications, mortality, or preoperative respiratory support, but we dropped 350 observations with a missing STAT mortality category. For

children with multiple surgeries throughout the study period, including children who received multiple surgeries within the same hospital admission, we treated each surgery as a discrete case. All patients who received preoperative care, including neonates, were managed within the Pediatric Cardiac ICU. The institutional review board at Duke University approved this study and waived the need for informed consent.

Study Outcomes and Exposure Definitions

The primary study outcome was postoperative hospital length of stay, defined as the number of days from surgery to hospital discharge. Secondary outcomes included major postoperative complications and mortality. We included the following major postoperative complications as previously defined by the Society of Thoracic Surgeons: renal failure requiring dialysis, neurologic deficit persisting at discharge, arrhythmia requiring permanent pacemaker, paralyzed diaphragm/phrenic nerve injury, mechanical circulatory support, and unplanned re-intervention (includes surgical or catheter-based unplanned interventions as well as cardiac and noncardiac procedures).⁸ Mortality was defined using the Society of Thoracic Surgeons definition of operative mortality, which includes all deaths occurring during the hospitalization in which the operation was performed and all deaths occurring after discharge from the hospital within 30 days after the operation.⁹

We separately evaluated the prevalence and adjusted odds of the primary and secondary outcomes based on three primary study exposures: highest mode of preoperative respiratory support, escalation of preoperative respiratory support, and invasive mechanical ventilation on the day of surgery. Postoperative length of stay and postoperative major complications were evaluated in survivors only. Preoperative respiratory support was categorized into four escalating modes: room air, nasal cannula, high flow nasal cannula or noninvasive ventilation, and invasive mechanical ventilation. Noninvasive ventilation included bi-level positive airway pressure and continuous positive airway pressure. Invasive mechanical ventilation included both invasive conventional mechanical ventilation and invasive high frequency ventilation. For children who received multiple modes of preoperative respiratory support during one hospital admission, the mode was defined as the highest mode of respiratory support received preoperatively. For children who underwent multiple surgeries within one hospital admission, the preoperative respiratory support mode was defined by the highest mode of support received prior to each surgery. We defined escalation of respiratory support as any progression of respiratory support from a lower to a higher mode during the preoperative period (e.g., from high flow nasal cannula/noninvasive ventilation to mechanical ventilation), regardless of any downward progression thereafter. To evaluate the timing of ventilation, we classified children based on whether or not they were on invasive mechanical ventilation on the day of surgery. The total duration of preoperative respiratory support was defined as the number of days on any preoperative respiratory support, and preoperative duration was defined as the number of days from hospital admission to surgery.

Statistical Analysis

We summarized the distribution of continuous and categorical variables using medians (25th and 75th percentiles) and counts (%), respectively. Patients with missing data were excluded from analysis. To compare study outcomes based on preoperative respiratory support

parameters, we performed the Kruskal–Wallis test for continuous variables and chi-square or Fischer’s exact tests for categorical variables. We reported the study outcomes based on the primary exposures as counts (%) or medians (25th, 75th percentiles). To assess the association between study exposures and study outcomes, we developed separate multivariable logistic regression models for the three primary predictors of interest: (1) highest mode of preoperative respiratory support, (2) escalation of preoperative respiratory support, and (3) invasive mechanical ventilation on the day of surgery. Logistic regression was performed for binary outcomes (operative mortality and occurrence of any postoperative complication) and Poisson regression for postoperative hospital length of stay. To build the regression models, we included variables determined a priori based on reported association with outcomes in prior studies and biological plausibility.^{4,10} The final models were adjusted for age, any preoperative cardiac or noncardiac risk factors, any noncardiac congenital anatomic or chromosomal abnormalities, any syndromes, surgical complexity, and cardiopulmonary bypass duration. In addition, a Poisson regression model was used to assess the association between preoperative duration and escalation in preoperative respiratory support. The regression model controlled for all of the same variables listed above, except for cardiopulmonary bypass duration. Preoperative cardiac or noncardiac risk factors included all factors coded in the Society of Thoracic Surgeons Congenital Heart Surgery Database under the heading “preoperative factors,” which comprises factors such as preoperative complete AV block or preoperative renal dysfunction.⁹ All syndromes, noncardiac congenital anatomic abnormalities, and chromosomal abnormalities coded in the Society of Thoracic Surgeons Congenital Heart Surgery Database under the headings “syndromes,” “noncardiac congenital anatomic abnormalities,” and “chromosomal abnormalities” were included.⁹ Surgical complexity was categorized using the Society of Thoracic Surgeons and European Association for Cardiothoracic Surgery (STAT) Mortality Categories, with category 1 representing the lowest mortality risk and category 5 representing the highest mortality risk.^{11,12}

The adjusted results from the logistic regression models are displayed as odds ratios and 95% confidence intervals. All analyses were repeated in a subcohort of neonates less than 30 days of age, with the addition of birthweight and gestational age as covariates in the neonatal multivariable model. Analyses were performed with R and R Studio software, version 1.1.453. A *p* value < 0.05 was considered statistically significant.

Results

Cohort Clinical Characteristics

We identified a total of 701 children who underwent 800 cardiac surgical procedures during the study period. A complete list of surgical procedures and fundamental diagnoses can be found in Supplementary Tables S1 and S2, respectively. The median (25th, 75th percentiles) age at surgery was 0.5 years (0.1, 3.2), and 47% of the cohort was female. In the preoperative period, the maximum respiratory support for 60% of children was room air, while 7% received nasal cannula, 12% received high flow nasal cannula or noninvasive ventilation, and 21% received invasive mechanical ventilation (Table 1). Preoperative duration was 0 days (0, 0) for those on room air, 7 days (3, 12) for those on nasal cannula, 6

days (3, 16) for those on high flow nasal cannula/noninvasive ventilation, and 14 days (6, 34) for those on mechanical ventilation. Approximately 10% of children underwent an escalation in preoperative respiratory support from the time of admission to the day of surgery. Escalation in preoperative respiratory support was found to correlate with preoperative duration, with those who experienced an escalation in preoperative respiratory support more likely to have a longer preoperative duration (odds ratio = 1.51, 95% CI 1.42–1.60). The total duration of preoperative respiratory support for children was 2 days (1, 5) on nasal cannula, 4 days (2, 10) on high flow nasal cannula/noninvasive ventilation, and 14 days (6, 31) on MV, and 13% of children were on invasive mechanical ventilation on the day of surgery. To address concern for variations between noninvasive ventilation and high flow nasal cannula and the impact of including these children in the same group, the analysis was repeated excluding the 6 children on noninvasive support. Comparing the adjusted odds ratios (with 95% CI) for high flow nasal cannula alone versus high flow nasal cannula and noninvasive ventilation together, mortality (3.53 [1.11–11.26] vs 3.69 [1.20–11.40]), postoperative length of stay (1.70 [1.61–1.80] vs 1.77 [1.68–1.86]), and major complications (1.09 [0.43–2.76] vs 1.15 [0.47–2.82]) varied slightly with no change in statistical significance.

Mortality

Operative mortality for the whole cohort was 5%, and it varied based on the maximum mode of preoperative respiratory support received. Operative mortality for children on nasal cannula (6%), high flow nasal cannula/noninvasive ventilation (12%), and invasive mechanical ventilation (12%) was greater than for those who remained on room air (1%, $p<0.001$) (Table 2). In addition, operative mortality was greater for children who experienced an escalation in preoperative respiratory support mode compared to those who did not (11% vs. 5%, $p=0.021$) and for those on invasive mechanical ventilation on the day of surgery (15% vs. 4%, $p<0.001$) (Table 2). After adjusting for covariates, multivariable analysis showed that only children on preoperative high flow nasal cannula/noninvasive ventilation had increased odds of mortality following cardiac surgery compared to those who remained on room air (odds ratio=3.69, 95% CI 1.2–11.4) (Table 3).

Postoperative Hospital Length of Stay

Among survivors, the median postoperative length of stay was 10 days (6, 28), and both the median and the interquartile range increased with higher maximum mode of preoperative respiratory support (Table 2). The median postoperative length of stay for the surviving children on nasal cannula (20 days [13, 39]), high flow nasal cannula or noninvasive ventilation (19 days [12, 40]), and invasive mechanical ventilation (42 days [23, 117]) were all longer than for children who remained on room air preoperatively (6 days [5, 10], $p<0.001$). In addition, median postoperative length of stay was longer for both children who experienced an escalation in preoperative respiratory support compared to those who did not (36 days [16, 75] vs. 9 days [5, 21], $p<0.001$) and for children who were on invasive mechanical ventilation on the day of surgery compared to those who were not (50 days [24, 131] vs. 9 days [5, 19], $p<0.001$). In multivariable analysis, we found that all modes of maximum preoperative respiratory support were associated with an increased postoperative length of stay compared to children who remained on room air (Figure 1). In addition, both

escalation in preoperative respiratory support (odds ratio=1.19, 95% CI 1.15–1.23) and invasive mechanical ventilation on the day of surgery (odds ratio=1.37, 95% CI 1.32–1.43) were associated with increased postoperative length of stay.

Major Complications

Approximately 10% of the surviving children experienced a major postoperative complication. Major postoperative complications were more common among those on preoperative nasal cannula (8%), high flow nasal cannula or noninvasive ventilation (12%), and mechanical ventilation (14%) compared to those who remained on room air (5%, $p=0.001$) (Table 2). Postoperative complications were also more common for children who experienced an escalation in preoperative respiratory support (19% v. 7%, $p<0.001$) but not for children who were on mechanical ventilation on the day of surgery (13% v. 7%, $p=0.051$). Multivariable analysis showed no association between maximum preoperative respiratory support, escalation in preoperative respiratory support, or invasive mechanical ventilation on the day of surgery and postoperative complications (Figure 1).

Neonate Cohort Clinical Characteristics

We identified a subcohort of 243 neonates who underwent 253 cardiac procedures at a median age of 11 days (6, 22). The median gestational age for neonates was 38 weeks (32, 39), and the median birth weight was 2900 g (1700, 3450). Among neonates, 79% received preoperative respiratory support. This included 12% who received nasal cannula, 19% who received high flow nasal cannula or noninvasive ventilation, and 47% who received mechanical ventilation as their maximum preoperative respiratory support. In addition, 23% of neonates experienced an escalation in preoperative respiratory support, and 34% of neonates were on invasive mechanical ventilation on the day of surgery.

Neonatal Outcomes

Among neonates, operative mortality was 10%, and 17% experienced a major postoperative complication. The median postoperative length of stay for neonates was 31 days (15, 71). Similar to the main cohort, median postoperative lengths of stay for neonates on preoperative nasal cannula (22 days [16, 41]), high flow nasal cannula or noninvasive ventilation (25 days [16, 51]), and invasive mechanical ventilation (47 days [24, 128]) were all longer compared to neonates who remained on room air (16 days [6, 33], $p<0.001$) (Table 2). Neonates on invasive mechanical ventilation on the day of surgery also experienced a greater postoperative length of stay than those who were not (56 days [24, 134] v. 23 days [14, 47], $p<0.001$). Multivariable analysis showed an association between all modes of preoperative respiratory support, escalation in preoperative respiratory support, and invasive mechanical ventilation on the day of surgery and an increased postoperative length of stay in neonates (Table 3). In addition, multivariable analysis showed an association between neonates on preoperative high flow nasal cannula or noninvasive ventilation and increased odds of mortality compared to those on room air (odds ratio=8.97, 95% CI 1.31–61.14) (Table 3). None of the other modes of preoperative respiratory support were associated with mortality, and there were no associations between any preoperative respiratory support modes and postoperative complications in neonates (Figure 2).

Discussion

Among survivors, children who received respiratory support prior to cardiac surgery had increased odds of prolonged postoperative length of stay compared to children on room air in the preoperative period. In addition, odds of prolonged postoperative hospital length of stay increased progressively with each increase in preoperative respiratory support mode, and children who required escalation in preoperative respiratory support compared to those who did not, as well as children on invasive mechanical ventilation on the day of surgery compared to those who were not, were more likely to experience prolonged postoperative length of stay. However, only preoperative high flow nasal cannula or noninvasive ventilation was associated with greater odds of operative mortality in children. These associations all held true in sensitivity analysis limited to neonates.

Prolonged length of stay is an important measure of morbidity as it captures the influence of several pre-, intra-, and postoperative factors and may serve as an indirect proxy of a patient's condition. Prolonged length of stay also has a negative impact on short- and long-term neurodevelopmental outcomes in children with CHD.¹³ In the Boston Circulatory Arrest Study of infants who underwent cardiac surgery, hospital length of stay was independently associated with lower full-scale, verbal, and performance IQ.¹⁴ Prolonged length of stay also increases resource utilization and costs. An analysis of 12,718 congenital heart surgery operations across 27 North American centers revealed an average excess cost per case of \$19,273 for each additional day of length of stay above the median.¹⁵ Identifying factors that predict or reduce postoperative length of stay in children undergoing cardiac surgery is therefore essential.

Our findings are consistent with single-center studies that analyzed the association between preoperative invasive mechanical ventilation and hospital length of stay. Among 355 children admitted to a cardiac ICU, preoperative mechanical ventilation was associated with two-fold greater odds of prolonged length of stay following cardiac surgery.⁶ Similarly, a retrospective study of 693 children after cardiac surgery found that preoperative mechanical ventilation was associated with an ICU stay above the 60th percentile of 3 days (odds ratio=5.23, 95% CI 1.71–16.13).⁷ Our study supplements these prior results by demonstrating that noninvasive preoperative respiratory support, escalation of preoperative respiratory support, and invasive respiratory support on the day of surgery are also associated with prolonged postoperative length of stay.

We did not find any associations between preoperative respiratory support and postoperative complications in either child or neonate survivors. This finding differs from past studies. An analysis of 2,557 neonates from 53 centers following the Norwood operation found that preoperative mechanical respiratory support was associated with 30% greater odds of experiencing one or more major postoperative complications (odds ratio=1.3, 95% CI 1.03–1.6).⁵ This significant finding may be related to the high morbidity associated with the Norwood operation. Our smaller, mixed cohort included children of all ages undergoing any cardiac surgery, which may have masked association between preoperative respiratory support and postoperative complications in higher complexity lesions. In addition, it is

possible that other complications, beyond those analyzed in this study, are associated with preoperative respiratory support.

Previous mortality risk factor studies in children undergoing cardiac surgery have demonstrated an association between preoperative invasive respiratory support and mortality. Among 25,476 pediatric cardiac operations recorded in the Society of Thoracic Surgeons Congenital Heart Surgery Database, discharge mortality was significantly greater in neonates (15.1% vs. 9.88%, $p<0.0001$), infants (11.5% vs. 2.91%, $p<0.0001$), and children (14.4% vs. 0.92%, $p<0.0001$) who received preoperative invasive mechanical ventilation compared to those who did not.⁴ Single-center studies have similarly identified preoperative invasive mechanical ventilation as a mortality risk factor.^{16–18} However, none of these studies analyzed noninvasive respiratory support. Our study builds on these findings by demonstrating that noninvasive preoperative respiratory support is also associated with operative mortality. In our cohort, mortality for children on noninvasive support (12%) and on invasive mechanical ventilation (12%) were comparable, and both were significantly greater than mortality among children who remained on room air before surgery (1%, $p<0.001$). However, mortality in neonates on noninvasive support (18%) was greater than mortality in neonates on invasive support (10%). Further, while invasive mechanical ventilation was not significantly associated with mortality in our multivariable analysis, children (odds ratio=3.69, 95% CI 1.2–11.4) and neonates (odds ratio=8.97, 95% CI 1.31–61.14) with preoperative noninvasive ventilation were found to have greater adjusted odds of operative mortality. These increased odds of mortality are similar to those observed in a previous study of 180 premature neonates with critical congenital heart disease receiving preoperative conventional invasive mechanical ventilation in 2002–2008 (odds ratio=5.1, 95% CI 1.4–18.4).¹⁸ We can only speculate as to why noninvasive ventilation was associated with mortality in our cohort, while invasive ventilation was not. Noninvasive ventilation does not reduce work of breathing and respiratory energy expenditure to the same extent as invasive mechanical ventilation.¹⁹ Thus, it is possible that children and neonates on noninvasive ventilation are inadequately supported prior to surgery, thereby worsening their postoperative outcomes. This may be particularly relevant in children with significant pulmonary pathology, who may have traditionally been supported with invasive mechanical ventilation, but who may now be receiving noninvasive support instead.²⁰ It is also possible that greater use of preoperative noninvasive respiratory support in children with CHD in recent years has simply shifted the mortality risk of preoperative invasive respiratory support to the noninvasive support group. Given the potential for noninvasive support to reduce intubation-related morbidities, such as preoperative sedation or neuromuscular blockade-associated complications, this finding alone may be beneficial. In either case, our findings suggest that careful consideration of preoperative patient selection for noninvasive support may be needed.

To our knowledge, this study is the first to provide a detailed analysis of how preoperative respiratory support affects outcomes in children undergoing cardiac surgery. While previous studies have reported mechanical ventilation as a risk factor for poor outcomes in this population, ours is the first to investigate associations between mode, timing, and escalation of preoperative respiratory support and outcomes. In addition, this study is the first to analyze these preoperative respiratory support parameters in neonates alone.

The key limitation of our study is our inability to control for the indication for preoperative respiratory support, which may vary in this population. For children with preoperative primary pulmonary disease, preoperative respiratory support may be a proxy for prolonged postoperative pulmonary convalescence, or worsened postoperative cardiopulmonary interactions.²¹ These children will need time for their lungs to recover before undergoing surgery. Among the neonates and children included in our cohort, some may have received preoperative ventilation to treat central apnea associated with prostaglandins administration, or to preempt loss of airway protection during transport or preoperative diagnostic procedures (e.g. brain MRI).^{22,23} Under these circumstances, preoperative respiratory support may contribute to pulmonary and overall morbidity, due to the possible need for sedation, the increased risk of pulmonary infections, the impaired ability to feed, and the potential for mechanical ventilation induced lung injury.²⁴ In the absence of underlying primary respiratory disease, children may receive preoperative respiratory support to reduce work of breathing and improve oxygen delivery in left heart failure.²⁵ By normalizing venous saturations, reducing work of breathing, eliminating negative intrathoracic pressures swings, and thereby decreasing systemic ventricular afterload, respiratory support may improve the preoperative state of children suffering from left heart failure by optimizing their oxygen supply-demand balance.²⁶ In children with parallel pulmonary and systemic circulations (e.g., hypoplastic left heart syndrome), respiratory support may be needed to control relative fractions of pulmonary and systemic blood flow and be indicative of worsened preoperative systemic cardiac output.²⁷ For children who receive ventilation to support their altered cardiovascular physiology, prolonged postoperative recovery may be due to the severity of their lesions, and respiratory support may actually improve their preoperative condition and subsequent outcomes.

Additional limitations of our study include several factors inherent in retrospective research. Although we controlled for a wide variety of patient, surgical, and respiratory support characteristics, there may be other risk factors contributing to variability in outcomes. For example, we were not able to obtain the duration of each type of preoperative respiratory support, which likely affects outcomes. In addition, this was a single-center study subject to the effects of local practice patterns and may not be generalizable to other care settings. Incorporation of preoperative respiratory support parameters into an existing registry for children undergoing cardiac surgery would provide useful information on how preoperative respiratory support affects outcomes. Used in combination with blood gas, hemodynamic, and chest x-ray data, research derived from this registry could provide greater insight into the associations between preoperative respiratory support, lung injury, and prolonged recovery.

In conclusion, we found that any preoperative respiratory support is associated with increased odds of prolonged postoperative length of stay following cardiac surgery compared to children who remain on room air preoperatively. Understanding how preoperative respiratory support affects children undergoing cardiac surgery may help guide preoperative management and surgical timing, inform prognostic conversations, and improve risk stratification models in this population.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors thank the Duke Clinical Research Institute and Duke University Medical Center for data acquisition and support.

Financial Support. This work was funded by the National Heart, Lung, and Blood Institute (grant 5R25HL13530402).

Conflicts of Interest. K.O.Z. receives financial support from Purdue Pharma. C.P.H. receives salary support for research from National Institute for Child Health and Human Development (NICHD) (K23HD090239), the U.S. government for his work in pediatric and neonatal clinical pharmacology (Government Contract HHSN267200700051C, PI: Benjamin, under the Best Pharmaceuticals for Children Act), and industry for drug development in children (www.dcri.duke.edu/research/coi.jsp).

References

1. The Society of Thoracic Surgeons. STS Congenital Heart Surgery Data Summary Children. 2018 Available from: https://www.sts.org/sites/default/files/documents/Congenital-STSExecSummary_Children_Spring2018.pdf
2. Jacobs JP, Mayer JE Jr, Pasquali SK, et al. The Society of Thoracic Surgeons Congenital Heart Surgery Database: 2019 Update on Outcomes and Quality. *Ann Thorac Surg* 2019; 107: 691–704. [PubMed: 30641069]
3. Jacobs ML, O'Brien SM, Jacobs JP, et al. An empirically based tool for analyzing morbidity associated with operations for congenital heart disease. *J Thorac Cardiovasc Surg* 2013; 145: 1046–1057.e1. [PubMed: 22835225]
4. Jacobs JP, O'Brien SM, Pasquali SK, et al. The importance of patient-specific preoperative factors: an analysis of the Society of Thoracic Surgeons Congenital Heart Surgery Database. *Ann Thorac Surg* 2014; 98: 1653–1658. [PubMed: 25262395]
5. Hornik CP, He X, Jacobs JP, et al. Complications after the Norwood operation: an analysis of The Society of Thoracic Surgeons Congenital Heart Surgery Database. *Ann Thorac Surg* 2011; 92: 1734–1740. [PubMed: 21937021]
6. Brown KL, Ridout DA, Goldman AP, Hoskote A, Penny DJ. Risk factors for long intensive care unit stay after cardiopulmonary bypass in children. *Crit Care Med* 2003; 31: 28–33. [PubMed: 12544989]
7. Pagowska-Klimek I, Pychynska-Pokorska M, Krajewski W, Moll JJ. Predictors of long intensive care unit stay following cardiac surgery in children. *Eur J Cardiothorac Surg* 2011; 40: 179–184. [PubMed: 21227714]
8. Jacobs JP, Jacobs ML, Austin EH 3rd, et al. Quality measures for congenital and pediatric cardiac surgery. *World J Pediatr Congenit Heart Surg* 2012; 3: 32–47. [PubMed: 23804682]
9. The Society of Thoracic Surgeons. STS Congenital Heart Surgery Database Data Specifications. Version 3.3. 6 26, 2015 Available from: https://www.sts.org/sites/default/files/documents/CongenitalDataSpecsV3_3_Updated.pdf.
10. O'Brien SM, Jacobs JP, Pasquali SK, et al. The Society of Thoracic Surgeons Congenital Heart Surgery Database Mortality Risk Model: Part 1-Statistical Methodology. *Ann Thorac Surg* 2015; 100: 1054–1062. [PubMed: 26245502]
11. O'Brien SM, Clarke DR, Jacobs JP, et al. An empirically based tool for analyzing mortality associated with congenital heart surgery. *J Thorac Cardiovasc Surg* 2009; 138: 1139–1153. [PubMed: 19837218]
12. Jacobs JP, Jacobs ML, Maruszewski B, et al. Initial application in the EACTS and STS Congenital Heart Surgery Databases of an empirically derived methodology of complexity adjustment to evaluate surgical case mix and results. *Eur J Cardiothorac Surg* 2012; 42: 775–779. [PubMed: 22700597]

13. Wernovsky G, Licht DJ. Neurodevelopmental outcomes in children with congenital heart disease-what can we impact? *Pediatr Crit Care Med* 2016; 17(8 Suppl 1): S232–S242. [PubMed: 27490605]
14. Newburger JW, Wypij D, Bellinger DC, et al. Length of stay after infant heart surgery is related to cognitive outcome at age 8 years. *J Pediatr* 2003; 143: 67–73. [PubMed: 12915826]
15. Pasquali SK, He X, Jacobs ML, et al. Excess costs associated with complications and prolonged length of stay after congenital heart surgery. *Ann Thorac Surg* 2014; 98: 1660–1666. [PubMed: 25201725]
16. Shamszad P, Gospin TA, Hong BJ, McKenzie ED, Petit CJ. Impact of preoperative risk factors on outcomes after Norwood palliation for hypoplastic left heart syndrome. *J Thorac Cardiovasc Surg* 2014; 147: 897–901. [PubMed: 23849462]
17. Stieh J, Fischer G, Scheewe J, et al. Impact of preoperative treatment strategies on the early perioperative outcome in neonates with hypoplastic left heart syndrome. *J Thorac Cardiovasc Surg* 2006; 131: 1122–1129.e2. [PubMed: 16678599]
18. Cheng HH, Almodovar MC, Laussen PC, et al. Outcomes and risk factors for mortality in premature neonates with critical congenital heart disease. *Pediatr Cardiol* 2011; 32: 1139–1146. [PubMed: 21713439]
19. Wysocki M, Antonelli M. Noninvasive mechanical ventilation in acute hypoxaemic respiratory failure. *Eur Respir J* 2001; 18: 209–220. [PubMed: 11510794]
20. Romans RA, Schwartz SM, Costello JM, et al. Epidemiology of noninvasive ventilation in pediatric cardiac ICUs. *Pediatr Crit Care Med* 2017; 18: 949–957. [PubMed: 28742724]
21. Alexander P Respiratory physiology for intensivists In: Ungerleider RM, Meliones JN, McMillan KN, Cooper DS, Jacobs JP (eds.) *Critical Heart Disease in Infants and Children* (3rd ed). Elsevier, Philadelphia, 2019, pp 134–149.
22. Gupta N, Leven L, Stewart M, Cheung M, Patel N. Transport of infants with congenital heart disease: benefits of antenatal diagnosis. *Eur J Pediatr* 2014; 173: 655–660. [PubMed: 24326382]
23. Meckler GD, Lowe C. To intubate or not to intubate? Transporting infants on prostaglandin E1. *Pediatrics* 2009; 123: e25–e30. [PubMed: 19064611]
24. Slutsky AS, Ranieri VM. Ventilator-induced lung injury. *N Engl J Med* 2013; 369: 2126–2136. [PubMed: 24283226]
25. Donnellan A, Justice L. Preoperative stabilization of infants with hypoplastic left heart syndrome before stage I palliation. *Crit Care Nurse* 2016; 36: 52–59.
26. Cooper DS, Jacobs JP, Chai PJ, et al. Pulmonary complications associated with the treatment of patients with congenital cardiac disease: consensus definitions from the Multi-Societal Database Committee for Pediatric and Congenital Heart Disease. *Cardiol Young* 2008; (18 Suppl 2): 215–221. [PubMed: 19063794]
27. Sarris GE, Balmer CS, Bonou PG, et al. Clinical guidelines for the management of patients with transposition of the great arteries with intact ventricular septum. *Eur J Cardiothorac Surg* 2017; 51(1): e1–e32 [PubMed: 28077506]

Figure 1.

Adjusted odds ratios for study outcomes based on maximum preoperative respiratory support mode, escalation in respiratory support, and mechanical ventilation on the day of surgery

Figure 2.

Adjusted odds ratios for study outcomes based on maximum preoperative respiratory support mode, escalation in respiratory support, and mechanical ventilation on the day of surgery in neonates

Table 1.

Patient and operative characteristics based on maximum respiratory support *

Characteristic	Maximum Respiratory Support			
	RA N=482 (60)	NC N=53 (7)	HFNC/NIV N=94 (12)	MV N=171 (21)
Age at surgery, days	560 (155, 1902)	14 (7, 170)	50 (8, 174)	22 (10, 68)
Female	238 (49)	21 (40)	37 (39)	69 (40)
Race				
White	271 (56)	31 (58)	46 (49)	84 (49)
Black	131 (27)	13 (25)	21 (22)	62 (36)
Other	80 (17)	9 (17)	27 (29)	25 (15)
Preoperative duration, days	0 (0, 0)	7 (3, 12)	6 (3, 16)	14 (6, 34)
Preoperative respiratory duration, days	NA	2 (1, 5)	4 (2, 10)	14 (6, 31)
Preoperative Factors				
Noncardiac congenital syndrome or chromosomal abnormality	116 (24)	15 (28)	37 (39)	43 (25)
Surfactant exposure	5 (1)	0 (0)	3 (3)	12 (7)
Cardiac complication	24 (5)	9 (17)	17 (18)	169 (99)
Noncardiac complication	126 (26)	25 (47)	55 (59)	137 (80)
Cyanotic Lesions	106 (22)	20 (38)	33 (35)	48 (28)
STAT mortality category				
1	203 (42)	4 (8)	11 (12)	11 (6)
2	168 (35)	12 (23)	27 (29)	73 (43)
3	34 (7)	3 (6)	12 (13)	16 (9)
4–5	77 (16)	34 (64)	44 (47)	71 (42)
CPB time, minutes	109 (54, 151)	148 (33, 210)	128 (16, 197)	0 (0, 182)

Abbreviations: RA, room air; NC, nasal cannula; HFNC, high flow nasal cannula; NIV, noninvasive ventilation; MV, mechanical ventilation; STAT, The Society of Thoracic Surgeons-European Association for Cardio-Thoracic Surgery; CPB, cardiopulmonary bypass.

* Counts (%) or median (25th, 75th percentiles).

Table 2.

Study outcomes based on maximum respiratory support mode, escalation in respiratory support, and mechanical ventilation on the day of surgery*

	Any Postoperative Complication			Postoperative Length of Stay		Operative Mortality		
Variable	No	Yes	p-value	Median (25 th , 75 th percentiles)	p-value	No	Yes	p-value
All Subjects								
Maximum support								
RA	452 (95)	23 (5)	0.001	6 (5, 10)	<0.001	475 (99)	7 (1)	<0.001
NC	46 (92)	4 (8)		20 (13, 39)		50 (94)	3 (6)	
HFNC/NIV	73 (88)	10 (12)		19 (12, 40)		83 (88)	11 (12)	
MV	129 (86)	21 (14)		42 (23, 117)		150 (88)	21 (12)	
Respiratory Support Escalation								
No	643 (93)	45 (7)	<0.001	9 (5, 21)	<0.001	688 (95)	33 (5)	0.021
Yes	57 (81)	13 (19)		36 (16, 75)		70 (89)	9 (11)	
MV on day of surgery								
No	622 (93)	46 (7)	0.051	9 (5, 19)	<0.001	668 (96)	26 (4)	<0.001
Yes	78 (87)	12 (13)		50 (24, 131)		90 (85)	16 (15)	
Neonates								
Maximum support								
RA	48 (92)	4 (8)	0.444	16 (6, 33)	<0.001	52 (96)	2 (4)	0.111
NC	26 (93)	2 (7)		22 (16, 41)		28 (90)	3 (10)	
HFNC/NIV	34 (85)	6 (15)		25 (16, 51)		40 (82)	9 (18)	
MV	91 (85)	16 (15)		47 (24, 128)		107 (90)	12 (10)	
Respiratory support escalation								
No	162 (91)	16 (9)	0.007	27 (15, 66)	0.299	178 (91)	18 (9)	0.416
Yes	37 (76)	12 (24)		34 (16, 57)		49 (86)	8 (14)	
MV on day of surgery								
No	135 (88)	18 (12)	0.873	23 (14, 47)	<0.001	153 (91)	15 (9)	0.439
Yes	64 (86)	10 (14)		56 (24, 134)		74 (87)	11 (13)	

Abbreviations: RA, room air; NC, nasal cannula; HFNC, high flow nasal cannula; NIV, noninvasive ventilation, MV, mechanical ventilation.

* Counts (%) or median (25th, 75th percentiles).

Table 3.

Adjusted odds ratios (95% CIs) for study outcomes based on maximum preoperative respiratory support mode, escalation in respiratory support, and mechanical ventilation on the day of surgery

Variable	Any Postoperative Complication	Postoperative Length of Stay	Operative Mortality
All Subjects			
Maximum support			
RA	Reference	Reference	Reference
NC	0.60 (0.18–1.96)	1.59 (1.49–1.69)	1.50 (0.32–7.11)
HFNC/NIV	1.15 (0.47–2.82)	1.77 (1.68–1.86)	3.69 (1.20–11.40)
MV	0.65 (0.23–1.80)	2.50 (2.35–2.67)	1.27 (0.39–4.13)
Respiratory support escalation			
No	Reference	Reference	Reference
Yes	1.26 (0.59–2.70)	1.19 (1.15–1.23)	0.87 (0.37–2.05)
MV on day of surgery			
No	Reference	Reference	Reference
Yes	0.94 (0.40–2.20)	1.37 (1.32–1.43)	1.43 (0.62–3.29)
Neonates			
Maximum support			
RA	Reference	Reference	Reference
NC	0.72 (0.10–5.13)	1.12 (1.03–1.22)	2.16 (0.26–17.73)
HFNC/NIV	2.19 (0.46–10.54)	1.32 (1.22–1.43)	8.97 (1.31–61.14)
MV	1.53 (0.29–8.03)	2.10 (1.92–2.29)	1.08 (0.17–6.93)
Respiratory support escalation			
No	Reference	Reference	Reference
Yes	2.29 (0.80–6.60)	1.35 (1.29–1.41)	1.31 (0.48–3.57)
MV on day of surgery			
No	Reference	Reference	Reference
Yes	0.89 (0.25–3.20)	1.18 (1.12–1.25)	1.18 (0.37–3.78)

Abbreviations: RA, room air; NC, nasal cannula; HFNC, high flow nasal cannula; NIV, noninvasive ventilation, MV, mechanical ventilation.