Published in final edited form as:

Arch Sex Behav. 2020 January ; 49(1): 275–286. doi:10.1007/s10508-019-01508-w.

Birth Cohort and Racial/Ethnic Differences in the Age of First Oral and Anal Sex among United States Men who have Sex with Men

Travis H. Sanchez, DVM, MPH¹, Mona Rai, MPH¹, Maria Zlotorzynska, PhD, MPH¹, Jeb Jones, PhD, MPH, MS¹, Patrick S. Sullivan, DMV, PhD¹

¹ Rollins School of Public Health, Emory University, Atlanta, Georgia, USA

Abstract

Earlier age of first sex has potential direct and indirect health effects later in life. Though there are multiple nationwide general population studies on ages of first sex, there is no such nationwide study of first male-male oral or anal sex among men who have sex with men (MSM). This may be important for understanding racial/ethnic disparities in HIV and sexually transmitted infection (STI) acquisition among young racial/ethnic minority MSM. Our study examined the birth cohort and racial/ethnic differences in ages of first male-male oral and anal sex using a diverse 2015 US nationwide sample of 10,217 sexually-active MSM. The mean age of first male-male oral sex was 18.0 years. Compared with older birth cohorts, those MSM born 1990–2000 were more likely to have younger age of first male-male oral sex. Compared to white MSM, Hispanic MSM and non-Hispanic black MSM were more likely to have younger age of first male-male oral sex with a man. The mean age of first male-male anal sex was 20.3 years. Compared with older birth cohorts, those MSM born 1990–2000 were more likely to have younger age of first male-male anal sex. Compared to white MSM, MSM of all other racial/ethnic groups were more likely to have younger age of first male-male anal sex. These findings emphasize the need for comprehensive and MSM-inclusive sexual health education for young teens and online sexual health resources for young gay, bisexual, queer and other MSM.

Keywords

MSM; gay; anal sex; oral sex; debut

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <https://www.springer.com/aam-terms-v1>

Corresponding Author and Reprint Request: Travis Sanchez, DVM, MPH, Rollins School of Public Health, Emory University, 1518 Clifton Road, Atlanta, GA 30322, Tel: +1 404 727 8403, Travis.Sanchez@emory.edu.

Conflict of Interest: The authors declare that they have no conflict of interest.

Publisher's Disclaimer: This Author Accepted Manuscript is a PDF file of an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Introduction

Earlier age of first sexual intercourse has potential direct health effects such as teen pregnancies and sexually transmitted infections (STI), and indirect health effects such as increased number of sexual partners, problem substance use, and mental health disorders later in life.(Heywood, Patrick, Smith, & Pitts, 2015; Kaestle, Halpern, Miller, & Ford, 2005; Sandfort, Orr, Hirsch, & Santelli, 2008; Vasilenko, Kugler, & Rice, 2016; Zimmer-Gembeck & Helfand, 2008) The average age of initiation of first sexual intercourse for US youth is approximately 17 years, with boys initiating sex less than a year before girls. (Abama JC, 2017; Cavazos-Rehg et al., 2009) The age of first sex is also younger for racial/ethnic minority youth and youth from economically disadvantaged backgrounds.(Abama JC, 2017; Biello, Ickovics, Niccolai, Lin, & Kershaw, 2013; Zimmer-Gembeck & Helfand, 2008) Studies have also shown that the prevalence of having ever engaged in sex is decreasing among some teens and age of first sex has been increasing for those in more recent birth cohorts.(Abama JC, 2017; Ethier, Kann, & McManus, 2018) When sexual intercourse is specified in these studies it is most typically defined as penetrative penile-vaginal sex. Other types of sexual intercourse, such as oral or anal intercourse, are either not included in the definition, not reported by gender of sex partner, or are not specifically reported in the literature for most studies. One study has reported initiation of oral or anal sex (though not by partner gender) and found that age of first oral sex occurs slightly later than penile-vaginal sex and approximately one year prior to penile-anal sex.(Halpern & Haydon, 2012)

There is even less published information on sexual identity differences in the age of initiation of sexual intercourse. Where data do exist, non-heterosexual males initiated sex at an earlier age than heterosexual males,(Brown, Masho, Perera, Mezuk, & Cohen, 2015; Glick et al., 2012) but these studies used broad definitions of sexual intercourse, did not differentiate initiation of sex with men, and conflated initiation of any type of sex among homosexual- or bisexual-identified respondents with age of first male-male sex. Other studies have focused more specifically on age of first sex among men who have sex with men (MSM), often for purposes of examining sexual health risks with highly specified samples such as MSM living with HIV infection or MSM in only a single major US city. (Glick & Golden, 2014; Outlaw et al., 2011) These studies also found that, similar to the general population studies, racial/ethnic minority MSM may also initiate sex with men at earlier ages. There have been no published nationwide studies with MSM that differentiated between age of initiation of oral and anal sex or have examined the potential birth cohort effect noted among youth in general.

Better understanding age of initiation of first male-male sexual intercourse is important to public health in several ways. MSM are at substantially higher risk of acquiring HIV and other STIs compared to others groups.(Centers for Disease Control and Prevention, 2016a, 2016b; Singh, Song, Johnson, McCray, & Hall, 2018) Condomless anal intercourse poses the greatest per-act sexual transmission risk for HIV infection.(Patel et al., 2014) In addition to the increased risk for HIV infection, MSM experience high rates of oropharyngeal and rectal STIs which are often asymptomatic and require substantial public health resources to diagnose and treat.(Centers for Disease Control and Prevention, 2017) The risk is magnified

for racial/ethnic minority MSM who have the highest rates of newly acquired HIV infection, mainly due to high HIV incidence among racial/ethnic minority MSM 13 to 24 years of age. (Singh et al., 2018) Earlier age of initiation of sexual intercourse with men may result in a longer period at risk of HIV/STI acquisition, and may partly explain these racial/ethnic disparities. (Maulsby et al., 2014) More recent agent-based models have been created to better understand HIV and STI transmission and acquisition, particularly for young racial and ethnic minority MSM. (Goodreau et al., 2017; Jones et al., 2019) These models are also being increasingly used to estimate potential population-level effects of various prevention strategies, such as increased screening and pre-exposure prophylaxis. (Goodreau et al., 2018; Hamilton et al., 2018; Jenness et al., 2016; Jenness et al., 2019) The models currently don't address the potential for demographic group variation in the cumulative time at risk of HIV infection. One of the first steps in better understanding whether the agent-based models are improved with these parameters is determining whether there are racial, ethnic, and birth cohort differences in age of first sex.

To better understand the ongoing sexual health needs of US MSM, we have conducted the annual American Men's Internet Survey (AMIS) which reaches a large and geographically diverse sample. (Sanchez, Sineath, Kahle, Tregear, & Sullivan, 2015) In 2015, we asked AMIS participants what age they were when they first had sex with a man. The purpose of the present study is to describe the ages of first oral and anal sex with a man for this diverse sample of sexually active US MSM, and determine whether these ages of first sex significantly differed by birth cohort or race/ethnicity.

Methods

Participants and Procedures

AMIS is conducted in annual rounds with a goal of at least 10,000 complete surveys from eligible US MSM each year. The AMIS methods have been previously reported in detail. (Sanchez et al., 2015) Briefly, participants are recruited each year through convenience sampling from a variety of websites and through social media applications using banner ads and email blasts to members (hereafter referred to as "ads"). Men who click on ads are taken directly to the survey website hosted on a secure server administered by SurveyGizmo (Boulder, CO, USA). Participants are eligible to participate if they are age 15 years or older, male sex at birth, reside in the US, and report oral or anal sex with a man at least once at any time in the past. Participants who meet the eligibility criteria and consent to participate in the study start the online survey immediately. The survey is self-administered, can be taken on a computer or mobile device and is comprised of questions on demographics, sexual behaviors, substance use, HIV and sexually transmitted infection testing and diagnosis, and use of HIV prevention services. No incentive is provided to participants.

Measures

The AMIS round of data collection conducted from September 2015 to April 2016 included questions on the age of first oral and anal sex with a man: "How old were you the first time you had oral sex (mouth on the penis) with a man?" and "How old were you the first time you had anal sex (penis in the butt) with a man?" For this study, only responses of age of

10+ years were used for age of initiation and implausible sex debut ages (e.g., where the age of sexual debut was reported to be greater than their age at the time of survey completion) were recoded as missing. Year of birth was calculated from self-reported age at the time of the survey and collapsed into a four-level birth cohort variable: those born between 1950–1969, between 1970–1979, between 1980–1989, and between 1990–2000. Race and ethnicity were asked using the standard 2-question US Office of Management and Budget 1997 standard format in which persons are asked to report their race with multiple selections possible and are asked separately whether they identify as Hispanic or Latino. (Office of Management and Budget, 1997) Responses to these questions are then categorized into a single race/ethnicity measure using the same scheme employed by federal HIV/STI case surveillance systems. (Centers for Disease Control and Prevention, 2018, 2019) Because of relatively few responses for some race categories, we used the 4-level categorization scheme typical in these studies. (An, Wejnert, Bernstein, & Paz-Bailey, 2017; Shadaker, Magee, Paz-Bailey, & Hoots, 2017; Singh, Mitsch, & Wu, 2017) Those who identify as Hispanic or Latino are categorized as Hispanic in the combined measure regardless of response to the race question. Those who identify their race as only black, African American or only white are categorized into black, non-Hispanic and white, non-Hispanic categories. All other responses (American Indian or Alaska Native, Asian, Native Hawaiian or Other Pacific Islander, other race, or multiple responses) were categorized into the other or multiple race category. Self-reported HIV status was determined from responses to questions about having ever had an HIV test, results of the most recent HIV test, and having ever had a positive HIV test. Participants were categorized as self-reported HIV-positive, HIV-negative, or unknown status. Residential county population density was categorized using the National Center for Health Statistics (NCHS) Rural-Urban classification scheme. (Ingram & Franco, 2014) We further collapsed these categories into a binary variable: urban (urban, suburban, medium metropolitan) and rural (small metropolitan, micropolitan, and non-core).

Analyses

Eligible participants were included in analyses if they were unduplicated by IP address, completed the survey, ever had sex with a man in the past 12 months, provided a valid US ZIP code, and were between the ages of 15–59 at the time of survey completion. Log-rank tests were used to compare the equality of survival until the first oral or anal sex with a man across race/ethnicity and birth cohort groups. These 2 independent measures were chosen because they are key measures of disparity in HIV and STI infection among MSM, and they were the only demographic factors collected in AMIS that were likely contemporary with adolescent sexual development. All other measures were collected for more recent or current timeframes and pose substantial temporarily concerns for a cross-sectional analysis of outcomes occurring in adolescence for most participants. Wilcoxon two-sample tests were also used to compare the mean ages of first oral or anal sex with a man across groups. For participants who reported ages of first oral and anal sex with a man, the age difference was also calculated as age at first anal sex minus age at first oral sex age, and mean age differences were compared between groups using Wilcoxon two-sample tests. Cox proportional hazard modeling was used to assess independence of the associations between race/ethnicity or birth cohort and age of first oral sex and anal sex. Interactions between race/ethnicity and birth cohort in relation to age of first oral and anal sex were assessed with

likelihood ratio tests. For significant interactions, results are presented as hazard ratios and 95% confidence intervals within the race/ethnicity by birth cohort strata. Analyses were all conducted using SAS 9.4 (SAS Institute, Cary, NC) with statistical significance denoted at $\alpha=0.05$.

Results

Of the 10,217 AMIS participants who had sex with a man in the past 12 months, 8,331 (81.5%) answered the questions on age of first oral or anal sex with a man and were included in these analyses. MSM participants were most commonly under 35 years of age (birth cohorts 1980–1989 and 1990–2000), from the Southern US, from urban or suburban counties, HIV negative, and non-Hispanic white (Table 1). Overall, 99.9% (8320/8331) of MSM had ever had oral sex with a man and 95.2% (7931/8331) had ever had anal sex with a man – 95.1% (7922/8331) of MSM had both oral and anal sex with a man.

Age of First Oral Sex with a Man

The mean age of first oral sex with a man was 18.0 years. Overall, 34.1% of MSM had oral sex with a man by age 15, 65.9% by age 18, and 92.8% by age 25 (Table 2). Compared with older birth cohorts, those MSM born 1990–2000 (age 15 to 20) were significantly more likely to have younger age of first oral sex with a man. The birth cohort differences in age of first oral sex show that though a greater percentage of the oldest cohort (1950–1969) may have had oral sex in pre-teen years, the younger cohorts (1980–1989 and 1990–2000) had greater cumulative percentage of having engaged in oral sex by age 18 (Figure 1). This resulted in those MSM born 1990–2000 (mean first oral sex = 16.8 years; Table 2) having first oral sex nearly 1 year younger than the next closest birth cohort (1980–1989) and 2.6 years younger than the oldest cohort (1950–1969).

Compared to non-Hispanic white MSM, Hispanic MSM and non-Hispanic black MSM were significantly more likely to have younger age of first oral sex with a man (Table 3). These racial/ethnic differences in age of first oral sex appear in pre-teen non-Hispanic black MSM and are present by the mid-teens for Hispanic MSM (Figure 2), resulting in both of these groups having first oral sex an average of 1 year earlier than non-Hispanic white MSM (means of 17.2 years versus 18.2 years, respectively; Table 3). Some of the birth cohort differences may partly explain the racial/ethnic differences in earlier age of first oral sex with a man (Table 4 and Supplemental Table). For instance, the black versus white MSM differences in age of first oral sex were only significant for those born in the 1950–1969 and 1980–1989 cohorts, whereas the Hispanic versus white MSM differences were only significant for those in the 2 youngest cohorts (1980–1989 and 1990–2000).

Age of First Anal Sex with a Man

The mean age of first anal sex with a man was 20.3 years. Overall, 9.6% of MSM had anal sex with a man by age 15, 46.1% by age 18, and 83.1% by age 25 (Table 5). Compared with older birth cohorts, those MSM born 1990–2000 were significantly more likely to have younger age of first anal sex with a man. These birth cohort differences in age of first anal sex appear in mid- to late-teen MSM (Figure 3). This resulted in those MSM born in 1990–

2000 (mean first anal sex = 17.8 years, Table 5) having first anal sex with a man 1.6 years younger than the next closest birth cohort (1980–1989) and 6.2 years younger than the oldest cohort (1950–1969).

Compared to non-Hispanic white MSM, MSM of all other racial/ethnic groups were significantly more likely to have younger age of first anal sex with a man (Table 6). Compared to non-Hispanic white MSM (mean first anal sex = 20.9 years), Hispanic MSM on average had anal sex 2.3 years younger (mean 18.3 years), black MSM had anal sex 1.9 years younger (mean 19.0 years), and MSM of other or multiple races had anal sex 1.3 years younger (mean 19.6 years). These racial/ethnic differences in age of first anal sex by race/ethnicity appear by the early teens (Figure 4). Even when accounting for birth cohort differences, racial/ethnic minority MSM remained significantly more likely to have engaged in anal sex at earlier ages compared to non-Hispanic white MSM (Table 7 and Supplemental Table).

Age Difference between First Oral Sex and First Anal Sex with a Man

On average, MSM first had oral sex with a man 2.5 years (SD = 4.9 years) before they first had anal sex with a man. Compared with white MSM (mean [SD] = 2.8 [5.4] years difference between first oral and anal sex), black MSM (1.8 [3.4] years), Hispanic/Latino MSM (1.5 [3.1] years), and MSM of other or multiple races (1.9 [3.7] years) had significantly (all $p < 0.001$) shorter time differences between first oral and anal sex. Compared with the youngest MSM born in 1990–2000 (mean [SD] = 1.1 [1.8] years difference between first oral and anal sex), those born 1980–1989 (1.8 [2.9] years), 1970–1979 (2.8 [4.5] years), and 1950–1969 (4.8 [7.7] years) had significantly (all $p < 0.001$) longer time differences between first oral and anal sex with a man.

Discussion

Most sexually active MSM in our study had oral sex with a man by age 18 and had anal sex with a man by age 20. Between 15% (anal sex) and 34% (oral sex) of MSM had initiated sex with a man by age 15. The mean ages of first oral and anal sex in our study were higher than those previously reported, but may be due to differences in the sampled populations and definitions of sexual initiation. For instance, a study of HIV-positive adolescent MSM found mean age of 14.5 for first sex of any type with male partners. (Outlaw et al., 2011) An earlier study with MSM from only a few major cities found mean age of 16.5 for first sex of any type with male partners and 19.6 for first anal sex. An even smaller study with adolescent MSM recruited to examine HIV prevention needs found mean age of 15.0 for first oral sex and age 16.6 for first anal sex with male partners. Finally, a nationwide online survey that was similar in design to ours but was conducted 3 years earlier and with only approximately 800 MSM, found mean age of 15.6 for first sex of any type with male partners and 17.6 for first anal sex. In addition to the larger scope and more recent sample of our study, some of the demographic group findings may partly explain the age differences between the present study and previously published studies.

MSM who were born in more recent years were significantly more likely than those born in earlier years to have younger age at first oral or anal sex with a man. The time period

between initiation of oral and anal sex was also significantly shorter for MSM born more recently. This could mean that studies which mainly or more substantially sampled younger MSM may have younger average ages of first sex with male partners. This birth cohort effect in younger age of first sex for younger MSM has been previously reported in a Swiss study, (Balthasar, Jeannin, & Dubois-Arber, 2009) but was not found in the one other small US study that examined this association.(Nelson, Gamarel, Pantalone, Carey, & Simoni, 2016) There are no predominant theories about why MSM who were born in the past few decades may be initiating sex at younger ages, but it could be due to de-criminalization and changing social norms about male-male sex, increase in male teens' ability to find male sex partners using social media, or increased willingness of youth to disclose their early male-male sexual experiences. It should also be noted that the trend we found of decreasing age of first sex is opposite that seen in general population studies of US adolescents in more recent years, which has been attributed to improvements in sexual health education programs in schools.(Lindberg & Maddow-Zimet, 2012) It's possible that MSM adolescents are not equally benefiting from these same programs which may struggle to provide comprehensive and inclusive sexual health information for non-heterosexual youth.(Hall, Jones, Witkemper, Collins, & Rodgers, 2019) Further study regarding the underlying processes that are driving this change is warranted, particularly because young MSM experience disproportionately high burden of new HIV infection.(Singh et al., 2018)

Racial/ethnic minority MSM were significantly more likely than white MSM to have younger age of first oral or anal sex with a man. These differences were not entirely explained by the overlapping birth cohort effects. The time period between initiation of oral and anal sex was also significantly shorter for racial/ethnic minority MSM than it was for white MSM. Other previous studies of US MSM that have examined racial/ethnic differences in age of first sex with male partners have not found a significant difference like we have, potentially due to their limited sample size and limited geographic scope compared to nationwide AMIS.(Glick & Golden, 2014; Outlaw et al., 2011; Sanchez et al., 2015; Warren et al., 2008) Our findings are consistent with large-scale general population studies that have found the racial/ethnic differences in age of first sex.(Alabama JC, 2017; Biello et al., 2013; Zimmer-Gembeck & Helfand, 2008) The mechanisms by which racial and ethnic identities are associated with early sexual histories are poorly understood, but identities are likely a proxy indicator for other cultural, familial, social, structural and socioeconomic factors.(Cavazos-Rehg et al., 2009; Corbie-Smith, Henderson, Blumenthal, Dorrance, & Estroff, 2008; Outlaw et al., 2011)

The approximately 2 year earlier age of initiation of anal sex for black MSM and Hispanic/Latino MSM compared to white MSM could also be playing a role in increased HIV and STI incidence seen in these groups of young MSM.(Garofalo, Hotton, Kuhns, Gratzner, & Mustanski, 2016; Kaestle et al., 2005; Sandfort et al., 2008; Singh et al., 2018) Recent agent-based modeling of HIV and STI transmission among young US MSM had not previously accounted for duration at risk of infection because there were no consistently available parameters, particularly those among young MSM by racial and ethnic groups.(Beck, Birkett, Armbruster, & Mustanski, 2015; Goodreau et al., 2017; Jones et al., 2019) The next step will be incorporate these parameters into those agent-based models to determine whether variation in duration at risk of infection plays a key role in racial and

ethnic disparities in HIV and STI infection. These updated models can also be used to re-estimate population-level effects of key prevention strategies. (Hamilton et al., 2018; Jenness et al., 2016)

Several limitations of this study should be noted. AMIS data are not generalizable to all MSM in the US or to all MSM online. The convenience sampling approach, although conducted with multiple types of websites and social networking applications intended to increase sample diversity, increases the potential for selection or enrollment biases. The surveys only involved self-report of behaviors. It is possible that earlier age of first sex may be less socially desirable and therefore may be misreported as being older. These behaviors potentially occurred many years prior to the interview and could be prone to differential recall bias where those with initiation of sex closer to the interview date may be more accurate. There were also many participants who reported having oral or anal sex at very young ages, possibly due to sexual abuse rather than consensual sex. Participants were not directly asked whether those first sexual encounters were consensual, but a subset were asked questions from the Adverse Childhood Experiences scale which assesses sexual abuse of minors by an adult family member, friend or other. (Anda, Butchart, Felitti, & Brown, 2010) We conducted a sensitivity analysis (data not reported) that excluded participants who reported these experiences. Average ages of first oral and anal sex increased overall and for all demographic groups by approximately 1 year. We found the racial/ethnic differences in age of oral sex were somewhat attenuated, but racial differences in initiation age of anal sex and birth cohort differences in oral and anal sex initiation age remained significant. Some studies of age of first sex have attempted to either restrict their definition to consensual sex or define consensual sex separately from non-consensual sexual abuse of minors, but none have been done with large samples of MSM. (Finer & Philbin, 2013; Glick & Golden, 2014; Holmes, 2009; Sandfort et al., 2008) There is also no published research on how coercion may factor into age-appropriate sexual relationships of adolescent MSM, but one study of young adult males showed a high prevalence (67%) of sexual violence with partners. (McKool, Stephenson, Winskell, Teten Tharp, & Parrott, 2017) A new adolescent couples HIV prevention trial is designed to capture some of this relationship data and may help inform better measurement of early sexual partnership dynamics. (Gamarel, Darbes, Hightow-Weidman, Sullivan, & Stephenson, 2019) Finally, the study was intended to identify age of initiation of sex with men, but we did not ask questions about the age of initiation of sex with females to allow us to determine an absolute age of first sex for comparability with other previous general population studies.

Conclusion

A substantial proportion of US MSM begin sexual activity with men in their teen years and younger birth cohorts are even more likely to first have sex at these earlier ages. This finding combined with younger age of sexual initiation for racial/ethnic minority MSM may partly explain disparities in HIV acquisition for young black and Hispanic/Latino MSM and should be further explored. Because the majority of sexually active MSM begin engaging in sex with males in their teens, our study also emphasizes the need for comprehensive and MSM-inclusive sexual health education for young teens, such as San Francisco's Healthy Me, Healthy Us middle school curriculum, (San Francisco Unified School District, 2018) and

online sexual health resources specifically for young gay, bisexual, queer and other MSM, such as Northwestern University's QueerSexEd.(Mustanski, Greene, Ryan, & Whitton, 2015; "Queer Sex Ed," 2018)

Human Subjects Statement

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. Informed consent was obtained from all individual participants included in the study.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding: The study was funded by a grant from the MAC AIDS Fund and by the National Institutes of Health [P30AI050409] – the Emory Center for AIDS Research.

References

- Abama JC, M. G. (2017). Sexual Activity and Contraceptive Use Among Teenagers in the United States, 2011–2015. Retrieved from
- An Q, Wejnert C, Bernstein K, & Paz-Bailey G (2017). Syphilis Screening and Diagnosis Among Men Who Have Sex With Men, 2008–2014, 20 U.S. Cities. *J Acquir Immune Defic Syndr*, 75 Suppl 3, S363–s369. doi:10.1097/qai.0000000000001412 [PubMed: 28604440]
- Anda RF, Butchart A, Felitti VJ, & Brown DW (2010). Building a framework for global surveillance of the public health implications of adverse childhood experiences. *Am J Prev Med*, 39(1), 93–98. doi:10.1016/j.amepre.2010.03.015 [PubMed: 20547282]
- Balthasar H, Jeannin A, & Dubois-Arber F (2009). First anal intercourse and condom use among men who have sex with men in Switzerland. *Arch Sex Behav*, 38(6), 1000–1008. doi:10.1007/s10508-008-9382-5 [PubMed: 18561013]
- Beck EC, Birkett M, Armbruster B, & Mustanski B (2015). A Data-Driven Simulation of HIV Spread Among Young Men Who Have Sex With Men: Role of Age and Race Mixing and STIs. *J Acquir Immune Defic Syndr*, 70(2), 186–194. doi:10.1097/qai.0000000000000733 [PubMed: 26102448]
- Biello KB, Ickovics J, Niccolai L, Lin H, & Kershaw T (2013). Racial differences in age at first sexual intercourse: residential racial segregation and the black-white disparity among U.S. adolescents. *Public Health Rep*, 128 Suppl 1, 23–32. doi:10.1177/00333549131282s103 [PubMed: 23450882]
- Brown MJ, Masho SW, Perera RA, Mezuk B, & Cohen SA (2015). Sex and sexual orientation disparities in adverse childhood experiences and early age at sexual debut in the United States: results from a nationally representative sample. *Child Abuse Negl*, 46, 89–102. doi:10.1016/j.chiabu.2015.02.019 [PubMed: 25804435]
- Cavazos-Rehg PA, Krauss MJ, Spitznagel EL, Schootman M, Bucholz KK, Peipert JF, . . . Bierut LJ (2009). Age of sexual debut among US adolescents. *Contraception*, 80(2), 158–162. doi:10.1016/j.contraception.2009.02.014 [PubMed: 19631791]
- Centers for Disease Control and Prevention. (2016a, 11/1/2016). Diagnoses of HIV Infection in the United States and Dependent Areas, 2015. Retrieved from <https://www.cdc.gov/hiv/pdf/library/reports/surveillance/cdc-hiv-surveillance-report-2015-vol-27.pdf>
- Centers for Disease Control and Prevention. (2016b, 10/1/2016). Sexually Transmitted Disease Surveillance 2015. Retrieved from <https://www.cdc.gov/std/stats15/STD-Surveillance-2015-print.pdf>

- Centers for Disease Control and Prevention. (2017). STDs in Men Who Have Sex with Men. Retrieved from <https://www.cdc.gov/std/stats17/msm.htm>
- Centers for Disease Control and Prevention. (2018). HIV Surveillance by Race/Ethnicity. Retrieved from <https://www.cdc.gov/hiv/pdf/library/slidesets/cdc-hiv-surveillance-race-ethnicity-2017.pdf>
- Centers for Disease Control and Prevention. (2019). HIV by Race/Ethnicity. Retrieved from <https://www.cdc.gov/hiv/group/raciaethnic/index.html>
- Corbie-Smith G, Henderson G, Blumenthal C, Dorrance J, & Estroff S (2008). Conceptualizing race in research. *J Natl Med Assoc*, 100(10), 1235–1243. doi:10.1016/s0027-9684(15)31470-x [PubMed: 18942287]
- Ethier KA, Kann L, & McManus T (2018). Sexual Intercourse Among High School Students - 29 States and United States Overall, 2005–2015. *MMWR Morb Mortal Wkly Rep*, 66(5152), 1393–1397. doi:10.15585/mmwr.mm665152a1 [PubMed: 29300723]
- Finer LB, & Philbin JM (2013). Sexual initiation, contraceptive use, and pregnancy among young adolescents. *Pediatrics*, 131(5), 886–891. doi:10.1542/peds.2012-3495 [PubMed: 23545373]
- Gamarel KE, Darbes LA, Hightow-Weidman L, Sullivan P, & Stephenson R (2019). The Development and Testing of a Relationship Skills Intervention to Improve HIV Prevention Uptake Among Young Gay, Bisexual, and Other Men Who Have Sex With Men and Their Primary Partners (We Prevent): Protocol for a Randomized Controlled Trial. *JMIR Res Protoc*, 8(1), e10370. doi: 10.2196/10370 [PubMed: 30602433]
- Garofalo R, Hotton AL, Kuhns LM, Gratzner B, & Mustanski B (2016). Incidence of HIV Infection and Sexually Transmitted Infections and Related Risk Factors Among Very Young Men Who Have Sex With Men. *J Acquir Immune Defic Syndr*, 72(1), 79–86. doi:10.1097/qai.0000000000000933 [PubMed: 26745827]
- Glick SN, & Golden MR (2014). Early male partnership patterns, social support, and sexual risk behavior among young men who have sex with men. *AIDS Behav*, 18(8), 1466–1475. doi:10.1007/s10461-013-0678-7 [PubMed: 24356869]
- Glick SN, Morris M, Foxman B, Aral SO, Manhart LE, Holmes KK, & Golden MR (2012). A comparison of sexual behavior patterns among men who have sex with men and heterosexual men and women. *J Acquir Immune Defic Syndr*, 60(1), 83–90. doi:10.1097/QAI.0b013e318247925e [PubMed: 22522237]
- Goodreau SM, Hamilton DT, Jenness SM, Sullivan PS, Valencia RK, Wang LY, . . . Rosenberg ES (2018). Targeting Human Immunodeficiency Virus Pre-Exposure Prophylaxis to Adolescent Sexual Minority Males in Higher Prevalence Areas of the United States: A Modeling Study. *J Adolesc Health*, 62(3), 311–319. doi:10.1016/j.jadohealth.2017.09.023 [PubMed: 29248392]
- Goodreau SM, Rosenberg ES, Jenness SM, Luisi N, Stansfield SE, Millett GA, & Sullivan PS (2017). Sources of racial disparities in HIV prevalence in men who have sex with men in Atlanta, GA, USA: a modelling study. *Lancet HIV*, 4(7), e311–e320. doi:10.1016/s2352-3018(17)30067-x [PubMed: 28431923]
- Hall WJ, Jones BLH, Witkemper KD, Collins TL, & Rodgers GK (2019). State Policy on School-based Sex Education: A Content Analysis Focused on Sexual Behaviors, Relationships, and Identities. *Am J Health Behav*, 43(3), 506–519. doi:10.5993/ajhb.43.3.6 [PubMed: 31046882]
- Halpern CT, & Haydon AA (2012). Sexual timetables for oral-genital, vaginal, and anal intercourse: sociodemographic comparisons in a nationally representative sample of adolescents. *Am J Public Health*, 102(6), 1221–1228. doi:10.2105/ajph.2011.300394 [PubMed: 22571710]
- Hamilton DT, Goodreau SM, Jenness SM, Sullivan PS, Wang LY, Dunville RL, . . . Rosenberg ES (2018). Potential Impact of HIV Preexposure Prophylaxis Among Black and White Adolescent Sexual Minority Males. *Am J Public Health*, 108(S4), S284–s291. doi:10.2105/ajph.2018.304471 [PubMed: 30383415]
- Heywood W, Patrick K, Smith AM, & Pitts MK (2015). Associations between early first sexual intercourse and later sexual and reproductive outcomes: a systematic review of population-based data. *Arch Sex Behav*, 44(3), 531–569. doi:10.1007/s10508-014-0374-3 [PubMed: 25425161]
- Holmes WC (2009). Potential Impact of Childhood Sexual Abuse Among Males, by Whether It Occurred at or After Sexual Debut. *J Interpers Violence*. doi:10.1177/0886260508329122

- Ingram DD, & Franco SJ (2014). 2013 NCHS Urban-Rural Classification Scheme for Counties. *Vital Health Stat* 2(166), 1–73.
- Jenness SM, Goodreau SM, Rosenberg E, Beylerian EN, Hoover KW, Smith DK, & Sullivan P (2016). Impact of the Centers for Disease Control's HIV Preexposure Prophylaxis Guidelines for Men Who Have Sex With Men in the United States. *J Infect Dis*, 214(12), 1800–1807. doi:10.1093/infdis/jiw223 [PubMed: 27418048]
- Jenness SM, Maloney KM, Smith DK, Hoover KW, Goodreau SM, Rosenberg ES, . . . Sullivan PS (2019). Addressing Gaps in HIV Preexposure Prophylaxis Care to Reduce Racial Disparities in HIV Incidence in the United States. *Am J Epidemiol*, 188(4), 743–752. doi:10.1093/aje/kwy230 [PubMed: 30312365]
- Jones J, Weiss K, Mermin J, Dietz P, Rosenberg ES, Gift TL, . . . Jenness SM (2019). Proportion of Incident Human Immunodeficiency Virus Cases Among Men Who Have Sex With Men Attributable to Gonorrhea and Chlamydia: A Modeling Analysis. *Sex Transm Dis*, 46(6), 357–363. doi:10.1097/olq.0000000000000980 [PubMed: 31095100]
- Kaestle CE, Halpern CT, Miller WC, & Ford CA (2005). Young age at first sexual intercourse and sexually transmitted infections in adolescents and young adults. *Am J Epidemiol*, 161(8), 774–780. doi:10.1093/aje/kwi095 [PubMed: 15800270]
- Lindberg LD, & Maddow-Zimet I (2012). Consequences of sex education on teen and young adult sexual behaviors and outcomes. *J Adolesc Health*, 51(4), 332–338. doi:10.1016/j.jadohealth.2011.12.028 [PubMed: 22999833]
- Maulsby C, Millett G, Lindsey K, Kelley R, Johnson K, Montoya D, & Holtgrave D (2014). HIV among Black men who have sex with men (MSM) in the United States: a review of the literature. *AIDS Behav*, 18(1), 10–25. doi:10.1007/s10461-013-0476-2 [PubMed: 23620241]
- McKool M, Stephenson R, Winskell K, Teten Tharp A, & Parrott D (2017). Peer Influence on IPV by Young Adult Males: Investigating the Case for a Social Norms Approach. *J Interpers Violence*, 886260517725735. doi:10.1177/0886260517725735
- Mustanski B, Greene GJ, Ryan D, & Whitton SW (2015). Feasibility, acceptability, and initial efficacy of an online sexual health promotion program for LGBT youth: the Queer Sex Ed intervention. *J Sex Res*, 52(2), 220–230. doi:10.1080/00224499.2013.867924 [PubMed: 24588408]
- Nelson KM, Gamarel KE, Pantalone DW, Carey MP, & Simoni JM (2016). Sexual Debut and HIV-Related Sexual Risk-Taking by Birth Cohort Among Men Who Have Sex with Men in the United States. *AIDS Behav*, 20(10), 2286–2295. doi:10.1007/s10461-016-1330-0 [PubMed: 26860630]
- Office of Management and Budget. (1997). Standards for Maintaining, Collecting, and Presenting Federal Data on Race and Ethnicity. Retrieved from <https://www.doi.gov/pmb/eco/Data-Standards>
- Outlaw AY, Phillips G 2nd, Hightow-Weidman LB, Fields SD, Hidalgo J, Halpern-Felsher B, & Green-Jones M (2011). Age of MSM sexual debut and risk factors: results from a multisite study of racial/ethnic minority YMSM living with HIV. *AIDS Patient Care STDS*, 25 Suppl 1, S23–29. doi:10.1089/apc.2011.9879 [PubMed: 21711140]
- Patel P, Borkowf CB, Brooks JT, Lasry A, Lansky A, & Mermin J (2014). Estimating per-act HIV transmission risk: a systematic review. *AIDS*, 28(10), 1509–1519. doi:10.1097/QAD.0000000000000298 [PubMed: 24809629]
- Queer Sex Ed. (2018). Retrieved from <http://www.queersexed.org/>
- San Francisco Unified School District. (2018). Healthy Me. Health Us. Retrieved from <https://sfusdhealtheducation.org/ms-healthy-healthy-us/>
- Sanchez TH, Sineath RC, Kahle EM, Tregear SJ, & Sullivan PS (2015). The Annual American Men's Internet Survey of Behaviors of Men Who Have Sex With Men in the United States: Protocol and Key Indicators Report 2013. *JMIR Public Health Surveill*, 1(1), e3. doi:10.2196/publichealth.4314 [PubMed: 27227126]
- Sandfort TG, Orr M, Hirsch JS, & Santelli J (2008). Long-term health correlates of timing of sexual debut: results from a national US study. *Am J Public Health*, 98(1), 155–161. doi:10.2105/ajph.2006.097444 [PubMed: 18048793]
- Shadaker S, Magee M, Paz-Bailey G, & Hoots BE (2017). Characteristics and Risk Behaviors of Men Who Have Sex With Men and Women Compared With Men Who Have Sex With Men-20 US

- Cities, 2011 and 2014. *J Acquir Immune Defic Syndr*, 75 Suppl 3, S281–s287. doi:10.1097/qai.0000000000001403 [PubMed: 28604429]
- Singh S, Mitsch A, & Wu B (2017). HIV Care Outcomes Among Men Who Have Sex With Men With Diagnosed HIV Infection - United States, 2015. *MMWR Morb Mortal Wkly Rep*, 66(37), 969–974. doi:10.15585/mmwr.mm6637a2 [PubMed: 28934185]
- Singh S, Song R, Johnson AS, McCray E, & Hall HI (2018). HIV Incidence, Prevalence, and Undiagnosed Infections in U.S. Men Who Have Sex With Men. *Ann Intern Med*, 168(10), 685–694. doi:10.7326/m17-2082 [PubMed: 29554663]
- Vasilenko SA, Kugler KC, & Rice CE (2016). Timing of First Sexual Intercourse and Young Adult Health Outcomes. *J Adolesc Health*, 59(3), 291–297. doi:10.1016/j.jadohealth.2016.04.019 [PubMed: 27265422]
- Warren JC, Fernandez MI, Harper GW, Hidalgo MA, Jamil OB, & Torres RS (2008). Predictors of unprotected sex among young sexually active African American, Hispanic, and White MSM: the importance of ethnicity and culture. *AIDS Behav*, 12(3), 459–468. doi:10.1007/s10461-007-9291-y [PubMed: 17721725]
- Zimmer-Gembeck M, & Helfand M (2008). Ten Years of Longitudinal Research on U.S. Adolescent Sexual Behavior: Developmental Correlates of Sexual Intercourse, and the Importance of Age, Gender and Ethnic Background. *Developmental Review*, 28, 153–224. doi:10.1016/j.dr.2007.06.001

Figure 1.

Age of Initiaion of Oral Sex with a Man by Birth Cohort of MSM Participants in the American Men's Internet Survey, United States, 2015

Figure 2.
Age of Initiaion of Oral Sex with a Man by Race/Ethnicity of MSM Participants in the
American Men’s Internet Survey, United States, 2015

Figure 3.
Age of Initiaion of Anal Sex with a Man by Birth Cohort of MSM Participants in the
American Men's Internet Survey, United States, 2015

Figure 4.
Age of Initiaion of Anal Sex with a Man by Race/Ethnicity of MSM Participants in the
American Men's Internet Survey, United States, 2015

Table 1.

Characteristics of MSM Participants in the American Men's Internet Survey by Race/Ethnicity, United States, 2015

Participant Characteristics	Total		Black, non-Hispanic		Hispanic		White, non-Hispanic		Other or multiple races	
	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
Age (years)										
15–19	907	(10.9)	29	(5.3)	184	(15.2)	591	(10.2)	103	(13.6)
20–24	1,786	(21.4)	72	(13.2)	320	(26.4)	1,194	(20.5)	200	(26.3)
25–29	1,477	(17.7)	102	(18.8)	249	(20.5)	972	(16.7)	154	(20.3)
30–34	789	(9.5)	70	(12.9)	133	(11.0)	517	(8.9)	69	(9.1)
35–39	565	(6.8)	63	(11.6)	101	(8.3)	348	(6.0)	53	(7.0)
40–44	498	(6.0)	50	(9.2)	64	(5.3)	338	(5.8)	46	(6.1)
45–49	673	(8.1)	49	(9.0)	58	(4.8)	516	(8.9)	50	(6.6)
50–54	884	(10.6)	59	(10.8)	61	(5.0)	714	(12.3)	50	(6.6)
55–59	752	(9.0)	50	(9.2)	44	(3.6)	623	(10.7)	35	(4.6)
Region										
Midwest	1,753	(21.0)	77	(14.2)	124	(10.2)	1,412	(24.3)	140	(18.4)
Northeast	1,669	(20.0)	95	(17.5)	196	(16.1)	1,240	(21.3)	138	(18.2)
South	3,040	(36.5)	305	(56.1)	476	(39.2)	2,026	(34.9)	233	(30.7)
West	1,863	(22.4)	66	(12.1)	414	(34.1)	1,135	(19.5)	248	(32.6)
U.S. dependent areas	6	(0.1)	1	(0.2)	4	(0.3)	0	(0.0)	1	(0.1)
Population density^a										
Rural	1,537	(18.4)	67	(12.3)	126	(10.4)	1,180	(20.3)	164	(21.6)
Urban/Suburban	6,785	(81.4)	476	(87.5)	1,083	(89.2)	4,631	(79.7)	595	(78.3)
Self-reported HIV status										
Positive	712	(8.5)	116	(21.3)	130	(10.7)	406	(7.0)	60	(7.9)
Negative	5,782	(69.4)	355	(65.3)	797	(65.7)	4,112	(70.7)	518	(68.2)
Unknown	1,837	(22.1)	73	(13.4)	287	(23.6)	1,295	(22.3)	182	(23.9)
TOTAL	8,331		544	(6.5)	1,214	(14.6)	5,813	(69.8)	760	(9.1)

^aThere were 9 participants missing information needed to determine the population density of the area where they lived.

Table 2.

Age of Initiation of Oral Sex with a Man by Birth Cohort of MSM Participants in the American Men's Internet Survey, United States, 2015

	Birth Cohort									
	All		1950–1969		1970–1979		1980–1989		1990–2000	
	(N=8,331)		(N=2,164)		(N=1,085)		(N=2,038)		(N=3,044)	
Had oral sex by age x ^a	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
10	156	(1.9)	76	(3.5)	20	(1.8)	33	(1.6)	27	(0.9)
11	321	(3.9)	140	(6.5)	46	(4.2)	70	(3.4)	65	(2.1)
12	873	(10.5)	346	(16.0)	125	(11.5)	194	(9.5)	208	(6.8)
13	1436	(17.2)	525	(24.3)	191	(17.6)	317	(15.6)	403	(13.2)
14	2118	(25.4)	694	(32.1)	284	(26.2)	476	(23.4)	664	(21.8)
15	2837	(34.1)	872	(40.3)	357	(32.9)	638	(31.3)	970	(31.9)
16	3734	(44.8)	1046	(48.3)	441	(40.6)	849	(41.7)	1398	(45.9)
17	4451	(53.4)	1158	(53.5)	502	(46.3)	1041	(51.1)	1750	(57.5)
18	5486	(65.9)	1321	(61.0)	601	(55.4)	1284	(63.0)	2280	(74.9)
19	6149	(73.8)	1433	(66.2)	679	(62.6)	1459	(71.6)	2578	(84.7)
20	6605	(79.3)	1515	(70.0)	743	(68.5)	1568	(76.9)	2779	(91.3)
21	7036	(84.5)	1624	(75.0)	808	(74.5)	1700	(83.4)	2904	(95.4)
22	7298	(87.6)	1689	(78.0)	863	(79.5)	1780	(87.3)	2966	(97.4)
23	7481	(89.8)	1731	(80.0)	901	(83.0)	1847	(90.6)	3002	(98.6)
24	7612	(91.4)	1760	(81.3)	925	(85.3)	1898	(93.1)	3029	(99.5)
25	7728	(92.8)	1800	(83.2)	953	(87.8)	1938	(95.1)	3037	(99.8)
Log-rank p-value ^b			<0.001		<0.001		<0.001		ref	
Mean [SD] age of first oral sex	18.0	[5.7]	19.4	[8.8]	18.9	[6.07]	17.7	[4.14]	16.8	[2.8]
Rank-sum p-value ^c			<0.001		<0.001		<0.001		ref	

^aNumber and percentages are cumulative. For example by 18 years of age, 5,486 (65.9%) total participants ever had oral sex with a male partner.

^bLog-rank p-value comparing equality of survival until first oral sex of one birth cohort to referent group of those MSM born 1990–2000.

^cRank-sum p-value by Wilcoxon two-sample test comparing mean age of first oral sex of one birth cohort to referent group of those MSM born 1990–2000.

Table 3.

Age of Initiation of Oral Sex with a Man by Race/Ethnicity of MSM Participants in the American Men's Internet Survey, United States, 2015

	Race/Ethnicity									
	All		Black, non-Hispanic		Hispanic		White, non-Hispanic		Other or multiple races	
	(N=8,331)		(N=544)		(N=1,214)		(N=5,813)		(N=760)	
Had oral sex by age x ^a	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
10	156	(1.9)	15	(2.8)	21	(1.7)	105	(1.8)	15	(2.0)
11	321	(3.9)	31	(5.7)	51	(4.2)	211	(3.6)	28	(3.7)
12	873	(10.5)	78	(14.3)	122	(10.0)	592	(10.2)	81	(10.7)
13	1436	(17.2)	118	(21.7)	202	(16.6)	983	(16.9)	133	(17.5)
14	2118	(25.4)	152	(27.9)	297	(24.5)	1483	(25.5)	186	(24.5)
15	2837	(34.1)	202	(37.1)	433	(35.7)	1948	(33.5)	254	(33.4)
16	3734	(44.8)	259	(47.6)	572	(47.1)	2569	(44.2)	334	(43.9)
17	4451	(53.4)	315	(57.9)	709	(58.4)	3030	(52.1)	397	(52.2)
18	5486	(65.9)	374	(68.8)	900	(74.1)	3715	(63.9)	497	(65.4)
19	6149	(73.8)	414	(76.1)	974	(80.2)	4200	(72.3)	561	(73.8)
20	6605	(79.3)	442	(81.3)	1026	(84.5)	4532	(78.0)	605	(79.6)
21	7036	(84.5)	468	(86.0)	1089	(89.7)	4842	(83.3)	637	(83.8)
22	7298	(87.6)	485	(89.2)	1117	(92.0)	5031	(86.5)	665	(87.5)
23	7481	(89.8)	502	(92.3)	1142	(94.1)	5146	(88.5)	691	(90.9)
24	7612	(91.4)	510	(93.8)	1149	(94.6)	5248	(90.3)	705	(92.8)
25	7728	(92.8)	516	(94.9)	1165	(96.0)	5326	(91.6)	721	(94.9)
Log-rank p-value ^b			<0.001		<0.001		ref		0.104	
Mean [SD] age of first oral sex	18.0	[5.7]	17.2	[4.6]	17.2	[4.4]	18.2	[6.1]	17.8	[5.3]
Rank-sum p-value ^c			0.004		<0.001		ref		0.555	

^aNumber and percentages are cumulative. For example by 18 years of age, 5,486 (65.9%) total participants ever had oral sex with a male partner.

^bLog-rank p-value comparing equality of survival until first oral sex of one racial/ethnic group to referent group of white MSM.

^cRank-sum p-value by Wilcoxon two-sample test comparing mean age of first oral sex of one racial/ethnic group to referent group of white MSM.

Table 4.

Cox Proportional Hazard Ratios and 95% Confidence Intervals of Younger Age of First Oral Sex with a Man, by Race/Ethnicity and Birth Cohort of MSM Participants in the American Men's Internet Survey, United States, 2015

Race/Ethnicity	Birth Cohort			
	1950–1969	1970–1979	1980–1989	1990–2000
Black, non-Hispanic	1.205 (1.021, 1.423)	1.173 (0.957, 1.438)	1.370 (1.164, 1.613)	1.123 (0.934, 1.350)
Hispanic	1.162 (0.980, 1.378)	1.054 (0.886, 1.253)	1.135 (1.010, 1.275)	1.237 (1.126, 1.360)
Other or multiple races	1.021 (0.851, 1.225)	0.921 (0.747, 1.136)	1.099 (0.944, 1.280)	0.978 (0.872, 1.096)
White, non-Hispanic	Ref	Ref	Ref	Ref

Table 5.

Age of Initiation of Anal Sex with a Man by Birth Cohort of MSM Participants in the American Men's Internet Survey, United States, 2015

	Birth Cohort									
	All		1950–1969		1970–1979		1980–1989		1990–2000	
	(N=8,331)		(N=2,164)		(N=1,085)		(N=2,038)		(N=3,044)	
Had oral sex by age x ^a	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
10	31	(0.4)	10	(0.5)	6	(0.6)	9	(0.4)	6	(0.2)
11	78	(0.9)	29	(1.3)	11	(1.0)	22	(1.1)	16	(0.5)
12	243	(2.9)	89	(4.1)	44	(4.1)	53	(2.6)	57	(1.9)
13	470	(5.6)	161	(7.4)	68	(6.3)	104	(5.1)	137	(4.5)
14	800	(9.6)	232	(10.7)	107	(9.9)	180	(8.8)	281	(9.2)
15	1286	(15.4)	333	(15.4)	159	(14.7)	297	(14.6)	497	(16.3)
16	1981	(23.8)	457	(21.1)	213	(19.6)	451	(22.1)	860	(28.3)
17	2677	(32.1)	586	(27.1)	257	(23.7)	620	(30.4)	1214	(39.9)
18	3843	(46.1)	785	(36.3)	371	(34.2)	941	(46.2)	1746	(57.4)
19	4655	(55.9)	902	(41.7)	468	(43.1)	1145	(56.2)	2140	(70.3)
20	5235	(62.8)	1021	(47.2)	542	(50.0)	1301	(63.8)	2371	(77.9)
21	5826	(69.9)	1148	(53.0)	640	(59.0)	1471	(72.2)	2567	(84.3)
22	6210	(74.5)	1251	(57.8)	704	(64.9)	1588	(77.9)	2667	(87.6)
23	6474	(77.7)	1320	(61.0)	758	(69.9)	1683	(82.6)	2713	(89.1)
24	6710	(80.5)	1375	(63.5)	810	(74.7)	1770	(86.8)	2755	(90.5)
25	6927	(83.1)	1472	(68.0)	854	(78.7)	1837	(90.1)	2764	(90.8)
Log-rank p-value ^b			<0.001		<0.001		<0.001		ref	
Mean (SD) age of first anal sex	20.3 [6.7]		24.0 [10.1]		21.5 [6.3]		19.4 [4.04]		17.8 [2.6]	
Rank-sum p-value ^c			<0.001		<0.001		<0.001		ref	

^aNumber and percentages are cumulative. For example by 18 years of age, 3,843 (46.1%) participants ever had anal sex with a male partner.

^bLog-rank p-value comparing equality of survival until first anal sex of one birth cohort to referent group of those MSM born 1990–2000.

^cRank-sum p-value by Wilcoxon two-sample test comparing mean age of first anal sex of one birth cohort to referent group of those MSM born 1990–2000.

Table 6.

Age of Initiation of Anal Sex with a Man by Race/Ethnicity of MSM Participants in the American Men's Internet Survey, United States, 2015

	Race/Ethnicity									
	All		Black, non-Hispanic		Hispanic		White, non-Hispanic		Other or multiple races	
	(N=8,331)		(N=544)		(N=1,214)		(N=5,813)		(N=760)	
Had oral sex by age x ^a	No.	(%)	No.	(%)	No.	(%)	No.	(%)	No.	(%)
10	31	(0.4)	5	(0.9)	7	(0.6)	18	(0.3)	1	(0.1)
11	78	(0.9)	13	(2.4)	13	(1.1)	46	(0.8)	6	(0.8)
12	243	(2.9)	28	(5.1)	44	(3.6)	147	(2.5)	24	(3.2)
13	470	(5.6)	43	(7.9)	85	(7.0)	299	(5.1)	43	(5.7)
14	800	(9.6)	62	(11.4)	144	(11.9)	507	(8.7)	87	(11.4)
15	1286	(15.4)	102	(18.8)	238	(19.6)	807	(13.9)	139	(18.3)
16	1981	(23.8)	157	(28.9)	362	(29.8)	1255	(21.6)	207	(27.2)
17	2677	(32.1)	202	(37.1)	494	(40.7)	1722	(29.6)	259	(34.1)
18	3843	(46.1)	303	(55.7)	713	(58.7)	2458	(42.3)	369	(48.6)
19	4655	(55.9)	343	(63.1)	814	(67.1)	3064	(52.7)	434	(57.1)
20	5235	(62.8)	371	(68.2)	897	(73.9)	3484	(59.9)	483	(63.6)
21	5826	(69.9)	410	(75.4)	978	(80.6)	3907	(67.2)	531	(69.9)
22	6210	(74.5)	438	(80.5)	1016	(83.7)	4189	(72.1)	567	(74.6)
23	6474	(77.7)	453	(83.3)	1049	(86.4)	4374	(75.2)	598	(78.7)
24	6710	(80.5)	467	(85.8)	1069	(88.1)	4547	(78.2)	627	(82.5)
25	6927	(83.1)	486	(89.3)	1089	(89.7)	4704	(80.9)	648	(85.3)
Log-rank p-value ^b			<0.001		<0.001		ref		<0.001	
Mean (SD) age of first anal sex	20.3	[6.7]	19.0	[4.8]	18.6	[4.8]	20.9	[7.3]	19.6	[5.8]
Rank-sum p-value ^c			<0.001		<0.001		ref		<0.001	

^aNumber and percentages are cumulative. For example by 18 years of age, 3,843 (46.1%) participants ever had anal sex with a male partner.

^bLog-rank p-value comparing equality of survival until first anal sex of one racial/ethnic group to referent group of white MSM.

^cRank-sum p-value by Wilcoxon two-sample test comparing mean age of first anal sex of one racial/ethnic group to referent group of white MSM.

Table 7.

Cox Proportional Hazard Ratios and 95% Confidence Intervals of Younger Age of First Anal Sex with a Man, by Race/Ethnicity and Birth Cohort of MSM Participants in the American Men's Internet Survey, United States, 2015

Race/Ethnicity	Birth Cohort			
	1950–1969	1970–1979	1980–1989	1990–2000
Black, non-Hispanic	1.685 (1.425, 1.992)	1.260 (1.026, 1.547)	1.457(1.237, 1.716)	1.165 (0.961, 1.411)
Hispanic	1.460 (1.228, 1.735)	1.241 (1.043, 1.477)	1.181 (1.050, 1.328)	1.282 (1.162, 1.414)
Other or multiple races	1.290 (1.074, 1.550)	0.966 (0.782, 1.194)	1.092 (0.936, 1.274)	0.901 (0.797, 1.017)
White, non-Hispanic	Ref	Ref	Ref	Ref