Published in final edited form as:

Neurosci Biobehav Rev. 2019 March ; 98: 122–134. doi:10.1016/j.neubiorev.2019.01.004.

Chronobiology of limbic seizures: Potential mechanisms and prospects of chronotherapy for mesial temporal lobe epilepsy

Daniel Leite Góes Gitai^{a,b}, Tiago Gomes de Andrade^c, Ygor Daniel Ramos dos Santos^b, Sahithi Attaluri^a, Ashok K. Shetty^{a,d,*}

^aInstitute for Regenerative Medicine, Department of Molecular and Cellular Medicine, Texas A&M University, College Station, Texas, USA

^bInstitute of Biological Sciences and Health, Federal University of Alagoas, Maceio, Alagoas, Brazil

^cFaculty of Medicine, Federal University of Alagoas, Maceió, Alagoas, Brazil

^dResearch Service, Olin E. Teague Veterans' Medical Center, Central Texas Veterans Health Care System, Temple, Texas, USA

Abstract

Mesial Temporal Lobe Epilepsy (mTLE) characterized by progressive development of complex partial seizures originating from the hippocampus is the most prevalent and refractory type of epilepsy. One of the remarkable features of mTLE is the rhythmic pattern of occurrence of spontaneous seizures, implying a dependence on the endogenous clock system for seizure threshold. Conversely, circadian rhythms are affected by epilepsy too. Comprehending how the circadian system and seizures interact with each other is essential for understanding the pathophysiology of epilepsy as well as for developing innovative therapies that are efficacious for better seizure control. In this review, we confer how the temporal dysregulation of the circadian clock in the hippocampus combined with multiple uncoupled oscillators could lead to periodic seizure occurrences and comorbidities. Unraveling these associations with additional research would help in developing chronotherapy for mTLE, based on the chronobiology of spontaneous seizures. Notably, differential dosing of antiepileptic drugs over the circadian period and/or strategies that resynchronize biological rhythms may substantially improve the management of seizures in mTLE patients.

Keywords

Seizure; Clock genes; Circadian rhythms; Epilepsy; Hippocampus; Temporal lobe epilepsy

*Corresponding author at: Institute for Regenerative Medicine, Texas A&M Health Science Center, College of Medicine, 1114 TAMU, 206 Olsen Boulevard, College Station, TX, 77843, USA. shetty@medicine.tamhsc.edu (A.K. Shetty).

Publisher's Disclaimer: Department of defense, department of veterans affairs, and United States government disclaimer
The contents of this article suggest the views of authors and do not represent the views of the Department of Defense, U.S. Department of Veterans Affairs, or the United States Government.

1. Introduction

Circadian rhythms are central properties of living organisms tailored to anticipate and dynamically respond to a cyclical environment. Changes enacted by alternation between day and night on factors such as light, temperature, sleep, food and other periodic ecological elements instituted selective pressures to the evolving life to develop intrinsic biological rhythms. The enormity of these interfaces is apparent from the widespread presence of circadian rhythms in nature, from bacteria to complex organisms, and its impact on many biological processes (Bhadra et al., 2017). Notably, the discovery of the molecular mechanism of circadian rhythms in flies in the mid 80's won the Nobel Prize in Physiology and Medicine (2017), showing how fast and substantially the role of circadian rhythms has been integrated to the management of human health.

Transcriptional-translational feedback loops (TTFL) modulated at different levels control the circadian clock in eukaryotes (Takahashi, 2017) (Fig. 1). Several components of the mammalian clock are highly conserved in flies. The core-clock transcription factors BMAL1 and CLOCK heterodimerize in the cytoplasm and form BMAL1:CLOCK complex, which translocates to the nucleus and instigates transcription of target genes by interacting with E-box promoters. The target genes comprise the core clock-controlled genes (CCGS), *Period* (*Per1*, *Per2*, *Per3*) and *Cryptochrome* (*Cry1*, *Cry2*). The protein products of PERs and CRYs fluctuate across the 24-hour cycle by inhibiting their promoters operating in a complex negative feedback loop. Specifically, the PER:CRY heterodimers translocate into the nucleus and restrain their transcription by reducing the activity of the BMAL1-CLOCK complex. The accumulation of PER:CRY proteins is modulated through phosphorylation by Casein Kinase 1 (CK1) and degradation by Skp/Cullin/F-box (SCF) ubiquitin ligases. An added regulatory loop is produced when BMAL1:CLOCK heterodimers activate the transcription of two retinoic acid-related orphan receptor response elements, RORA (activator of BMAL1 transcription) and REV-ERBA (repressor of BMAL1 transcription). Both RORA and REV-ERBA partake in binding to rev responsive element (RRE) present in BMAL1 promoter to regulate the expression of BMAL1. Thus, RORA and REV-ERBA are respectively involved in positive and negative regulation of the circadian oscillation of BMAL1.

Furthermore, the Nuclear Factor, Interleukin 3 Regulated (NFIL3) gene represses the CCG, D-box binding PAR bZIP transcription factor (DBP), to regulate the rhythmic transcription of D-box elements containing genes, RORA, and ERBA (McDaniel et al., 2011). The occupancy of promoter sites by transcription factors (TFs) and RNA polymerase present a circadian pattern, regulating the transcription globally (Koike et al., 2012). It is estimated that up to 50% of the genome is transcribed in a circadian manner in different tissues, with high specificity for each tissue (Zhang et al., 2014). The other post-transcriptional/translational mechanisms also regulate circadian rhythms (Kojima and Green, 2015).

Circadian rhythms in mammals comprise a multi-oscillatory system, in which practically every single cell displays an autonomous genetically programmed rhythm, coordinated by a master oscillator, the hypothalamic suprachiasmatic nucleus (SCN) (Honma, 2018; Takahashi et al., 2008). The SCN receives photic inputs from the retinohypothalamic tract (RHT) and nonphotic cues from other afferent inputs, to orchestrate the endogenous rhythms

to environmental signals (LeGates et al., 2014) (Ramkisoensing and Meijer, 2015). Intrinsically photosensitive retinal ganglion cells (ipRGC) expressing melanopsin transduce light signals, independent of the canonical forming image pathways, to the SCN, inducing the expression of Per1 and Per2 genes (LeGates et al., 2014). This photic input resets the circadian molecular clock in the SCN with consequent coordination of several oscillators in the brain and other organs, synchronizing the internal body clocks to the environmental light. Misalignments of these external signals and the internal clock, as occurs in shift workers, or alterations in the clock genes have been correlated with disruptions in the circadian rhythms and many metabolic and neuropsychiatric disorders (Ramkisoensing and Meijer, 2015); (Smolensky et al., 2016).

The circadian activity of several brain regions has been described in humans and animal models (Frank et al., 2013; Li et al., 2013; Rath et al., 2014). In the hippocampus, the inherent circadian timekeeping capacity is fundamental for rhythmic fluctuation of hippocampus-dependent memory and learning (Snider et al., 2018) as well as its plasticity, including patterning, spine density, generation of new neurons (adult neurogenesis) and long-term potentiation (Chaudhury et al., 2005; Ikeno et al., 2013, 2014; O'Callaghan et al., 2012; Tamai et al., 2008). Therefore, structural and physiological alterations in the hippocampus in certain disorders may be linked to complex interactions with the circadian system.

Epilepsy, typified by spontaneous recurrent seizures (SRS), is one of the most common neurological disorders (Devinsky et al., 2018). Mesial temporal lobe epilepsy (mTLE), characterized by the progressive development of the complex partial seizures originating from the temporal lobe foci such as the hippocampus, is seen in ~30% of epilepsy patients (Fisher et al., 1998). Cognitive, memory and mood impairments are the behavioral comorbidities in mTLE, which are associated with multiple adverse changes and plasticity in the hippocampus. These include substantially diminished normal hippocampal neurogenesis, continued aberrant migration of newly born neurons into the dentate hilus, dispersion of dentate granule cells, considerable loss of GABA-ergic interneurons, aberrant mossy fiber sprouting, chronic neuroinflammation, gene expression reorganization (Shetty and Turner, 1999; Hattiangady et al., 2004; Shetty et al., 2005; Romcy-Pereira et al., 2008; Vannest et al., 2008; Kuruba et al., 2009; Waldau et al., 2010; Hattiangady and Shetty, 2010; Long et al., 2017; Shetty, 2014). The interaction between circadian rhythms and epileptic seizure occurs in a bi-directional manner as evidenced by the influence of environmental zeitgebers (food and light) on both onset and severity of seizures (Cho, 2012; Mirzoev et al., 2012; Roberts and Keith, 1994; Stewart et al., 2001; Weiss et al., 1993), and by the influence of seizures on temporal oscillation of body temperature (Quigg et al., 1999), locomotor activity (Stewart and Leung, 2003), sleep architecture (Kothare and Zarowski, 2011) and hormones (Molina-Carballo et al., 1994; van Campen et al., 2015; Yalín et al., 2006). A remarkable feature of this crosstalk is that the seizures, although unpredictable, present a rhythmic pattern of occurrence in mTLE and animal models of mTLE (Hofstra and de Weerd, 2009). Discerning how the time control system and seizures influence each other is of great interest for not only understanding the pathophysiology of epilepsy but also for developing new therapeutic strategies.

In this review, we focused on mTLE to provide an overview of the temporal remapping of the hippocampus as a potential contributor to the rhythmic pattern of limbic seizures. Moreover, we discuss several studies that investigated the effects of circadian rhythm disturbances on the epileptogenic process to bring insights regarding the role of the circadian system on limbic seizures and its implication for chronotherapy in epilepsy. The key terminologies and phrases used in this review are defined in Box 1.

2. Potential mechanisms for the temporal pattern of limbic seizures

Circadian regulatory system modulates the occurrence of partial complex seizures in certain types of epilepsy, which is evidenced from observations that seizures have a time-of-day preference, depending on its origin (Durazzo et al., 2008; Spencer et al., 2016). Patients afflicted with mTLE display a 24-hour non-uniform distribution of seizure occurrence with either a unimodal (afternoon) or bimodal (early morning and afternoon) time peak (Kalevias et al., 2011; Karafin et al., 2010; Mirzoev et al., 2012; Nzwalo et al., 2016). Since sex and age modulate circadian rhythms in both humans and animal models (De Nobrega and Lyons, 2018; Bailey and Silver, 2014; Yan and Silver, 2016; Díaz-Morales and Parra-Robledo, 2018), it is essential to consider their potential interference in the temporal pattern of seizures. Age and sex discrepancies influence the highest occurrence of seizures in mesial temporal sclerosis (Passarelli and Castro, 2015). However, in a recent study using a large dataset of unspecified epilepsy, no significant differences in temporal patterns were seen between men and women for seizure cycles at different time scales (Karoly et al., 2018). Thus, additional studies are necessary to examine the effects of sex and age on circadian rhythms and temporal pattern of seizures.

Spontaneous seizures in kainic acid and pilocarpine models of mTLE have been found to occur in a daily pattern with a peak incidence befalling between 2–4 PM in multiple studies (Arida et al., 1999; Bertram and Cornett, 1994; Cavalheiro et al., 1991; Matzen et al., 2012; Tchekalarova et al., 2010, 2011; Van Nieuwenhuyse et al., 2015). Although the comparison across species is complicated because of nocturnal habits of rodents, it is remarkable that most time peaks in both humans and animal models occur in the daytime period. Unraveling the components involved in ictogenic mechanisms related to circadian rhythmicity across species would likely suggest new strategies for epilepsy management. On the other hand, it is important to note that the investigations for detecting the seizure periodicity in humans were performed by using a population-based approach, reflecting the overlapping seizure peaks across heterogeneous populations. When transferred to the individual level, the comparison between time peaks from humans and experimental models is not straightforward. Indeed, the endogenous and environmental factors composing mTLE pathophysiology give rise to individual variability, ultimately leading to patient-specific seizure cycle patterns that deviate from the mean in some cases (Karoly et al., 2018). Studies using long-term 24/7 video-EEG combined with algorithms for periodicity analysis will allow a better characterization of biological rhythms of spontaneous seizures. A study in an experimental model demonstrated that the circadian periodicity of seizures persists in a constant darkness condition (Quigg et al., 2000), suggesting a dependence of the endogenous clock system for epilepsy threshold. As the temporal pattern of seizures progresses uniquely depending on the localization of epileptogenic focus, determining the

temporal pattern of seizures in different conditions has considerable significance. In mTLE, it is likely that increased electrical activity in the hippocampus modifies the oscillation pattern of clock genes and uncouples central and peripheral oscillators (Fig. 2).

2.1. Dysregulation of clock genes and clock-controlled genes in the hippocampus

A recent study has shown that the circadian expression of the core clock genes *Bmal1*, *Cry1*, and *Cry2* persist in the hippocampus of epileptic animals (Matos et al., 2018), although with phase and amplitude changes. The dysregulation of core clock genes can affect the 24 h availability of molecules that play roles directly or indirectly in the electrical activity of the hippocampus with consequences on the occurrence, frequency, and duration of seizures. A higher and lower circadian amplitude of excitatory and inhibitory CCGS can cause rhythmic fluctuations in the inhibitory/excitatory balance (Matzen et al., 2012; Talathi et al., 2009). Accordingly, several studies have pointed out that seizure threshold and severity (Golombek et al., 1992; Roberts and Keith, 1994; Stewart et al., 2001; Weiss et al., 1993), as well as the anticonvulsant efficacy of antiepileptic drugs (Khedhaier et al., 2017), depends on the daytime or dark/light conditions. Essential regulatory systems involved in the modulation of seizure thresholds present circadian activity in the hippocampus including the melatoninergic (Mushhoff et al., 2002) and the mammalian/mechanistic target of rapamycin (mTOR) pathways (Saraf et al., 2014).

Melatonin levels present higher amplitude in patients with epilepsy (Molina-Carballo et al., 1994) and the limbic seizures occur most frequently 6h before the melatonin onset (Hofstra et al., 2011). Patients with intractable epilepsy have low baseline interictal melatonin levels that increase dramatically following seizures (Bazil et al., 2000). Melatonin receptors (MT1 and MT2), which have an inhibitory effect on CNS, show a significant amplitude change (decrease at ZT0 for MT1 and increase at ZT18 for MT2) in the hippocampus of epileptic rats (Rocha et al., 2017). Both melatonin and agomelatine (an agonist of MT1 and MT2 receptors) have been reported to have anticonvulsant actions in models of acute and chronic seizures (Aguilar et al., 2012; Dastgheib and Moezi, 2014). Also, RORA, a binding site of melatonin, has 24-h expression pattern in the hippocampus of control and epileptic rats (Rocha et al., 2016). Both mRNA and protein expression for RORA showed a decline in the acute and latent phases in the pilocarpine model. This pattern of expression may be involved in the epileptogenic processes. The ROR mRNA was also reduced in the chronic phase of epilepsy (Rocha et al., 2016). It is highly likely that RORA contributes to the temporal pattern of limbic seizures as RORA is an essential component in the regulation of circadian rhythms (Solt and Burris, 2012) and has anti-inflammatory (Delerive et al., 2001) and antioxidant activities (Boukhtouche et al., 2006).

The mTOR signaling, engaged on a broader reorganization of gene expression (Laplanche and Sabatini, 2012; Solt and Burris, 2012), is overactivated in the hippocampus of both mTLE patients and animal models of mTLE (Talós et al., 2018; Zeng et al., 2009). The inhibition of mTOR by Rapamycin results in both antiepileptic and antiepileptogenic effect in the mTLE model (Broekaart et al., 2017; Citraro et al., 2016; Huang et al., 2012). As the mTOR activity and its key products display circadian fluctuation in several tissues, with a higher peak during the day (Cao et al., 2010; Cao and Obrietan, 2010), the overstimulation caused

by epilepsy could change the circadian oscillation of some downstream genes that act in excitatory/inhibitory balance. Investigating the temporal expression of these molecules in epilepsy models should bring valuable insights (Cho, 2012). Furthermore, a series of recent publications have shown that mTOR, which is regulated by the circadian clock (Hughes et al., 2009; Zhang et al., 2014), plays a significant role in regulating central and peripheral clock function including period and amplitude (Liu et al., 2018; Ramanathan et al., 2018). Interestingly, at night, when the limbic seizures are less frequent, there are high levels of melatonin and low activity of mTOR, suggesting that these systems may contribute to the temporal occurrence of seizures.

Chiang and colleagues conducted a quantitative proteome analysis of the normal hippocampus throughout a 24 h cycle from C57BL/6/J male mice maintained in continuous darkness regimen (Chiang et al., 2017). They used JTK_CYCLE, an algorithm that identifies and characterizes cycling variables in large data sets (Hughes et al., 2010), to identify significant patterns of rhythmic expression at six time-points. The authors reported that 51 proteins out of 3052 displayed circadian rhythmicity. An individual functional annotation with QuickGO (Binns et al., 2009) revealed that Gene Ontology (GO) terms corresponded to central pathways in epileptogenesis, including *action potential* (GO:0001508), *positive regulation of neuron apoptotic process* (GO:0043525), *positive regulation of synapse structural plasticity* (GO:0051835), *positive regulation of cytosolic calcium ion concentration* (GO:0007204) and *neuron fate determination* (GO:0048664). Verification of these proteins against two epilepsy genetic databases (CarpeDB and epiGAD) and Pubmed (using the search terms “epilepsy OR seizure”) showed that 5 of them (*Cdc42*, *Chga*, *Gng3*, *Scarb2*, and *Sptan1*) have links to epileptogenic processes.

Remarkably, both *Cdc42* and *Gng3* have been shown to affect the seizure threshold. *Cdc42* encodes for cell division cycle 42 GTP-binding protein, which is involved in the regulation of neuronal structure (Bustelo et al., 2007; Govek et al., 2011). Moreover, *cdc42* protein is overexpressed in the hippocampus of mTLE patients (Xiao et al., 2007) and *cdc42* inhibition reduces seizure severity in the lithium/pilocarpine rat model of epilepsy (Zhang et al., 2015). The *cdc42* shows a lower expression peak at CT18 in the hippocampus, the time window displaying the lowest seizure frequency (early night) within 24 h. Furthermore, analysis with STRING database version 10.5, a repository that gathers information about physical and functional (indirect) protein-protein interactions (Szklarczyk et al., 2017), showed that *cdc42* protein has a functional interaction with mTOR protein (Combined score: 0.974). Indeed, mTOR is a downstream signaling target of *cdc42* (Chou and Blenis, 1996; Fang et al., 2003) and *cdc42* depletion inhibits mTOR activation (Endo et al., 2009). The anticonvulsant effect from both *cdc42* and mTOR inhibition (Citraro et al., 2016; Zhang et al., 2015) indicate a functional link between these proteins in the seizure threshold. *Gng3* encodes for G-protein gamma 3 subunit, which contributes to the specificity of receptor signaling pathways associated with a G-protein function (Schwindinger et al., 2004). Furthermore, *Gng3* knockout mice (*Gng3*^{-/-}) displays increased susceptibility to seizures (Schwindinger et al., 2004), due to defective GABA_B receptor signaling (Schwindinger et al., 2011). The *Gng3* temporal expression profile reaches a peak at CT22, which are inserted in time window with the lowest seizure frequency (at night) within 24 h, implying the anticonvulsant activity of the *Gng3* protein in the hippocampus. Thus, changes in the expression of core clock genes in

the hippocampus would result in modifications of the oscillatory expression patterns of CCGS acting on inhibitory or excitatory pathways, leading to a reduced threshold for seizure occurrence at precise times. The identification of harmonics of gene transcription in the brain (Hughes et al., 2009, 2012) may help in elucidating the mechanism behind the bimodal pattern of seizures in some TLE patients, with a reported ultradian seizure periodicity at 6–8 AM and 3–5PM (Karafin et al., 2010). A recent study, using NeuroPace, Inc. RNS® System to monitor human interictal epileptiform discharges and seizures at protracted timescales, further demonstrated that all subjects exhibited a 12-h harmonic of the circadian rhythm and a multitude of other rhythms (Baud et al., 2018).

In mice maintained in continuous darkness regimen, 22 proteins were found with ultradian oscillation in the normal hippocampus (Chiang et al., 2017). Four of them (*Ncdn*, *Ptprzl*, *Stx1b*, and *Calbl*) had been associated with epileptogenic processes in the hippocampus following status epilepticus. Nonetheless, analyzing the epileptic hippocampus with a higher temporal resolution is essential to confirm the ultradian pattern in gene expression. Evaluation of the temporal distribution of SRS through continuous video-EEG recordings in epileptic animals with knocked down of specific genes of interest would be ideal. Such analysis and in-depth characterization of the temporal expression of CCGS in an epileptic hippocampus would directly examine the potential time-dependent functional roles of genes in the generation/maintenance of SRS.

2.2. Uncoupled oscillators

The normal functioning of a multi-oscillatory system relies on the synchronization of the central and peripheral oscillators. Disruptions in this internal temporal order have been linked to multiple diseases (Smolensky et al., 2016). Hippocampus has intense connections with other brain structures, including direct neural pathways to SCN and other peripheral oscillators. As the clock gene *Per1* is responsive to electrical activity (Eun et al., 2011), the hyperactivity coming from the hippocampus could affect several circadian outputs leading to changes in amplitude or phase shifts. The clinical consequences of altered *Per1* expression (and other clock genes) in epileptic individuals are enormous as this could lead to extensive metabolism alterations and comorbidities linked with mTLE. Indeed, patients with epilepsy do demonstrate a wide variety of macrostructural sleep abnormalities (Derry and Duncan, 2013; Walker and Eriksson, 2011), such as sleep fragmentation and reduced sleep efficiency. Moreover, several functions controlled by the circadian clock are altered in epileptic individuals, including the hippocampus-dependent learning and memory.

The onset of spontaneous locomotor activity (SLA) has been shown to be unaltered in the mTLE model under light-dark conditions, suggesting that the SCN is appropriately responding to the light input (Matos et al., 2018; Stewart and Leung, 2003). However, the changes observed for some core clock genes in the hippocampus along with fragmented SLA (Matos et al., 2018) and body temperature (Quigg et al., 1999) rhythms may indicate a decoupling mechanism, in which the hippocampus and other oscillators are in different phase relationships compared to the healthy brain. The alterations in the hippocampus to SLA is likely linked to the activation of hippocampal-accumbens pathway after PILO-SE induction as demonstrated previously (Stewart and Leung, 2003). Alternatively, the seizures

could modulate SCN-related functions, affecting SLA downstream (Quigg, 2000). If interacting oscillators are uncoupled, the input signals can reach their targets out of phase, modifying the structure in many ways. Inhibitory molecules may not have their receptors expressed at the right time, or excitatory signals can reach the hippocampus in a low threshold condition (Fig. 2). Another possibility is a *de novo* circadian output effect, in which new genes could be rhythmically expressed due to the temporal changes in signal pathways. That is, the temporal inputs from other oscillators would not be in synchrony with the hippocampal timekeeping causing dysfunction in transcription of immediate early genes with a gain of oscillation for some targets (Fig. 2). A recent study using large-scale approaches provided an inventory of 1650 transcripts and 237 proteins undergoing diurnal expression in the hippocampus of a mouse model of mTLE (Debski et al., 2017). Notably, the authors observed that ~30% of the transcripts gain circadian fluctuation only when the brain becomes epileptic, which can contribute to *de novo* features of the limbic seizure rhythmicity.

A scenario combining multiple uncoupled oscillators superimposed in different phases may also lead to the arousal of harmonics in circadian seizure occurrence. Nevertheless, the potential contribution of circadian uncoupling to epilepsy still need to be addressed in future studies. In particular, assessment of circadian outputs in the hippocampus as well as in other brain oscillators would bring insights on the role of seizures as a disrupting cue for desynchrony with clinical consequences. Indeed, recent advances in techniques would likely help in the quantification of the oscillatory output *in vivo* and to determine a phase shift, amplitude of variation and periodicity over days. Such measurements have immense value to the epileptology field, especially for the implementation of chronotherapy in the future.

3. Chronotherapy for limbic seizures

Nearly 35% of mTLE patients acquire medically intractable epilepsy despite the availability of many medications over the last decade and adequate antiepileptic drug (AED) treatment (Kwan and Brodie, 2000). Furthermore, AED therapy is mostly focused on seizure reduction, which can lead to side effects even with a reduced frequency of seizures. Indeed, most mTLE patients display memory and mood impairments (Eddy et al., 2011; Schmitz, 2006). Only a minority of patients with intractable epilepsy qualify for surgical resection of the epileptic hippocampus, but this surgery is often associated with significant cognitive dysfunction (Helmstaedter et al., 2008). Hence, there is an urgent medical need for therapy that is efficient for both reducing the frequency of SRS and improving functions that are disrupted by seizures. Advanced knowledge on the involvement of the endogenous clock in the generation and sustenance of limbic seizures might shed light on the management of mTLE, a condition that has been recognized to be suitable for the application of chronotherapy. From these perspectives, it is plausible to implement an AED treatment regimen based on differential time dosing. Also, resetting the circadian rhythms in epileptic conditions may also be useful for reducing seizures.

3.1. Differential time dosing approach

By discerning the temporal pattern of seizure occurrence, one may design a drug administration regimen that would enable drug availability at times of higher seizure susceptibility, increased drug efficacy, and reduced toxicity. Such goals can be achieved by controlling both the time of drug administration and pharmacokinetic parameters under circadian control, including metabolism, excretion, absorption and distribution of the drug. By taking these variables into account, one may tailor drug schedules to improve their effectiveness by decreasing medication tolerance due to receptor holidays (Ramgopal et al., 2013). This kind of therapy is being investigated for several diseases with a periodic pattern of manifestation including some types of epilepsy (Manganaro et al., 2017; Tekade and Gattani, 2010). Overall, antiepileptic chronotherapy with a differential dose of carbamazepine, phenytoin, clobazam has been shown to be more useful for seizure control without increasing adverse side effects, in comparison to conventional AED therapy (Amengual-Gual et al., 2018; Thome-Souza et al., 2016; Yegnaranayan et al., 2006). While such studies have not been performed widely, they are nonetheless promising for suggesting guidelines that ensure the timing of drug activity during periods of maximal seizures. In this context, the development and the use of time-dependent drug delivery systems may gain priority for the pharmacotherapy of epilepsy (Singh et al., 2012). Moreover, studies on chronopharmacokinetic properties of AEDs in animal models of mTLE are critical for identifying optimal windows for dosing appropriate concentration of AEDs (Khedhaier et al., 2017). For example, it has been shown that carbamazepine level doubled around noon in rodents (Ramgopal et al., 2013). Furthermore, the identification of molecular rhythms in the mTLE model may provide new targets for AEDs (Debski et al., 2017).

3.2. Approaches for resetting the circadian rhythms

Since the potential mechanism for a periodic pattern of limbic seizures involves dysregulation of clock genes and uncoupling among the hippocampus and other oscillators, implementation of strategies that minimize harm from clock dysregulation/desynchronization could provide a new target in the management of epilepsy. In this context, application of external cues that reset endogenous rhythms may help in controlling the seizures and adverse effects. At the molecular level, this approach would induce changes in the amplitude or phase of core clock genes or their specific targets. Pharmacotherapy as a resetting approach has been applied in the circadian field to treat jet-lag, shift-workers and other sleep problems (Cuesta et al., 2015; Emens and Eastman, 2017; Lockley et al., 2015). The capability of this therapy to modulate phase, amplitude or period has the potential to be used in different conditions such as hypertension, immune disorders, mood dysfunction and aging (Chen et al., 2018; Portaluppi and Smolensky, 2010).

Melatonin is an endogenous chronobiotic molecule able to adjust altered circadian oscillation in several conditions perhaps due to its capacity to align the phase among peripheral and central oscillators (Sánchez-Barceló et al., 2011; Tchekalarova et al., 2015). Indeed, exogenous melatonin has been shown to have dose-dependent anticonvulsant actions and neuroprotective effects in several mTLE models and human patients (Petkova et al., 2014; Tchekalarova et al., 2013). However, Studies evaluating circadian phase effect of melatonin or its agonists are scarce. Costa-Lotufo et al. (2002) examined melatonin

treatment at two-time points (8:30 h and 17:00h) in female rats. The authors suggested that the anticonvulsant effect of melatonin seemed to be more intense at the light-dark transition, but the differences were not significant statistically. Moreover, a few studies have shown a dose-dependent pro-convulsant effect of melatonin (Banach et al., 2011). These conflicting results reinforce the necessity for chron-opharmacological studies that consider phase differences in melatonin responses. The mechanism of melatonin effects on epilepsy is likely multifactorial because melatonin can exert antioxidant and GABA-potential effects, in addition to its ability to synchronize disrupted circadian rhythms (Gupta et al., 2004; Kabuto et al., 1998; Niles et al., 1987).

Rapamycin, an mTOR pathway inhibitor, can both reduce seizures and modulate resetting of circadian rhythms occurring in SCN and peripheral oscillators in rodents (Liu et al., 2018; Ostendorf and Wong, 2015; Ramanathan et al., 2018). Clinical trials and case studies revealed a reduction in seizures with the use of rapamycin and other mTOR inhibitors, such as everolimus, in patients with tuberous sclerosis complex-associated epilepsy (Curatolo, 2015; French et al., 2016). The pre-clinical findings indicate that seizure-suppressive effect of rapamycin is maintained in the TLE model as long as rapamycin blood levels are sufficiently high (Drion et al., 2016). High doses and long-term treatment are likely necessary to achieve substantial anticonvulsant activity in human mTLE. Such regimens may give rise to unwanted side effects, however (Drion et al., 2016; Sliwa et al., 2012; van Vliet et al., 2012). Further clinical trials will be necessary to define the efficacy and safety profile of rapamycin and other mTOR inhibitors for treating mTLE.

It has been shown that diazepam and valproic acid, two of the most used AEDs, change the phase of the core clock genes by increasing or decreasing their expression (Griggs et al., 2018; Johansson et al., 2011; Oggier et al., 2010). Moreover, based on the analysis of pharmaco-transcriptome data available in DrugBank database version 5.0.6 (Wishart et al., 2018), it appears that phenobarbital treatment results in increased expression of *Per2* and *Per3* mRNA (Lambert et al., 2009). Additionally, several molecules have been discovered in screening strategies as clock modulators and could be used in the near future as new “clock drugs,” with the co-adjuvant potential of resetting oscillators in epileptic patients (Chen et al., 2018; Wallach and Kramer, 2015). Whether or to what extent the chronobiological effects of AEDs contribute to the amelioration of limbic seizure frequency and severity remains an interesting topic for future studies. Besides, it is interesting to note that several *zeitgebers* have been reported to have an antiepileptogenic role, including diet, light and physical exercise.

3.2.1. Diet—In mammals, feeding is a relevant cue for peripheral clocks including those in liver, muscle, pancreas and brain regions outside the SCN, regardless of the central clock (Damiola et al., 2000; Stokkan et al., 2001). This function is related to the role of specific nutrients (such as high-fat and high salt diet) and food factors (e.g., resveratrol) in resetting the cellular circadian clocks (Oike, 2017). The ketogenic diet (KD), a high fat, adequate protein, and low-carbohydrate diet, advances the onset of behavioral rhythms and disrupts the circadian clock (Challet et al., 1998; Oishi et al., 2009). Inhibition of mTOR pathway is likely one the mechanisms by which KD mediates these effects (McDaniel et al., 2011), which also involves the activation of peroxisomal proliferator-activated receptor γ (PPAR γ)

(Simeone et al., 2017). PPAR γ is also linked to the control of the circadian clock (Chen and Yang, 2014) and hence a target for epilepsy treatment (San et al., 2015). Indeed, KD has been used as an antiepileptic treatment for some patients with drug-resistant epilepsy, but the underlying mechanisms of the KD action are still largely unknown.

Resveratrol, a grape polyphenol, can delay the phase of the circadian clock in cultured fibroblasts (Oike, 2017). Resveratrol also displays anticonvulsant activity (Ethemoglu et al., 2017) and has shown efficacy for easing SE-induced epileptogenesis (Mishra et al., 2015; Castro et al., 2017). Nonetheless, additional studies are required to elucidate the extent to which nutritional signals can reset peripheral clocks related to the antiepileptic action. From this perspective, studies in mTLE models assessing the anticonvulsant effects of specific nutrients and food factors by using a time-restricted approach would be helpful.

3.2.2. Light—Light is one of the most important cues that synchronizes the endogenous rhythms with the cyclic environmental changes occurring within 24 h. This role is likely due to its capacity for resetting the clock system even by quick flash exposures (Kaladchibachi and Fernandez, 2018). For example, artificial light has been used as an approach to realign endogenous rhythms within a therapeutic perspective, especially in sleep (van Maanen et al., 2016) and mood disorders (Geoffroy et al., 2018; Kaladchibachi and Fernandez, 2018; San et al., 2015). The mTLE could be a good candidate for testing this kind of therapy (Baxendale, 2011) since it has been shown that in individuals with this condition the biological rhythms are altered probably due to dysregulation of the clock system. Based on a double-blinded randomized control trial (Baxendale et al., 2012), it was observed that a periodic bright light exposure over 12 weeks reduced the seizure frequency during the treatment in patients with intractable epilepsy and the response was more effective in patients with hippocampal sclerosis. In another controlled trial, the same group observed that the bright light resulted in a significant reduction in symptoms of anxiety and depression in patients with medically intractable epilepsy (Baxendale et al., 2013). Further clinical and preclinical studies are however required to evaluate the realignment of brain rhythms by the light as a therapeutic possibility for mTLE.

3.2.3. Physical exercise—Studies in humans and animal models have shown that physical exercise affects the SCN's firing rate and synchronizes circadian rhythms, acting as a non-photoc cue (Mistlberger and Skene, 2005; Ramkisoensing and Meijer, 2015). Furthermore, several other studies have demonstrated that epileptic rats under different physical exercise regimen present a reduced number of seizures (Arida et al., 2004, 2007; de Almeida et al., 2017; Peixinho-Pena et al., 2012). Recently, it was shown that the antiepileptic effect mediated by physical exercise is related to elevated BDNF and decreased TrkB levels (de Almeida et al., 2018) in the hippocampus. Interestingly, the BDNF-TrkB has a vital role in the synchronization of endogenous rhythms to the 24-h light/dark by regulating the daily ultrastructural rearrangements of the neuronal-glial network in the master clock (Girardet et al., 2013) but not in the hippocampus (Ikeno et al., 2013).

4. Future directions

Collectively, the approaches mentioned above are supportive for investigating other ways of resetting the biological rhythms as a potential therapy for improving seizure control and ameliorating comorbidities observed in epilepsy patients. As mentioned above, many phase-resetting molecules have been uncovered by high-throughput screening *in vitro* (Hirota et al., 2008; Oike, 2017). Overall, they interact with clock proteins to alter period length or phase at the cellular level. Thus, the screening of these clock modulators as potential antiepileptic agents can bring new perspectives for the treatment of mTLE. Alternatively, light, diet, exercise and other *zeitgebers* which potentially include cognitive plasticity (Gritton et al., 2012, 2013) could modulate the rhythms in the hippocampus and related structures with clinical benefits. Cognitive behavioral therapy has been applied for the treatment of epilepsy, with inconsistent results (Leeman-Markowski and Schachter, 2017; Tang et al., 2014). It would be interesting to investigate whether cognitive interventions at different phases could affect the rhythm in the hippocampus and modulate seizures more effectively.

Studies investigating the application of chronotherapy in mTLE or animal models of mTLE are still scarce to evaluate the suitability of time-specific therapeutic intervention regimens. However, this is not a straightforward approach, as the individual nature of epilepsy can influence its efficacy. From the perspective of inter-individual variability in seizure occurrence, a clinical protocol for chronotherapy must include a patient-based tracking of seizure clusters during the day. Such-tracking would help in incorporating the temporal patterns of seizures into individualized medication management.

Furthermore, transplantation of neural stem cells (NSCs) is being considered as a therapy for epilepsy and other conditions (Waldau et al., 2010; Shetty, 2011, 2014; Hattiangady and Shetty, 2012; Shetty and Hattiangady, 2016). Coupling of circadian rhythms to the cell cycle has been observed in NSCs, and clock genes have been shown to influence the proliferation and differentiation of NSCs (Malik et al., 2015; Akle et al., 2017; Draijer et al., 2018; Shimozaki, 2018). Therefore, the timing of NSC grafting or the use of synchronizers for aligning the clock system of grafted NSCs with the host NSCs in the neurogenic region may enhance the yield of graft-derived cells by stimulating the proliferation and differentiation of grafted NSCs. Indeed, a study demonstrated that melatonin-supported stem cell therapy is superior to the administration of stem cells alone (Lee et al., 2017). Regarding the application of NSCs for the treatment of epilepsy, grafting of the medial ganglionic eminence-derived NSCs into the hippocampus of rats exhibiting chronic TLE resulted in ~50% reductions in the frequency and intensity of spontaneous seizures with no improvements in the memory function (Waldau et al., 2010). Analyses of grafts, however, revealed that the overall yield of graft-derived cells was ~28% of injected cells. From this perspective, grafting of NSCs into the chronically epileptic hippocampus along with circadian synchronizers, or phase-dependent grafting approach, may enhance graft cell survival and differentiation as well as host hippocampal neurogenesis, which may further diminish the frequency and intensity of spontaneous seizures, and alleviate memory and mood impairments seen in chronic TLE. Also, the use of circadian synchronizers in cell therapy may enhance the survival, differentiation and seizure-suppressing effects of other

donor cell types considered for treating epilepsy. The cells include the GABA-ergic precursor cells from the embryonic brain, and GABA-ergic progenitors derived from the human embryonic stem cells and human induced pluripotent stem cells (Hattiangady et al., 2008; Hunt et al., 2013; Cunningham et al., 2014; Shetty and Upadhyaya, 2016; Upadhyaya et al., 2018).

5. Conclusions

Learning how the clock system modulates the seizure threshold is essential for comprehending the circadian occurrence of seizures and comorbidities as well as for developing innovative therapeutic approaches for mTLE. The temporal dysregulation of core clock genes in the epileptic hippocampus could affect the phase and amplitude of different genes that play roles in the electrical activity of neurons with consequences on the rhythmic fluctuation in the inhibitory/excitatory imbalance. As examples, the melatonergic and the mTOR pathways are two essential regulatory systems in the modulation of seizure thresholds that present circadian activity in the hippocampus. Moreover, at the system level, the epileptic hippocampus could be uncoupled to other oscillators contributing in several ways to the occurrence of seizures as a de novo circadian output and to the extensive metabolic alterations linked with mTLE. From these perspectives, mTLE is a suitable condition for the application of chronotherapy for reducing seizures and comorbidities. Differential time dosing of AEDs and resetting mechanisms by light, food, physical exercise or other approaches could be valuable strategies for the management of mTLE.

Acknowledgements

The authors are supported by grants from the Department of Defense (CDMRP W81XWH-14-1-0558 to A.K.S.), Department of Veterans Affairs (VA Merit Award grant I01BX000883 and VA-BLR&D Research Career Scientist award, 1IK6BX003612 to A.K.S.). Daniel Gtai and Tiago Andrade were supported by a Visiting Scientist Award from CAPES Foundation, Ministry of Education, Government of Brazil and by grants from Fundacao de Amparo a Pesquisa do Estado de Alagoas (FAPEAL) and Conselho Nacional de Desenvolvimento Cientifico e Tecnologico (CNPq), Brazil.

Abbreviations:

AED	anti epileptic drug
BDNF	brain derived neurotropic factor
CCGS	clock-controlled genes
CK1	casein kinase 1
CLOCK	circadian locomotor output cycles kaput
Cry	cryptochrome
GO	gene ontology
ipRGC	intrinsically photosensitive retinal ganglion cells
KD	ketogenic diet

MT	melatonin receptors
mTLE	mesial temporal lobe epilepsy
mTOR	mammalian/mechanistic target of rapamycin
NFIL3	nuclear factor, interleukin 3 regulated
NPAS2	neuronal PAS domain protein 2
Per	period
RHT	retinohypothalamic tract
RORs	retinoid-related orphan receptors
RRE	rev responsive element
SCF	skp1/cullin/F-box protein
SCN	suprachiasmatic nucleus
SLA	spontaneous locomotor activity
SRS	spontaneous recurrent seizures
TFs	transcription factors
TrKB	tropomyosin receptor kinase B
TTFL	transcriptional-translational feedback loops

References

- Aguiar CCT, Almeida AB, Araújo PVP, Vasconcelos GS, Chaves EMC, do Vale OC, Macedo DS, de Sousa FCF, Viana GS, de B, Vasconcelos SMM, 2012 Anticonvulsant effects of agomelatine in mice. *Epilepsy Behav.* 24, 324–328. 10.1016/j.yebeh.2012.04.134. [PubMed: 22658946]
- Akle V, Stankiewicz AJ, Kharchenko V, Yu L, Kharchenko PV, Zhdanova IV, 2017 Circadian kinetics of cell cycle progression in adult neurogenic niches of diurnal vertebrate. *J. Neurosci* 37, 1900–1909. 10.1523/JNEUROSCI.3222-16.2017. [PubMed: 28087763]
- Amengual-Gual M, Sánchez Fernández I, Loddenkemper T, 2018 Patterns of epileptic seizure occurrence. *Brain Res.* 10.1016/j.brainres.2018.02.032.
- Arida RM, Scorza FA, Peres CA, Cavalheiro EA, 1999 The course of untreated seizures in the pilocarpine model of epilepsy. *Epilepsy Res.* 34, 99–107. [PubMed: 10210024]
- Arida RM, Sanabria ERG, da Silva AC, Faria LC, Scorza FA, Cavalheiro EA, 2004 Physical training reverts hippocampal electrophysiological changes in rats submitted to the pilocarpine model of epilepsy. *Physiol. Behav* 83, 165–171. 10.1016/j.physbeh.2004.08.008. [PubMed: 15501504]
- Arida RM, Scorza CA, Scorza FA, Gomes da Silva S, da Graça Naffah-Mazzacoratti M, Cavalheiro EA, 2007 Effects of different types of physical exercise on the staining of parvalbumin-positive neurons in the hippocampal formation of rats with epilepsy. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 31, 814–822. 10.1016/j.pnpbp.2007.01.021. [PubMed: 17331634]
- Bailey M, Silver R, 2014 Sex differences in circadian timing systems: implications for disease. *Front. Neuroendocrinol* 35, 111–139. 10.1016/j.yfrne.2013.11.003. [PubMed: 24287074]
- Banach M, Gurdziel E, J drych M, Borowicz KK, 2011 Melatonin in experimental seizures and epilepsy. *Pharmacol. Rep* 63, 1–11. [PubMed: 21441606]

- Baud MO, Kleen JK, Mirro EA, Andrechak JC, King-Stephens D, Chang EF, Rao VR, 2018 Multi-day rhythms modulate seizure risk in epilepsy. *Nat. Commun* 9, 88 10.1038/s41467-017-02577-y. [PubMed: 29311566]
- Baxendale SA, 2011 Light therapy as a treatment for epilepsy. *Med. Hypotheses* 76, 661–664. 10.1016/j.mehy.2011.01.025. [PubMed: 21333455]
- Baxendale S, O'Sullivan J, Heaney D, 2012 Bright light therapy as an add on treatment for medically intractable epilepsy. *Epilepsy Behav.* 24, 359–364. 10.1016/j.yebeh.2012.04.123. [PubMed: 22658437]
- Baxendale S, O'Sullivan J, Heaney D, 2013 Bright light therapy for symptoms of anxiety and depression in focal epilepsy: randomized controlled trial. *Br. J. Psychiatry* 202, 352–356. 10.1192/bjp.bp.112.122119. [PubMed: 23520221]
- Bazil CW, Short D, Crispin D, Zheng W, 2000 Patients with intractable epilepsy have low melatonin, which increases following seizures. *Neurology* 55, 1746–1748. [PubMed: 11113238]
- Bertram EH, Cornett JF, 1994 The evolution of a rat model of chronic spontaneous limbic seizures. *Brain Res.* 661, 157–162. [PubMed: 7834366]
- Bhadra U, Thakkar N, Das P, Pal Bhadra M, 2017 Evolution of circadian rhythms: from bacteria to human. *Sleep Med.* 35, 49–61. 10.1016/j.sleep.2017.04.008. [PubMed: 28619182]
- Binns D, Dimmer E, Huntley R, Barrell D, O'Donovan C, Apweiler R, 2009 QuickGO: a web-based tool for Gene Ontology searching. *Bioinformatics* 25, 3045–3046. 10.1093/bioinformatics/btp536. [PubMed: 19744993]
- Boukhtouche F, Vodjdani G, Jarvis CI, Bakouche J, Staels B, Mallet J, Mariani J, Lemaigre-Dubreuil Y, Brugg B, 2006 Human retinoic acid receptor-related orphan receptor alpha1 overexpression protects neurons against oxidative stress-induced apoptosis. *J. Neurochem* 96, 1778–1789. 10.1111/jM471-4159.2006.03708.x. [PubMed: 16539693]
- Broekaart DWM, van Scheppingen J, Geijtenbeek KW, Zuidberg MRJ, Anink JJ, Baayen JC, Muhlebnner A, Aronica E, Gorter JA, van Vliet EA, 2017 Increased expression of (immuno)proteasome subunits during epileptogenesis is attenuated by inhibition of the mammalian target of rapamycin pathway. *Epilepsia* 58, 1462–1472. 10.1111/epi.13823. [PubMed: 28643873]
- Bustelo XR, Sauzeau V, Berenjeno IM, 2007 GTP-binding proteins of the Rho/Rac family: regulation, effectors and functions in vivo. *Bioessays* 29, 356–370. 10.1002/bies.20558. [PubMed: 17373658]
- Cao R, Obrietan K, 2010 mTOR signaling and entrainment of the mammalian circadian clock. *Mol. Cell. Pharmacol* 2, 125–130. 10.4255/mcpharmacol.10.17. [PubMed: 21274417]
- Cao R, Li A, Cho H-Y, Lee B, Obrietan K, 2010 Mammalian target of rapamycin signaling modulates photic entrainment of the suprachiasmatic circadian clock. *J. Neurosci* 30, 6302–6314. 10.1523/JNEUROSCI.5482-09.2010. [PubMed: 20445056]
- Castro OW, Upadhy D, Kodali M, Shetty AK, 2017 Resveratrol for easing status epilepticus induced brain injury, inflammation, epileptogenesis, and cognitive and memory dysfunction- are we there yet? *Front. Neurol* 8, 603 10.3389/fneur.2017.00603. [PubMed: 29180982]
- Cavalheiro EA, Leite JP, Bortolotto ZA, Turski WA, Ikonomidou C, Turski L, 1991 Longterm effects of pilocarpine in rats: structural damage of the brain triggers kindling and spontaneous recurrent seizures. *Epilepsia* 32, 778–782. [PubMed: 1743148]
- Challet E, Solberg LC, Turek FW, 1998 Entrainment in calorie-restricted mice: conflicting zeitgebers and free-running conditions. *Am. J. Physiol* 274, R1751–61. [PubMed: 9841486]
- Chaudhury D, Wang LM, Colwell CS, 2005 Circadian regulation of hippocampal long-term potentiation. *J. Biol. Rhythms* 20, 225–236. 10.1177/0748730405276352. [PubMed: 15851529]
- Chen L, Yang G, 2014 PPARs integrate the mammalian clock and energy metabolism. *PPAR Res*, 653017 10.1155/2014/653017. [PubMed: 24693278]
- Chen Z, Yoo S-H, Takahashi JS, 2018 Development and therapeutic potential of small-molecule modulators of circadian systems. *Annu. Rev. Pharmacol. Toxicol* 58, 231–252. 10.1146/annurev-pharmtox-010617-052645. [PubMed: 28968186]
- Chiang C-K, Xu B, Mehta N, Mayne J, Sun WYL, Cheng K, Ning Z, Dong J, Zou H, Cheng H-YM, Figeys D, 2017 Phosphoproteome profiling reveals circadian clock regulation of posttranslational modifications in the murine Hippocampus. *Front. Neurol* 8, 110 10.3389/fneur.2017.00110. [PubMed: 28382018]

- Cho C-H, 2012 Molecular mechanism of circadian rhythmicity of seizures in temporal lobe epilepsy. *Front. Cell. Neurosci* 6, 55 10.3389/fncel.2012.00055. [PubMed: 23189039]
- Chou MM, Blenis J, 1996 The 70 kDa S6 kinase complexes with and is activated by the Rho family G proteins Cdc42 and Rac1. *Cell* 85, 573–583. [PubMed: 8653792]
- Citraro R, Leo A, Constanti A, Russo E, De Sarro G, 2016 mTOR pathway inhibition as a new therapeutic strategy in epilepsy and epileptogenesis. *Pharmacol. Res* 107, 333–343. <https://doi.org/10.1016/j.phrs.2016.03.039>. [PubMed: 27049136]
- Costa-Lotufo LV, Fonteles MM, de F, Lima ISP, de Oliveira AA, Nascimento VS, de Bruin VMS, Viana GSB, 2002 Attenuating effects of melatonin on pilocarpine-induced seizures in rats. *Comp. Biochem. Physiol. C Toxicol. Pharmacol* 131, 521–529. [PubMed: 11976067]
- Cuesta M, Cermakian N, Boivin DB, 2015 Glucocorticoids entrain molecular clock components in human peripheral cells. *FASEB J* 29, 1360–1370. 10.1096/fj.14-265686. [PubMed: 25500935]
- Cunningham M, Cho JH, Leung A, Savvidis G, Ahn S, Moon M, Lee PK, Han JJ, Azimi N, Kim KS, Bolshakov VY, Chung S, 2014 hPSC-derived maturing GABAergic interneurons ameliorate seizures and abnormal behavior in epileptic mice. *Cell Stem Cell* 15, 559–573. 10.1016/j.stem.2014.10.006. [PubMed: 25517465]
- Curatolo P, 2015 Mechanistic target of rapamycin (mTOR) in tuberous sclerosis complex-associated epilepsy. *Pediatr. Neurol* 52, 281–289. 10.1016/j.pediatrneurol.2014.10.028. [PubMed: 25591831]
- Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U, 2000 Restricted feeding uncouples circadian oscillators in peripheral tissues from the central pacemaker in the suprachiasmatic nucleus. *Genes Dev* 14, 2950–2961. [PubMed: 11114885]
- Dastgheib M, Moezi L, 2014 Acute and chronic effects of agomelatine on intravenous pentylenetetrazol-induced seizure in mice and the probable role of nitric oxide. *Eur. J. Pharmacol* 736, 10–15. 10.1016/j.ejphar.2014.04.039. [PubMed: 24803306]
- de Almeida AA, Gomes da Silva S, Lopim GM, Vannucci Campos D, Fernandes J, Cabral FR, Arida RM, 2017 Resistance exercise reduces seizure occurrence, attenuates memory deficits and restores BDNF signaling in rats with chronic epilepsy. *Neurochem. Res* 42, 1230–1239. 10.1007/s11064-016-2165-9. [PubMed: 28078614]
- de Almeida AA, Gomes da Silva S, Lopim GM, Vannucci Campos D, Fernandes J, Cabral FR, Arida RM, 2018 Physical exercise alters the activation of downstream proteins related to BDNF-TrkB signaling in male Wistar rats with epilepsy. *J. Neurosci. Res* 96, 911–920. 10.1002/jnr.24196. [PubMed: 29098710]
- De Nobrega AK, Lyons LC, 2018 Aging and the clock: perspective from flies to humans. *Eur. J. Neurosci* 10.1111/ejn.14176.
- Debski K, Ceglia N, Ghestem A, Ivanov A, Brancati GE, Broer S, Bot A, Muller A, Schoch S, Becker A, Loscher W, Sassone-Corsi P, Lukasiuk K, Baldi P, Bernard C, 2017 The Circadian Hippocampus and Its Reprogramming in Epilepsy: Impact for Chronotherapeutics. 10.1101/199372.
- Delerive P, Monté D, Dubois G, Trottein F, Fruchart-Najib J, Mariani J, Fruchart JC, Staels B, 2001 The orphan nuclear receptor ROR alpha is a negative regulator of the inflammatory response. *EMBO Rep* 2, 42–48. 10.1093/embo-reports/kve007. [PubMed: 11252722]
- Derry CP, Duncan S, 2013 Sleep and epilepsy. *Epilepsy Behav* 26, 394–404. 10.1016/j.yebeh.2012.10.033. [PubMed: 23465654]
- Devinsky O, Vezzani A, O'Brien TJ, Jette N, Scheffer IE, de Curtis M, Perucca P, 2018 Epilepsy. *Nat. Rev. Dis. Primers* 4, 18024 10.1038/nrdp.2018.24. [PubMed: 29722352]
- Díaz-Morales JF, Parra-Robledo Z, 2018 Age and sex differences in Morningness/Eveningness along the life span: a cross-sectional study in Spain. *J. Genet. Psychol* 179, 71–84. 10.1080/00221325.2018.1424706. [PubMed: 29424669]
- Dictionary of Circadian Physiology [WWW Document], n.d. URL <https://www.circadian.org/dictionary.html> (accessed 10.30.18).
- Draijer S, Chaves I, Hoekman MFM, 2018 The circadian clock in adult neural stem cell maintenance. *Prog. Neurobiol* 10.1016/j.pneurobio.2018.05.007. 6 7 pii: S0301-0082(18)30020-0. [Epub ahead of print].

- Drion CM, Borm LE, Kooijman L, Aronica E, Wadman WJ, Hartog AF, van Vliet EA, Gorter JA, 2016 Effects of rapamycin and curcumin treatment on the development of epilepsy after electrically induced status epilepticus in rats. *Epilepsia* 57, 688–697. 10.1111/epi.13345. [PubMed: 26924447]
- Durazzo TS, Spencer SS, Duckrow RB, Novotny EJ, Spencer DD, Zaveri HP, 2008 Temporal distributions of seizure occurrence from various epileptogenic regions. *Neurology* 70, 1265–1271. 10.1212/01.wnl.0000308938.84918.3f. [PubMed: 18391158]
- Eddy CM, Rickards HE, Cavanna AE, 2011 The cognitive impact of antiepileptic drugs. *Ther. Adv. Neurol. Disord* 4, 385–407. 10.1177/1756285611417920. [PubMed: 22164192]
- Emens JS, Eastman CI, 2017 Diagnosis and treatment of non-24-h sleep-wake disorder in the blind. *Drugs* 77, 637–650. 10.1007/s40265-017-0707-3. [PubMed: 28229310]
- Endo M, Antonyak MA, Cerione RA, 2009 Cdc42-mTOR signaling pathway controls Hes5 and Pax6 expression in retinoic acid-dependent neural differentiation. *J. Biol. Chem* 284, 5107–5118. 10.1074/jbc.M807745200. [PubMed: 19097998]
- Ethemoglu MS, Seker FB, Akkaya H, Kilic E, Aslan I, Erdogan CS, Yilmaz B, 2017 Anticonvulsant activity of resveratrol-loaded liposomes in vivo. *Neuroscience* 357, 12–19. 10.1016/j.neuroscience.2017.05.026. [PubMed: 28577913]
- Eun B, Kim HJ, Kim SY, Kim TW, Hong ST, Choi KM, Shim JK, Moon Y, Son GH, Kim K, Kim H, Sun W, 2011 Induction of Per1 expression following an experimentally induced epilepsy in the mouse hippocampus. *Neurosci. Lett* 498, 110–113. 10.1016/j.neulet.2011.03.039. [PubMed: 21420468]
- Fang Y, Park I-H, Wu A-L, Du G, Huang P, Frohman MA, Walker SJ, Brown HA, Chen J, 2003 PLD1 regulates mTOR signaling and mediates Cdc42 activation of S6K1. *Curr. Biol* 13, 2037–2044. [PubMed: 14653992]
- Fisher PD, Sperber EF, Moshé SL, 1998 Hippocampal sclerosis revisited. *Brain Dev.* 20, 563–573. [PubMed: 9865538]
- Frank E, Sidor MM, Gamble KL, Cirelli C, Sharkey KM, Hoyle N, Tikotzky L, Talbot LS, McCarthy MJ, Hasler BP, 2013 Circadian clocks, brain function, and development. *Ann. N. Y. Acad. Sci* 1306, 43–67. 10.1111/nyas.12335. [PubMed: 24329517]
- French JA, Lawson JA, Yapici Z, Ikeda H, Polster T, Nabbout R, Curatolo P, de Vries PJ, Dlugos DJ, Berkowitz N, Voi M, Peyrard S, Pelov D, Franz DN, 2016 Adjunctive everolimus therapy for treatment-resistant focal-onset seizures associated with tuberous sclerosis (EXIST-3): a phase 3, randomised, double-blind, placebo-controlled study. *Lancet* 388, 2153–2163. 10.1016/S0140-6736(16)31419-2. [PubMed: 27613521]
- Geoffroy PA, Schroder CM, Bourgin P, 2018 Light treatment in depression: an antique treatment with new insights. *Sleep Med. Rev* 10.1016/j.smrv.2018.03.002.
- Girardet C, Lebrun B, Cabirol-Pol M-J, Tardivel C, François-Bellan A-M, Becquet D, Bosler O, 2013 Brain-derived neurotrophic factor/TrkB signaling regulates daily astroglial plasticity in the suprachiasmatic nucleus: electron-microscopic evidence in mouse. *Glia* 61, 1172–1177. 10.1002/glia.22509. [PubMed: 23640807]
- Glossary | Centre for Chronobiology [WWW Document], n.d. URL <http://www.chronobiology.ch/glossary/> (accessed 10.30.18).
- Golombek DA, Fernandez Duque D, De Brito Sanchez MG, Burin L, Cardinali DP, 1992 Time-dependent anticonvulsant activity of melatonin in hamsters. *Eur. J. Pharmacol* 210, 253–258. [PubMed: 1612101]
- Govek E-E, Hatten ME, Van Aelst L, 2011 The role of Rho GTPase proteins in CNS neuronal migration. *Dev. Neurobiol* 71, 528–553. 10.1002/dneu.20850. [PubMed: 21557504]
- Griggs CA, Malm SW, Jaime-Frias R, Smith CL, 2018 Valproic acid disrupts the oscillatory expression of core circadian rhythm transcription factors. *Toxicol. Appl. Pharmacol* 339, 110–120. 10.1016/j.taap.2017.12.005. [PubMed: 29229235]
- Gritton HJ, Kantorowski A, Sarter M, Lee TM, 2012 Bidirectional interactions between circadian entrainment and cognitive performance. *Learn. Mem* 19, 126–141. 10.1101/lm.023499.111. [PubMed: 22383380]

- Gritton HJ, Stasiak AM, Sarter M, Lee TM, 2013 Cognitive performance as a zeitgeber: cognitive oscillators and cholinergic modulation of the SCN entrain circadian rhythms. *PLoS One* 8, e56206 10.1371/journal.pone.0056206. [PubMed: 23441168]
- Gupta M, Gupta YK, Agarwal S, Aneja S, Kalaivani M, Kohli K, 2004 Effects of add-on melatonin administration on antioxidant enzymes in children with epilepsy taking carbamazepine monotherapy: a randomized, double-blind, placebo-controlled trial. *Epilepsia* 45, 1636–1639. 10.1111/j.0013-9580.2004.17604.x. [PubMed: 15571523]
- Hafner M, Koepl H, Gonze D, 2012 Effect of network architecture on synchronization and entrainment properties of the circadian oscillations in the suprachiasmatic nucleus. *PLoS Comput. Biol* 8, e1002419 10.1371/journal.pcbi.1002419. [PubMed: 22423219]
- Hattiangady B, Shetty AK, 2010 Decreased neuronal differentiation of newly generated cells underlies diminished neurogenesis in chronic temporal lobe epilepsy. *Hippocampus* 20, 97–112. 10.1002/hipo.20594. [PubMed: 19309040]
- Hattiangady B, Shetty AK, 2012 Neural stem cell grafting counteracts hippocampal injury-mediated impairments in mood, memory, and neurogenesis. *Stem Cells Transl. Med* 1, 696–708. 10.5966/sctm.2012-0050. [PubMed: 23197876]
- Hattiangady B, Rao MS, Shetty AK, 2004 Chronic temporal lobe epilepsy is associated with severely declined dentate neurogenesis in the adult hippocampus. *Neurobiol. Dis* 17, 473–490. 10.1016/j.nbd.2004.08.008. [PubMed: 15571983]
- Hattiangady B, Rao MS, Shetty AK, 2008 Grafting of striatal precursor cells into hippocampus shortly after status epilepticus restrains chronic temporal lobe epilepsy. *Exp. Neurol* 212, 468–481. 10.1016/j.expneurol.2008.04.040. [PubMed: 18579133]
- Helmstaedter C, Loer B, Wohlfahrt R, Hammen A, Saar J, Steinhoff BJ, Quiske A, Schulze-Bonhage A, 2008 The effects of cognitive rehabilitation on memory outcome after temporal lobe epilepsy surgery. *Epilepsy Behav.* 12, 402–409. 10.1016/j.yebeh.2007.11.010. [PubMed: 18155965]
- Hirota T, Lewis WG, Liu AC, Lee JW, Schultz PG, Kay SA, 2008 A chemical biology approach reveals period shortening of the mammalian circadian clock by specific inhibition of GSK-3 β . *Proc. Natl. Acad. Sci. U. S. A* 105, 20746–20751. 10.1073/pnas.0811410106. [PubMed: 19104043]
- Hofstra WA, de Weerd AW, 2009 The circadian rhythm and its interaction with human epilepsy: a review of literature. *Sleep Med. Rev* 13, 413–420. 10.1016/j.smrv.2009.01.002. [PubMed: 19398353]
- Hofstra WA, Gordijn MCM, van der Palen J, van Regteren R, Grootemarsink BE, de Weerd AW, 2011 Timing of temporal and frontal seizures in relation to the circadian phase: a prospective pilot study. *Epilepsy Res.* 94, 158–162. 10.1016/j.epilepsyres.2011.01.015. [PubMed: 21388790]
- Honma S, 2018 The mammalian circadian system: a hierarchical multi-oscillator structure for generating circadian rhythm. *J. Physiol. Sci* 68, 207–219. 10.1007/s12576-018-0597-5. [PubMed: 29460036]
- Huang X, McMahon J, Huang Y, 2012 Rapamycin attenuates aggressive behavior in a rat model of pilocarpine-induced epilepsy. *Neuroscience* 215, 90–97. 10.1016/j.neuroscience.2012.04.011. [PubMed: 22522471]
- Hughes ME, DiTacchio L, Hayes KR, Vollmers C, Pulivarthy S, Baggs JE, Panda S, Hogenesch JB, 2009 Harmonics of circadian gene transcription in mammals. *PLoS Genet.* 5, e1000442 10.1371/journal.pgen.1000442. [PubMed: 19343201]
- Hughes ME, Hogenesch JB, Kornacker K, 2010 JTK_CYCLE: an efficient nonpara-metric algorithm for detecting rhythmic components in genome-scale data sets. *J. Biol. Rhythms* 25, 372–380. 10.1177/0748730410379711. [PubMed: 20876817]
- Hughes ME, Hong H-K, Chong JL, Indacochea AA, Lee SS, Han M, Takahashi JS, Hogenesch JB, 2012 Brain-specific rescue of Clock reveals system-driven transcriptional rhythms in peripheral tissue. *PLoS Genet.* 8, e1002835 10.1371/journal.pgen.1002835. [PubMed: 22844252]
- Hunt RF, Girsakis KM, Rubenstein JL, Alvarez-Buylla A, Baraban SC, 2013 GABA progenitors grafted into the adult epileptic brain control seizures and abnormal behavior. *Nat. Neurosci* 16, 692–697. 10.1038/nn.3392. [PubMed: 23644485]

- Ikeno T, Weil ZM, Nelson RJ, 2013 Photoperiod affects the diurnal rhythm of hippocampal neuronal morphology of Siberian hamsters. *Chronobiol. Int* 30, 1089–1100. 10.3109/07420528.2013.800090. [PubMed: 23879697]
- Ikeno T, Weil ZM, Nelson RJ, 2014 Timing of light pulses and photoperiod on the diurnal rhythm of hippocampal neuronal morphology of Siberian hamsters. *Neuroscience* 270, 69–75. 10.1016/j.neuroscience.2014.04.002. [PubMed: 24726983]
- Johansson A-S, Brask J, Owe-Larsson B, Hetta J, Lundkvist GBS, 2011 Valproic acid phase shifts the rhythmic expression of period2: luciferase. *J. Biol. Rhythms* 26, 541–551. 10.1177/0748730411419775. [PubMed: 22215612]
- Kabuto H, Yokoi I, Ogawa N, 1998 Melatonin inhibits iron-induced epileptic discharges in rats by suppressing peroxidation. *Epilepsia* 39, 237–243. [PubMed: 9578039]
- Kaladchibachi S, Fernandez F, 2018 Precision light for the treatment of psychiatric disorders. *Neural Plast.*, 5868570 10.1155/2018/5868570. [PubMed: 29593784]
- Kaleyias J, Loddenkemper T, Vendrame M, Das R, Syed TU, Alexopoulos AV, Wyllie E, Kothare SV, 2011 Sleep-wake patterns of seizures in children with le-sional epilepsy. *Pediatr. Neurol* 45, 109–113. 10.1016/j.pediatrneurol.2011.03.006. [PubMed: 21763951]
- Karafin M, St Louis EK, Zimmerman MB, Sparks JD, Granner MA, 2010 Bimodal ultradian seizure periodicity in human mesial temporal lobe epilepsy. *Seizure* 19, 347–351. 10.1016/j.seizure.2010.05.005. [PubMed: 20580271]
- Karoly PJ, Goldenholz DM, Freestone DR, Moss RE, Grayden DB, Theodore WH, Cook MJ, 2018 Circadian and circaseptan rhythms in human epilepsy: a retrospective cohort study. *Lancet Neurol.* 17, 977–985. 10.1016/S1474-4422(18)30274-6. [PubMed: 30219655]
- Khedhaier WBC, Dridi I, Aouam K, Ben-Attia M, Reinberg A, Boughattas NA, 2017 Circadian variation in anticonvulsant activity of valproic acid in mice. *Biomed. Pharmacother* 95, 25–30. 10.1016/j.biopha.2017.08.047. [PubMed: 28826093]
- Koike N, Yoo S-H, Huang H-C, Kumar V, Lee C, Kim T-K, Takahashi JS, 2012 Transcriptional architecture and chromatin landscape of the core circadian clock in mammals. *Science* 338, 349–354. 10.1126/science.1226339. [PubMed: 22936566]
- Kojima S, Green CB, 2015 Circadian genomics reveal a role for post-transcriptional regulation in mammals. *Biochemistry* 54, 124–133. 10.1021/bi500707c. [PubMed: 25303020]
- Kothare SV, Zarowski M, 2011 Sleep and epilepsy: common bedfellows. *J. Clin. Neurophysiol* 28, 101–102. 10.1097/WNP.0b013e3182120d30. [PubMed: 21399510]
- Kumaki Y, Ukai-Tadenuma M, Uno K-ID, Nishio J, Masumoto K-H, Nagano M, Komori T, Shigeyoshi Y, Hogenesch JB, Ueda HR, 2008 Analysis and synthesis of high-amplitude Cis-elements in the mammalian circadian clock. *Proc. Natl. Acad. Sci. U. S. A* 105, 14946–14951. 10.1073/pnas.0802636105. [PubMed: 18815372]
- Kuruba R, Hattiangady B, Shetty AK, 2009 Hippocampal neurogenesis and neural stem cells in chronic temporal lobe epilepsy. *Epilepsy Behav.* 14 (Suppl 1), 65–73. [PubMed: 18796338]
- Kwan P, Brodie MJ, 2000 Early identification of refractory epilepsy. *N. Engl. J. Med* 342, 314–319. 10.1056/nejm200002033420503. [PubMed: 10660394]
- Lambert CB, Spire C, Claude N, Guillouzo A, 2009 Dose- and time-dependent effects of phenobarbital on gene expression profiling in human hepatoma HepaRG cells. *Toxicol. Appl. Pharmacol* 234, 345–360. 10.1016/j.taap.2008.11.008. [PubMed: 19084549]
- Laplanche M, Sabatini DM, 2012 mTOR signaling in growth control and disease. *Cell* 149, 274–293. 10.1016/j.cell.2012.03.017. [PubMed: 22500797]
- Lee MS, Yin T-C, Sung P-H, Chiang JY, Sun C-K, Yip H-K, 2017 Melatonin enhances survival and preserves functional integrity of stem cells: a review. *J. Pineal Res.* 62 10.1111/jpi.12372.
- Leeman-Markowski BA, Schachter SC, 2017 Cognitive and behavioral interventions in epilepsy. *Curr. Neurol. Neurosci. Rep* 17, 42 10.1007/s11910-017-0752-z. [PubMed: 28382495]
- LeGates TA, Fernandez DC, Hattar S, 2014 Light as a central modulator of circadian rhythms, sleep and affect. *Nat. Rev. Neurosci* 15, 443–454. 10.1038/nrn3743. [PubMed: 24917305]
- Li JZ, Bunney BG, Meng F, Hagenauer MH, Walsh DM, Vawter MP, Evans SJ, Choudary PV, Cartagena P, Barchas JD, Schatzberg AF, Jones EG, Myers RM, Watson SJ Jr., Akil H, Bunney WE, 2013 Circadian patterns of gene expression in the human brain and disruption in major

- depressive disorder. *Proc. Natl. Acad. Sci. U. S. A* 110, 9950–9955. 10.1073/pnas.1305814110. [PubMed: 23671070]
- Liu D, Stowie A, de Zavalía N, Leise T, Pathak SS, Drewes LR, Davidson AJ, Amir S, Sonenberg N, Cao R, 2018 mTOR signaling in VIP neurons regulates circadian clock synchrony and olfaction. *Proc. Natl. Acad. Sci. U. S. A* 115, E3296–E3304. 10.1073/pnas.1721578115. [PubMed: 29555746]
- Lockley SW, Dressman MA, Licamele L, Xiao C, Fisher DM, Flynn-Evans EE, Hull JT, Torres R, Lavedan C, Polymeropoulos MH, 2015 Tasimelteon for non-24-hour sleep-wake disorder in totally blind people (SET and RESET): two multicentre, randomised, double-masked, placebo-controlled phase 3 trials. *Lancet* 386, 1754–1764. 10.1016/S0140-6736(15)60031-9. [PubMed: 26466871]
- Long Q, Upadhyaya D, Hattiangady B, Kim DK, An SY, Shuai B, Prockop DJ, Shetty AK, 2017 Intranasal MSC-derived A1-exosomes ease inflammation and prevent abnormal neurogenesis and memory dysfunction after status epilepticus. *Proc. Natl. Acad. Sci. U. S. A* 114, E3536–E3545. 10.1073/pnas.1703920114. [PubMed: 28396435]
- Malik A, Kondratov RV, Jamasbi RJ, Geusz ME, 2015 Circadian clock genes are essential for normal adult neurogenesis, differentiation, and fate determination. *PLoS One* 10, e0139655 10.1371/journal.pone.0139655. [PubMed: 26439128]
- Manganaro S, Loddenkemper T, Rotenberg A, 2017 The need for antiepileptic drug chronotherapy to treat selected childhood epilepsy syndromes and avert the harmful consequences of drug resistance. *J. Cent. Nerv. Syst. Dis* 9, 1179573516685883 10.1177/1179573516685883. [PubMed: 29308021]
- Matos H, de C, Koike BDV, Pereira WDS, de Andrade TG, Castro OW, Duzzioni M, Kodali M, Leite JP, Shetty AK, Gitali DLG, 2018 Rhythms of core clock genes and spontaneous locomotor activity in post- model of mesial temporal lobe epilepsy. *Front. Neurol* 9, 632 10.3389/fneur.2018.00632. [PubMed: 30116220]
- Matzen J, Buchheim K, Holtkamp M, 2012 Circadian dentate gyrus excitability in a rat model of temporal lobe epilepsy. *Exp. Neurol* 234, 105–111. 10.1016/j.expneurol.2011.12.029. [PubMed: 22226597]
- McDaniel SS, Rensing NR, Thio LL, Yamada KA, Wong M, 2011 The ketogenic diet inhibits the mammalian target of rapamycin (mTOR) pathway. *Epilepsia* 52, e7–e11. 10.1111/j.1528-1167.2011.02981.x. [PubMed: 21371020]
- Mirzoev A, Bercovici E, Stewart LS, Cortez MA, Snead OC, Desrocher M 3rd, 2012 Circadian profiles of focal epileptic seizures: a need for reappraisal. *Seizure* 21, 412–416. 10.1016/j.seizure.2012.03.014. [PubMed: 22537415]
- Mishra V, Shuai B, Kodali M, Shetty GA, Hattiangady B, Rao X, Shetty AK, 2015 Resveratrol treatment after status epilepticus restrains neurodegeneration and abnormal neurogenesis with suppression of oxidative stress and inflammation. *Sci. Rep* 5, 17807 10.1038/srep17807. [PubMed: 26639668]
- Mistlberger RE, Skene DJ, 2005 Nonphotic entrainment in humans? *J. Biol. Rhythms* 20, 339–352. 10.1177/0748730405277982. [PubMed: 16077153]
- Molina-Carballo A, Muñoz-Hoyos A, Rodríguez-Cabezas T, Acuña-Castroviejo D, 1994 Daynight variations in melatonin secretion by the pineal gland during febrile and epileptic convulsions in children. *Psychiatry Res.* 52, 273–283. [PubMed: 7991721]
- Musshoff U, Riewenherm D, Berger E, Fauteck J-D, Speckmann E-J, 2002 Melatonin receptors in rat hippocampus: molecular and functional investigations. *Hippocampus* 12, 165–173. 10.1002/hipo.1105. [PubMed: 12000116]
- Niles LP, Pickering DS, Arciszewski MA, 1987 Effects of chronic melatonin administration on GABA and diazepam binding in rat brain. *J. Neural Transm.* 70, 117–124. [PubMed: 3668517]
- Nzwalo H, Menezes Cordeiro I, Santos AC, Peralta R, Paiva T, Bentes C, 2016 24-hour rhythmicity of seizures in refractory focal epilepsy. *Epilepsy Behav.* 55, 75–78. 10.1016/j.yebeh.2015.12.005. [PubMed: 26773673]

- O'Callaghan EK, Anderson ST, Moynagh PN, Coogan AN, 2012 Long-lasting effects of sepsis on circadian rhythms in the mouse. *PLoS One* 7, e47087 10.1371/journal.pone.0047087. [PubMed: 23071720]
- Oggier DM, Weisbrod CJ, Stoller AM, Zenker AK, Fent K, 2010 Effects of diazepam on gene expression and link to physiological effects in different life stages in zebrafish *Danio rerio*. *Environ. Sci. Technol* 44, 7685–7691. 10.1021/es100980r. [PubMed: 20804179]
- Oike H, 2017 Modulation of circadian clocks by nutrients and food factors. *Biosci. Biotechnol. Biochem* 81, 863–870. 10.1080/09168451.2017.1281722. [PubMed: 28114877]
- Oishi K, Uchida D, Ohkura N, Doi R, Ishida N, Kadota K, Horie S, 2009 Ketogenic diet disrupts the circadian clock and increases hypofibrinolytic risk by inducing expression of plasminogen activator inhibitor-1. *Arterioscler. Thromb. Vasc. Biol* 29, 1571–1577. 10.1161/ATVBAHA.109.190140. [PubMed: 19628783]
- Ostendorf AP, Wong M, 2015 mTOR inhibition in epilepsy: rationale and clinical perspectives. *CNS Drugs* 29, 91–99. 10.1007/s40263-014-0223-x. [PubMed: 25633849]
- Passarelli V, Castro LHM, 2015 Gender and age influence in daytime and nighttime seizure occurrence in epilepsy associated with mesial temporal sclerosis. *Epilepsy Behav.* 50, 14–17. <https://doi.org/10.1016/j.yebeh.2015.05.028>. [PubMed: 26093513]
- Peixinho-Pena LF, Fernandes J, de Almeida AA, Gomes FGN, Cassilhas R, Venancio DP, de Mello MT, Scorza FA, Cavaleiro EA, Arida RM, 2012 A strength exercise program in rats with epilepsy is protective against seizures. *Epilepsy Behav.* 25, 323–328. 10.1016/j.yebeh.2012.08.011. [PubMed: 23103304]
- Petkova Z, Tchekalarova J, Pechlivanova D, Moyanova S, Kortenska L, Mitreva R, Popov D, Markova P, Lozanov V, Atanasova D, Lazarov N, Stoynev A, 2014 Treatment with melatonin after status epilepticus attenuates seizure activity and neuronal damage but does not prevent the disturbance in diurnal rhythms and behavioral alterations in spontaneously hypertensive rats in kainate model of temporal lobe epilepsy. *Epilepsy Behav.* 31, 198–208. 10.1016/j.yebeh.2013.12.013. [PubMed: 24440891]
- Portaluppi F, Smolensky MH, 2010 Perspectives on the chronotherapy of hypertension based on the results of the MAPEC study. *Chronobiol. Int* 27, 1652–1667. 10.3109/07420528.2010.510788. [PubMed: 20854140]
- Quigg M, Clayburn H, Straume M, Menaker M, Bertram EH, 1999 Hypothalamic neuronal loss and altered circadian rhythm of temperature in a rat model of mesial temporal lobe epilepsy. *Epilepsia* 40, 1688–1696. 10.1111/jM528-1157.1999.tb01585.x. [PubMed: 10612331]
- Quigg M, Clayburn H, Straume M, Menaker M, Bertram EH 3rd, 2000 Effects of circadian regulation and rest-activity state on spontaneous seizures in a rat model of limbic epilepsy. *Epilepsia* 41, 502–509. [PubMed: 10802754]
- Ramanathan C, Kathale ND, Liu D, Lee C, Freeman DA, Hogenesch JB, Cao R, Liu AC, 2018 mTOR signaling regulates central and peripheral circadian clock function. *PLoS Genet.* 14, e1007369 10.1371/journal.pgen.1007369. [PubMed: 29750810]
- Ramgopal S, Thome-Souza S, Loddenkemper T, 2013 Chronopharmacology of anti-convulsive therapy. *Curr. Neurol. Neurosci. Rep* 13, 339 10.1007/s11910-013-0339-2. [PubMed: 23456771]
- Ramkisoensing A, Meijer JH, 2015 Synchronization of biological clock neurons by light and peripheral feedback systems promotes circadian rhythms and health. *Front. Neurol* 6, 128 10.3389/fneur.2015.00128. [PubMed: 26097465]
- Rath MF, Rovsing L, Møller M, 2014 Circadian oscillators in the mouse brain: molecular clock components in the neocortex and cerebellar cortex. *Cell Tissue Res.* 357, 743–755. 10.1007/s00441-014-1878-9. [PubMed: 24842045]
- Reinberg AE, Ashkenazi IE, 1993 Interindividual differences in chronopharmacologic effects of drugs: a background for individualization of chronotherapy. *Chronobiol. Int* 10, 449–460. [PubMed: 8111870]
- Roberts AJ, Keith LD, 1994 Sensitivity of the circadian rhythm of kainic acid-induced convulsion susceptibility to manipulations of corticosterone levels and miner-allocorticoid receptor binding. *Neuropharmacology* 33, 1087–1093. 10.1016/0028-3908(94)90147-3. [PubMed: 7838321]

- Rocha AKA, de A, de Lima E, do Amaral FG, Peres R, Cipolla-Neto J, Amado D, 2016 Pilocarpine-induced epilepsy alters the expression and daily variation of the nuclear receptor ROR α in the hippocampus of rats. *Epilepsy Behav.* 55, 38–46. 10.1016/j.yebeh.2015.11.026. [PubMed: 26731717]
- Rocha AKA, de A, de Lima E, Amaral F, Peres R, Cipolla-Neto J, Amado D, 2017 Altered MT1 and MT2 melatonin receptors expression in the hippocampus of pilocarpine-induced epileptic rats. *Epilepsy Behav.* 71, 23–34. 10.1016/j.yebeh.2017.01.020. [PubMed: 28460319]
- Roenneberg T, Merrow M, 2016 The circadian clock and human health. *Curr. Biol* 26, R432–43. 10.1016/j.cub.2016.04.011. [PubMed: 27218855]
- Romcy-Pereira RN, Gitaí DLG, Gitaí LLG, Leite JP, Garcia-Cairasco N, Paçó-Larson ML, 2008 [Genes and epilepsy II: differential gene expression in epilepsy]. *Rev. Assoc. Med. Bras* 54, 461–466. [PubMed: 18989569]
- San Y-Z, Liu Y, Zhang Y, Shi P-P, Zhu Y-L, 2015 Peroxisome proliferator-activated receptor- γ agonist inhibits the mammalian target of rapamycin signaling pathway and has a protective effect in a rat model of status epilepticus. *Mol. Med. Rep* 12, 1877–1883. 10.3892/mmr.2015.3641. [PubMed: 25891824]
- Sánchez-Barceló EJ, Mediavilla MD, Reiter RJ, 2011 Clinical uses of melatonin in pediatrics. *Int. J. Pediatr.* 892624. 10.1155/2011/892624.
- Saraf A, Luo J, Morris DR, Storm DR, 2014 Phosphorylation of eukaryotic translation initiation factor 4E and eukaryotic translation initiation factor 4E-binding protein (4EBP) and their upstream signaling components undergo diurnal oscillation in the mouse hippocampus: implications for memory persistence. *J. Biol. Chem* 289, 20129–20138. 10.1074/jbc.M114.552638. [PubMed: 24895127]
- Schmitz B, 2006 Effects of antiepileptic drugs on mood and behavior. *Epilepsia* 47 (Suppl 2), 28–33. 10.1111/j.1528-1167.2006.00684.x. [PubMed: 17105456]
- Schwindinger WF, Giger KE, Betz KS, Stauffer AM, Sunderlin EM, Sim-Selley LJ, Selley DE, Bronson SK, Robishaw JD, 2004 Mice with deficiency of G protein 3 are lean and have seizures. *Mol. Cell. Biol* 24, 7758–7768. 10.1128/mcb.24.17.7758-7768.2004. [PubMed: 15314181]
- Schwindinger WF, Mirshahi UL, Baylor KA, Sheridan KM, Stauffer AM, Usef S, Stecker MM, Mirshahi T, Robishaw JD, 2011 Synergistic roles for G-protein $\gamma 3$ and $\gamma 7$ subtypes in seizure susceptibility as revealed in double knock-out mice. *J. Biol. Chem* 287, 7121–7133. 10.1074/jbc.M111.308395. [PubMed: 22207761]
- Shetty AK, 2011 Progress in cell grafting therapy for temporal lobe epilepsy. *Neurotherapeutics* 8, 721–735. 10.1007/s13311-011-0064-y. [PubMed: 21892793]
- Shetty AK, 2014 Hippocampal injury-induced cognitive and mood dysfunction, altered neurogenesis, and epilepsy: can early neural stem cell grafting intervention provide protection? *Epilepsy Behav.* 38, 117–124. 10.1016/j.yebeh.2013.12.001. [PubMed: 24433836]
- Shetty AK, Hattiangady B, 2016 Grafted subventricular zone neural stem cells display robust engraftment and similar differentiation properties and form new neurogenic niches in the young and aged hippocampus. *Stem Cells Transl. Med* 5, 1204–1215. 10.5966/sctm.2015-0270. [PubMed: 27194744]
- Shetty AK, Turner DA, 1999 Aging impairs axonal sprouting response of dentate granule cells following target loss and partial deafferentation. *J. Comp. Neurol* 414, 238–254. [PubMed: 10516594]
- Shetty AK, Upadhy D, 2016 GABA-ergic cell therapy for epilepsy: advances, limitations and challenges. *Neurosci. Biobehav. Rev* 62, 35–47. 10.1016/j.neubiorev.2015.12.014. [PubMed: 26748379]
- Shetty AK, Zaman V, Hattiangady B, 2005 Repair of the injured adult hippocampus through graft-mediated modulation of the plasticity of the dentate gyrus in a rat model of temporal lobe epilepsy. *J. Neurosci* 25, 8391–8401. [PubMed: 16162921]
- Shimozaki K, 2018 Involvement of nuclear receptor REV-ERB β in formation of neurites and proliferation of cultured adult neural stem cells. *Cell. Mol. Neurobiol* 38, 1051–1065. 10.1007/s10571-018-0576-7. [PubMed: 29397477]

- Simeone TA, Matthews SA, Samson KK, Simeone KA, 2017 Regulation of brain PPARgamma2 contributes to ketogenic diet anti-seizure efficacy. *Exp. Neurol* 287, 54–64. 10.1016/j.expneurol.2016.08.006. [PubMed: 27527983]
- Singh R, Sharma PK, Malviya R, 2012 CIRCADIAN cycle and chronotherapeutics: recent trend for the treatment of various biological disorders. *Recent Pat. Drug Deliv. Formul* 6, 80–91. [PubMed: 22044257]
- Sliwa A, Plucinska G, Bednarczyk J, Lukasiuk K, 2012 Post-treatment with rapamycin does not prevent epileptogenesis in the amygdala stimulation model of temporal lobe epilepsy. *Neurosci. Lett* 509, 105–109. 10.1016/j.neulet.2011.12.051. [PubMed: 22227620]
- Smolensky MH, Hermida RC, Reinberg A, Sackett-Lundeen L, Portaluppi F, 2016 Circadian disruption: new clinical perspective of disease pathology and basis for chronotherapeutic intervention. *Chronobiol. Int* 33, 1101–1119. 10.1080/07420528.2016.1184678. [PubMed: 27308960]
- Snider KH, Sullivan KA, Obrietan K, 2018 Circadian regulation of hippocampal-dependent memory: circuits, synapses, and molecular mechanisms. *Neural Plast.*, 7292540. 10.1155/2018/7292540.
- Solt LA, Burris TP, 2012 Action of RORs and their ligands in (patho)physiology. *Trends Endocrinol. Metab* 23, 619–627. 10.1016/j.tem.2012.05.012. [PubMed: 22789990]
- Spencer DC, Sun FT, Brown SN, Jobst BC, Fountain NB, Wong VSS, Mirro EA, Quigg M, 2016 Circadian and ultradian patterns of epileptiform discharges differ by seizure-onset location during long-term ambulatory intracranial monitoring. *Epilepsia* 57, 1495–1502. 10.1111/epi.13455. [PubMed: 27396544]
- Stewart LS, Leung LS, 2003 Temporal lobe seizures alter the amplitude and timing of rat behavioral rhythms. *Epilepsy Behav.* 4, 153–160. 10.1016/S1525-5050(03)00006-4. [PubMed: 12697140]
- Stewart LS, Leung LS, Persinger MA, 2001 Diurnal variation in pilocarpine-induced generalized tonic-clonic seizure activity. *Epilepsy Res.* 44, 207–212. 10.1016/S0920-1211(01)00192-9. [PubMed: 11325576]
- Stokkan KA, Yamazaki S, Tei H, Sakaki Y, Menaker M, 2001 Entrainment of the circadian clock in the liver by feeding. *Science* 291, 490–493. 10.1126/science.291.5503.490. [PubMed: 11161204]
- Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, Santos A, Doncheva NT, Roth A, Bork P, Jensen LJ, von Mering C, 2017 The STRING database in 2017: quality-controlled protein-protein association networks, made broadly accessible. *Nucleic Acids Res.* 45, D362–D368. 10.1093/nar/gkw937. [PubMed: 27924014]
- Takahashi JS, 2017 Transcriptional architecture of the mammalian circadian clock. *Nat. Rev. Genet* 18, 164–179. 10.1038/nrg.2016.150. [PubMed: 27990019]
- Takahashi JS, Hong H-K, Ko CH, McDearmon EL, 2008 The genetics of mammalian circadian order and disorder: implications for physiology and disease. *Nat. Rev. Genet* 9, 764–775. 10.1038/nrg2430. [PubMed: 18802415]
- Talathi SS, Hwang D-U, Ditto WL, Mareci T, Sepulveda H, Spano M, Carney PR, 2009 Circadian control of neural excitability in an animal model of temporal lobe epilepsy. *Neurosci. Lett* 455, 145–149. 10.1016/j.neulet.2009.03.057. [PubMed: 19368864]
- Talos DM, Jacobs LM, Gourmaud S, Coto CA, Sun H, Lim K-C, Lucas TH, Davis KA, Martinez-Lage M, Jensen FE, 2018 Mechanistic target of rapamycin complex 1 and 2 in human temporal lobe epilepsy. *Ann. Neurol* 83, 311–327. 10.1002/ana.25149. [PubMed: 29331082]
- Tamai S-I, Sanada K, Fukada Y, 2008 Time-of-day-dependent enhancement of adult neurogenesis in the hippocampus. *PLoS One* 3, e3835 10.1371/journal.pone.0003835. [PubMed: 19048107]
- Tang V, Michaelis R, Kwan P, 2014 Psychobehavioral therapy for epilepsy. *Epilepsy Behav.* 32, 147–155. 10.1016/j.yebeh.2013.12.004. [PubMed: 24418662]
- Tchekalarova J, Pechlivanova D, Itzev D, Lazarov N, Markova P, Stoynev A, 2010 Diurnal rhythms of spontaneous recurrent seizures and behavioral alterations of Wistar and spontaneously hypertensive rats in the kainate model of epilepsy. *Epilepsy Behav.* 17, 23–32. 10.1016/j.yebeh.2009.11.001. [PubMed: 20006556]
- Tchekalarova J, Pechlivanova D, Atanasova T, Markova P, Lozanov V, Stoynev A, 2011 Diurnal variations in depression-like behavior of Wistar and spontaneously hypertensive rats in the

- kainate model of temporal lobe epilepsy. *Epilepsy Behav.* 20, 277–285. 10.1016/j.yebeh.2010.12.021. [PubMed: 21277833]
- Tchekalarova J, Petkova Z, Pechlivanova D, Moyanova S, Kortenska L, Mitreva R, Lozanov V, Atanasova D, Lazarov N, Stoynev A, 2013 Prophylactic treatment with melatonin after status epilepticus: effects on epileptogenesis, neuronal damage, and behavioral changes in a kainate model of temporal lobe epilepsy. *Epilepsy Behav.* 27, 174–187. 10.1016/j.yebeh.2013.01.009. [PubMed: 23435277]
- Tchekalarova J, Moyanova S, Fusco AD, Ngomba RT, 2015 The role of the melatonergic system in epilepsy and comorbid psychiatric disorders. *Brain Res. Bull.* 119, 80–92. 10.1016/j.brainresbull.2015.08.006. [PubMed: 26321393]
- Tekade AR, Gattani SG, 2010 Dual cross-linked pulsatile beads for chronotherapy of asthma: development and evaluation. *Drug Deliv.* 17, 581–586. 10.3109/10717544.2010.501460. [PubMed: 20626253]
- Thome-Souza S, Klehm J, Jackson M, Kadish NE, Manganaro S, Fernandez IS, Loddenkemper T, 2016 Clobazam higher-evening differential dosing as an add-on therapy in refractory epilepsy. *Seizure* 40, 1–6. 10.1016/j.seizure.2016.05.014. [PubMed: 27281712]
- Tilson HA, Mitchell CL, 1984 Neurobehavioral techniques to assess the effects of chemicals on the nervous system. *Annu. Rev. Pharmacol. Toxicol.* 24, 425–450. 10.1146/annurev.pa.24.040184.002233. [PubMed: 6375549]
- Upadhyay D, Hattiangady B, Castro OW, Shuai B, Kodali M, Attaluri S, Bates A, Dong Y, Zhang S-C, Prockop DJ, Shetty AK, 2018 Human induced pluripotent stem cell-derived grafting after status epilepticus attenuates chronic epilepsy and comorbidities via synaptic integration. *Proc. Natl. Acad. Sci. U. S. A.* 10.1073/pnas.1814185115. 12 17 pii: 2018 14185 [Epub ahead of print].
- van Campen JS, Valentijn FA, Jansen FE, Joëls M, Braun KPJ, 2015 Seizure occurrence and the circadian rhythm of cortisol: a systematic review. *Epilepsy Behav.* 47, 132–137. 10.1016/j.yebeh.2015.04.071. [PubMed: 25982883]
- van Maanen A, Meijer AM, van der Heijden KB, Oort FJ, 2016 The effects of light therapy on sleep problems: a systematic review and meta-analysis. *Sleep Med. Rev.* 29, 52–62. 10.1016/j.smr.2015.08.009. [PubMed: 26606319]
- Van Nieuwenhuyse B, Raedt R, Sprengers M, Dauwe I, Gadeyne S, Carrette E, Delbeke J, Wadman WJ, Boon P, Vonck K, 2015 The systemic kainic acid rat model of temporal lobe epilepsy: long-term EEG monitoring. *Brain Res.* 1627, 1–11. 10.1016/j.brainres.2015.08.016. [PubMed: 26381287]
- van Vliet EA, Forte G, Holtman L, den Burger JCG, Sinjewel A, de Vries HE, Aronica E, Gorter JA, 2012 Inhibition of mammalian target of rapamycin reduces epileptogenesis and blood-brain barrier leakage but not microglia activation. *Epilepsia* 53, 1254–1263. 10.1111/j.1528-1167.2012.03513.x. [PubMed: 22612226]
- Vannest J, Szaflarski JP, Privitera MD, Schefft BK, Holland SK, 2008 Medial temporal fMRI activation reflects memory lateralization and memory performance in patients with epilepsy. *Epilepsy Behav.* 12, 410–418. 10.1016/j.yebeh.2007.11.012. [PubMed: 18162441]
- Waldau B, Hattiangady B, Kuruba R, Shetty AK, 2010 Medial ganglionic eminence-derived neural stem cell grafts ease spontaneous seizures and restore GDNF expression in a rat model of chronic temporal lobe epilepsy. *Stem Cells* 28, 1153–1164. 10.1002/stem.446. [PubMed: 20506409]
- Walker MC, Eriksson SH, 2011 Epilepsy and sleep disorders. *Eur. Neurol. Rev.* 6, 60 10.17925/ENR.2011.06.01.60.
- Wallach T, Kramer A, 2015 Chemical chronobiology: toward drugs manipulating time. *FEBS Lett.* 589, 1530–1538. <https://doi.org/10.1016Zj.febslet.2015.04.059>. [PubMed: 25957770]
- Weiss G, Lucero K, Fernandez M, Karnaze D, Castillo N, 1993 The effect of adrenalectomy on the circadian variation in the rate of kindled seizure development. *Brain Res.* 612, 354–356. 10.1016/0006-8993(93)91686-M. [PubMed: 8330212]
- Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, Sajed T, Johnson D, Li C, Sayeeda Z, Assempour N, Iynkkaran I, Liu Y, Maciejewski A, Gale N, Wilson A, Chin L, Cummings R, Le D, Pon A, Knox C, Wilson M, 2018 DrugBank 5.0: a major update to the DrugBank database for 2018. *Nucleic Acids Res.* 46, D1074–D1082. 10.1093/nar/gkx1037. [PubMed: 29126136]

- Wright KP, Burke TM, Lee-Chiong T, 2009 A glossary of circadian rhythm terminology for the researcher and clinician. *Sleep Med. Clin* 4, 301–304. 10.1016/j.jsmc.2009.03.002.
- Xiao F, He M, Wang X-F, Xi Z-Q, Li J-M, Wu Y, Zhong Z-F, Zhu D, Lu Y, Li H-W, 2007 [Overexpression of the Cdc42 in the brain tissue of human with intractable temporal epilepsy]. *Zhonghua Yi Xue Za Zhi* 87, 2030–2032. [PubMed: 17925172]
- Yalın O, Arman F, Erdoğan F, Kula M, 2006 A comparison of the circadian rhythms and the levels of melatonin in patients with diurnal and nocturnal complex partial seizures. *Epilepsy Behav.* 8, 542–546. 10.1016/j.yebeh.2005.12.015. [PubMed: 16524783]
- Yan L, Silver R, 2016 Neuroendocrine underpinnings of sex differences in circadian timing systems. *J. Steroid Biochem. Mol. Biol* 160, 118–126. 10.1016/j.jsbmb.2015.10.007. [PubMed: 26472554]
- Yegnanarayan R, Mahesh SD, Sangle S, 2006 Chronotherapeutic dose schedule of phenytoin and carbamazepine in epileptic patients. *Chronobiol. Int* 23, 1035–1046. 10.1080/07420520600921112. [PubMed: 17050216]
- Zeng L-H, Rensing NR, Wong M, 2009 The mammalian target of rapamycin signaling pathway mediates epileptogenesis in a model of temporal lobe epilepsy. *J. Neurosci* 29, 6964–6972. 10.1523/JNEUROSCI.0066-09.2009. [PubMed: 19474323]
- Zhang R, Lahens NF, Ballance HI, Hughes ME, Hogenesch JB, 2014 A circadian gene expression atlas in mammals: implications for biology and medicine. *Proc. Natl. Acad. Sci. U. S. A* 111, 16219–16224. 10.1073/pnas.1408886111. [PubMed: 25349387]
- Zhang Y, Liu J, Luan G, Wang X, 2015 Inhibition of the small GTPase Cdc42 in regulation of epileptic-seizure in rats. *Neuroscience* 289, 381–391. 10.1016/j.neuroscience.2014.12.059. [PubMed: 25595978]

Box 1**Key terms in Chronobiology.****Amplitude:**

The difference between the peak or trough and the mean value of a wave. It is a measure of how the variable intensity changes in the circadian time.

Central or master oscillator:

In mammals, the hypothalamic suprachiasmatic nucleus (SCN), which receives photic inputs from the retinohypothalamic tract (RHT) and nonphotic cues from other afferent inputs, systemically orchestrates the multiple endogenous rhythms in the brain and other tissues to environmental signals, particularly the light (LeGates et al., 2014; Ramkisoensing and Meijer, 2015).

Chronobiotic:

An agent capable of influencing parameters of biological rhythms (e.g., the phase setting) (Reinberg and Ashkenazi, 1993).

Chronotherapy:

A treatment regimen developed based on the circadian rhythms of the patient. It can be related to chronopharmacology or other manipulations of the body's clock or external cues (*zeitgebers*).

Chronopharmacology:

Pharmacotherapy adjusted according to the internal biological time to maximize treatment efficacy and minimize side effects.

Circadian rhythm:

A rhythm of physiological, metabolic or behavioral oscillations occurring in ~24 h. Circadian rhythms persist in constant conditions.

Circadian Time (CT):

CT is a quantification of time as defined by an organism's endogenous circadian clock, without reference to any environmental regulators or *zeitgebers*.

(Core) Clock genes:

Clock genes are components of the circadian clock, which intricately interact with each other and generate oscillations of gene expression. The protein products of clock genes are essential components for the generation and regulation of circadian rhythms.

Clock-controlled genes (CCGS):

CCGS are genes regulated by the core clock genes (Wright et al., 2009). Rhythmic translation and function of CCGS ultimately underlie daily oscillations at a cellular and organismal level.

Constant darkness:

It is an experimental procedure employed to estimate the circadian period and phase-shifting agents under environmental conditions of continuous darkness exposure (Wright et al., 2009).

Coupling:

Coupling is a process by which two or more oscillators interact in a stable phase relationship. Synchronization occurs through coupling mechanisms. In this sense, the SCN is composed of a network of coupled oscillators (Hafner et al., 2012). Uncoupling occurs when two or more oscillators drive out of this phase stability (e.g., peripheral oscillators uncoupled from the SCN).

D-box:

D-box is a transcriptional regulatory element (TTATG[T/C]AA) that participates in circadian rhythm regulation (Kumaki et al., 2008). It is a DBP/E4BP4 binding element.

E-box:

E-box is a transcriptional regulatory element (CACGTG) that participates in circadian rhythm regulation (Kumaki et al., 2008). It is a CLOCK/BMAL1-binding element.

Harmonic periods:

These are period lengths harmonic with the circadian rhythms (e.g., 24 h, 12 h, 8 h). Harmonics are cycles that fit an exact number of times into a major cycle, such as the 24-hour cycle.

Internal temporal order:

It is an order of maintenance of internal synchronization among the central and peripheral oscillators within an organism.

Misalignment:

A state in which the individual's sleep/wake cycle is misaligned to the biological night, or there is a misalignment between central and peripheral rhythms (Roenneberg and Merrow, 2016).

Period:

The period is an amount of time for a cycle to be completed, returning to the same phase. For example, the distance in time between two consecutive peaks (or troughs, etc.) of a wave.

Peripheral oscillators:

Refer to the circadian oscillators located in cells, tissues or organs outside of the SCN.

Phase:

The phase is an instantaneous state of oscillation within a period (reference point), generally the acrophase (peak of a rhythm).

Phase shift:

The phase shift is a displacement of oscillation along the time axis. Can be a phase advance or a phase delay.

Resetting:

Resetting is a phase shift in clock gene expression due to a synchronizing agent able to restart the cycle.

Shift work:

Shift work is a schedule of work involving non-traditional working hours, usually during the evening and night (“Dictionary of Circadian Physiology,” n.d.).

Spontaneous Locomotor Activity (SLA):

SLA is a spontaneous motor behavior resulting from the integrated neural activity, controlled by the circadian rhythm and measured as a rest-activity pattern (Tilson and Mitchell, 1984). SLA is a behavioral marker for circadian rhythm alterations.

Synchronization:

Synchronization is a state in which two or more oscillators assume the same frequency due to mutual or unilateral influences (“Glossary | Centre for Chronobiology,” n.d.).

Ultradian rhythm:

Ultradian rhythm is an oscillatory biological process with a periodicity of fewer than 24 h.

Zeitgebers:

Zeitgebers are inputs to circadian clocks that alter the timing of rhythmic biological oscillations (Wright et al., 2009).

Zeitgeber Time (ZT):

It is a quantification of time defined with reference to environmental regularities or *zeitgebers* (Wright et al., 2009).

Fig. 1.

A general overview of the molecular control of circadian rhythms. Synchronizing inputs such as light, food, and exercise modulate the endogenously generated circadian clock present in the central oscillator (SCN) or the peripheral oscillators throughout the body. The middle panel shows the cellular clock pathways. By interacting with E-box promoters, the BMAL1: CLOCK complex instigates transcription of target genes, period (PER1/2) and cryptochrome (CRY1/2), which leads to the formation of PER: CRY heterodimers in the cytoplasm. The activity of BMAL1: CLOCK complex is regulated through translocation of PER: CRY heterodimers into the nucleus. BMAL1: CLOCK heterodimers also activate the transcription of two retinoic acid-related orphan receptor response elements, RORA (activator of BMAL1 transcription) and REV-ERBA (repressor of BMAL1 transcription). Both RORA and REV-ERBA bind to rev responsive element (RRE) present in BMAL1 promoter to regulate the expression of BMAL1. The rhythmic outputs may also feedback to the cellular clock adding another level of temporal regulation in a multicellular system.

Fig. 2. Potential effects of epileptogenesis and chronotherapy on circadian rhythm parameters. At level of genes (A), distinct genes could be affected in phase and amplitude, changing the temporal transcriptome profile. At the system level (B), different oscillators may lose the stability in phase relationship due to changes in the temporal outputs from the hippocampus, uncoupling functional interconnected structures. A possible contribution of chronotherapy relies on resetting mechanisms that could restore the synchrony between the central (SCN) and peripheral oscillators.