CORRESPONDENCE

Central Pontine Myelinolysis and Osmotic Demyelination Syndrome

by Dr. med. Johann Lambeck, Dr. med. Maren Hieber, Dr. med. Andrea Dreßing, and Dr. med. Wolf-Dirk Niesen in issue 35–36/2019

Inadequate Therapy

The authors emphasize the role of magnetic resonance imaging (MRI) and present some new therapeutic approaches, the effectiveness of which cannot yet be conclusively determined (1). Even though MRI has led to a deepening of the pathophysiological understanding of these syndromes, I would argue that neither MRI nor “the considerable progress achieved by modern intensive care medicine” has led to a decisive advancement.

Osmotic demyelination syndromes can be prevented by avoiding too rapid changes in serum osmolality (<1 mosmol/h, corresponding to 0.5 mmol/L sodium/h if only changes in sodium concentration are considered). However, diseases are not themselves the cause of osmotic demyelination syndromes, as all changes in serum osmolality resulting from diseases are slow (with the exception of rapid-onset central diabetes insipidus, with increased sodium). Rather, inadequate therapy causes demyelination syndromes. With the goal of quickly correcting hyponatremia, the brain tissue is exposed to osmotic stress.

Careful planning of the infusion therapy and tightly controlled monitoring of the osmotically active substances, followed by an adequate adjustment of the infusion therapy, can prevent rapid fluctuations in osmolality and thus prevent demyelination syndromes. It is important to consider the cause of the electrolyte imbalance. In this light, it is also necessary to determine whether hypovolemia, normovolemia, or hypervolemic hyponatremia is present. For this, the technical requirements (point-of-care devices, to determine sodium and potassium levels, among other things) and knowledge (osmolality should not change by more than 1 mosmol/h) are well-known and widely available. If osmotic demyelination syndromes nevertheless occur, it is due to inadequate care during therapy. Work overload of doctors may be one of the causes.

DOI: 10.3238/arztebl.2020.0041a

Reference

1. Lambeck J, Hieber M, Dreßing A, Niesen WD: Central pontine myelinolysis and osmotic demyelination syndrome. *Dtsch Arztebl Int* 2019; 116: 600–6.

Reiner Klick
Iserlohn, Germany
reiner-klick@gmx.de

Open Questions

The article (1) provides an excellent overview of a difficult, often somewhat neglected problem that also plays a role in intensive care medicine. However, the proposed preparation of a 3% sodium chloride (NaCl) solution is sometimes not easy to carry out. I think measuring exactly 318 mL during an emergency situation is not always feasible. My simpler solution is based on the calculation that 500 mL of a 3% NaCl solution must contain 15 g NaCl. So, if one lets 150 mL run out of an infusion bottle containing 500 mL of 5% glucose, and replaces it with 150 mL of

a 10% NaCl solution, a 3% NaCl solution is prepared. This procedure prevents an excessive chloride concentration. Are there any objections to this?

What should be the initial rate of infusion of the 3% solution in patients with or without additional fluid deficiency in the case of severe neurological symptoms?

In my opinion, the recommendation to initially compensate for a possibly existing potassium deficit before therapy of hyponatremia is often hardly practicable in case of severe neurological symptoms. Should a confused or convulsing patient, whose cerebral condition could probably be improved by administering a NaCl solution, really receive an intravenous potassium replacement—possibly over several hours—before beginning the actual treatment of the underlying disease? Here, 30 mmol/h is the maximum permissible rate of potassium replacement via central venous catheter. I remember patients with hyponatremia who would not have tolerated this for long. What would you recommend in this case?

DOI: 10.3238/arztebl.2020.0041b

Reference

1. Lambeck J, Hieber M, Dreßing A, Niesen WD: Central pontine myelinolysis and osmotic demyelination syndrome. *Dtsch Arztebl Int* 2019; 116: 600–6.

Prof. Dr. med. Friedrich Lübbecke
Uelzen, Germany

Etiology Falls Short

It is alarming that the authors have shortened the etiology in their article (1) in an unacceptable manner, and not only from an intensive care perspective. After all, osmotic myelinolysis and demyelination are much more frequently due to pharmacogenomics than to a singular consequence of alcohol abuse and/or liver transplantation (LTX). Therefore, in such a review, it is indispensable to include at least a summary of typical iatrogenic hyponatremia syndromes as well as the risk of syndrome of inappropriate antidiuretic hormone secretion (SIADH) as an adverse drug effect.

In addition to or together with diuretics, the following can be considered as causal (2, 3):

- Neuroleptics
- Antidepressants (e.g. selective serotonin reuptake inhibitors [SSRI])
- Coxibs
- Nonsteroidal anti-inflammatory drugs (NSAIDs)
- Proton pump inhibitors (PPI)
- Cardiac agents
- Anticonvulsants (including pregabalin)
- Immunologicals
- Opiates
- Oncologics
- Ecstasy and other party drugs

An accurate medication history and high therapy vigilance are crucial for the course, especially as the pharmacogenomic adverse effects (e.g., paresis, signs of Parkinson's, delirium) often overlap the respective primary complaint. Patients with reduced ability to cooperate in gerontology, intensive care, neurology, pediatrics, and psychiatry (4) are therefore particularly often and severely affected.

DOI: 10.3238/arztebl.2020.0041c

References

1. Lambeck J, Hieber M, Dreßing A, Niesen WD: Central pontine myelinolysis and osmotic demyelination syndrome. *Dtsch Arztebl Int* 2019; 116: 600–6.
2. Liamis G, Milionis H, Elisaf M: A review of drug-induced hyponatraemia. *Am J Kidney Dis* 2008; 52: 144–53.
3. Spasovski G, Vanholder R, Allolio B, et al: Clinical practice guideline on diagnosis and treatment of hyponatraemia. *Nephrol Dial Transplant* 2014; 29 (Suppl. 2): ii1–ii39.
4. Post B, van Gool W A, Tijssen MAJ: Transient parkinsonism in isolated extrapontine myelinolysis. *Neurol Sci* 2009, 30: 325–8.

Dr. med. Burger Lichtenstein

Stuttgart, Germany
helensson@t-online.de

In Reply:

We thank Mr. Klick for his comments regarding the prognosis and prevention of osmotic demyelination syndromes (ODS). We agree that earlier and better detection of ODS using magnetic resonance imaging (MRI) has not been the decisive basis for improving the disease prognosis. As discussed critically in our article (1), the increased use of MRI has led to the detection of many ODS patients who are oligosymptomatic and/or have symptoms only visible by morphological imaging, and their inclusion in the overall prognosis makes it appear to be better than in previous years. Nevertheless, modern intensive care medicine has an increasing number of early rehabilitation approaches that also play a part in improving the prognosis. Having a good long-term outcome is now more frequent, even for patients with initial severe clinical impairment (2). The most relevant influence on prognosis, however, is the careful preventive therapy in the form of a slow electrolyte and osmolality correction, as described in our section Therapy, and we would also like to thank Mr. Klick for highlighting the importance of prevention.

We thank Prof. Lübbecke for his practice-oriented questions. With respect to producing the 3% NaCl solution, it makes sense

from our point of view to carefully measure the quantities mentioned. While an emergency preparation is usually not necessary, there is nothing to be said against the proposed formula if this is the case. With respect to the initial infusion rate, a correction of 1.5–2 mmol/L/h can be considered for severely affected patients for a maximum of 3 h. The infusion rate can then be adjusted to the sodium deficit (for correction up to 120 mmol/L) ($0.5 \times \text{kg body weight} \times [120 \text{ minus current serum sodium mmol/L}]$). This correction can be continued for a maximum of 12 mmol/L/24 h. Overall, however, it is usually better to do nothing than to do too much if necessary. Therefore, a quick potassium correction that exceeds the maximum infusion rate mentioned is not necessary. Any neurological complications (e.g. epileptic seizures) must of course also be treated with an anticonvulsant in addition to sodium and/or potassium correction.

We would also like to thank Dr. Lichtenstein for his critical remarks on the overly-brief etiology of ODS. We agree about the importance of clarifying the underlying cause in patients with ODS, as we addressed in the article (1). While we would have liked to have intensively examined and discussed the etiological background of hyponatremia, this would have required its own review. Here it is important to note the frequency of hyponatremia and to refer to the various reviews on this topic (3–5), as well as the article mentioned by Dr. Lichtenstein.

DOI: 10.3238/arztebl.2020.0042

References

1. Lambeck J, Hieber M, Dreßing A, Niesen WD: Central pontine myelinolysis and osmotic demyelination syndrome. *Dtsch Arztebl Int* 2019; 116: 600–6.
2. de Souza A: Movement disorders and the osmotic demyelination syndrome. *Parkinsonism Relat Disord* 2013; 19: 709–16.
3. Sterns RH: Disorders of plasma sodium - causes, consequences, and correction. *N Engl J Med* 2015; 372: 55–65.
4. Knepper MA, Kwon TH, Nielsen S: Molecular physiology of water balance. *N Engl J Med* 2015; 372: 1349–58.
5. Adrogue HJ, Madias NE: Hyponatremia. *N Engl J Med* 2000; 342: 1581–9.

On behalf of the authors

Dr. med. Wolf-Dirk Niesen

Klinik für Neurologie und Neurophysiologie, Universitätsklinikum Freiburg, Germany
wolf-dirk.niesen@uniklinik-freiburg.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.