THE AURICULAR WAVE (P) OF THE ELECTRO-CARDIOGRAM

CLINICAL OBSERVATIONS WITH ESPECIAL REFERENCE TO PULMONIC
AND MITRAL STENOSES

A. A. ALEXANDER, M.D.

OAKLAND, CALIF.

H. F. KNIGHT, M.D.

ROCHESTER, N. Y.

AND

PAUL D. WHITE, M.D.

BOSTON

LITERATURE

Studies concerning the diagnostic significance of changes in the auricular or P wave of the electrocardiogram have not been numerous. Increase in height and prolongation of the wave, occurring most often in mitral stenosis, have been noted by Einthoven,¹ Kraus and Nicolai² and Samojloff;³ Goddard⁴ assumed a more or less definite quantitative relationship to exist between such augmentation of the P wave and auricular hypertrophy. He considered as increased all P waves when they were greater than T in the same curve, that is $\frac{P}{T} = 1 +$. Checking this relationship, he found that there was some relative increase in height (of the P wave) in 54 per cent. of the cases of mitral disease studied. However, in five cases that came to necropsy, one with very slight hypertrophy of the auricle showed marked exaggeration of the P wave while another with decided hypertrophy showed only a slight relative alteration of the P. On the other hand a check of all the exaggerated P waves against the histories of the patients showed that 45 per cent. had mitral disease and that a small percentage had clinically normal hearts. Goddard then concludes that the association of P wave exaggeration with auricular hypertrophy, though suggestive, is not a constant one.

Experiments seeking to determine the nature and source of the P wave by Samojloff,⁵ Kraus and Nicolai,⁶ Lewis, Meakins and White;⁷

1. Einthoven, W.: *Le Telecardiogramme*, Arch. internat. de physiol. **4**:148, 1906-1907.

2. Kraus, F., and Nicolai, G. F.: *Berl. klin. Wchnschr.* **44**:812, 1907.

3. Samojloff, A.: *München. med. Wchnschr.* **56**:1943, 1909.

4. Goddard, C. H.: *Arch. Int. Med.* **16**:633 (Oct.) 1915.

5. Samojloff, A.: *Festschrift Herman*, 1908, p. 171.

6. Kraus, F., and Nicolai, G. F.: *Das Elektrokardiogramm des Gesunden und Kranken Menschen*, Leipzig, 1910.

7. Lewis, T.; Meakins, J., and White, P. D.: *Phil. Tr. Roy Soc. London*, **205**:375, 1914.

and Eyster and Meek⁸ have shown that it is a composite representation of the excitation wave as it spreads from the region of the sinus, the sino-auricular node, over the auricle. Recently, the interesting and fundamentally important work of Einthoven⁹ has shown that the electrical and mechanical expressions of muscular activity are synchronous and parallel in degree. This applies to heart muscle as well as to skeletal muscle. The myocardiogram if accurately obtained corresponds exactly with the electrocardiogram in time of onset, duration and force. Thus the auricular or P wave of the electrocardiogram does not represent simply the brief early expression of the excitatory process in the auricle but in fact the entire auricular contraction. Instead of measuring in time only one third of the duration of auricular action it really coincides with it. The T wave of the auricle—the so-called Ta wave of the electrocardiogram—had been pointed out years ago by several observers but it has now been correlated to auricular systole by Einthoven.⁹ Sprague and White¹⁰ have recently shown the frequent clinical occurrence of this Ta wave of the electrocardiogram. Thus, the larger and more active the auricles, the larger and longer should be the auricular deflections of the electrocardiogram.

Attention has also been called to the influence of the position of the heart in the body, as determined by respiration or other factors, and of the periodic shift of the pacemaker within the node, as shown by Lewis, Meakins and White.⁸

Finally, the influence of the extrinsic cardiac nerves on the P wave has been studied. Einthoven¹¹ demonstrated that simple section of the vagus in dogs produced a marked rise in the P wave. In such an experiment of course changes in the auricle could play no rôle. Rothberger and Winterberg¹² repeated and confirmed this, and further showed that section of the cardiac accelerators (increased vagus effect) was followed by a pronounced decrease in P. Selenin,¹³ however, saw no such effect on the dog's heart in situ, following full therapeutic dosage of digitalis.

PRESENT INVESTIGATION

The reason for the present study was the desire to determine the significance of high P waves with reference to the diagnosis of pulmonary stenosis and mitral stenosis.

8. Eyster, J. A. E., and Meek, W. J.: *Arch. Int. Med.* **11**:204 (Feb.) 1913.

9. Einthoven, W.: *Goddard Lectures*, Harvard Medical School, 1924.

10. Sprague, H. B., and White, P. D.: *J. Clin. Invest.* **1**:389 (April) 1925.

11. Einthoven, W.: *Arch. f. d. ges. Physiol.* **122**:537, 1908.

12. Rothberger, J., and Winterberg, H.: *Arch. f. d. ges. Physiol.* **135**:506, 1910.

13. Selenin, W. P.: *Arch. f. d. ges. Physiol.* **143**:137, 1912.

The study has included three distinct parts as follows: First, the analysis of (a) the amplitude of the P wave, (b) the values of angle and index of electrical axis deviation, and (c) the clinical diagnosis in 132 unselected cases seen at the Massachusetts General Hospital from 1914 to 1924 in whom electrocardiograms showed an amplitude of P in Lead I greater than 0.1 millivolt (1 mm.), or of P in Lead II equal to or greater than 0.3 millivolt (3 mm.). Second, a similar analysis of all cases definitely diagnosed pulmonic stenosis and coming to electro-

Fig. 1.—Leads I, II and III, normal, for comparison with Figures 2 and 3; time, 0.2 second; potential, 10^{-4} volts per millimeter.

cardiographic study at the Massachusetts General Hospital from 1914 to 1924. Third, the same study of 100 unselected cases of "pure" mitral stenosis electrocardiographed at the Massachusetts General Hospital.

The height of the P wave has been studied rather than its length or duration for several reasons: First, it is more evident and more easily measured; second, the occurrence of the T wave of the auricle, the so-called Ta wave, sometimes appearing as a direct prolongation

of the P wave, makes it difficult to determine the limits of the P, and third, the height of the P wave may be abnormally increased with little if any increase in duration while the reverse is rarely true. Almost invariably if the P is longer than normal it is also higher than normal in either Lead I, Lead II, or both. For example, out of the fifty-eight cases of mitral stenosis with normal rhythm (listed in Table 7) only nine failed to show high P waves in either Lead I or Lead II and none of these nine showed any abnormal length of the P waves; incidentally, the P wave in Lead III was low in all of these nine cases.

Rarely if ever is the P wave in Lead III as large as in the other leads; usually it is of very low potential, either upright, inverted or iso-electric. Thus P_3 has not been included in the present analysis because of its unimportance.

With the abnormal increase in height of the P waves in Lead I and Lead II there is often abnormal notching in addition to increased length. A duration of the P wave (not including the Ta wave) beyond 0.1 second in any lead and a height above 1 mm. (0.1 millivolt) in Lead I, or a height equal to or above 3 mm. (0.3 millivolt) in Lead II, we have considered abnormal.

The formula used for determining the index was that outlined by White and Bock¹⁴ and the angles were arrived at by the use of the triangle of Einthoven, following the method described by Carter, Richter and Greene.¹⁵ By the index method figures of -12 or more negative indicate abnormal right axis deviation, and of $+20$ or more positive abnormal left axis deviation; -12 to $+20$ are within normal limits. By the angle method figures beyond $+90$ degrees represent abnormal right axis deviation and on the negative (upper) side of 0 degrees abnormal left axis deviation; 0 to $+90$ degrees (the right lower quadrant) are within normal limits.

PART I

Tables 1, 2, 3, 4, 5 and 6 include the cases examined in the first part of the study. The clinical diagnosis has determined the inclusion of the various cases under one or another of these tables, wherein pulmonic stenosis, mitral stenosis pure, and mitral stenosis with other valve defects are each given separately, while other diagnoses encountered are embodied in Table 6.

Examination of Tables 2, 3, 4 and 5 shows that the amplitude of P_2 was equal to or greater than 3 mm. in 68 per cent. of all of the seventy-two cases of mitral stenosis encountered, and an interesting

14. White, P. D., and Bock, A. V.: *Am. J. M. Sc.* **156**:17 (July) 1918.

15. Carter, E. P.; Richter, C. P., and Greene, C. H.: *Bull. Johns Hopkins Hosp.* **30**:162 (June) 1919.

TABLE 1.—*Pulmonic Stenosis* *

Plate No.	Index, Mm.	Angle, Degrees	P ₁ , Mm.	P ₂ , Mm.
6413.....	-20	150	2.5	4
7465.....	-30	90	1	5
6838.....	-34	125	2	4
1994.....	-12	165	1.8	3.8
7919.....	-42	159	4	4
745.....	-43	143	1.5	1.2
3940.....	-18	112	2	2.5
6448.....	-36	134	2	3
5805.....	-28	150	2	4.5
974.....	-25	140	2	2.5
4057.....	-23	142	2.5	3
6573.....	-32	132	1	3
7465.....	-30	90	1	5
7438.....	-24	118	2	3.5
5879.....	-30	143	1.7	4
7050.....	-18	149	2	2.5
6590.....	-28	118	2	3
8870.....	-16	116	1.5	4
6448.....	-36	134	3	4
8098.....	-29	118	0.5	6

* Values throughout these tables are positive unless otherwise noted.
 Summary: Total cases, nineteen (twenty plates). Sixteen cases had an amplitude of P₂ equal to or greater than 3 mm. The other three cases had an amplitude of P₁ greater than 1 mm. Eighteen had an index below -15. One had an index of -12. All had angles of 90 degrees or more.

TABLE 2.—*Mitral Stenosis (Pure)* *

Plate No.	Index, Mm.	Angle, Degrees	P ₁ , Mm.	P ₂ , Mm.
905.....	-15	120	3	3
1917.....	-11	69	2	2
2209.....	-17	102	2	2
2275.....	-15	130	1.8	2
1605.....	-8.5	81	1.5	2.7
3095.....	-13	95	1	4
2352.....	-19.5	120	2.2	2.2
2362.....	2	56	1.5	3.5
6653.....	-1	59	2	3
6685.....	-20	90	1	3
6675.....	-1	62	2	3.5
6801.....	-7	75	1.5	3.5
6774.....	-4.1	164	1	3
6865.....	-6	90	2.2	1.8
6879.....	-23	102	1.2	2.5
6959.....	-10	72	1.5	3
6948.....	5.5	25	2	4
8518.....	-12	98	2	5
7151.....	-5	69	1	4
6525.....	-10	90	3	3
6471.....	-16	117	2	4
7106.....	-6	76	1	5
6898.....	-20	112	1.5	5
1324.....	-13	121	1.6	2.5
2683.....	-10	78	1.5	3.2
6556.....	-20	122	2	2.5
5562.....	-32	122	2	2.5
8518.....	-12	98	3	5
6435.....	-9	68	2.5	4
7057.....	-11	99	2.5	5
6308.....	-3	67	2	3
6325.....	-23	98	2	2.5
6535.....	-13	99	2	3
6831.....	15	-18	1.5	3.5
6935.....	-9	108	2	2.5
7104.....	-13	92	2	2.5
5023.....	0	60	1.5	3
6233.....	-1	62	2	3
4476.....	2	43	2	3
4501.....	-9	73	2	2
7191.....	-31	118	3	2.5
1355.....	-21	122	2.5	4
1396.....	-19	105	3	2.5

* Among the cases with electrocardiograms showing high P waves these had been diagnosed as pure mitral stenosis.

Summary: Total cases, forty-three. Thirty-eight had an amplitude of P₁ greater than 1 mm. (88 per cent.); thirty-eight had an amplitude of P₂ greater than 2 mm. (88 per cent.), and twenty-seven had an amplitude of P₂ equal to or greater than 3 mm. (63 per cent.). Indexes varied from +15 to -41 (four had positive values). Fourteen had indexes of -15 or below (33 per cent.); twenty-five had indexes of -10 or below (58 per cent.), and twenty-five had angles of +90 degrees or more (58 per cent.).

TABLE 3.—*Mitral Stenosis and Regurgitation*

Plate No.	Index, Mm.	Angle, Degrees	P ₁ , Mm.	P ₂ , Mm.
2173.....	-25	102	3	6
547.....	-25	127	2	5
948.....	-12	140	2	4
21.....	-22.5	96	2.5	5
332.....	-28.5	114	1	3
987.....	-18	112	1.5	2.1
1383.....	-25	107	1	3
1742.....	-14.5	88	1.5	2
2475.....	-32	109	3	2.5
3197.....	-6	85	1.5	1.3
4355.....	-21	132	2	2.5
4103.....	-3	68	2	3
50.....	-25	90	1.5	3.5
6689.....	-11	71	4	7

Summary: Total cases, fourteen. Twelve had an amplitude of P₁ greater than 1 mm. (85 per cent.); twelve had an amplitude of P₂ greater than 2 mm. (85 per cent.), and nine had an amplitude of P₂ equal to or greater than 3 mm. (64 per cent.). Nine had indexes below -15; twelve had indexes of -10 or below (85 per cent.), and ten had angles equal to or greater than +90 degrees (71 per cent.).

TABLE 4.—*Mitral Stenosis and Aortic Regurgitation*

Plate No.	Index, Mm.	Angle, Degrees	P ₁ , Mm.	P ₂ , Mm.
270.....	-15	90	1.8	2.5
4170.....	-12	73	3.5	5.5
7182.....	10	-47	1	3
7786.....	5	53	1.5	3
8017.....	-10	90	1.5	3.5
2189.....	-11.5	76	1.3	3.3
4171.....	2	58	4	3
2419.....	19	-4	1.2	1.5
3620.....	-9	72	1	3
6647.....	42	-32	2	4

Summary: Total cases, ten. Eight had an amplitude of P₁ over 1 mm. (80 per cent.); nine had an amplitude of P₂ over 2 mm., and eight had an amplitude of P₂ of 3 mm. or over (80 per cent.). Two had an angle of only 90 degrees or over. Four cases only had indexes of -10 or below (40 per cent.), and one case only had an index of -15.

TABLE 5.—*Mitral Stenosis and Regurgitation and Aortic Stenosis and Regurgitation*

Plate No.	Index, Mm.	Angle, Degrees	P ₁ , Mm.	P ₂ , Mm.
7682.....	-8	75	1.5	4
7721.....	-17	100	2	4.5
3974.....	0	62	3	5
3968.....	0.5	60	3	3
6800.....	-26	110	1.5	3.5

Summary: Total cases, five. All had amplitude of P₁ greater than 1 mm.; all had amplitude of P₂ of 3 mm. or over. All had angles of 60 degrees or over; two had angles of 90 degrees or over (40 per cent.), and two had indexes of -10 or below.

Summary of Tables 3, 4 and 5: Total cases, twenty-nine. Twenty-five had amplitude of P₁ greater than 1 mm. (86 per cent.); twenty-six had amplitude of P₂ equal to or over 2 mm., and twenty-two had amplitude of P₂ equal to or over 3 mm. (76 per cent.). Eighteen had indexes of -10 or below. Twenty-four had angles of 60 degrees or more, and fourteen had angles of 90 degrees or more (48 per cent.).

TABLE 6.—*Miscellaneous Conditions* *

Plate No.	Index, Mm.	Angle, Degrees	P ₁ , Mm.	P ₂ , Mm.	Clinical Diagnoses
4166	10	48	3	2	Arteriosclerosis; cerebral embolism; sino-auricular tachycardia
6148	-17	170	2	4	Effort syndrome; sino-auricular tachycardia
2057	-10	72	1	3	Arteriosclerosis; sino-auricular tachycardia
928	-15	123	1.5	2	Cardiac neurosis; sino-auricular tachycardia
809	0.5	61	1.6	5	Sino-auricular tachycardia
1313	-13	78	0.5	3	Pericarditis (?); mitral stenosis (?); sino-auricular tachycardia
1256	1	57	1.5	5	Goiter; toxemia of pregnancy; sino-auricular tachycardia
908	3	56	2.1	5	Nephritis; hypertension; arteriosclerosis
3295	-0.5	65	1.5	2	Nephritis; hypertension; arteriosclerosis; sino-auricular tachycardia
4358	5	83	2.5	2.5	Nephritis; hypertension; sino-auricular tachycardia
6410	19	23	2	4	Hypertension; congestive heart failure
6476	31	4	2.5	4	Hypertension; congestive heart failure; sino-auricular tachycardia
6601	1.5	59	1	6	Exophthalmic goiter; sino-auricular tachycardia
6716	0	54	1.5	4	Coronary sclerosis; sino-auricular tachycardia
6771	-27	110	1	4.5	Sino-auricular tachycardia
7001	14	5	2	1.5	Nephritis; hypertension; coronary sclerosis
7031	22	7	2	4	Hypertension; cardiac hypertrophy; arteriosclerosis
7042	45	35	1.5	3	Hypertension; arteriosclerosis; sino-auricular tachycardia
7162	-1	63	1	4.5	Essential hypertension
8281	19	30	2.1	4	Hypertension; arteriosclerosis; hemiplegia
7156	10	30	1	5	No disease
6575	-36	128	2	4	Mitral stenosis (?); sino-auricular tachycardia
6588	-11	73	1	4.6	Mitral stenosis (?); sino-auricular tachycardia
7063	9	28	1.5	2.5	Rheumatic history; normal heart
2954	-35	85	2	1	Arteriosclerotic and syphilitic heart; sino-auricular tachycardia
4654	-7	58	1.5	2	Perinephritic abscess; sino-auricular tachycardia
3854	-7	94	1	4	Hypertension; varicocele
5480	15	10	2	3.2	Chorea (?); mitral stenosis (?); sino-auricular tachycardia
3730	2	35	3	3	Nephritis; hypertension
1445	-8	75	2	3	Phthisis (?); mitral stenosis (?)
6820	-7	78	1	4	Mitral stenosis (?); sino-auricular tachycardia
6857	-8	100	2	2	Hypertension; uremia; prostatitis
6487	6	46	1	3	Nephritis; hypertension; sino-auricular tachycardia
7068	-2	65	2	4	Congenital heart (?); ventricular septal defect; sino-auricular tachycardia
7654	8	59	2	6.5	Congenital heart; septal defect; heart block
6590	-28	118	2	4	Congenital heart; septal defect; patent ductus arteriosus; sino-auricular tachycardia; pulmonic stenosis (?)
7170	-12	110	2	1	Congenital heart; ventricular septal defect (?); pulmonic stenosis (?); sino-auricular tachycardia
6459	-20	86	2.5	4	Congenital heart; pulmonic stenosis (?); sino-auricular tachycardia
8121	2	57	2	3.5	Nephritis; hypertension; uremia
8155	5	53	1.5	5	Nephritis; hypertension; uremia; sino-auricular tachycardia
8975	0	60	2.5	3	Sino-auricular tachycardia; curvature of spine

* All cases having other than definite pulmonic stenosis or mitral stenosis encountered in the review of cases with electrocardiograms showing high P waves.

Summary: Total cases, forty-one; doubtful mitral stenosis, six; doubtful pulmonic stenosis, three; probable interventricular septal defect without pulmonic stenosis, two; total congenital heart cases, five. Sino-auricular tachycardia (heart rate of 100 or more) without question of mitral stenosis or congenital heart defects, eighteen. Hypertension with or without nephritis, not included in foregoing listing, ten. Rheumatic history without evidence of heart disease, one, and no disease, one.

approximation was shown by the percentage of right axis deviation, in that 60 per cent. of all cases had indexes equal to or below -10 , and 54 per cent. had angles equal to or greater than 90 degrees.

A further interesting and expected finding in this series of combined mitral lesions (Tables 3, 4 and 5) is that the mechanism involved causes an increase in the amplitude of P but a decrease in the negativity of the indexes and in the positive angles (that is a lessened tendency to right axis deviation), as compared with the pure mitral stenosis cases.

Fig. 2.—Leads I, II and III in well marked mitral stenosis; large P waves in Leads I and II and pronounced right axis deviation (right ventricular preponderance).

It is of much interest to note that twenty-nine of these forty-one miscellaneous cases (Table 6) showed conditions that could easily explain the abnormally high P waves. Mitral stenosis and pulmonic stenosis were probably present in the cases listed doubtful though the signs were not well enough marked to make a positive diagnosis. There were two other cases of congenital heart disease showing large P waves probably with interventricular defects; it is likely that the auri-

cles as well as the ventricles in some patients of this type are overworked with resultant hypertrophy or increased activity. Sino-auricular tachycardia is usually the result of sympathetic nerve stimulation, and it is a well recognized fact physiologically that such stimulation may result in increase in amplitude of the auricular deflection (and also of the T wave) of the electrocardiogram. The frequent finding of hypertension in some of the cases with high P waves may be significant—here again there is a possibility that the auricles as well as the ventricles

Fig. 3.—Leads I, II and III in congenital pulmonic stenosis; large P waves in Leads I and II and marked right axis deviation.

may be overworked with resultant evidence of this increased activity, although large P waves are not the usual finding in hypertension. Only two of the whole group of the forty-one miscellaneous cases, and in fact only two of the entire series of 132 cases showing high P waves, remain unexplained in the presence of definite or doubtful mitral stenosis, definite or doubtful pulmonic stenosis, other congenital heart defects, sino-auricular tachycardia, or the possible influence of hypertension.

PART II

When we came to undertake the second part of the study mentioned above, that is, an analysis of the P waves as to their amplitude and of the angles and indexes in all cases of definite pulmonic stenosis in which electrocardiograms were made at the Massachusetts General Hospital since 1914, we discovered, as has previously been mentioned, that as a result of their increase in the amplitude of P they were all covered by Table 1, and had therefore all been analyzed. Seventy-nine per cent. of all these cases showed an amplitude of P_2 equal to or greater than 3 mm. Seventy-nine per cent. had an amplitude of P_1 greater than 1 mm., and 100 per cent. had definite right axis deviation with angles of over 90 degrees and indexes of —10 or below. All cases had either P_1 over 1 mm. or P_2 over or equal to 3 mm.

PART III

Turning to the third study mentioned we examined the plates of 100 unselected cases that had been diagnosed clinically as pure mitral stenosis (Table 7).

It is evident from Table 7 that in this series, 45 per cent. of those showing P waves (that is, without fibrillation) had an amplitude of P_2 equal to or greater than 3 mm., and 78 per cent. had an amplitude of P_1 greater than 1 mm., whereas 84 per cent. had P_1 or P_2 equaling these high figures. Furthermore, one half of these cases showed the angles and indexes of right axis deviation. With regard to the angles in the entire series (with or without fibrillation) they were found to equal or exceed 90 degrees in 47 per cent. of all cases, while the indexes in the total series were equal to or less than —10 in 52 per cent. of the cases.

SUMMARY AND CONCLUSIONS

1. A study has been made of all the patients (132) who have shown by electrocardiogram during nine and one-half years at the Massachusetts General Hospital auricular or P waves greater than 0.1 millivolt (1 mm.) in amplitude in Lead I or equal to or greater than 0.3 millivolt (3 mm.) in Lead II. Of these cases nineteen (14 per cent.) had definite congenital pulmonic stenosis; three (2 per cent.) had doubtful congenital pulmonic stenosis; seventy-two (55 per cent.) had definite mitral stenosis; six (5 per cent.) had doubtful mitral stenosis; two (2 per cent.) had probable interventricular septal defect, and eighteen (14 per cent.) had sino-auricular tachycardia without question of mitral stenosis or congenital heart defects—a total of 120, or 91 per cent. of the 132 cases. Only twelve cases (9 per cent.) are not readily accounted for. Ten of these (8 per cent.) had hypertension with or without nephritis, not included in previous groups.

TABLE 7.—*Mitral Stenosis (Pure) **

Plate No.	Index, Mm.	Angle, Degrees	P ₁ , Mm.	P ₂ , Mm.
905.....	-15	120	3	3
1917.....	-11	69	2	2
2209.....	-17	102	2	2
2275.....	-15	130	1.3	2
1805.....	-8.5	81	1.5	2.7
3095.....	-13	95	1	4
3252.....	-19.5	120	2.2	2.2
2362.....	2	56	1.5	3.5
6653.....	-1	59	2	3
6685.....	-20	90	1	3
6675.....	-1	62	2	3.5
6801.....	-7	75	1.5	3.5
6865.....	-6	90	2.2	1.8
6959.....	-10	72	1.5	3
6948.....	5.5	25	2	4
8518.....	-12	98	2	5
7151.....	-5	69	1	4
6471.....	-16	117	2	4
7106.....	-6	76	1	5
6898.....	-20	112	1.5	5
1324.....	-13	121	1.6	2.5
2683.....	-10	78	1.5	3.2
6569.....	-20	122	2	2.5
1355.....	-21	122	2.5	4
7859.....	-3	64	Auricular fibrillation	
6620.....	-14	126	Auricular fibrillation	
1396.....	-19	105	3	2.5
6763.....	-10	133	Auricular fibrillation	
6064.....	-8	90	Auricular fibrillation	
6765.....	0	—	1.5	2
6818.....	-9	140	1.5	1.2
6884.....	-17	98	Auricular fibrillation	
6890.....	-2	70	Auricular fibrillation	
6906.....	-20	112	Auricular fibrillation	
7166.....	4	-35	Auricular fibrillation	
7169.....	10	10	1	2
6411.....	-17	90	Auricular fibrillation	
6430.....	-20	97	Auricular fibrillation	
6440.....	-11	127	Auricular fibrillation	
6945.....	-30	90	1	1
5311.....	-19	88	Auricular fibrillation	
6879.....	-23	102	1.2	2.5
6450.....	-24	121	Auricular fibrillation	
5608.....	-14	80	Auricular fibrillation	
5822.....	5	52	Auricular fibrillation	
5903.....	4	0	Auricular fibrillation	
3863.....	-20	87	Auricular fibrillation	
5119.....	-10	90	Auricular fibrillation	
5907A.....	-15	97	Auricular fibrillation	
5275.....	-15	109	Auricular fibrillation	
4685.....	0	58	Auricular fibrillation	
5545.....	-8	108	Auricular fibrillation	
6863.....	-5	55	Auricular fibrillation	
5792.....	4	40	Auricular fibrillation	
5907.....	-18	110	Auricular fibrillation	
3640.....	-15	115	Auricular fibrillation	
5849.....	-10	158	Auricular fibrillation	
6031.....	8	-15	Auricular fibrillation	
4183.....	2	-32	Auricular fibrillation	
4531.....	-16	115	Auricular fibrillation	
4853.....	-2	60	Auricular fibrillation	
4063.....	-11	60	Auricular fibrillation	
4376.....	4	-11	1	2
4200.....	-8	76	Auricular fibrillation	
4529.....	-2	68	Auricular fibrillation	
3780.....	-12	115	Auricular fibrillation	
4717.....	12	7	Auricular fibrillation	
4685.....	2	55	Auricular fibrillation	
4710.....	5	-17	1	2
3608.....	16	5	1	2
3716.....	3	46	1	2
5562.....	-32	122	2	2.5
8518.....	-12	98	3	5
6435.....	-9	68	2.5	4
6064.....	-8	90	Auricular fibrillation	
7057.....	-11	99	2.5	5
6308.....	-3	67	2	3

TABLE 7.—*Mitral Stenosis (Pure) *—Continued*

Plate No.	Index, Mm.	Angle, Degrees	P ₁ , Mm.	P ₂ , Mm.
6826.....	-23	98	2	2.5
6835.....	-13	99	2	3
6831.....	15	-18	1.5	3+
6774.....	-44	164	2	2
6935.....	-9	108	2	2.5
7104.....	-13	92	2	2.5
5023.....	0	60	1.5	2
6233.....	-1	62	2	3
4478.....	2	42	2	3
4501.....	-9	73	2	2
6525.....	-10	90	3	3
2159.....	8	0	Auricular fibrillation	
4491.....	5	47	Auricular fibrillation	
8160.....	-5	74	Auricular fibrillation	
2450.....	-16	82	Auricular fibrillation	
3339.....	-18	142	2	2
3370.....	9	-16	1	1
2308.....	4	52	1.5	2.2
3182.....	-10	80	1.2	3
1742.....	-15	86	1.5	2.5
3319.....	1	57	1	1
3252.....	-20	120	2	2.5
3287.....	-6	69	1	1.5

* All were patients in the Massachusetts General Hospital electrocardiographed from 1918 to January, 1924.

Summary: Total cases, 100, including forty-two cases of auricular fibrillation. Of the remaining fifty-eight, forty-five had P₁ greater than 1 mm. amplitude (78 per cent.) and thirty-nine had P₂ greater than 2 mm. Twenty-six had P₂ equal to or greater than 3 mm. (45 per cent.); forty-nine had an abnormally large P₁ or P₂ (84 per cent.). Of the total (100) cases, forty-seven had angles equal to or greater than +90 degrees; seventy-four had angles equal to or greater than +60 degrees. Fifty-two had indexes equal to or below -10; thirty-one had indexes equal to or below -15. Of the fifty-eight cases with normal rhythm the indexes and angles were as follows: twenty-six had indexes equal to or below -10; sixteen had indexes equal to or below -15; forty-three had angles equal to or greater than 60 degrees, and twenty-seven had angles equal to or greater than 90 degrees.

2. Of all the nineteen patients with congenital pulmonic stenosis electrocardiographed at the Massachusetts General Hospital from 1914 to 1924 (nine and one-half years) 79 per cent. had an abnormally large P wave in Lead I, greater than 1 mm. in amplitude; 84 per cent. had an abnormally large P wave in Lead II, equal to or greater than 3 mm. in amplitude; 68 per cent. had both, and 100 per cent. had one or the other. Also, all showed abnormal right axis deviation (right ventricular preponderance).

3. Of the 100 unselected consecutive patients with mitral stenosis electrocardiographed at the Massachusetts General Hospital fifty-eight had normal rhythm. Forty-two showed auricular fibrillation. Of the fifty-eight cases with normal rhythm forty-nine (84 per cent.) showed either an abnormally large P₁, P₂ or both.

4. An amplitude of P₁ greater than 1 mm., of P₂ equal to or greater than 3 mm., an electrical axis of 90 degrees or more, and an axis index of -10 or less are consistently found in congenital pulmonic stenosis.

5. A P₁ consistently greater in amplitude than 1 mm. or a P₂ equal to or greater than 3 mm. points to the possibility of an existent mitral stenosis; if both of these conditions are present there is greater likelihood of the condition; and if either of these findings is accompanied by right axis deviation, in the absence of pulmonic stenosis, the diagnosis of mitral stenosis is probable.