Published in final edited form as:

Arch Womens Ment Health. 2017 June ; 20(3): 363–372. doi:10.1007/s00737-016-0707-4.

Treatment options for hyperemesis gravidarum

Amy Abramowitz¹, Emily S. Miller², Katherine L. Wisner³

¹UIC Department of Psychiatry, University of Illinois at Chicago, 912 S. Wood Street, Chicago, IL 60612, USA

²Department of Obstetrics and Gynecology, Division of Maternal Fetal Medicine, Department of Psychiatry and Behavioral Sciences, Northwestern University Feinberg School of Medicine, 303 E. Chicago Ave, Chicago, IL 60611, USA

³Department of Obstetrics and Gynecology, Division of Maternal Fetal Medicine, Department of Psychiatry and Behavioral Sciences, Northwestern University Feinberg School of Medicine, 303 E. Chicago Ave, Chicago, IL 60611, USA

Abstract

Hyperemesis gravidarum (HG) is a severe and prolonged form of nausea and/or vomiting during pregnancy. HG affects 0.3–2% of pregnancies and is defined by dehydration, ketonuria, and more than 5% body weight loss. Initial pharmacologic treatment for HG includes a combination of doxylamine and pyridoxine. Additional interventions include ondansetron or dopamine antagonists such as metoclopramide or promethazine. The options are limited for women who are not adequately treated with these medications. We suggest that mirtazapine is a useful drug in this context and its efficacy has been described in case studies. Mirtazapine acts on noradrenergic, serotonergic, histaminergic, and muscarinic receptors to produce antidepressant, anxiolytic, antiemetic, sedative, and appetite-stimulating effects. Mirtazapine is not associated with an independent increased risk of birth defects. Further investigation of mirtazapine as a treatment for HG holds promise to expand treatment options for women suffering from HG.

Keywords

Hyperemesis gravidarum; Mirtazapine; Gastrointestinal disorders in pregnancy; Nausea and vomiting in pregnancy; Hyperemesis

Introduction

During pregnancy, up to 80% of women experience nausea and/or vomiting (N/V) (Gazmararian et al. 2002) (Gadsby et al. 1993). For the majority of women, N/V resolves by the 16th week of pregnancy (Eliakim et al. 2000) (Gadsby et al. 1993). A severe and prolonged form of N/V, hyperemesis gravidarum (HG), affects 0.3–2% of pregnancies (Hod

Amy Abramowitz, abramowitz@gmail.com.

Conflict of interest The Department of Psychiatry at Northwestern University received contractual fees for Dr. Wisner's consultation to Quinn Emanuel Urquhart & Sullivan, LLP (New York City), who represent Pfizer Pharmaceutical Company, in 2015. Dr. Miller and Dr. Abramowitz declare that they have no conflict of interest.

et al. 1994). The criteria that define HG include dehydration, ketonuria, and more than 5% weight loss (Nelson-Piercy 1998) (McCarthy et al. 2014). HG is the most common indication for hospital admission in the first 20 weeks of pregnancy (Klebanoff et al. 1985) (Eliakim et al. 2000). Although it usually occurs between weeks 4 and 9 of pregnancy and resolves by mid-gestation (Eliakim et al. 2000), between 15 and 20% of women continue to experience symptoms until the third trimester and 5% through delivery (Goodwin 2008) (Fell et al. 2006).

Women with HG often present with increased blood urea nitrogen and hematocrit and, in 15–25% of cases, hyponatremia, hypokalemia, and hypochloremia (Goodwin 1998). Electrolyte abnormalities are corrected with intravenous replacement fluids; however, in severe cases, sustained weight loss requires enteral or total parenteral nutrition (Ayyavoo et al. 2014; Levine and Esser 1988). Between 15 and 50% of patients with HG have elevated serum aminotransferases and total bilirubin (Morali and Braverman 1990) (Wallstedt et al. 1990).

HG is a clinical diagnosis; however, tools such as the Pregnancy Unique Quantification of Emesis (PUQE) score or Rhodes index have been used to assess the severity of symptoms in research studies (Koren et al. 2002; Rhodes and McDaniel 1999). HG is a diagnosis of exclusion, and alternative diagnoses should be considered when the N/V begins after 9 weeks of gestation or if bilious emesis, fever, abdominal pain, headache, focal neurological findings, leukocytosis, or hypertension are present. Comorbid conditions, most commonly reflux, should also be evaluated and treated (Gideon et al. 2009). Women presenting with HG are evaluated with serial measures of maternal weight, orthostatic blood pressures, heart rate, electrolytes, urine ketones, and an obstetric ultrasound if not previously obtained (2015). The differential diagnosis for HG is provided in Table 1.

Women of African-American, Indian, or Pakistani descent have higher rates of HG compared to European women (Klebanoff et al. 1985) (Price et al. 1996). Primiparous women; adolescents; women with increased BMI, multiple gestations, and gestational trophoblastic disease; and women with HG during a previous pregnancy also have increased risk for HG (Klebanoff et al. 1985) (Depue et al. 1987; Eliakim et al. 2000). The risk of recurrent HG is 15–20% (Dodds et al. 2006; Trostad et al. 2005). Women with a sister or mother who had HG are also at increased risk, which suggests that genetic factors play a role (Zhang et al. 2011) (Fejzo et al. 2008). Female fetuses are more commonly born to women with HG (Veenendaal et al. 2011). Additionally, smoking appears to be a protective factor (Fell et al. 2006; Källén et al. 2003; Vikanes et al. 2010), although the mechanism has not been elucidated. Taking a multivitamin in early pregnancy or prior to conception was associated with decreased N/V (Czeizel et al. 1992; Emelianova et al. 1999; Källén et al. 2003).

Pathogenesis

Hormonal factors have been evaluated as etiologic contributors to HG. Several investigators have reported an increased level of free β -hCG in women with HG compared to matched controls (Goodwin et al. 1994). Additionally, hCG peaks in the first trimester when N/V is

typically the most severe. HG is also more common in conditions characterized by high hCG levels such as multiple gestations and gestational trophoblastic diseases (Goodwin et al. 1992; Kimura et al. 1993).

There is homology between the beta-subunit of hCG and TSH which causes hCG to have thyroid-stimulating activity (Ballabio et al. 1991). Initial studies found an increased free thyroxine in 40 and 73% of women with HG (Boullion et al. 1982) (Bober et al. 1986); however, these findings were not confirmed in another investigation (Wilson et al. 1992). Patients with hyperthyroidism rarely experience N/V, which suggests that thyrotoxicosis is an unlikely etiology of HG.

Additionally, total estradiol and sex hormone binding globulin were 26 and 37% higher, respectively, in women with HG (Depue et al. 1987). As progesterone increases throughout gestation, it is associated with decreased esophageal and gastric motility as well as lower esophageal sphincter pressure. However, progesterone concentrations are most pronounced in the third trimester, not early in pregnancy when HG is most common (Lagiou et al. 2003; Van Thiel et al. 1977).

Several systematic reviews and meta-analyses have shown a significant correlation between *Helicobacter pylori* infection and N/V of pregnancy (Golberg et al. 2007; Li et al. 2015; Niemeijer et al. 2014; Sandven et al. 2009). Treatment of *H. pylori* has been shown to improve N/V in infected women (El Younis et al. 1998; Jacoby and Porter 1999); however, the overall sensitivity and specificity for screening for *H. pylori* in women presenting with HG were only 73 and 55% respectively (Niemeijer et al. 2014).

Historically, HG was thought to be a response to stress or ambivalence about pregnancy (Buckwalter and Simpson 2002); however, no differences in psychological or personality characteristics between women with and without HG have been demonstrated (Orazio et al. 2011; Sheehan 2007; Simpson et al. 2001). An analysis of insurance data of 11,016 women who gave birth found that a pre-pregnancy psychiatric diagnosis doubled the risk of developing HG and pre-pregnancy psychiatric and somatic conditions quadrupled the risk, which suggest that these disorders, or a common etiologic factor, may play a role in the development of HG in some women (Seng et al. 2007). Women with HG have been found to have significantly higher levels of plasma serotonin compared to women with N/V of pregnancy and pregnant controls (Cengiz et al. 2015).

Maternal, fetal, and child outcomes associated with HG

Maternal outcomes

Women with HG are at risk for complications from excessive vomiting including hematemesis and from dehydration including dizziness and syncope (Mullin et al. 2012). More severe complications include Wernicke's encephalopathy, central pontine myelinolysis, and peripheral neuropathy due to vitamin B6 or B12 deficiency (Gardian et al. 1999) (Peeters et al. 1993) (Goodwin 1998).

Women with HG are at increased risk for developing secondary depression and anxiety (McCarthy et al. 2011) (Hizli et al. 2012) (Annagür et al. 2014) (Tan et al. 2010b) which typically resolve with remission of N/V by the third trimester of pregnancy (Tan et al. 2014). In a survey of women's experiences with HG, 15.2% women had voluntarily terminated at least one pregnancy (Poursharif et al. 2007).

Fetal outcomes

Decreased rates of early pregnancy loss in women with HG (2000; Bashiri et al. 1995; Hinkle et al. 2016; Weigel and Weigel 1989) have been reported. One speculation is that N/V eliminates harmful substances from food or alters hormones that facilitate placental development, thereby improving embryologic health (Huxley 2000; Nulman et al. 2009; Sherman and Flaxman 2002).

A Norwegian cohort study of 2,270,363 births found a decreased risk of very preterm birth (<32 weeks gestation; odds ratio 0.79, 95% CI, 0.67–0.93) born to women with HG (Vandraas et al. 2013). HG was associated with slight reductions in 21.4 g and gestational length (0.5 days) compared with unexposed infants (Vandraas et al. 2013). HG has been associated with increased rates of low birth weight and small for gestational age infants (Bailit 2005) (Dodds et al. 2006) (McCarthy et al. 2011; Veenendaal et al. 2011) whereas other studies have not confirmed these findings (Vikanes et al. 2013) (Depue et al. 1987) (Tsang et al. 1996). Further, some studies found that stratification by severity of HG (defined by losing greater than 5% of body weight) demonstrated that women with severe HG were more likely to have babies with smaller birth weight and in the <10th percentile at birth compared to women who maintained 95% of their body weight (Gross et al. 1989), although other authors have not replicated this finding (Hallak et al. 1996).

No association was found between HG and decreased Apgar scores, or perinatal mortality (Veenendaal et al. 2011). Boneva et al. (1999) observed lower rates of cardiac malformations in women with severe N/V (Boneva et al. 1999); however, an association between HG and other anomalies has not been found (Veenendaal et al. 2011).

Child outcomes

Few data have been published on long-term outcomes for children born to mothers with HG. One study showed 20% lower insulin sensitivity in pre-pubertal children of mothers with severe HG compared to children with mothers without HG (Ayyavoo et al. 2013). A retrospective case control study of adults showed an increase in psychological and behavioral disorders in a composite mental health outcome measure, but individual analyses did not show increases in depression, anxiety, or bipolar disorders (Mullin et al. 2011).

Treatment

Most women who experience N/V are counseled on the self-limited course of symptoms and on avoiding foods, odors, or activities that exacerbate symptoms (Davis 2004; Matthews et al. 2014). Eating small meals frequently (Bischoff and Renzer 2006) and snacks of carbohydrate-rich foods, such as soda crackers, may reduce nausea (Jednak et al. 1999).

Indications for hospital admission include significant weight loss, electrolyte abnormalities, and severe or persistent vomiting after rehydration. Inpatient treatment includes intravenous hydration with appropriate electrolyte replacement. Symptoms typically improve within 1 to 2 days of rehydration (Summers 2012).

Alternative therapies

Non-pharmaceutical approaches to treating N/V include herbs, such as ginger and chamomile, acupuncture, and massage (Davis 2004; Murphy 1998; Niebyl and Goodwin 2002). Capsules of ginger decreased episodes of N/V within 1 week compared to placebo tablets (Keating and Chez 2002; Nasrin et al. 2011; Ozgoli et al. 2009). A systematic review and meta-analysis found that ginger reduced nausea but not vomiting when compared to placebo (Viljoen et al. 2014).

Treatment with acupuncture was associated with decreased severity of N/V compared to placebo in one study (Aghadam and Mahfoozi 2010). However, three other studies have not confirmed improvement associated with acupuncture compared to placebo (Belluomini et al. 1994; Can Gürkan and Arslan 2008; Norheim et al. 2001).

Compared to placebo, pyridoxine or vitamin B6 supplementation reduced nausea but not vomiting (Sahakian et al. 1991; Vutyavanich et al. 1995). Vitamin B6 supplementation resulted in similar reductions in nausea and vomiting compared to ginger and acupuncture (Chittumma et al. 2007; Ensiyeh and Sakineh 2009; Jamigorn and Phupong 2007; Smith et al. 2004; Sripramote and Lekhyananda 2003). The mechanism of therapeutic effect is unclear and no correlation between vitamin B6 level and severity of nausea was observed (Schuster et al. 1985). Vitamin B6 has minimal side effects and is not associated with fetal malformations (ACOG 2015; Shrim et al. 2006).

Psychotherapeutic interventions

There are few studies examining psychotherapy treatment for HG, including no randomized trials (2015). However, there are several case studies reporting the efficacy of hypnotherapy (Fuchs et al. 1980; McCormack 2010; Simon and Schwartz 1999). Psychological support from family and the medical team has been shown to reduce symptoms of HG (Faramarzi et al. 2015; Liu et al. 2014; Tamay and Kusu 2011).

Pharmacologic therapies

Antihistamines (H1 antagonists), including doxylamine used in combination with pyridoxine, meclizine, dimenhydrinate, and diphenhydramine, are effective in reducing N/V. None of these agents has been associated with fetal malformations (Magee et al. 2002; Seto et al. 1997).

The combination of doxylamine and pyridoxine (Diclegis®) is the only FDA-indicated agent for the treatment of N/V in pregnancy. This formulation, previously sold under the name Bendectin®, was voluntarily withdrawn from the market in 1983 due to alleged teratogenic effects, although multiple studies demonstrated both its efficacy (Geiger et al. 1959; Koren

et al. 2010; Magee et al. 2002; Maltepe and Koren 2013b; Wheatley 1977) and safety (Brent 1995; Einarson et al. 1988; Kutcher et al. 2003; McKeigue et al. 1994; Neutel and Johansen 1995). After Bendectin® was withdrawn from the US market, a threefold increase in hospitalizations of women with HG was observed (Neutel and Johansen 1995). Diclegis® received FDA approval in 2013 with a Pregnancy Category A safety classification (Slaughter et al. 2014). Diclegis® is considered first-line pharmacotherapy for N/V in pregnancy by the American Professors in Gynecology and Obstetrics and the American College of Obstetricians and Gynecologists (2015). However, 97.7% of prescriptions for N/V during pregnancy in the USA are for medications other than Diclegis®, the only drug with an FDA indication for treatment (Koren 2014).

No adverse effects on cognitive development in children ages 3 to 7 with mothers who had HG both with and without treatment with pyridoxine/doxylamine were observed (Nulman et al. 2009), although another larger study of 8-year-old children of mothers with HG, regardless of treatment received, showed significantly increased rates of neurodevelopmental delay (Fejzo et al. 2015).

Ondansetron, a serotonin antagonist acting at the 5-HT₃ receptor, is the most commonly prescribed medication for the treatment of N/V during pregnancy, and its use is rapidly increasing (Koren 2014). In 2008, the rate of prescriptions per month was 50,000 and in 2013, 110,000 per month. Ondansetron is effective in reducing N/V during pregnancy, and a double-blind randomized controlled study of 36 women found that ondansetron was significantly more effective than combined pyridoxine and doxylamine treatment in reducing nausea and vomiting (Oliveira et al. 2014). Side effects include headache; fatigue; constipation; QT prolongation; and, rarely, serotonin syndrome (Freedman et al. 2013).

The safety of ondansetron during pregnancy was assessed in a Danish study of 1970 exposed infants, which did not show an increased risk of fetal malformations or adverse pregnancy outcomes (Pasternak et al. 2013). However, other studies have found that ondansetron was associated with birth defects. A Swedish cohort with 1349 exposed infants found a significantly increased risk of cardiac septal defects (Danielsson et al. 2014), and a US cohort found an increased risk of cleft palate (Anderka et al. 2012). Additionally, a second Danish study using the same registry but including more infants over more years found a two-fold increase in cardiac malformations (Andersen et al. 2013).

Dopamine antagonists, such as metoclopramide and promethazine and droperidol, have been used to treat HG (Fadi et al. 2003). Metoclopramide was found to be more effective at reducing N/V than other dopamine antagonists. For example, intravenous metoclopramide with diphenhydramine was more effective at reducing vomiting compared to droperidol with diphenhydramine (Lacasse et al. 2009). Also, metoclopramide with pyridoxine was also more effective than were other dopamine antagonists, prochlorperazine and promethazine (Fadi et al. 2003). When metoclopramide was compared to ondansetron, they were comparable in reducing nausea, although ondansetron had a greater effect on reducing vomiting (Kashifard et al. 2013). Both medications resulted in similar improvement in N/V, although metoclopramide was associated with sedation, dry mouth, and dystonias (Abas et

al. 2014; Pasricha et al. 2006). Cohort studies have shown that metoclopramide does not increase the risk of fetal malformations (Matok et al. 2009; Pasternak et al. 2014).

Promethazine is primarily an antihistaminergic medication, and it also acts as a weak dopamine antagonist. It is effective in relieving N/V in pregnancy but has significant maternal side effects including sedation, dystonia, and decreased seizure threshold (Braude and Crandall 2008; Fitzgerald 1955; Magee et al. 2002; Seto et al. 1997; Tan et al. 2010a). Droperidol was studied in combination with diphenhydramine and was found to reduce days in the hospital for HG (Nageotte et al. 1996). It has not been associated with fetal malformations, though it is associated with QTc prolongation in some women (Jackson et al. 2007).

Treatments for refractory HG

Many women do not experience resolution of their symptoms with existing treatments. Studies demonstrating the efficacy of ondansetron, for example, show that most women continue to have at least one episode of vomiting per day even after initiating therapy (Abas et al. 2014). The impairment associated with HG is demonstrated by women who pursue elective abortion due to the severity of their symptoms in otherwise desired pregnancies (Maltepe and Koren 2013a; Mazzota et al. 1997). One third of women surveyed with HG chose not to become pregnant again because of their experience with HG (Fejzo et al. 2011). In patients refractory to these treatments, several other options have been studied.

Glucocorticoids did not reduce rates of re-hospitalization when compared to placebo (Yost et al. 2003). Furthermore, they have been associated with a possible increased risk of oral cleft when used in the early first trimester (Carmichael and Shaw 1999; Park-Wyllie et al. 2000; Pradat et al. 2003; Shepard et al. 2002). Other therapeutic options for refractory cases include transdermal clonidine to reduce symptoms in women who cannot tolerate oral therapies. A randomized placebo-controlled clonidine trial with 13 patients showed significant decreases in symptoms and reduced need for enteral or parenteral nutrition (Maina et al. 2014). A study of 70 patients hospitalized with HG showed reduced rehospitalizations with diazepam and intravenous fluid compared to intravenous fluid alone (Tasci et al. 2009). Additionally, several case reports have suggested the efficacy of gabapentin (Guttuso et al. 2010; Spiegel and Webb 2012).

Women who do not respond to any of these interventions and continue to lose weight should be supported with enteral or parenteral nutrition in addition to any medication that provides improvement in symptoms (Stokke et al. 2015). Enteral nutrition provides more relief from N/V compared to parenteral nutrition (Hsu et al. 1996). Some women decline pharmacologic treatments due to concern about the risk of congenital defects associated with medications taken during early pregnancy.

Mirtazapine

We have included the antidepressant drug mirtazapine in this review to suggest consideration of its use for women who have not responded to other therapies. This drug is not included in recommendations for the treatment of N/V or HG from obstetrical publications (ACOG

2015). However, mirtazapine is used to treat patients undergoing cancer treatment with rapid onset of efficacy for nausea (day 1) and sleep (within the first 5 days) (Cardona 2006; Dupuis and Nathan 2010; Kim et al. 2008). Its pharmacologic profile of actions is similar to other drugs used to treat HG.

Pharmacology

Mirtazapine acts on noradrenergic, serotonergic, histaminergic, and muscarinic receptors. It is an indirect agonist of the 5-HT_{1A} receptor, an antagonist at the 5-HT₂ receptors, and an inverse agonist at the 5-HT_{2C} receptor to produce the antidepressant effect (Blier and Abbott 2001). Antagonism of the 5-HT₂ receptor also contributes to the anxiolytic, sedative, and appetite-stimulating effects (Fawcett and Barkin 1998). It is an inverse agonist on the H₁ receptor which contributes to the sedative effect. The antiemetic properties come from antagonism of the 5-HT₃ receptor, which is the same receptor that ondansetron affects (Anttila and Leinonen 2001). Mirtazapine exerts its antiemetic effect regardless of whether the patient has a psychiatric disorder, though a patient with depression or anxiety may benefit from both of these actions.

Mirtazapine's side effects were rated as average compared to other antidepressants and commonly include sedation, weight gain, dry mouth, and constipation (Cipriani et al. 2009). Similar to other antidepressants, patients should undergo screening for bipolar disorder using an instrument such as the Mood Disorder Questionnaire (Hirschfeld et al. 2000) before starting mirtazapine and educated about the rare possibility of anxiety, agitation, and suicidal ideation at the beginning of treatment.

Mirtazapine is metabolized in the liver and its elimination half-life is between 20 and 40 h. Mirtazapine is a weak inhibitor of CYP-isoenzymes and is unlikely to cause many clinically relevant drug interactions (Anttila and Leinonen 2001). Mirtazapine has been associated with serotonin syndrome in multiple case reports, typically when multiple serotonergic drugs are combined (Ansermot et al. 2014). Ondansetron and metoclopramide have also been associated with serotonin syndrome when used as monotherapy and when combined with other serotonergic agents. Though no case reports of serotonin syndrome have been found with the combination of mirtazapine and metoclopramide or ondansetron, it remains a theoretical risk. Signs of serotonin syndrome are diarrhea, diaphoresis, fever, hyperreflexia, confusion, ataxia, myoclonus, tremor, and rhabdomyolysis (Ables and Nagubilli 2010).

Winterfeld and colleagues (Winterfeld et al. 2015) conducted a prospective cohort study comparing birth defects in infants of women treated antenatally with mirtazapine to two comparison groups, each with 357 subjects. The comparison groups were (1) mothers treated with selective serotonin reuptake inhibitor antidepressants and (2) a general population of women without antidepressant treatment or exposure to known teratogenic agents. No statistically significant difference in the rate of major birth defects after first-trimester exposure between mirtazapine, SSRI-exposed, and non-exposed pregnancies was found. A marginally higher rate of birth defects was observed in the mirtazapine and SSRI groups compared with the low rate of birth defects in the general control subjects, which is consistent with other studies that evaluated the risk factors related to the underlying

depressive disorder (Huybrechts et al. 2014). Rates of preterm birth did not significantly differ across groups.

We present case series of mirtazapine for HG to summarize the published clinical data. A total of 15 reports of successful treatment of HG with mirtazapine have been published. Six women requested termination of their pregnancy due to the severity of their HG symptoms. Seven patients presented with symptoms of depression and five reported a history of depression. These patients had previously failed other treatments for HG, including patients who did not respond to the following drugs: metoclopramide ($n = 8$), ondansetron ($n = 5$), promethazine ($n = 4$), dimenhydrinate ($n = 2$), and pyridoxine ($n = 1$). Patients began treatment with mirtazapine between gestational weeks 6 and 25. Eight of the patients were treated with 15 mg/day of mirtazapine, four with 30 mg/day, and two with 45 mg/day. In eight patients for whom the time to resolution was reported, all responded within 1 week and five responded in 3 days. Mirtazapine was continued for between 6 days and 22 weeks. Delivery of healthy babies was reported in 12 of the patients and pregnancy outcomes were not reported in the other 3 cases (Dorn et al. 2002; Guclu et al. 2005; Rohde et al. 2003; Saks 2001; Schwarzer et al. 2008; Uguz, 2014).

Future directions

In a recent Cochrane review, Boelig, et al. (2016) concluded that little evidence existed to support the superiority of one intervention over another in the treatment of HG, and the authors recommended research comparing the side-effect profiles, safety, economic costs, and benefits of treatments to inform selection. Comparative evaluation of current treatments and expansion of therapies for severe or refractory HG, which has been associated with requests for termination of desired pregnancies, is imperative. Mirtazapine has highly promising but efficacy and side effect data are limited to multiple case studies. Mirtazapine may expand the options for treating HG, especially in women who are also experiencing psychiatric symptoms, such as depression and anxiety. Such studies should determine characteristics of women who would benefit most from each treatment option to personalize care.

References

- Abas MN, Tan PC, Azmi N, Omar SZ (2014) Ondansetron compared with metoclopramide for hyperemesis gravidarum: a randomized controlled trial. *Obstet Gynecol* 123(6):1272 [PubMed: 24807340]
- Ables AZ, Nagubilli R (2010) Prevention, recognition, and management of serotonin syndrome. *Am Fam Physician* 81(9):1139–1142 [PubMed: 20433130]
- ACOG (2015) Practice bulletin 153: nausea and vomiting of pregnancy. *Obstet Gynecol* 126:687–688 [PubMed: 26287781]
- Aghadam SKZ, Mahfoozi B (2010) Evaluation of the effects of acupressure by sea band on nausea and vomiting of pregnancy. *Iranian Journal of Obstetrics, Gynecology and Infertility* 13(2):39
- Anderka M, Mitchell AA, Louik C, Werler MM, Hernández-Díaz S, Rasmussen SA (2012) Medications used to treat nausea and vomiting of pregnancy and the risk of selected birth defects. Birth defects research. Part A, *Clinical And Molecular Teratology* 94(1):22
- Andersen J, Jimenez-Solem E, Andersen N, Poulsen H (2013) Ondansetron use in early pregnancy and the risk of congenital malformations—a register based Nationwide cohort study. *Pharmacoepidemiol Drug Saf* 22:13

- Annagür BB, Kerimo lu ÖS, Gündüz , Tazegül A (2014) Are there any differences in psychiatric symptoms and eating attitudes between pregnant women with hyperemesis gravidarum and healthy pregnant women? *J Obstet Gynaecol Res* 40(4):1009 [PubMed: 24320704]
- Ansermot N, Hodel PF, Eap CB (2014) Serotonin toxicity after addition of mirtazapine to escitalopram. *J Clin Psychopharmacol* 34(4):540–541 [PubMed: 24977717]
- Anttila SAK, Leinonen EVJ (2001) A review of the pharmacological and clinical profile of mirtazapine. *CNS Drug Reviews* 7(3):249–264 [PubMed: 11607047]
- Ayyavoo A, Derraik JGB, Hofman PL, Biggs J, Bloomfield FH, Cormack BE, Stone P, Cutfield WS (2013) Severe hyperemesis gravidarum is associated with reduced insulin sensitivity in the offspring in child-hood. *J Clin Endocrinol Metab* 98(8):3263 [PubMed: 23750032]
- Ayyavoo A, Derraik JGB, Hofman PL, Cutfield WS (2014) Hyperemesis gravidarum and long-term health of the offspring. *Am J Obstet Gynecol* 210(6):521 [PubMed: 24280249]
- Bailit JL (2005) Hyperemesis gravidarium: epidemiologic findings from a large cohort. *Am J Obstet Gynecol* 193(3):811 [PubMed: 16150279]
- Ballabio M, Poshychinda M, Ekins RP (1991) Pregnancy-induced changes in thyroid function: role of human chorionic gonadotropin as putative regulator of maternal thyroid. *J Clin Endocrinol Metab* 73(4):824 [PubMed: 1909707]
- Bashiri A, Newmann L, Maymon E, Katz M (1995) Hyperemesis gravidarum: epidemiologic features, complications and outcome. *Eur J Obstet Gynecol Reprod Biol* 63:135–138 [PubMed: 8903768]
- Belluomini J, Litt RC, Lee KA, Katz M (1994) Acupressure for nausea and vomiting of pregnancy: a randomized, blinded study. *Obstet Gynecol* 84(2):245 [PubMed: 8041539]
- Bischoff SC, Renzer C (2006) Nausea and nutrition. *Autonomic Neuroscience: Basic And Clinical* 129(1):22 [PubMed: 16935033]
- Blier P, Abbott FV (2001) Putative mechanisms of action of antidepressant drugs in affective and anxiety disorders and pain. *Journal Of Psychiatry & Neuroscience : Jpn* 26(1):37
- Bober SA, McGill AC, Tunbridge WM (1986) The thyroid function in hyperemesis gravidarum. *Acta Endocrinol* 111:404–410 [PubMed: 3083627]
- Boelig Rupsa C., Barton Samantha J., Saccone Gabriele, Kelly Anthony J., Edwards Steve J., and Berghella Vincenzo (2016) Interventions for treating hyperemesis gravidarum.” *Cochrane Database of Systematic Reviews*
- Boneva RS, Moore CA, Botto L, Wong LY, Erickson JD (1999) Nausea during pregnancy and congenital heart defects: a population-based case–control study. *Am J Epidemiol* 149(8):717 [PubMed: 10206621]
- Boullion R, Naesens M, van Assche FA (1982) Thyroid function in patients with hyperemesis gravidarum. *Am J Obstet Gynecol* 143: 922–926 [PubMed: 7102767]
- Braude D, Crandall C (2008) Ondansetron versus promethazine to treat acute undifferentiated nausea in the emergency department: a randomized, double-blind, noninferiority trial. *Acad Emerg Med Off J Soc Acad Emerg Med* 15(3):209
- Brent RL (1995) Bendectin: review of the medical literature of a comprehensively studied human nonteratogen and the most prevalent tortogen-litigen. *Reprod Toxicol* 9(4):337 [PubMed: 7496090]
- Buckwalter JG, Simpson SW (2002) Psychological factors in the etiology and treatment of severe nausea and vomiting in pregnancy. *Am J Obstet Gynecol* 186(5):S210 [PubMed: 12011888]
- Can Gürkan Ö, Arslan H (2008) Effect of acupressure on nausea and vomiting during pregnancy. *Complement Ther Clin Pract* 14(1):46 [PubMed: 18243942]
- Cardona D (2006) Pharmacological therapy of cancer anorexia-cachexia. *Nutricion hospitalaria* 21(Suppl 3):17–26 [PubMed: 16768027]
- Carmichael SL, Shaw GM (1999) Maternal corticosteroid use and risk of selected congenital anomalies. *Am J Med Genet* 86(3):242 [PubMed: 10482873]
- Cengiz H, Dagdeviren H, Caypinar S, Kanawati A, Yildiz S, Ekin M (2015) Plasma serotonin levels are elevated in pregnant women with hyperemesis gravidarum. *Arch Gynecol Obstet* 291(6):1271 [PubMed: 25502185]

- Chittumma P, Kaewkiattikun K, Wiriyasiriwach B (2007) Comparison of the effectiveness of ginger and vitamin B6 for treatment of nausea and vomiting in early pregnancy: a randomized double-blind controlled trial. *Journal Of The Medical Association Of Thailand = Chotmaihet Thangphaet* 90(1):15 [PubMed: 17621727]
- Cipriani A, Furukawa TA, Salanti G, Geddes JR, Higgins JPT, Churchill R, Watanabe N, Nakagawa A, Omori IM, McGuire H, Tansella M, Barbui C (2009) Comparative efficacy and acceptability of 12 new-generation antidepressants: a multiple-treatments meta-analysis. *Lancet* 373(9665):746–758 [PubMed: 19185342]
- Czeizel A, Dudas I, Fritz G, Técsői A, Hanck A, Kunovits G (1992) The effect of periconceptional multivitamin-mineral supplementation on vertigo, nausea and vomiting in the first trimester of pregnancy. *Arch Gynecol Obstet* 251(4):181 [PubMed: 1503509]
- Danielsson B, Wikner BN, Källén B (2014) Use of ondansetron during pregnancy and congenital malformations in the infant. *Reprod Toxicol* 50:134 [PubMed: 25450422]
- Davis M (2004) Nausea and vomiting of—an evidence-based review. *Journal Of Perinatal & Neonatal Nursing* 18(4):312
- Depue RH, Bernstein L, Ross RK, Judd HL, Henderson BE (1987) Hyperemesis gravidarum in relation to estradiol levels, pregnancy outcome, and other maternal factors: a seroepidemiology study. *Am J Obstet Gynecol* 156:1137–1141 [PubMed: 3578425]
- Dodds L, Fell DB, Joseph KS, Allen VM, Butler B (2006) Outcomes of pregnancies complicated by hyperemesis gravidarum. *Obstet Gynecol* 107(2 Pt 1):285 [PubMed: 16449113]
- Dorn C, Pantlen A, Rohde A (2002) Mirtazapine (Remergil (R)/Remeron (R)): treatment option of therapy resistant hyperemesis gravidarum? A case report. *Geburtshilfe Frauenheilkd* 62(7):677
- Dupuis LL, Nathan PC (2010) Optimizing emetic control in children receiving antineoplastic therapy: beyond the guidelines. *Paediatric drugs* 12(1):51–61 [PubMed: 20034341]
- Einarson TR, Leeder JS, Koren G (1988) A method for meta-analysis of epidemiological studies. *Drug Intelligence & Clinical Pharmacy* 22(10):813
- El Younis CM, Abulafia O, Sherer DM (1998) Rapid marked response of severe hyperemesis gravidarum to oral erythromycin. *Am J Perinatol* 15(9):533 [PubMed: 9890250]
- Eliakim R, Abulafia O, Sherer DM (2000) Hyperemesis gravidarum: a current review. *Am J Perinatol* 17(4):207 [PubMed: 11041443]
- Emelianova S, Mazzotta P, Einarson A, Koren G (1999) Prevalence and severity of nausea and vomiting of pregnancy and effect of vitamin supplementation. *Clinical and investigative medicine. Médecine Clinique Et Experimentale* 22(3):106 [PubMed: 10410832]
- Ensiyeh J, Sakineh M-AC (2009) Comparing ginger and vitamin B6 for the treatment of nausea and vomiting in pregnancy: a randomized controlled trial. *Midwifery* 25(6):649 [PubMed: 18272271]
- Fadi AB, Despina EH, David ES (2003) Comparison of three outpatient regimens in the Management of Nausea and Vomiting in pregnancy. *J Perinatol* 23(7):531 [PubMed: 14566347]
- Faramarzi M, Yazdani S, Barat S (2015) A RCT of psychotherapy in women with nausea and vomiting of pregnancy. *Hum Reprod* 30(12):2764–2773 [PubMed: 26466913]
- Fawcett J, Barkin RL (1998) Review of the results from clinical studies on the efficacy, safety and tolerability of mirtazapine for the treatment of patients with major depression. *J Affect Disord* 51(3):267 [PubMed: 10333982]
- Fejzo MS, Ingles SA, Wilson M, Wang W, MacGibbon K, Romero R, Goodwin TM (2008) High prevalence of severe nausea and vomiting of pregnancy and hyperemesis gravidarum among relatives of affected individuals. *European Journal Of Obstetrics And Gynecology* 141(1):13
- Fejzo MS, MacGibbon KW, Romero R, Goodwin TM, Mullin PM (2011) Recurrence risk of hyperemesis gravidarum. *The Journal of Midwifery & Women's Health* 56(2):132–136
- Fejzo MS, Magtira A, Schoenberg FP, Macgibbon K, Mullin PM (2015) Neurodevelopmental delay in children exposed in utero to hyperemesis gravidarum. *European Journal of Obstetrics and Gynecology* 189:79
- Fell DB, Dodds L, Joseph KS, Allen VM, Butler B (2006) Risk factors for hyperemesis gravidarum requiring hospital admission during pregnancy. *Obstet Gynecol* 107(2 Pt 1):277 [PubMed: 16449112]

- Fitzgerald JP (1955) The effect of promethazine in nausea and vomiting of pregnancy. *N Z Med J* 54(300):215 [PubMed: 14384015]
- Freedman SB, Uleryk E, Rumantir M, Finkelstein Y (2013) Ondansetron and the risk of cardiac arrhythmias: a systematic review and postmarketing analysis. *Ann Emerg Med* 64(1):19 [PubMed: 24314899]
- Fuchs K, Paldi E, Abramovici H, Peretz BA (1980) Treatment of hyperemesis gravidarum by hypnosis. *Int J Clin Exp Hypn* 28(4):313 [PubMed: 7409926]
- Gadsby R, Barnie-Adshead AM, Jagger C (1993) A prospective study of nausea and vomiting during pregnancy. *British Journal Of General Practice : The Journal Of The Royal College Of General Practitioners* 43(371):245
- Gardian G, Voros E, Jardanhazy T, Ungurean A, Vecsei L (1999) Wernicke's encephalopathy induced by hyperemesis gravidarum. *Acta Neurol Scand* 99:196–198 [PubMed: 10100965]
- Gazmararian JA, Petersen R, Jamieson DJ, Schild L, Adams MM, Deshpande AD, Franks AL (2002) Hospitalizations during pregnancy among managed care enrollees. *Obstetrics & Gynecology* 100(1):94 [PubMed: 12100809]
- Geiger CJ, Fahrenbach DM, Healey FJ (1959) Bendectin in the treatment of nausea and vomiting in pregnancy. *Obstet Gynecol* 14:688 [PubMed: 13827111]
- Gideon K, Katayoon M, Caroline M, Simerpal Kaur G (2009) The effect of acid-reducing pharmacotherapy on the severity of nausea and vomiting of pregnancy. *Obstet Gynecol Int* 2009(2009)
- Golberg D, Szilagyi A, Graves L (2007) Hyperemesis gravidarum and helicobacter pylori infection: a systematic review. *Obstet Gynecol* 110(3):695 [PubMed: 17766620]
- Goodwin TM (1998) Hyperemesis gravidarum. *Clin Obstet Gynecol* 41: 597–605 [PubMed: 9742356]
- Goodwin TM (2008) Hyperemesis Gravidarum. *Obstet Gynecol Clin N Am* 35(3):401
- Goodwin TM, Hershman JM, Cole L (1994) Increased concentration of the free beta-subunit of human chorionic gonadotropin in hyperemesis gravidarum. *Acta Obstet Gynecol Scand* 73:770–772 [PubMed: 7817726]
- Goodwin TM, Montoro M, Mestman JH, Pekary AE, Hershman JM (1992) The role of chorionic gonadotropin in transient hyperthyroidism of hyperemesis gravidarum. *J Clin Endocrinol Metab* 75(5):1333 [PubMed: 1430095]
- Gross S, Librach C, Cecutti A (1989) Maternal weight loss associated with hyperemesis gravidarum: a predictor of fetal outcome. *Am J Obstet Gynecol* 160:906–909 [PubMed: 2712122]
- Guclu S, Gol M, Dogan E, Saygili U (2005) Mirtazapine use in resistant hyperemesis gravidarum: report of three cases and review of the literature. *Arch Gynecol Obstet* 272(4):298 [PubMed: 16007504]
- Guttuso T, Robinson LK, Amankwah KS (2010) Gabapentin use in hyperemesis gravidarum: a pilot study. *Early Hum Dev* 86(1):65 [PubMed: 20015600]
- Hallak M, Tsalamandris K, Drombrowski MP, Isada NB, Pryde PG, Evans MI (1996) Hyperemesis gravidarum: effects on fetal outcome. *J Reprod Med* 41:871–874 [PubMed: 8951141]
- Herrell HE (2014) Nausea and vomiting of pregnancy. *Am Fam Physician* 89(12):965 [PubMed: 25162163]
- Hinkle Stefanie N., Mumford Sunni L., Grantz Katherine L., Silver Robert M., Mitchell Emily M., Sjaarda Lindsey A., Radin Rose G., Perkins Neil J., Galai Noya, Schisterman Enrique F. (2016) Association of Nausea and Vomiting During Pregnancy With Pregnancy Loss. *JAMA Inter Med*
- Hirschfeld RMA, Williams JBW, Spitzer RL, Calabrese JR, Flynn L, Keck PE, Lewis L, McElroy SL, Post RM, Rappaport DJ, Russell JM, Sachs GS, Zajecka J (2000) Development and validation of a screening instrument for bipolar spectrum disorder: the mood disorder questionnaire. *Am J Psychiatr* 157(11):1873–1875 [PubMed: 11058490]
- Hizli D, Kosus A, Kosus N, Kamalak Z, Akkurt G (2012) Hyperemesis gravidarum and depression in pregnancy: is there an association? *J Psychosom Obstet Gynecol* 33(4):171
- Hod M, Orvieto R, Kaplan B, Friedman S, Ovadia J (1994) Hyperemesis GRAVIDARUM—a review. *J Reprod Med* 39(8):605 [PubMed: 7996524]

- Hsu JJ, Rene C-G, Nelson DK, Kim CH (1996) Nasogastric enteral feeding in the management of hyperemesis gravidarum. *Obstetrics & Gynecology* 88(3):343 [PubMed: 8752236]
- Huxley RR (2000) Nausea and vomiting in early pregnancy: its role in placental development. *Obstetrics & Gynecology* 95(5):779 [PubMed: 10775746]
- Huybrechts KF, Hernández-Díaz S, Avorn J (2014) Antidepressant use in pregnancy and the risk of cardiac defects. *N Engl J Med* 371 (12): 1168–1169 [PubMed: 25229932]
- Jackson CW, Sheehan AH, Reddan JG (2007) Evidence-based review of the black-box warning for droperidol.(SPECIAL FEATURE)(drug overview)(clinical report). *Am J Health Syst Pharm* 64(11):1174 [PubMed: 17519460]
- Jacoby EB, Porter KB (1999) Helicobacter pylori infection and persistent hyperemesis gravidarum. *Am J Perinatol* 16(2):85 [PubMed: 10355915]
- Jamigorn M, Phupong V (2007) Acupressure and vitamin B6 to relieve nausea and vomiting in pregnancy: a randomized study. *Arch Gynecol Obstet* 276(3):245 [PubMed: 17318558]
- Jednak MA, Shadigian EM, Kim MS, Woods ML, Hooper FG, Owyang C, Hasler WL (1999) Protein meals reduce nausea and gastric slow wave dysrhythmic activity in first trimester pregnancy. *Am J Physiol* 277(4 Pt 1):G855 [PubMed: 10516152]
- Källén B, Lundberg G, Aberg A (2003) Relationship between vitamin use, smoking, and nausea and vomiting of pregnancy. *Acta Obstet Gynecol Scand* 82(10):916 [PubMed: 12956841]
- Kashifard M, Basirat Z, Kashifard M, Golsorkhtabar-Amiri M, Moghaddamnia A (2013) Ondansetron or metoclopramide? Which is more effective in severe nausea and vomiting of pregnancy? A randomized trial double-blind study. *Clinical and experimental obstetrics & gynecology* 40(1):127–130 [PubMed: 23724526]
- Keating A, Chez RA (2002) Ginger syrup as an antiemetic in early pregnancy. *Altern Ther Health Med* 8(5):89 [PubMed: 12233808]
- Sw Kim, Is Shin, Jm Kim, Yc Kim, Ks Kim, Km Kim, Sj Yang, Js Yoon (2008) Effectiveness of mirtazapine for nausea and insomnia in cancer patients with depression. *Psychiatry Clin Neurosci* 62(1):75 [PubMed: 18289144]
- Kimura M, Amino N, Tamaki H, Ito E, Mitsuda N, Miyai K, Tanizawa O (1993) Gestational thyrotoxicosis and hyperemesis gravidarum: possible role of hCG with higher stimulating activity. *Clin Endocrinol* 38(4):345
- Klebanoff MA, Koslowe PA, Kaslow R, Rhodes GG (1985) Epidemiology of vomiting in early pregnancy. *Obstet Gynecol* 66:612–616 [PubMed: 3903578]
- Koren G (2014) Treating morning sickness in the United States—changes in prescribing are needed. *Am J Obstet Gynecol* 211(6):602–606 [PubMed: 25151184]
- Koren G, Boskovic R, Hard M, Maltepe C, Navioz Y, Einarson A (2002) Motherisk-PUQE (pregnancy-unique quantification of emesis and nausea) scoring system for nausea and vomiting of pregnancy. (Understanding and treating nausea and vomiting of pregnancy). *Am J Obstet Gynecol* 186(5):S228 [PubMed: 12011891]
- Koren G, Clark S, Hankins GDV, Caritis SN, Miodovnik M, Umans JG, Mattison DR (2010) Effectiveness of delayed-release doxylamine and pyridoxine for nausea and vomiting of pregnancy: a randomized placebo controlled trial. *Am J Obstet Gynecol* 203(6):571.e571 [PubMed: 20843504]
- Kutcher JS, Engle A, Firth J, Lamm SH (2003) Bendectin and birth defects. II: ecological analyses. Birth defects research. Part A, Clinical And Molecular Teratology 67(2):88
- Lacasse A, Lagoutte A, Ferreira E, Bérard A (2009) Metoclopramide and diphenhydramine in the treatment of hyperemesis gravidarum: effectiveness and predictors of rehospitalisation. *European Journal Of Obstetrics And Gynecology* 143(1):43
- Lagiou P, Tamimi R, Mucci LA, Trichopoulos D, Adami H-O, Hsieh C-C (2003) Nausea and vomiting in pregnancy in relation to prolactin, estrogens, and progesterone: a prospective study. *Obstetrics & Gynecology* 101(4):639 [PubMed: 12681864]
- Levine MG, Esser D (1988) Total parenteral nutrition for the treatment of severe hyperemesis gravidarum: maternal nutritional effects and fetal outcome. *Obstet Gynecol* 72(1):102 [PubMed: 3132667]

- Li L, Li L, Zhou X, Xiao S, Gu H, Zhang G, 2015 Infection is associated with an increased risk of hyperemesis gravidarum: a meta-analysis 2015.
- Liu M-C, Kuo S-H, Lin C-P, Yang Y-M, Chou F-H, Yang Y-H (2014) Effects of professional support on nausea, vomiting, and quality of life during early pregnancy. *Biological Research For Nursing* 16(4):378 [PubMed: 24113384]
- Magee LA, Mazzotta P, Koren G (2002) Evidence-based view of safety and effectiveness of pharmacologic therapy for nausea and vomiting of pregnancy (NVP). *Am J Obstet Gynecol* 186(5):S256 [PubMed: 12011897]
- Maina A, Arrotta M, Cicogna L, Donvito V, Mischinelli M, Todros T, Rivolo S (2014) Transdermal clonidine in the treatment of severe hyperemesis. A pilot randomised control trial: CLONEMESI. (Clinical report). *BJOG: An International Journal of Obstetrics and Gynaecology* 121(12):1556 [PubMed: 24684734]
- Maltepe C, Koren G, 2013a The management of nausea and vomiting of pregnancy and hyperemesis gravidarum—a 2013 update. *Journal Of Population Therapeutics And Clinical Pharmacology = Journal De La Thérapeutique Des Populations Et De La Pharmacologie Clinique* 20(2), e184. [PubMed: 23863612]
- Maltepe C, Koren G, 2013b Preemptive treatment of nausea and vomiting of pregnancy: results of a randomized controlled trial. 2013b.
- Matok I, Gorodischer R, Koren G, Sheiner E, Wiznitzer A, Levy A (2009) The safety of metoclopramide use in the first trimester of pregnancy. *N Engl J Med* 360(24):2528 [PubMed: 19516033]
- Matthews A, Haas DM, Mathúna DPO, Dowswell T, Doyle M (2014) Interventions for nausea and vomiting in early pregnancy. *Cochrane Database Syst Rev* 3:CD007575
- Mazzota P, Magee L, Koren G (1997) Therapeutic abortions due to severe morning sickness unacceptable combination Canadian family physician *Médecin de famille canadien* 43:1055
- McCarthy FP, Khashan AS, North RA, Moss-Morris R, Baker PN, Dekker G, Poston L, Kenny LC, Wang H (2011) A prospective cohort study investigating associations between hyperemesis gravidarum and cognitive, behavioural and emotional well-being in pregnancy. *PLoS One* 6(11)
- McCarthy FP, Lutonski JE, Greene RA (2014) Hyperemesis gravidarum: current perspectives. *International Journal Of Women's Health* 6:719
- McCormack D (2010) Hypnosis for hyperemesis gravidarum. *Journal Of Obstetrics And Gynaecology : The Journal Of The Institute Of Obstetrics And Gynaecology* 30(7):647 [PubMed: 20925602]
- McKeigue P, Lamm S, Linn S, Kutcher J (1994) Bendectin and birth-defects. I. A metaanalysis of the epidemiologic studies. *Teratology* 50(1):27 [PubMed: 7974252]
- Morali JA, Braverman DZ (1990) Abnormal liver enzymes and ketonuria in hyperemesis gravidarum: a retrospective review of eighty patients. *J Clin Gastroenterol* 12:303–305 [PubMed: 2362099]
- Mullin PM, Ching C, Schoenberg F, Macgibbon K, Romero R, Goodwin TM, Fejzo MS (2012) Risk factors, treatments, and outcomes associated with prolonged hyperemesis gravidarum. *Journal Of Maternal-Fetal & Neonatal Medicine: The Official Journal Of The European Association Of Perinatal Medicine, The Federation Of Asia And Oceania Perinatal Societies, The International Society Of Perinatal Obstetricians* 25(6):632
- Mullin PM, Goodwin A, Fejzo F, Bray KW, Schoenberg R, MacGibbon TM, Romero MS, Fejzo MS, 2011 Prenatal exposure to hyperemesis gravidarum linked to increased risk of psychological and behavioral disorders in adulthood. p 200.
- Murphy PA (1998) Alternative therapies for nausea and vomiting of pregnancy. *Obstetrics & Gynecology* 91(1):78 [PubMed: 9464725]
- Nageotte MP, Briggs GG, Towers CV, Asrat T (1996) Droperidol and diphenhydramine in the management of hyperemesis gravidarum. *Am J Obstet Gynecol* 174(6):1801 [PubMed: 8678143]
- Nasrin S, Sholeh S, Robabeh M, Masoumeh R, Fariba F (2011) Comparing the effects of ginger and metoclopramide on the treatment of pregnancy nausea. *Pak J Biol Sci* 14(16):817 [PubMed: 22545357]
- Nelson-Piercy C (1998) Treatment of nausea and vomiting in pregnancy. *Drug Saf* 19(2):155 [PubMed: 9704251]

- Neutel CI, Johansen HL (1995) Measuring drug effectiveness by default: the case of Bendectin. *Canadian Journal Of Public Health/Revue Canadienne De Sante&Apos;E Publique* 86(1):66
- Niebyl JR, Goodwin TM (2002) Overview of nausea and vomiting of pregnancy with an emphasis on vitamins and ginger. *Am J Obstet Gynecol* 186(5):S253 [PubMed: 12011896]
- Niemeijer M, Grooten I, Vos N, Bais J, van Der Post J, Mol B, Roseboom T, Leeftang M, Painter RC (2014) Diagnostic markers for hyperemesis gravidarum: a systematic review and metaanalysis. *Am J Obstet Gynecol* 211(2)
- Norheim AJ, Pedersen EJ, Fønnebø V, Berge L, 2001 Acupressure treatment of morning sickness in pregnancy. A randomised, double-blind, placebo-controlled study. *Scand J Prim Health Care* 19(1), 43. [PubMed: 11303547]
- Nulman I, Rovet J, Barrera M, Knittel-Keren D, Feldman BM, Koren G (2009) Long-term neurodevelopment of children exposed to maternal nausea and vomiting of pregnancy and diclectin. *J Pediatr* 155(1):45 [PubMed: 19394042]
- Oliveira LG, Capp SM, You WB, Riffenburgh RH, Carstairs SD (2014) Ondansetron compared with doxylamine and pyridoxine for treatment of nausea in pregnancy: a randomized controlled trial. *Obstet Gynecol* 124(4):735 [PubMed: 25198265]
- Orazio LM, Meyerowitz BE, Korst LM, Goodwin TM, Romero R (2011) Evidence against a link between hyperemesis gravidarum and personality characteristics from an ethnically diverse sample of pregnant women: a pilot study. *Journal of Women's Health* 20(1):137
- Ozgoli G, Goli M, Simbar M (2009) Effects of ginger capsules on pregnancy, nausea, and vomiting. *Journal Of Alternative And Complementary Medicine (New York, NY)* 15(3):243
- Park-Wyllie L, Mazzotta P, Pastuszak A, Moretti ME, Beique L, Hunnisett L, Friesen MH, Jacobson S, Kasapinovic S, Chang D, Diav-Citrin O, Chitayat D, Nulman I, Einarson TR, Koren G (2000) Birth defects after maternal exposure to corticosteroids: prospective cohort study and meta-analysis of epidemiological studies. *Teratology* 62(6):385 [PubMed: 11091360]
- Pasricha PJ, Nonko P, Aravind S, Joseph J (2006) Drug insight: from disturbed motility to disordered movement—a review of the clinical benefits and medicolegal risks of metoclopramide. *Nature Clinical Practice Gastroenterology & Hepatology* 3(3):138
- Pasternak B, Svanström H, Hviid A (2013) Ondansetron in pregnancy and risk of adverse fetal outcomes. *N Engl J Med* 368(9):814 [PubMed: 23445092]
- Pasternak B, Svanström H, Mølgaard-Nielsen D, Melbye M, Hviid A (2014) Metoclopramide in pregnancy and risk of major congenital malformations and fetal death. *Obstet Anesth Dig* 34(4):211
- Peeters A, van de Wyngaert F, Van Lierde M, Sindic CJ, Laterre EC (1993) Wernicke's encephalopathy and central pontine myelinolysis induced by hyperemesis gravidarum. *Acta Neurol Belg* 93:276–282 [PubMed: 8296523]
- Pradat P, Robert-Gnansia E, Di Tanna GL, Rosano A, Lisi A, Mastroiacovo P (2003) First trimester exposure to corticosteroids and oral clefts. Birth defects research. Part A, Clinical And Molecular Teratology 67(12):968 [PubMed: 14745915]
- Price A, Davies R, Heller SR, Milford-Ward A, Weetman AP (1996) Asian women are at risk of gestational thyrotoxicosis. *J Clin Endocrinol Metab* 81:1160–1163 [PubMed: 8772593]
- Poursharif B, Korst LM, Fejzo MS, Macgibbon KW, Romero R, Goodwin TM (2007) The psychosocial burden of hyperemesis gravidarum. *J Perinatol* 28(3):176–181 [PubMed: 18059463]
- Quigley EMM, Hasler WL, Parkman HP (2001) AGA technical review on nausea and vomiting. *Gastroenterology* 120(1):263 [PubMed: 11208736]
- Rhodes VA, McDaniel RW (1999) The index of nausea, vomiting, and retching: a new format of the index of nausea and vomiting. *Oncol Nurs Forum* 26(5):889 [PubMed: 10382187]
- Rohde A, Dembinski J, Dorn C (2003) Mirtazapine (Remergil) for treatment resistant hyperemesis gravidarum: rescue of a twin pregnancy. *Arch Gynecol Obstet* 268(3):219 [PubMed: 12819986]
- Sahakian V, Rouse D, Sipes S, Rose N, Niebyl J (1991) Vitamin B6 is effective therapy for nausea and vomiting of pregnancy: a randomized, double-blind placebo-controlled study. *Obstet Gynecol* 78(1):33 [PubMed: 2047064]

- Saks BR (2001) Mirtazapine: treatment of depression, anxiety, and hyperemesis gravidarum in the pregnant patient. A report of 7 cases. *Archives Of Women's Mental Health* 3(4):165
- Sandven I, Abdelnoor M, Nesheim B-I, Melby KK (2009) *Helicobacter pylori* infection and hyperemesis gravidarum: a systematic review and meta-analysis of case-control studies. *Acta Obstet Gynecol Scand* 88(11):1190 [PubMed: 19900137]
- Schuster K, Bailey LB, Dimperio D, Mahan CS (1985) Morning sickness and vitamin B 6 status of pregnant women. *Human Nutrition: Clinical Nutrition* 39(1):75 [PubMed: 3997551]
- Schwarzer V, Heep A, Gembruch U, Rohde A (2008) Treatment resistant hyperemesis gravidarum in a patient with type 1 diabetes mellitus: neonatal withdrawal symptoms after successful antiemetic therapy with mirtazapine. *Arch Gynecol Obstet* 277(1):67 [PubMed: 17628816]
- Seng JS, Schrot JA, Van De Ven C, Liberzon I (2007) Service use data analysis of pre-pregnancy psychiatric and somatic diagnoses in women with hyperemesis gravidarum. *J Psychosom Obstet Gynaecol* 28(4):209 [PubMed: 17852658]
- Seto A, Einarson T, Koren G (1997) Pregnancy outcome following first trimester exposure to antihistamines: meta-analysis. *Am J Perinatol* 14(3):119 [PubMed: 9259911]
- Sheehan P (2007) Hyperemesis gravidarum: assessment and management. *Aust Fam Physician* 36(9):698 [PubMed: 17885701]
- Shepard TH, Brent RL, Friedman JM, Jones KL, Miller RK, Moore CA, Polifka JE (2002) Update on new developments in the study of human teratogens. *Teratology* 65(4):153 [PubMed: 11948561]
- Sherman PW, Flaxman SM (2002) Nausea and vomiting of pregnancy in an evolutionary perspective. *Am J Obstet Gynecol* 186(5):S190 [PubMed: 12011885]
- Shrim A, Boskovic R, Maltepe C, Navios Y, Garcia-Bournissen F, Koren G, Shrim G (2006) Pregnancy outcome following use of large doses of vitamin B 6 in the first trimester. *J Obstet Gynaecol* 26(8):749 [PubMed: 17130022]
- Simon EP, Schwartz J (1999) Medical Hypnosis for Hyperemesis Gravidarum *Birth* 26(4):248 [PubMed: 10655831]
- Simpson SW, Goodwin TM, Robins SB, Rizzo AA, Howes RA, Buckwalter DK, Buckwalter JG (2001) Psychological factors and hyperemesis gravidarum. *Journal Of Women's Health & Gender-Based Medicine* 10(5):471
- Slaughter SR, Hearn-Stokes R, Van Der Vlugt T, Joffe HV (2014) FDA approval of doxylamine-pyridoxine therapy for use in pregnancy. *N Engl J Med* 370(12):1081 [PubMed: 24645939]
- Smith C, Crowther C, Willson K, Hotham N, McMillian V (2004) A randomized controlled trial of ginger to treat nausea and vomiting in pregnancy. *Obstet Gynecol* 103(4):639 [PubMed: 15051552]
- Spiegel DR, Webb K (2012) A case of treatment refractory hyperemesis gravidarum in a patient with comorbid anxiety, treated successfully with adjunctive gabapentin: a review and the potential role of neurogastroenterology in understanding its pathogenesis and treatment. *Innovations In Clinical Neuroscience* 9(11-12):31
- Sripramote M, Lekhyananda N (2003) A randomized comparison of ginger and vitamin B6 in the treatment of nausea and vomiting of pregnancy. *Journal Of The Medical Association Of Thailand = Chotmaihet Thangphaet* 86(9):846 [PubMed: 14649969]
- Stokke G, Gjelsvik BL, Flaatten KT, Birkeland E, Flaatten H, Trovik J (2015) Hyperemesis gravidarum, nutritional treatment by nasogastric tube feeding: a 10-year retrospective cohort study. *Acta Obstet Gynecol Scand* 94(4):359-367 [PubMed: 25581215]
- Summers A (2012) Emergency management of hyperemesis gravidarum: AnthonySummersdiscussesthecauses,signsand symptomsofsevere morning sickness, and explains how the condition should be treated in emergency care settings. (Report). *Emergency Nurse* 20(4):24
- Tamay AG, Kescu N (2011) Hyperemesis gravidarum: current aspect. *J Obstet Gynaecol* 31(8):708 [PubMed: 22085059]
- Tan PC, Khine PP, Vallikkannu N, Omar SZ (2010a) Promethazine compared with metoclopramide for hyperemesis gravidarum: a randomized controlled trial. *Obstet Gynecol* 115(5):975 [PubMed: 20410771]

- Tan PC, Vani S, Lim BK, Omar SZ (2010b) Anxiety and depression in hyperemesis gravidarum: prevalence, risk factors and correlation with clinical severity. *Eur J Obstet Gynecol Reprod Biol* 149(2):153 [PubMed: 20097465]
- Tan PC, Zaidi SN, Azmi N, Omar SZ, Khong SY (2014) Depression, anxiety, stress and hyperemesis gravidarum: temporal and case controlled correlates. *PLoS One* 9(3)
- Tasci Y, Demir B, Dilbaz S, Haberal A (2009) Use of diazepam for hyperemesis gravidarum. *Journal Of Maternal-Fetal & Neonatal Medicine: The Official Journal Of The European Association Of Perinatal Medicine, The Federation Of Asia And Oceania Perinatal Societies, The International Society Of Perinatal Obstetricians* 22(4):353
- Trogstad LIS, Stoltenberg C, Magnus P, Skjaerven R, Irgens LM (2005) Recurrence risk in hyperemesis gravidarum. *Bjog : An International Journal Of Obstetrics And Gynaecology* 112(12):1641 [PubMed: 16305568]
- Tsang IS, Katz VL, Wells SD (1996) Maternal and fetal outcomes in hyperemesis gravidarum. *Int J Gynecol Obstet* 55:231–223
- Uguz F (2014) Low-dose mirtazapine in treatment of major depression developed following severe nausea and vomiting during pregnancy: two cases. *Gen Hosp Psychiatry* 36(1):125.e125
- Van Thiel DH, Gravalier JS, Joshi SN (1977) Heartburn of pregnancy. *Gastroenterol* 72:666–668
- Vandraas K, Vikanes A, Vangen S, Magnus P, Stoer N, Grjibovski A (2013) Hyperemesis gravidarum and birth outcomes—a population-based cohort study of 2.2 million births in the Norwegian birth registry. *BJOG: An International Journal of Obstetrics and Gynaecology* 120(13):1654 [PubMed: 24021026]
- Veenendaal M, van Abeelen A, Painter RC, van Der Post J, Roseboom T (2011) Consequences of hyperemesis gravidarum for offspring: a systematic review and meta-analysis. *Bjog-An International Journal Of Obstetrics And Gynaecology* 118(11):1302 [PubMed: 21749625]
- Vikanes Å, Grjibovski AM, Vangen S, Gunnes N, Samuelsen SO, Magnus P (2010) Maternal body composition, smoking, and hyperemesis gravidarum. *Ann Epidemiol* 20(8):592 [PubMed: 20609338]
- Vikanes ÅV, Stør NC, Magnus P, Grjibovski AM (2013) Hyperemesis gravidarum and pregnancy outcomes in the Norwegian mother and child cohort—a cohort study. *Bmc Pregnancy And Childbirth* 13:169 [PubMed: 24004605]
- Viljoen E, Visser J, Koen N, Musekiwa A, 2014 A systematic review and meta-analysis of the effect and safety of ginger in the treatment of pregnancy-associated nausea and vomiting. *Nutrition Journal* 13.
- Vutyavanich T, Wongtra-ngan S, Ruangsri R-A (1995) Pyridoxine for nausea and vomiting of pregnancy: a randomized, double-blind, placebo-controlled trial. *Am J Obstet Gynecol* 173(3):881 [PubMed: 7573262]
- Wallstedt A, Riely CA, Shaver D (1990) Prevalence and characteristics of liver dysfunction in hyperemesis gravidarum. *Clin Res* 38:970–972
- Wegrzyniak LJ, Repke JT, Ural SH (2012) Treatment of hyperemesis gravidarum. *Reviews In Obstetrics & Gynecology* 5(2):78
- Weigel RM, Weigel MM (1989) Nausea and vomiting of early-pregnancy and pregnancy outcome—a meta-analytical review. *Br J Obstet Gynaecol* 96(11):1312 [PubMed: 2611170]
- Wheatley D (1977) Treatment of pregnancy sickness. *Br J Obstet Gynaecol* 84(6):444 [PubMed: 329860]
- Wilson R, McKillop JH, MacLean M (1992) Thyroid function test are rarely abnormal in patients with severe hyperemesis gravidarum. *Clin Endocrinol* 37:331–334
- Winterfeld U, Klinger G, Panchaud A, Stephens S, Arnon J, Malm H, Te Winkel B, Clementi M, Pistelli A, Ma áková E, Eleftheriou G, Merlob P, Kaplan YC, Buclin T, Rothuizen LE (2015) Pregnancy outcome following maternal exposure to mirtazapine: a multicenter, prospective study. *J Clin Psychopharmacol* 35(3):250 [PubMed: 25830592]
- Yost NP, McIntire DD, Wians FH, Ramin SM, Balko JA, Leveno KJ (2003) A randomized, placebo-controlled trial of corticosteroids for hyperemesis due to pregnancy. *Obstetrics & Gynecology* 102(6):1250 [PubMed: 14662211]

Zhang Y, Cantor RM, MacGibbon K, Romero R, Goodwin TM, Mullin PM, Fejzo MS (2011) Familial aggregation of hyperemesis gravidarum. Am J Obstet Gynecol 204(3):230.e231 [PubMed: 20974461]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1**Differential diagnosis for nausea and vomiting in pregnancy**

Gastrointestinal disorders
Gastroenteritis
Cholecystitis
Gastroesophageal reflux disease
Pancreatitis
Appendicitis
Peptic ulcer disease
Hepatitis
Gastroparesis
Bowel obstruction
Endocrine and metabolic disorders
Hyperthyroidism
Hyperparathyroidism
Hypercalcemia
Diabetic ketoacidosis
Genitourinary and renal disorders
Nephrolithiasis
Pyelonephritis
Uremia
Ovarian torsion
Neurological disorders
Migraines
Pseudotumor cerebri
Tumors of the central nervous system
Vestibular disease
Psychiatric disorders
Substance use disorders—alcohol intoxication and withdrawal, opioid withdrawal, sedative/hypnotic/anxiolytic withdrawal, stimulant intoxication
Cannabinoid hyperemesis syndrome
Eating disorders—anorexia nervosa, bulimia
Antidepressant discontinuation syndrome (due to abrupt discontinuation of medication with confirmation of pregnancy)
Other
Medications—antiarrhythmic, antihypertensives, narcotics, anticonvulsants, antibiotics, iron supplementation
Cyclic vomiting syndrome

Compiled from Goodwin (1998), Herrell (2014), Quigley et al. (2001), Wegrzyniak et al. (2012)