Published in final edited form as:

Nurs Res. 2020 ; 69(1): 42–50. doi:10.1097/NNR.0000000000000397.

Inpatient Maternal Mortality in the United States, 2002-2014

Mulubrhan F. Mogos, PhD, MSc [Assistant Professor],

Department of Women, Children and Family Health Science, College of Nursing, University of Illinois at Chicago, Chicago, IL, USA

Kylea L. Liese, PhD, CNM, RN [Assistant Professor],

Department of Women, Children and Family Health Science, College of Nursing, University of Illinois at Chicago, Chicago, IL, USA

Patrick D. Thornton, PhD, CNM [Clinical Assistant Professor],

Department of Women, Children and Family Health Science, College of Nursing, University of Illinois at Chicago, Chicago, IL, USA

Tracy A. Manuck, MD [Associate Professor],

Department of Obstetrics and Gynecology, University of North Carolina, Chapel Hill, NC, USA

William D. O'Brien Jr., PhD [Professor],

Department of Electrical and Computer Engineering, University of Illinois at Urbana-Champaign, Urbana, IL, USA

Barbara L. McFarlin, PhD, CNM, RDMS, FACNM, FAAN [Professor]

Department of Women, Children and Family Health Science, College of Nursing, University of Illinois at Chicago, Chicago, IL, USA

Abstract

Background: Although prior studies of inpatient maternal mortality in the U.S provides data on the overall rate and trend in inpatient maternal mortality, there are no published reports of maternal mortality data stratified by timing of its occurrence across the pregnancy continuum (antepartum, intrapartum, and postpartum).

Objective: The study objective was to determine whether the maternal mortality rate, trends over time, self-reported race/ethnicity, and associated factors vary based on the timing of the occurrence of death during pregnancy.

Corresponding author: Mulubrhan F. Mogos, PhD, MSc, Department of Women Children and Family Health Science, College of Nursing, University of Illinois, 845 S. Damen Ave, Chicago, IL 60612, Phone: 312.996.5801, Fax: 312.996.8871, mmogos@uic.edu. Author Note

Mulubrhan F. Mogos, PhD, MSc, Kylea L. Liese, PhD, CNM, RN, and Patrick D. Thornton, PhD, CNM, RN are Assistant Professors at the Department of Women, Children and Family Health Science, College of Nursing, University of Illinois at Chicago, Chicago, IL.

Tracy A. Manuck, MD is an Associate Professor at the Department of Obstetrics and Gynecology, University of North Carolina, Chapel Hill, NC.

William D. O'Brien, Jr., PhD, is a Professor at the Department of Electrical and Computer Engineering, University of Illinois at Urbana-Champaign, Urbana, IL.

Barbara L. McFarlin, PhD, CNM RDMS, FACNM, FAAN is a professor at the Department of Women, Children and Family Health Science, College of Nursing, University of Illinois at Chicago, Chicago, IL.

The authors have no conflicts of interest to report.

Ethical Conduct of Research: The study has been approved by the IRB of University of Illinois at Chicago.

Methods: We conducted a cross-sectional analysis of the Nationwide Inpatient Sample database to identify pregnancy-related inpatient stays stratified by timing. Among women in the sample, we determined in-hospital mortality and used International Classification of Diseases, Ninth Revision, Clinical Modification codes to identify comorbidities and behavioral characteristics associated with mortality including alcohol, drug, and tobacco use. Joinpoint regression was used to calculate rates and trends of in-hospital maternal mortality.

Results: During the study period, there were 7,411 inpatient maternal mortalities among an estimated 58,742,179 hospitalizations of women 15 – 49 years of age. In-hospital maternal mortality rate stratified by race showed that African Americans died at significantly higher rates during antepartum, intrapartum, and postpartum periods compared to hospitalizations for whites or Hispanics during the same time period. Although the postpartum hospitalization represents only 2% of pregnancy-related hospitalizations among women age 15 – 49 years, hospitalization during this time period accounted for 27.2% of all maternal deaths during pregnancy-related hospitalization.

Discussion: Most in-hospital maternal mortalities occur after hospital discharge from child birth (postpartum period). Yet, the postpartum period continues to be the time period with the least maternal healthcare surveillance in the pregnancy continuum. African American women experience three times more in-hospital mortality when compared to their white counterparts.

Keywords

inpatient; maternal mortality; United States; childbirth; postpartum

In the United States (U.S.), the maternal mortality rate, defined as the number of maternal deaths per 100,000 live births has increased from 7.2 to 18 per 100,000 (a 150% increase) between the years 1987 and 2014 (Centers for Disease Control and Prevention, 2014). A recent study reported an increase in the maternal mortality rate from 18.8 per 100,000 live births in 2000 to 23.8 per 100,000 live births in 2014, a 26.6% increase (MacDorman, Declercq, & Thoma, 2017). In contrast, the maternal mortality rate has decreased in most developed countries during the same period of time (Alkema et al., 2016; Li, Luo, Deng, Jacoby, & de Klerk, 2007; Sullivan, Ford, Chambers, & Slaytor, 2004). Although prior studies of inpatient maternal mortality in the U.S. provided data on the overall rate and trend in inpatient maternal mortality, to our knowledge, no study has provided maternal mortality data stratified by timing of its occurrence across the pregnancy continuum (antepartum, intrapartum, and postpartum). Scholarly and popular media reports (Creanga, 2018; Curtin & Hoyert, 2017; Martin & Montagne, 2017; Metcalfe, Wick, & Ronksley, 2018) have drawn attention to racial disparities in rates of maternal mortality, particularly among African American women, who are 3-5 times as likely to die of maternal causes than white women (Martin & Montagne, 2017; Tucker, Berg, Callaghan, & Hsia, 2007). While some attribute the disparity to increasing risk factors in this population (obesity, advanced maternal age, hypertension, diabetes), others suggest structural racism as an underlying etiology (Jain et al., 2018; Martin & Montagne, 2017). Structural racism refers to the totality of ways in which society enacts racial discrimination through institutions, policies and systems (Bailey, et al., 2017; Krieger, 2014).

The study objective was to determine whether the maternal mortality rate, trends over time, self-reported race/ethnicity and associated factors vary based on the timing of its occurrence during pregnancy.

METHODS

Using the National Inpatient Sample (NIS) from the Healthcare Cost and Utilization Project (HCUP) (HCUP, 2011, 2015), we conducted a retrospective analysis of all pregnancy-related inpatient hospitalizations from January 1, 2002 to December 31, 2014. The NIS is the largest publicly available all-payer inpatient hospitalization database in the United States. To create the NIS, HCUP uses a cluster sampling method and stratifies hospitalizations from participating non-federal community hospitals using five characteristics (rural/urban location, number of beds, geographic region, teaching status, and ownership). Then, using a simple random sampling approach, 20% of hospitals are selected from each stratum (HCUP, 2011) and all inpatient discharges from hospitals selected using the simple random sampling approach are included in the NIS. In 2012, the NIS sampling approach was revised to include a sample of discharges from all hospitals instead of only the 20% selected using a stratified sampling approach. The NIS has been validated against other databases and is representative of the population of all hospitalizations in the United States on critical hospital and patient characteristics (Hashimoto, Brodt, Skelly, & Dettori, 2014; Murphy, Alavi, & Maykel, 2013).

Study Population

The study population consisted of pregnancy-related inpatient hospitalizations 15-49 years of age, identified using an HCUP-created “NEOMAT” variable. The NEOMAT indicator identifies maternal diagnosis records on the basis of International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) diagnosis and procedure codes for pregnancy (HCUP, 2015; Weiss & Elixhauser, 2006).

We categorized each maternal pregnancy-related hospitalization as antepartum, intrapartum, or postpartum (HCUP, 2015). Antepartum hospitalizations reflect the admission of a pregnant person with no record of childbirth during the same hospitalization. An intrapartum hospitalization reflects all admissions for intrapartum management (birth of fetus) through initial discharge (approximately 1-3 days) after birth. Postpartum hospitalizations are any readmission within the first 42 days after birth of the baby. We used ICD-9-CM diagnosis codes to determine whether each woman had a hospitalization during the AP, IP, or PP period using algorithm previously published by Kuklina and colleagues (Callaghan, Creanga, & Kuklina, 2012; Kuklina et al., 2008). IP hospitalizations were identified using delivery-specific ICD-9-CM diagnosis, procedure, and diagnosis-related group (DRG) codes. PP hospitalizations (within 42 days of delivery) were identified using V24 code, postpartum DRG codes and presence of the number “4” in the fifth digit of selected ICD-9-CM codes. Pregnancy-related discharges that did not have a code for IP or PP conditions were classified as AP hospitalizations (Callaghan et al., 2012; Kuklina et al., 2008). Supplemental Table 1 (Table S1) provides the complete list of ICD-9-CM diagnosis,

procedure, and DRG codes used in this study. Mortality was identified using the variable disposition status provided by HCUP.

Covariates

Individual-level sociodemographic and behavioral characteristics were also extracted from the NIS databases. Patient age in years was classified into the following categories: 15-24, 25-34, and 35-49. We used median household income in quartiles, calculated by HCUP, as a proxy for participants' socioeconomic status. Type of insurance used was grouped into three categories: government (Medicare and Medicaid), private (commercial carrier, private health maintenance organization, and preferred provider organization), and other sources (e.g., self-pay and charity).

In addition, hospital characteristics such as geographic region (northeast, midwest, south, or west), location and teaching status (rural, urban non-teaching, urban teaching) were considered. We considered the effect of several individual-level behavioral characteristics, including tobacco, alcohol, and drug use. We also examined distribution of all comorbidities included in the Elixhauser comorbidity index among those who died by timing of death (AP, IP, and PP) (Elixhauser, Steiner, Harris, & Coffey, 1998).

Statistical Analyses

The distribution of inpatient maternal mortality diagnoses among AP, IP, and PP discharges were described by sociodemographic, behavioral, hospital, insurance, clinical, and pregnancy-related comorbidities using descriptive statistics such as frequencies and rates. Joinpoint regression was used to examine temporal trends of inpatient maternal mortality during the study period by timing of hospitalization (AP, IP, and PP). We further examined trends in inpatient mortality across the pregnancy continuum stratified by self-reported race/ethnicity (white, African American, and Hispanic). The annual percent change (APC) was calculated to describe the trend within each time interval, as previously described by Clegg and colleagues (Clegg, Hankey, Tiwari, Feuer, & Edwards, 2009).

All statistical analyses accounted for the complex sampling design of the NIS and were weighted to facilitate the generation of national frequency and prevalence estimates. To account for NIS sampling design changes, we used the NIS-Trends files, so that trend weights and data elements would be consistently defined over time (Houchens & Elixhauser, 2006). All analyses were conducted by the primary author using software (SAS 9.4; SAS Institute Inc., Cary, NC) using a 5% type I error rate and two-sided hypothesis tests. The study was approved by the institutional review board of the primary author's institution. Data used in this study are available for purchase from HCUP for interested researchers. Upon request, authors are willing to share information related to analytic approach used in this study.

RESULTS

In the United States from 2002 to 2014, there were 7,411 inpatient maternal mortalities among an estimated 58,742,179 hospitalizations among women 15 – 49 years of age (Table 1). The overall inpatient maternal mortality rate during the study period was 34, 7, and 226

per 100,000 for antepartum, intrapartum, and postpartum hospitalizations, respectively. The mean age of women who died versus women who survived during pregnancy-related hospitalization was (28.3 vs 27.1 years, $p < 0.01$) for antepartum; (29.9 vs 27.7 years, $p < 0.01$) for intrapartum; and (30 vs 27.9 years, $p < 0.01$) for postpartum hospitalizations. Table 1 presents the distribution of inpatient maternal mortality across the pregnancy continuum by selected demographic, behavioral, and hospital characteristics. Across all pregnancy time periods (antepartum, intrapartum, and postpartum), the proportion of women who died during pregnancy-related hospitalization was higher among women who are older (age >35 years), non-Hispanic (NH)-African Americans, users of alcohol and abused drugs, and from southern region of the U.S.

Table 2 summarizes the distribution of inpatient maternal mortality across racial groups by selected demographic, behavioral, and hospital characteristics. The distributions of maternal age, alcohol use, and hospital size (occupancy) among women who died during pregnancy-related hospitalization were not statistically different across maternal race groups. However, there was statistically significant difference in rates of tobacco use, drug abuse, household income, and insurance status.

In-hospital maternal mortality rate stratified by race showed that African Americans die at significantly higher rates of 47, 17, and 379 per 100,000 maternal hospitalizations during antepartum, intrapartum, and postpartum periods, respectively, compared to 30, 5, 160 per 100,000 hospitalizations for whites and 28, 6 and 181 per 100,000 hospitalizations for Hispanics during the same time period ($p < 0.001$). In-hospital mortality rate during IP hospitalization among African Americans is three times higher than that of whites and Hispanics. When compared to women 15 – 34 years of age, advanced maternal age women (age ≥ 35 years) experienced significantly higher rates of in-hospital maternal mortality during antepartum (32 vs 44 per 100,000), intrapartum (6 vs 13 per 100,000), and postpartum (206 vs 319 per 100,000) hospitalizations ($p < 0.001$). However, it is important to highlight that the average age at the time of in-hospital maternal death during IP hospitalization was 29.7 years for whites, 29.7 years for African Americans and 30.3 years for Hispanics, indicating that age is not a factor for the observed 3-fold high mortality among African Americans. Although the postpartum hospitalization represents only 2% of pregnancy-related hospitalizations among women age 15 – 49 years, hospitalization during this time period accounted for 27.2% of all maternal deaths during pregnancy-related hospitalization (Figure 1).

The rate of cesarean section (CS) varied among women who survived to intrapartum discharge (31.7%) compared to those who died during hospitalization (72.8%). The rate of CS among those who died during hospitalization was 77.1% for whites, 73.8 (African Americans) and 68% among Hispanics. Our data did not allow us to pinpoint the indication(s) for the CS procedures.

Among all in-hospital maternal deaths during the study period, 19.2%, 19.0%, and 20.5% of antepartum, intrapartum, and postpartum hospitalization involved three or more indicators for severe maternal morbidity respectively (Table S2). Among maternal deaths that occurred during antepartum, intrapartum, and postpartum hospitalizations, acute myocardial

infarction was present in 41.3%, 52.4%, and 62.9% respectively. Next to acute myocardial infarction, the most frequently reported indicator of severe maternal morbidity indicator during antepartum, intrapartum, and postpartum hospitalizations were: amniotic fluid embolism (8.75%), conversion of cardiac rhythm (4.54%), and pulmonary edema/acute heart failure (7.08%) respectively. Our data, seen in Table 3) indicate that comorbidities such as fluid and electrolyte imbalance, coagulopathy, anemia related to blood loss, hypertension, and neurological disorders are the most commonly reported comorbidities among women who died during pregnancy-related hospitalization.

Between 2002 and 2014, we found a statistically significant 3.6% annual reduction ($APC = -3.6 [-6.1, -1.2]$) in inpatient maternal mortality during the intrapartum period. However, the inpatient maternal mortality rate remained steady among antepartum and postpartum hospitalizations (Figure 2). Further trend analyses stratified by race showed that inpatient maternal mortality during intrapartum hospitalization decreased more among African Americans ($APC = -4.7 [-8.0, -1.3]$) when compared to whites ($APC = -3.3 [-6.0, -0.5]$) (Figures S1 and S2). Hispanic women experienced a statistically significant 8.7% annual reduction in maternal mortality during postpartum hospitalization from 2002 to 2014 ($APC = -8.7, [-16.2, -0.6]$) (Figure S3).

DISCUSSION

The major new findings of this study are: 1) although intrapartum deaths during hospitalization decreased by 3.6% annually between 2002 and 2014 for all women, and by 4.7% annually for NH African-American women, African-American women are still three times more likely to die during intrapartum hospitalization than white women; 2) the postpartum period, a time with the least maternal healthcare surveillance, had the highest rate (226/100,000) of maternal mortality compared with the antepartum period (34/100,000) and intrapartum period (7/100,000); and 3) in-hospital maternal mortality rates significantly vary across racial groups, with African-American women experiencing the highest rate of mortality during antepartum, intrapartum, and postpartum hospitalizations.

Our study assessed rates of maternal mortality that occurred in an in-patient hospital setting only. Findings in this study underscore that the postpartum period is the deadliest time for pregnant women with 226 maternal deaths occurring per 100,000 postpartum hospitalizations. The recent guideline by the American College of Obstetricians and Gynecologists (American College of Obstetricians and Gynecologists, 2018) emphasized the need for postpartum care, urging providers to focus on individualized early fourth trimester care. Previously we have shown that women were at increased risk of cardiac pregnancy complications that are deadly during the postpartum period (Mogos, et al., 2018). Thus, the recent guideline regarding early fourth trimester care has the potential to save maternal lives by providing care during a time of least healthcare surveillance for women at increased risk of postpartum mortality.

On the other hand, the relatively lower rate of mortality during intrapartum hospitalization, a time when the mother and fetus are intensely monitored, is encouraging. The observed decline in inpatient maternal mortality during the intrapartum period could be due to

improved intrapartum monitoring, presence of obstetric in-house healthcare providers and availability of in-house anesthesia service in most labor and delivery units (Mogos, Salemi, Spooner, McFarlin, & Salihu, 2016). However, surveillance of high-risk women during intrapartum hospitalization before discharge needs to be emphasized as one of the focal points for intervention to decrease maternal mortality during postpartum period. Improved patient education is also needed so that women and their families can recognize early warning signs of severe morbidity such as fatigue, myocardial infarction and heart failure.

The three-fold high mortality rate experienced by African-American women during the intrapartum period is consistent with several studies that show significant racial disparities in maternal morbidity and mortality among US women (Creanga, Bateman, Kuklina, & Callaghan, 2014; Creanga, et al., 2015; Liese, et al., 2019; Louis, Menard, & Gee, 2015). Demonstrated black-white differences in maternal death could be the result of multifaceted factors (Moaddab et al., 2018; Rosenberg, Geller, Studee, & Cox, 2006; Tucker et al., 2007). Due to the limited nature of our dataset, we were not able to test the effect of specific risk factors in the observed racial disparity.

Hispanic women experienced a significant decline in inpatient maternal mortality during postpartum hospitalization. Because of the nature of our data, we are not able to elucidate reasons behind the observed reduction in inpatient maternal mortality among Hispanic women. Previously it has been reported that strong social support was associated with better postpartum outcomes (Clark, Christmas, Frye, Meyers, & Perlin, 2014; Keller et al., 2014; Mann, Mannan, Quinones, Palmer, & Torres, 2010; Negron, Martin, Almog, Balbierz, & Howell, 2013). Overall, the observed racial disparity in inpatient maternal mortality suggests that specifically targeted quality improvement efforts are needed to reduce inpatient maternal mortality and eliminate associated disparity (Adams & Thomas, 2018; Negron et al., 2013). In the era of personalized healthcare, different interventions may be needed based upon racial, ethnic, and health risk factors, rather than using generalized care plans to be used in all circumstances.

Since we used the NIS data to estimate prevalence, our data can only be generalized to maternal mortality in an inpatient setting. We also acknowledge that our prevalence estimates of maternal mortality may be underestimated since our data do not include mortalities that might have occurred in an emergency department, home or outpatient setting; however, the use of a large, multi-year, nationally-representative hospital discharge database is the primary strength of the current study. The NIS data contain hospital discharge data containing demographic, behavioral, hospital, and clinical variables from roughly 1,000 U.S. community hospitals with approximately 8 million discharge records annually, allowing us to adjust for multiple potential confounders in a multivariate model. The current study has other limitations that relate to our use of cross-sectional administrative data. The first limitation is that the databases lack data on some important variables including, gestational age and any treatment received by patients. Therefore, for example, we did not assess when exactly during the antepartum period (first, second, or third trimester) the death occurred. Also, we could not assess effectiveness of various treatments or procedures that might have occurred during the hospitalization. Second, identification of cases, behaviors, and clinical diagnoses using ICD-9-CM codes is subject to errors in

coding. Third, not all states report race-ethnicity data, and there is variation in reporting. The dataset used in this study has missing race-ethnicity for one fourth of the study population. Although racial-ethnic disparities were not considered in this study, our ability to fully control for the variable race was limited. Finally, the NIS data do not provide information about the main cause of death.

CONCLUSION

This study provides an estimate of the prevalence of inpatient maternal mortality in relation to the timing of delivery and indicates that trends in inpatient maternal mortality vary across racial groups. We have assessed these trends over a 13 year period and noted persistence of racial disparity in maternal mortality. In the past five years, there has been increased media attention on issues of maternal mortality and associated racial disparity. The authors hope that such an increased attention on the topic combined with ongoing efforts by perinatal quality collaboratives and maternal care providers from various states will lead to reducing maternal mortality and associated racial disparity. Future studies focusing on in-hospital mortality should try to disentangle unique factors contributing to maternal mortality and associated racial disparity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgement:

This study was partially funded by National Institutes of Health R01 HD089935 (BLM & WDO), and R01 MD011609 (TAM)

The content is solely the responsibility of the authors and does not necessarily represent the official views of the author's institutions. Part of this work was presented at the 2018 Council for the advancement of nursing science (CANS) in Washington DC on September 14th, 2018

References

- American College of Obstetricians and Gynecologists (2018). Committee Opinion No. 736: Optimizing Postpartum Care. *Obstetrics and Gynecology*, 131, e140–e150. doi:10.1097/aog.0000000000002633. [PubMed: 29683911]
- Adams C, & Thomas SP (2018). Alternative prenatal care interventions to alleviate Black-White maternal/infant health disparities. *Sociology Compass*, 12, e12549. doi:doi:10.1111/soc4.12549.
- Alkema L, Chou D, Hogan D, Zhang S, Moller AB, Gemmill A, ... & Say L (2016). Global, regional, and national levels and in maternal mortality between 1990 and 2015, with scenario-based projections to 2030: a systematic analysis by the UN Maternal Mortality Estimation Inter-Agency Group. *Lancet*, 387, 462–474. doi:10.1016/s0140-6736(15)00838-7. [PubMed: 26584737]
- Bailey ZD, Krieger N, Agenor M, Graves J, Linos N, & Bassett MT (2017). Structural racism and health inequities in the USA: evidence and interventions. *Lancet*, 389, 1453–1463. doi:10.1016/s0140-6736(17)30569-x. [PubMed: 28402827]
- Callaghan WM, Creanga AA, & Kuklina EV (2012). Severe maternal morbidity among delivery and postpartum hospitalizations in the United States. *Obstetrics and Gynecology*, 120, 1029–1036. doi:<https://doi.org/10.1097/AOG.0b013e31826d60c5>. [PubMed: 23090519]
- Centers for Disease Control and Prevention (2014). Pregnancy Mortality Surveillance System. Retrieved on September 9, 2018 from: <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pregnancy-mortality-surveillance-system.htm>.

- Clark SL, Christmas JT, Frye DR, Meyers JA, & Perlin JB (2014). Maternal mortality in the United States: predictability and the impact of protocols on fatal postcesarean pulmonary embolism and hypertension-related intracranial hemorrhage. *American Journal of Obstetrics and Gynecology*, 211, 32 e31–39. doi:10.1016/j.ajog.2014.03.031. [PubMed: 24631705]
- Clegg LX, Hankey BF, Tiwari R, Feuer EJ, & Edwards BK (2009). Estimating average annual per cent change in trend analysis. *Statistics in Medicine*, 28, 3670–3682. doi:10.1002/sim.3733. [PubMed: 19856324]
- Creanga AA (2018). Maternal mortality in the United States: A review of contemporary data and their limitations. *Clinical Obstetrics and Gynecology*, 61, 296–306. doi:10.1097/grf.0000000000000362. [PubMed: 29561285]
- Creanga AA, Bateman BT, Kuklina EV, & Callaghan WM (2014). Racial and ethnic disparities in severe maternal morbidity: a multistate analysis, 2008–2010. *American Journal of Obstetrics and Gynecology*, 210, 435e431–438. doi:10.1016/j.ajog.2013.11.039. [PubMed: 24295922]
- Creanga AA, Berg CJ, Syverson C, Seed K, Bruce FC, & Callaghan WM (2015). Pregnancy-related mortality in the United States, 2006–2010. *Obstetrics and Gynecology*, 125, 5–12. doi:10.1097/aog.0000000000000564. [PubMed: 25560097]
- Curtin SC, & Hoyert DL (2017). Maternal morbidity and mortality: Exploring racial/ethnic differences using new data from birth and death certificates In Hoque MN, Pecotte B, & McGehee MA (Eds.), *Applied Demography and Public Health in the 21st Century* (pp. 95–113). Cham: Springer International Publishing.
- Elixhauser A, Steiner C, Harris DR, & Coffey RM (1998). Comorbidity measures for use with administrative data. *Medical Care*, 36, 8–27. [PubMed: 9431328]
- Hashimoto RE, Brodt ED, Skelly AC, & Dettori JR (2014). Administrative database studies: Goldmine or goose chase? *Evidence-Based Spine-Care Journal*, 5, 74–76. doi:10.1055/s-0034-1390027. [PubMed: 25278880]
- Cost Healthcare and Project Utilization (2011). Introduction to the HCUP Nationwide Inpatient Sample (NIS) 2011. Retrieved from https://www.hcup-us.ahrq.gov/db/nation/nis/NIS_Introduction_2011.pdf.
- Cost Healthcare and Project Utilization (HCUP) (2015). NIS description of data elements. Retrieved from <https://www.hcup-us.ahrq.gov/db/nation/nis/nisdde.jsp>.
- Houchens RL, Ross D, & Elixhauser A. (2015). Using the HCUP National Inpatient Sample to Estimate Trends. HCUP Methods Series Report # 2006-05ONLINE, January 4, 2016. U.S. Agency for Healthcare Research and Quality. Retrieved from: <http://www.hcup-us.ahrq.gov/reports/methods/methods.jsp>.
- Jain JA, Temming LA, D'Alton ME, Gyamfi-Bannerman C, Tuuli M, Louis JM, ... Riley LE (2018). SMFM Special Report: Putting the “M” back in MFM: Reducing racial and ethnic in maternal morbidity and mortality: A call to action. *American Journal of Obstetrics and Gynecology*, 218, B9–B17. doi:10.1016/j.ajog.2017.11.591
- Keller C, Ainsworth B, Records K, Todd M, Belyea M, Vega-Lopez S, ... Nagle-Williams A (2014). A comparison of a social support physical activity intervention in weight management among postpartum Latinas. *BMC Public Health*, 14, 971. doi:10.1186/1471-2458-14-971 [PubMed: 25233867]
- Krieger N (2014). Discrimination and health inequities. *International Journal of Health Services*, 44(4), 643–710. doi:10.2190/HS.44.4.b [PubMed: 25626224]
- Kuklina EV, Whiteman MK, Hillis SD, Jamieson DJ, Meikle SF, Posner SF, & Marchbanks PA (2008). An enhanced method for identifying obstetric deliveries: implications for estimating maternal morbidity. *Maternal and Child Health Journal*, 12, 469–477. doi:10.1007/s10995-007-0256-6 [PubMed: 17690963]
- Li J, Luo C, Deng R, Jacoby P, & de Klerk N (2007). Maternal mortality in Yunnan, China: Recent trends and associated factors. *BJOG*, 114, 865–874. doi:10.1111/j.1471-0528.2007.01362.x. [PubMed: 17506792]
- Liese KL, Mogos M, Abboud S, Decocker K, Koch AR, & Geller SE (2019). Racial and ethnic disparities in severe maternal morbidity in the United States. *Journal of Racial and Ethnic Health Disparities*. doi:10.1007/s40615-019-00577-w.

- Louis JM, Menard MK, & Gee RE (2015). Racial and ethnic disparities in maternal morbidity and mortality. *Obstetrics and Gynecology*, 125, 690–694. doi:10.1097/aog.0000000000000704 [PubMed: 25730234]
- MacDorman MF, Declercq E, & Thoma ME (2017). Trends in maternal mortality by sociodemographic characteristics and cause of death in 27 states and the District of Columbia. *Obstetrics and Gynecology*, 129, 811–818. doi:10.1097/aog.0000000000001968 [PubMed: 28383383]
- Mann JR, Mannan J, Quinones LA, Palmer AA, & Torres M (2010). Religion, spirituality, social support, and perceived stress in pregnant and postpartum Hispanic women. *Journal of Obstetrics Gynecologic and Neonatal Nursing*, 39, 645–657. doi:10.1111/j.1552-6909.2010.01188.x
- Metcalfe A, Wick J, & Ronksley P (2018). Racial disparities in comorbidity and severe maternal morbidity/mortality in the United States: an analysis of temporal trends. *Acta Obstet Gynecol Scand*, 97, 89–96. doi:10.1111/aogs.13245 [PubMed: 29030982]
- Moaddab A, Dildy GA, Brown HL, Bateni ZH, Belfort MA, Sangi-Haghpeykar H, & Clark SL (2018). Health care disparity and pregnancy-related mortality in the United States, 2005–2014. *Obstetrics and Gynecology*, 131, 707–712. doi:10.1097/aog.0000000000002534 [PubMed: 29528919]
- Mogos MF, Piano MR, McFarlin BL, Salemi JL, Liese KL, & Briller JE (2018). Heart failure in pregnant women: A concern across the pregnancy continuum. *Circulation: Heart Failure*, 11, e004005. doi:10.1161/circheartfailure.117.004005. [PubMed: 29330153]
- Mogos MF, Salemi JL, Spooner KK, McFarlin BL, & Salihu HM (2016). Differences in mortality between pregnant and nonpregnant women after cardiopulmonary resuscitation. *Obstetrics and Gynecology*, 128, 880–888. doi:10.1097/aog.0000000000000000 [PubMed: 27607874]
- Murphy M, Alavi K, & Maykel J (2013). Working with existing databases. *Clinics in Colon and Rectal Surgery*, 26, 5–11. doi:10.1055/s-0033-1333627 [PubMed: 24436641]
- Negron R, Martin A, Almog M, Balbierz A, & Howell EA (2013). Social support during the postpartum period: Mothers' views on needs, expectations, and mobilization of support. *Maternal and Child Health Journal*, 17, 616–623. doi:10.1007/s10995-012-1037-4 [PubMed: 22581378]
- Martin N & Montagne R (2017). "Nothing protects black women from dying in pregnancy and childbirth," ProPublica, 12 7, 2017 Retrived from: <https://www.propublica.org/article/nothing-protects-black-women-from-dying-in-pregnancy-and-childbirth>.
- Rosenberg D, Geller SE, Studee L, & Cox SM (2006). Disparities in mortality among high risk pregnant women in Illinois: a population based study. *Annals of Epidemiology*, 16, 26–32. doi:10.1016/j.annepidem.2005.04.007 [PubMed: 16023371]
- Sullivan EA, Ford JB, Chambers G, & Slaytor EK (2004). Maternal mortality in Australia, 1973–1996. *Australian and New Zealand Journal of Obstetrics and Gynaecology*, 44, 452–457; discussion 377. doi:10.1111/j.1479-828X.2004.00313.x [PubMed: 15387869]
- Tucker MJ, Berg CJ, Callaghan WM, & Hsia J (2007). The black-white disparity in pregnancy-related mortality from 5 conditions: Differences in prevalence and case-fatality rates. *American Journal of Public Health*, 97, 247–251. doi:10.2105/AJPH.2005.072975 [PubMed: 17194867]
- Weiss AJ, & Elixhauser A (2006). Overview of hospital stays in the United States, 2012: Statistical brief #180 Healthcare Cost and Utilization Project (HCUP) Statistical Briefs. Rockville (MD): Agency for Healthcare Research and Quality (US).

Figure 1:
Distribution of pregnancy related hospitalizations and In-hospital mortality in the United States (NIS: 2002-2014)

Figure 2:

Trend in In-patient maternal mortality by timing in the pregnancy continuum in the United States: NIS (2002-2014)

APC = Annual Percent Change

Demographics, behavioral, and hospital characteristics of pregnancy-related deaths stratified by timing of pregnancy: NIS (2002-2014)

Table 1

Characteristics	a_n	C%	In-hospital pregnancy-related mortality by timing of pregnancy								
			Antepartum			Intrapartum			Postpartum		
			a_n	C%	R%	a_n	C%	R%	a_n	C%	R%
All	58,742,179	100%	1,798	100%	0.03	3,600	100%	0.01	2,013	100%	0.23
Age											
13-24	19,918,381	33.91	618	34.37	0.03	872	24.21	0.00	462	22.98	0.15
25-34	30,208,480	51.43	842	46.83	0.03	1,705	47.34	0.01	1,051	52.20	0.24
35-49	8,615,318	14.67	338	18.80	0.04	1,024	28.45	0.01	500	24.83	0.32
Race											
NH-white	24,716,530	42.08	598	33.29	0.03	1,082	30.04	0.00	572	28.44	0.16
NH-African American	6,997,714	11.91	468	26.06	0.05	961	26.70	0.02	639	31.72	0.38
Hispanic	11,029,179	18.78	267	14.86	0.03	623	17.32	0.01	251	12.46	0.18
Other	5,024,250	8.55	166	9.24	0.05	346	9.60	0.01	187	9.30	0.29
Alcohol											
Yes	100,779	0.17	55	3.07	0.14	25	0.70	0.04	14	0.71	0.30
No	58,641,400	99.83	1,743	96.93	0.03	3,575	99.30	0.01	1,999	99.29	0.23
Tobacco											
Yes	2,843,226	4.84	145	8.07	0.04	119	3.30	0.00	124	6.16	0.19
No	55,898,953	95.16	1,653	91.93	0.03	3,482	96.70	0.01	1,889	93.84	0.23
Drug abuse											
Yes	945,170	1.61	208	11.59	0.10	162	4.49	0.02	122	6.05	0.43
No	57,797,010	98.39	1,589	88.41	0.03	3,439	95.51	0.01	1,891	93.95	0.22
Household Income											
Highest quartile	12,979,032	22.09	194	10.80	0.02	554	15.39	0.00	317	15.74	0.18
Lowest quartile	15,971,946	27.19	658	36.61	0.04	1,157	32.13	0.01	784	38.93	0.29

Characteristics	a_n	C%	In-hospital pregnancy-related mortality by timing of pregnancy											
			Antepartum				Intrapartum				Postpartum			
			a_n	C%	R%	a_n	C%	R%	a_n	C%	R%	a_n	C%	R%
2nd quartile	14,554,198	24.78	534	29.70	0.04	991	27.51	0.01	485	24.12	0.22			
3rd quartile	14,133,460	24.06	366	20.35	0.03	792	22.00	0.01	382	18.98	0.19			
Primary payer														
Gov't	25,280,366	43.04	842	46.84	0.03	1,830	50.84	0.01	1,080	53.68	0.25			
Private	29,628,368	50.44	612	34.02	0.03	1,460	40.54	0.01	726	36.08	0.19			
Other	3,833,446	6.53	344	19.14	0.07	310	8.62	0.01	206	10.24	0.28			
Region of hospital														
Northeast	9,758,119	16.61	251	13.98	0.03	591	16.41	0.01	325	16.16	0.21			
Midwest	12,511,650	21.30	346	19.26	0.03	548	15.21	0.00	339	16.83	0.18			
South	22,259,017	37.89	807	44.87	0.04	1,663	46.19	0.01	999	49.61	0.28			
West	14,213,394	24.20	393	21.89	0.03	798	22.18	0.01	350	17.40	0.19			
Bed size of hospital														
Large	36,224,039	61.67	1,254	69.78	0.04	2,389	66.36	0.01	1,507	74.89	0.26			
Small	6,720,614	11.44	118	6.57	0.02	227	6.32	0.00	104	5.18	0.11			
Medium	15,544,115	26.46	409	22.77	0.03	968	26.90	0.01	390	19.38	0.17			
Location/teaching status of hospital														
Rural	6,649,507	11.32	99	5.52	0.02	210	5.84	0.00	82	4.06	0.10			
Urban, non-teaching	23,630,765	40.23	565	31.42	0.03	1,081	30.02	0.01	465	23.12	0.14			
Urban, teaching	28,208,496	48.02	1,118	62.17	0.04	2,294	63.71	0.01	1,455	72.28	0.30			

^aWeighted to estimate the national frequency; sum of all groups may not add up to the total because of missing data. NIS = Nationwide Inpatient Sample; C% = Column Percent; R% = Row Percent

Table 2

Demographics, behavioral, and hospital characteristics of In-hospital maternal mortality stratified by race group: NIS (2002-2014)

Characteristics	a_N	%	In-hospital Maternal Mortality by Race Group			p -value
			White a_n (%)	African-American a_n (%)	Hispanic a_n (%)	
All	58,742,179	100%	2,252 (30.4)	2,068 (27.9)	1,141 (15.4)	
Age						
13-24	19,918,381	33.91	576 (25.6)	588 (28.5)	283 (24.8)	0.34
25-34	30,208,480	51.43	1083 (48.1)	995 (48.1)	599 (52.5)	
35-49	8,615,318	14.67	594 (26.4)	485 (23.4)	259 (22.7)	
Alcohol (Yes)	100,779	0.17	41 (1.8)	25 (1.2)	14 (1.3)	0.67
Tobacco (Yes)	2,843,226	4.84	214 (9.5)	96 (4.7)	16 (1.4)	<0.01
Drug abuse (Yes)	945,170	1.61	222 (9.9)	88 (4.2)	55 (4.8)	0.01
Household Income						
Highest quartile	12,979,032	22.09	585 (26.0)	953 (46.1)	444 (38.9)	<0.01
Lowest quartile	15,971,946	27.19	631 (28.0)	507 (24.5)	356 (31.2)	
2nd quartile	14,554,198	24.78	511 (22.7)	384 (18.6)	194 (17.0)	
3rd quartile	14,133,460	24.06	477 (21.1)	155 (7.5)	108 (9.4)	
Primary payer						
Gov't	25,280,366	43.04	954 (42.4)	1251 (60.5)	707 (61.9)	<0.01
Private	29,628,368	50.44	1036 (46.0)	638 (30.9)	260 (22.7)	
Other	3,833,446	6.53	262 (11.6)	179 (8.7)	175 (15.3)	
Region of hospital						
Northeast	9,758,119	16.61	360 (16.0)	451 (21.8)	160 (14.0)	<0.01
Midwest	12,511,650	21.30	339 (15.0)	185 (8.9)	42 (3.7)	
South	22,259,017	37.89	1069 (47.4)	1248 (60.4)	447 (39.2)	
West	14,213,394	24.20	485 (21.5)	184 (8.9)	492 (43.1)	
Bed size of hospital						
Large	36,224,039	61.67	129 (0.9)	101 (1.0)	79 (1.6)	0.71
Small	6,720,614	11.44	543 (2.0)	524 (2.6)	260 (3.0)	
Medium	15,544,115	26.46	1576 (2.2)	1429 (2.7)	791 (3.4)	
Location/teaching status of hospital						
Rural	6,649,507	11.32	193	46	6	<0.01
Urban, non-teaching	23,630,765	40.23	712	490	445	
Urban, teaching	28,208,496	48.02	1343	1518	679	

^aWeighted to estimate the national frequency; sum of all groups may not add up to the total because of missing data. NIS = Nationwide Inpatient Sample; N= Sample Size; P = 0.05 considered statistically significant.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

Summary of percent distribution of Elixhauser comorbidities by timing of pregnancy among women age 15 – 49 years that died during pregnancy-related hospitalization in the United States (NIS 2002 – 2014).

Elixhauser Comorbidity	Timing of pregnancy-related hospitalization at the time of in-hospital mortality		
	AP (%)	IP (%)	PP (%)
AIDS/HIV	2.3	1.5	0.2
Alcohol Abuse	2.8	0.7	0.7
Anemia Deficiency	11.1	9.8	14.8
Depression	5.0	1.1	3.9
Diabetes without complication	4.8	3.7	6.4
Diabetes with complication	1.0	0.5	1.4
Drug Abuse	10.5	4.1	6.3
Hypertension	10.0	12.4	21.6
Hypothyroidism	3.0	1.6	3.4
Liver disease	2.4	1.6	1.6
Lymphoma	-	0.6	0.2
Fluid and Electrolyte Disorders	39.7	30.3	43.5
Metastatic Cancer	1.6	1.6	2.1
Neurological	13.0	12.2	15.6
Obesity	7.2	6.3	12.2
Paralysis	2.1	1.2	4.9
Peripheral Vascular Disease	0.6	2.0	1.5
Psychoses	3.5	0.8	1.2
Pulmonary circulation Disorder	5.3	3.6	5.1
Renal failure	3.0	1.9	6.7
Tumor without Metastasis	0.9	1.3	-
Peptic Ulcer Disease	-	-	-
Valvular Disease	3.3	1.8	5.7
Weight Loss	4.7	3.8	6.5
Rheumatoid Arthritis	1.6	1.7	2.9
Blood Loss Anemia	11.4	23.3	17.0
Congestive Heart Failure	5.0	3.7	20.4
Chronic Lung Disease	8.2	6.1	6.3
Coagulopathy	23.1	31.8	26.4

NIS = Nationwide Inpatient Sample; AP = Antepartum; IP = Intrapartum; PP = Postpartum. AIDS = Acquired Immune Deficiency Syndrome; HIV = Human Immunodeficiency Disease.