Published in final edited form as:

European J Org Chem. 2019 February 21; 2019(7): 1646–1648. doi:10.1002/ejoc.201801804.

Benylation of Imines with Activated Boronate Nucleophiles

Michael R. Hollerbach, Jacob C. Hayes, Timothy J. Barker

Department of Chemistry and Biochemistry, College of Charleston, Charleston, South Carolina 29424, USA

Abstract

Benzylation reactions of *N*-tosyl imines and *N*-*tert*-butanesulfinyl imines using benzylboronic acid pinacol ester are reported. *s*-Butyllithium was used to activate the boronic ester, rendering it nucleophilic. The reaction was compatible with electronically diverse substituents on the imine in both substrate classes. Good diastereoselectivity was observed in additions to *N*-*tert*-butylsulfinylaldehydes. The diastereoselectivity observed in these reactions is consistent with an open transition state for the addition. Examples of a secondary alkylboronic ester nucleophile and an *N*-*tert*-butanesulfinyl trifluoromethylketimine electrophile are also included.

Keywords

diastereoselectivity; boron; nucleophilic addition; chiral auxiliaries

Introduction

Organoboron reagents have been used extensively to form carbon-carbon bonds.^[1] The reaction between allylboron reagents and carbonyls or imines with and without catalysts has provided access to homoallylic alcohols and amines in high enantioselectivity and has been a very successful way to perform this transformation.^[2] Despite the success of allylboron reagents in 1,2-addition reactions, the use of benzylboron reagents and other alkylboron reagents in a similar reaction is less common.^{[3],[4],[5]}

Imines are an attractive class of electrophiles, because addition reactions between organometallic reagents and imines provide α -branched amines in a straightforward manner.^[6] Additionally, the diastereoselective addition to a chiral imine provides access to enantiomerically enriched amines, a direct approach to a medically relevant class of amines. Specifically, homobenzyl amines are present in a number of HIV protease inhibitors, including Fosamprenavir and Darunavir. Previously, benzyl Grignard reagents,^[7] benzylzinc reagents,^[7b, 8] benzyltrimethylsilanes,^[9] and tetrabenzylhafnium^[10] have been employed as nucleophiles with imines. A new synthetic approach, starting from a benchtop stable organometallic reagent such as an organosilicon or organoboron that upon activation can be rendered nucleophilic is attractive. A Rh-catalyzed addition of secondary benzylic trifluoroborates salts to *N*-sulfonylimines has also been reported.^[3c] Notably, *N*-*tert*-butanesulfinylimine were unreactive under these conditions. We sought to develop a 1,2-addition of benzylboronic esters to several classes of imines including *N*-tosylimines and *N*-*tert*-butanesulfinyl imines.

Alkylboronic esters can be activated with organolithium reagents to form the more nucleophilic alkylboronate species.^[11] Aggarwal has previously employed aryllithium reagents to activate alkylboronic esters and their subsequent reactions in numerous enantiospecific transformations.^[12] Previously, we have demonstrated benzylboronic acid pinacol ester (BnBpin) can be activated by an alkylolithium reagent, rendering it nucleophilic toward aldehydes.^[13]

Results and Discussion

Using our previously developed conditions for activation of BnBpin with *s*-BuLi employing *N*-tosylimines, good yields were observed of the amine products (Table 1). The reaction proved insensitive to substituents on the aromatic ring with both electron-donating and electron-withdrawing substituents well-tolerated, as seen in products 5–7. Additionally, the *o*-tolyl imine reacted well, affording product 8 in 72% yield. A reaction with the imine derived from furfural provided a 72% yield of the heteroaromatic amine product 9. A reaction using BnBF₃K in place of BnBpin under the standard conditions did not provide any of the desired product.

Next, (*R*)-*N*-*tert*-butanesulfinyl imines were examined as electrophiles, providing the opportunity for a diastereoselective reaction. As the *N*-*tert*-butanesulfinyl imines are not as activated as the *N*-tosyl imines, the reactions proceeded slower but provided the products in good yields across electronically diverse substrates (Table 2). This reaction with *N*-*tert*-butanesulfinyl imines was found to be more sensitive to substituent effects, with the imine derived from 4-trifluoromethylbenzaldehyde providing the highest yield of product 12 (83%) with a diastereomeric ratio of 91:9. The imine derived from furfural provided the desired amine product in 73% yield (86:14 d.r.). With the six *N*-*tert*-butanesulfinyl imines examined, the diastereomeric ratio observed ranged from 83:17–90:10. The diastereoselectivity observed was very close to that previously reported with benzyl zinc^[8a] and benzyl silicon^[9b] nucleophiles when using common substrates. The absolute configuration of the products was determined by comparison to previously reported data of the products that had rigorously established the configurations and by analogy for other examples.^[8a] With a proposed tetracoordinate boron nucleophile, the reaction is expected to proceed through an open transition state which is consistent with the observed stereochemistry of the products.^[8a] The previously mentioned Rh-catalyzed addition of alkylboron reagents did not work with *N*-*tert*-butanesulfinyl imines, suggesting these boronate are a more reactive nucleophile.^[3c]

To expand the scope of nucleophiles, the branched benzylboronic ester was also examined with an *N*-tosyl imine (Scheme 1). The product was isolated in a modest yield (51%, 2:1 d.r.). Previously this branched product had been made in a Rh-catalyzed reaction that also gave a 2:1 dr, however the major diastereomer is different under our reaction conditions.^[3c] This observed difference in selectivity can be attributed to a closed vs open transition state in the respective mechanisms.

Finally, an *N*-*tert*-butanesulfinyltrifluoromethylketamine was examined as a substrate to examine this nucleophile with a more sterically hindered imine electrophile (Scheme 2). The

reaction provided a 62% yield of the desired product. The diastereomeric ratio for the reaction was 63:37. While this reaction was less diastereoselective than in the additions to the aldimines, this reaction is one of the first examples of a benzyl addition to this class of ketamines to the best of our knowledge.

Conclusion

In summary, the nucleophilic 1,2-addition of benzylic boronic esters to imines has been demonstrated. Electron-donating and electron-withdrawing groups were well-tolerated with both imine classes. The diastereoselectivity observed in the reaction with the *N-tert*-butanesulfinylimines was consistent with that previously observed with other benzyl nucleophiles going through an open transition state. Branched alkylboronic esters was a competent nucleophile in the reaction, and a reaction with a trifluoromethylketimine was also demonstrated. Further studies on the use of alkylboronic esters as nucleophiles are underway in our laboratory.

Experimental Section

General Procedure for Benzyl Addition to *N*-Tosyl Imines: An oven dried round-bottom flask was charged with 4 Å molecular sieves and a magnetic stir bar, flame dried under vacuum, covered with a new septum and sealed thoroughly with Parafilm. The flask was allowed to cool under vacuum and purged three times with argon before adding benzyboronic acid pinacol ester (223 µL, 1.0 mmol, 2.0 equiv.) and 4 mL of THF. The contents were cooled to –78 °C under increased argon flow before addition of sec-butyllithium (0.75 mmol, 1.5 equiv., 1.2 M in hexanes) and allowed to stir at that temperature for 30 minutes. Subsequently, tosylimine (0.50 mmol, 1 equiv.) was added at –78 °C as a solution in THF (2 mL). The reaction was removed from the bath after an additional 15 minutes and allowed to warm to room temperature while stirring for 3 hours. The reaction was quenched with 0.2 ml sat. NH₄Cl and concentrated under reduced pressure and isolated by column chromatography, eluting with hexanes/ethyl acetate.

General Procedure for Benzyl Addition to *N-tert*-Butanesulfinyl Imines: An oven dried round-bottom flask was charged with 4 Å molecular sieves and a magnetic stir bar, flame dried under vacuum, covered with a new septum and sealed thoroughly with Parafilm. The flask was allowed to cool under vacuum and purged three times with argon before adding benzyboronic acid pinacol ester (223 µL, 1.0 mmol, 2.0 equiv.) and 4 mL of THF. The contents were cooled to –78 °C under increased argon flow before addition of sec-butyllithium (0.75 mmol, 1.5 equiv., 1.2 M in hexanes) and allowed to stir at that temperature for 30 minutes. Subsequently, *tert*-butylsulfinyl imine (0.50 mmol, 1 equiv.) was added at –78 °C as a solution in THF (2 mL). The reaction was removed from the bath after an additional 15 minutes and allowed to warm to room temperature while stirring for 4 hours. The reaction was quenched with 0.2 mL sat. NH₄Cl and concentrated under reduced pressure. The diastereomeric ratio of the product was determined by inverse-gated no NOE ¹³C NMR^[14] and ¹H NMR of the unpurified material before being isolated by column chromatography, eluting with hexanes/ethyl acetate.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

MRH acknowledges support for a summer stipend funded by the Howard Hughes Medical Institute to the College of Charleston as part of their 2012 Undergraduate Science Education Competition and the National Center for Research Resources (5 P20 RR016461) and the National Institute of General Medical Sciences (8 P20 GM103499) from the NIH. JCH acknowledges support for a summer stipend funded by the College of Charleston School of Science and Math Dean's Research Fund. The students enrolled in Advanced Synthesis Lab 371 during the fall of 2017 are acknowledged for synthesizing the imines used in this paper. The College of Charleston is acknowledged for additional financial support. The NMR spectrometer at the College of Charleston is supported by the National Science Foundation under Grant No. 1429308.

References

- [1]. Suzuki A, Angew. Chem. Int. Ed 2011, 50, 6722–6737.
- [2]. a)Hall DG, Lachance H, Allylboration of Carbonyl Compounds, Wiley, Hoboken, N. J., 2012;b)Diner C, Szabó KJ, J. Am. Chem. Soc 2017, 139, 2–14; [PubMed: 27966922] c)Yus M, Gonzalez-Gomez JC, Foubelo F, Chem. Rev 2013, 113, 5595–5698; [PubMed: 23540914] d)Hall DG, Lee JCH, Ding J, Pure Appl. Chem 2012, 84, 2263–2277;e)Huo H-X, Duvall JR, Huang M-Y, Hong R, Org. Chem. Front 2014, 1, 303–320;f)Hatano M, Ishihara K, Synthesis 2008, 1647–1675;g)Kennedy JWW, Hall DG, Angew. Chem. Int. Ed 2003, 42, 4732–4739;h)Ramachandran PV, Burghardt TE, Pure Appl. Chem 2006, 78, 1397–1406;i)Ramachandran PV, Aldrichimica Acta 2002, 35, 23–35;j)Hall DG, Synlett 2007, 1644–1655.k)Jonnalagadda SC, Suman P, Patel A, Jampana G, Colfer A, in Boron Reagents in Synthesis, Vol. 1236, American Chemical Society, 2016, pp. 67–122.
- [3]. a)Ros A, Aggarwal VK, Angew. Chem. Int. Ed 2009, 48, 6289–6292;b)Zhang C, Yun J, Org. Lett 2013, 15, 3416–3419; [PubMed: 23773146] c)Lee S, Lee WL, Yun J, Adv. Synth. Catal 2015, 357, 2219–2222.
- [4]. Reichle Markus A, Breit B, Angew. Chem. Int. Ed 2012, 51, 5730–5734.
- [5]. a)Li Y, Zhou K, Wen Z, Cao S, Shen X, Lei M, Gong L, J. Am. Chem. Soc 2018, 140, 15850–15858; [PubMed: 30372057] b)Plasko DP, Jordan CJ, Ciesa BE, Merrill MA, Hanna JM Jr., Photochem. Photobiol. Sci 2018, 17, 534–538. [PubMed: 29722411]
- [6]. a)Ellman JA, Owens TD, Tang TP, Acc. Chem. Res 2002, 35, 984–995; [PubMed: 12437323] b)Robak MT, Herbage MA, Ellman JA, Chem. Rev 2010, 110, 3600–3740. [PubMed: 20420386]
- [7]. a)Harried SS, Croghan MD, Kaller MR, Lopez P, Zhong W, Hungate R, Reider PJ, J. Org. Chem 2009, 74, 5975–5982; [PubMed: 19586010] b)Hatano M, Suzuki S, Ishihara K, J. Am. Chem. Soc 2006, 128, 9998–9999. [PubMed: 16881613]
- [8]. a)Buesking AW, Baguley TD, Ellman JA, Org. Lett 2011, 13, 964–967; [PubMed: 21306151] b)Almansa R, Guijarro D, Yus M, Tetrahedron: Asymmetry 2008, 19, 2484–2491.
- [9]. a)Zhang WX, Ding CH, Luo ZB, Hou XL, Dai LX, Tetrahedron Lett 2006, 47, 8391–8393;b)Das M, O'Shea Donal F, Chem. Eur. J 2015, 21, 18717–18723. [PubMed: 26558552]
- [10]. Mei HB, Ji XY, Han JL, Pan Y, Eur. J. Org. Chem 2011, 5783–5786.
- [11]. For an example of alkyllithium activation of alkylboronic esters for Pd cross-coupling, see: Zou G, Falck JR, Tetrahedron Lett 2001, 42, 5817–5819.
- [12]. a)Sandford C, Aggarwal VK, Chem. Commun 2017, 53, 5481–5494;b)Feeney K, Berionni G, Mayr H, Aggarwal VK, Org. Lett 2015, 17, 2614–2617. [PubMed: 25973673] c)Larouche-Gauthier R, Elford TG, Aggarwal VK, J Am Chem Soc 2011, 133, 16794–16797; [PubMed: 21939203] d)Sandford C, Rasappan R, Aggarwal VK, J. Am. Chem. Soc 2015, 137, 10100–10103; [PubMed: 26244235] e)Garcia-Ruiz C, Chen JLY, Sandford C, Feeney K, Lorenzo P, Berionni G, Mayr H, Aggarwal VK, J. Am. Chem. Soc 2017, 139, 15324–15327. [PubMed: 29028321]
- [13]. Hollerbach MR, Barker TJ, Organometallics 2018, 37, 1425–1427. [PubMed: 31680714]

- [14]. Otte DAL, Borchmann DE, Lin C, Weck M, Woerpel KA, Org. Lett 2014, 16, 1566–1569.
[PubMed: 24601654]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Scheme 1.
Addition with a Branched Nucleophile.

Scheme 2.
N-*tert*-Butanesulfinyl Ketimine Electrophile.

Table 1.

N-Tosyl Imines Substrate Scope.

Table 2.

N-*tert*-Butanesulfinylimine Substrate Scope.