Published in final edited form as:

Adv Exp Med Biol. 2020 ; 1131: 681–697. doi:10.1007/978-3-030-12457-1_27.

Readily releasable stores of calcium in neuronal endolysosomes: Physiological and pathophysiological relevance

Koffi L. Lakpa, Peter Halcrow, Xuesong Chen, Jonathan D. Geiger¹

Department of Biomedical Sciences, University of North Dakota School of Medicine and Health Sciences, 504 Hamline Street, Grand Forks, North Dakota 58203

Abstract

Neurons are long-lived post-mitotic cells that possess an elaborate system of endosomes and lysosomes (endolysosomes) for protein quality control. Relatively recently, endolysosomes were recognized to contain high concentrations (400 to 600 μM) of readily releasable calcium. The release of calcium from this acidic organelle store contributes to calcium-dependent processes of fundamental physiological importance to neurons including neurotransmitter release, membrane excitability, neurite outgrowth, synaptic remodeling, and cell viability. Pathologically, disturbance of endolysosome structure and/or function have been noted in a variety of neurodegenerative disorders including Alzheimer's disease (AD) and HIV-1 associated neurocognitive disorder (HAND). And, dysregulation of intracellular calcium has been implicated in the neuropathogenesis of these same neurological disorders. Thus, it is important to better understand mechanisms by which calcium is released from endolysosomes as well as the consequences of such release to inter-organelle signaling, physiological functions of neurons, and possible pathological consequences. In doing so, a path forward towards new therapeutic modalities might be facilitated.

Keywords

Endosomes; Lysosomes; Endolysosomes; Calcium; Store-operated calcium entry; N-type calcium channels; Neurodegenerative diseases; HIV-1 associated neurocognitive disorder; Alzheimer's disease; Neurons

An evolutionary perspective on calcium, intracellular organelles and endolysosomes

Intracellular calcium regulates many essential functions of neurons including neurotransmitter release, excitability, synaptic plasticity, and cell viability¹. Levels of intraneuronal calcium are very tightly regulated both temporally and spatially by various mechanisms including calcium release from intracellular stores, calcium influx across

¹Please address correspondence to: Jonathan D. Geiger, Ph.D., Chester Fritz Distinguished Professor, Department of Biomedical Sciences, University of North Dakota School of Medicine and Health Sciences, 504 Hamline Street, Room 110, Grand Forks, North Dakota 58203, (701) 777-2183, Jonathan.geiger@med.und.edu.

plasma membranes, and its association with a whole host of calcium binding proteins. Because of its importance both physiologically and pathologically, we start our story about the presence and functional significance of readily releasable stores of calcium in neuronal endolysosomes with a brief evolutionary perspective about calcium and intracellular organelles.

Calcium is well-known to be important for signal transduction in most cells including neurons. Indeed, calcium has been referred to as a universal second messenger in eukaryotic cells. The approximate 10,000-fold gradient of extracellular to intracellular calcium originated evolutionarily because of the gradual rise in calcium levels from about 100 nM during the period when the basic building blocks of life developed in thermal ducts under the ocean floor to about 1 mM during the Pre-Cambrian period when multicellular life evolved^{2,3}. Due to the toxic nature of millimolar levels of calcium, evolutionary pressure was applied such that cellular survival dictated that semipermeable membranes appeared and a variety of mechanisms were formed to maintain appropriate calcium gradients across plasma membranes³. Simultaneously, embedded in the plasma membranes were newly developed calcium pumps and calcium binding proteins which helped with calcium homeostasis³. Together, in neurons, these evolutionary changes provide unique and complex spatial and temporal handling of calcium that is essential for not only proper cellular signaling but also neuronal cell life and death.

It was also during this billion-year evolutionary period that intracellular organelles began appearing including mitochondria resulting from symbiotic relationships with bacteria and the development of functional endocytic machinery⁴. Mitochondria are integral to the maintenance of cellular energetics and they are important 'sinks' for intracellular calcium⁵. However, when too much calcium is up-taken into mitochondria cellular energetics are compromised and the resulting calcium overload can lead to a cascade of events including increased oxidative stress and cell death. It has also become increasingly appreciated that organelles including endoplasmic reticulum, endosomes and lysosomes (hereafter referred to as endolysosomes) have readily releasable and functionally important pools of intracellular calcium. Although less well known, the approximate 500 μ M levels of calcium in endolysosomes are similar to the calcium concentrations present in endoplasmic reticulum⁶. This is a very important concept because endoplasmic reticulum is commonly referred to as the principal intracellular store of readily releasable calcium. Furthermore, as the field of inter-organellar signaling as well as physical and chemical crosstalk between organelles has grown over the past decade it is prudent of us to now posit that this relatively new and highly complicated area of modern cell biology is key to our understanding of the regulation and dysregulation of calcium⁷.

With this as a very quick trip across 1 billion years of evolutionary biology, here we embark on a brief but focused summary of findings that neuronal endolysosomes contain readily releasable stores of calcium and once released this calcium can lead to calcium influx into cells, calcium release from other organelles, and calcium dysregulation-induced neurotoxicity. The relevance of such an important upstream store of calcium to the regulation of physiological functions and pathophysiological events is obvious and will be

discussed with particular relevance to the pathogenesis of two neurodegenerative disorders; Alzheimer's disease (AD) and HIV-1 associated neurocognitive disorder (HAND).

Endolysosomes contain readily releasable pools of calcium

Neurons are long-lived post-mitotic cells that possess an elaborate endolysosome system for quality control especially for proteins. Endolysosomes are well known to be acidic organelles that contain high levels of cations including calcium, iron, zinc and copper. However, for the cation calcium it was not until fairly recently that these organelles were described as being 'acidic calcium stores' because the luminal pH of endolysosomes is acidic and endolysosomes contain high (400 to 600 μM) levels of readily releasable calcium^{8,9}.

Neuronal calcium signals display spectacular spatiotemporal complexity and understanding how calcium signals are generated spatially and temporally is necessary to understand calcium-dependent cellular processes. Endolysosome calcium levels are maintained by a variety of uptake and efflux mechanisms. Essential for uptake of calcium into endolysosomes, proton gradients are established mainly by vacuolar H^+ -ATPase (v-ATPase) that pumps H^+ into the lumen and this drives Ca^{2+} uptake by $\text{Ca}^{2+}/\text{H}^+$ exchange^{9,10,11}. Four main mechanisms for calcium release from endolysosomes have been described including: (1) Calcium release through two-pore channels (TPCs) triggered by nicotinic acid adenine dinucleotide phosphate (NAADP)^{12,13,14,15,16,17}; (2) Elevation of endolysosome pH with, for example, the selective v-ATPase inhibitor bafilomycin (BAF) or the alkaline lysosomotropic agents NH_4Cl and chloroquine^{8,18,19}; (3) Involvement of TRPML1 mucolipin-type channels and P2X4 receptors^{20,21,22}; and (4) Selective disruption of endolysosome membranes with Gly-Phe- β -naphthylamide (GPN)^{23,24}. Of physiological significance, calcium released from endolysosomes has been shown to contribute to a variety of calcium-dependent neuronal processes including neurotransmitter release, neuronal excitability, neurite outgrowth, synaptic remodeling, and cell viability^{25,26,27}.

Endolysosomes can release calcium transiently and in a highly localized and distinct fashion^{17,28,29}. Endolysosome calcium can affect the release of calcium from organellar stores as well as through plasma membrane-based calcium influx mechanisms. The inter-organellar signaling and signaling with the plasma membrane is explained at least in part by findings that endolysosomes are highly mobile in cells, are highly dynamic metabolically, have high rates of biogenesis, and can interact physically and functionally with other intracellular organelles.

At least three models of acidic store-induced calcium signaling mechanisms have been described⁹. (1) Acidic stores of calcium might communicate with endoplasmic reticulum calcium stores such that calcium released from endolysosomes can enhance endoplasmic reticulum calcium loading³⁰ and calcium-induced calcium release^{13,15}. (2) Changes in endolysosome pH may release calcium from a subgroup of acidic calcium stores and the released calcium may affect other subgroups of acidic stores through mechanisms such as vesicular fusion of late endosomes and lysosomes^{9,15,31}. (3) Calcium released from acidic

calcium stores might depolarize plasma membranes, evoke calcium-dependent currents, and stimulate calcium influx across plasma membranes¹².

Acidic store-operated calcium entry in neurons

Acidic store-operated calcium entry (aSOCE) is a unique mechanism that links readily releasable calcium in endolysosomes with influx of extracellular calcium into neurons. This is a novel means by which intraneuronal stores of calcium can contribute to spatial and temporal integration of calcium signaling. In support of this novel mechanism, we found that calcium could be released from endolysosomes following stimulation of a number of different mechanisms, that the calcium release could be independent of other organellar stores of calcium, that release of calcium from endolysosomes triggered calcium influx, and that the calcium influx was regulated by N-type calcium channels and lysosome exocytosis.

Capacitative influx of calcium into cells was described over 30 years ago³². Such calcium influx mechanisms, that are now commonly referred to as store-operated calcium entry (SOCE), are principally initiated by a reduction in endoplasmic reticulum calcium stores followed by influx of extracellular calcium in a variety of cells including neurons in order to refill the depleted stores of calcium. Mechanistically, depleting endoplasmic reticulum calcium stores drives the oligomerization and translocation of stromal interaction molecule 1 (STIM1) proteins to endoplasmic reticulum junctions close to the plasma membrane. Such STIM1 translocation induces the clustering of calcium release-activated calcium modulator 1 (Orai1) channels and/or transient receptor potential (TRP) cation channels into plasma membranes thereby enabling extracellular calcium entry³³.

Conceptually, but not mechanistically, we observed similar store-operated calcium entry involving endolysosomes in neurons. Using primary cultures of rat cortical neurons, we found that calcium was released from endolysosomes following treatment with the two-pore channel agonist NAADP-AM, the v-ATPase inhibitor BAF, and the lysosomotropic agent GPN; all of which de-acidify endolysosomes³⁴. However, when these experiments were conducted in the absence of extracellular calcium, de-acidification of endolysosomes with NAADP-AM, BAF and GPN increased only slightly levels of free cytosolic calcium. When these same experiments were conducted in the presence of extracellular calcium, NAADP-AM, BAF and GPN all increased significantly the levels of free cytosolic calcium. Although it is not well understood currently, the relatively small release of calcium from endolysosomes causes a much larger influx of extracellular calcium and this might be due to plasma membrane depolarization as is accompanied by NAADP-induced endolysosome calcium release^{12,34}. Besides neurons, phenomena similar to aSOCE have been described in other cell types where NAADP has been found to induce endolysosome calcium release and large influxes of calcium across plasma membranes^{12,36,37,38,39}. These observations suggested to us that endolysosome de-acidification by three completely different mechanisms led directly or indirectly to an enhanced influx of calcium into neurons. Accordingly, we next tested more specifically the extent to which a store-operated mechanism might control the observed calcium influx across the plasma membrane.

Using approaches similar to those used by others and us, we began studying store-operated calcium mechanisms including the classical endoplasmic reticulum-based capacitative SOCE. Indeed, we confirmed that in the absence of extracellular calcium and following depletion of endoplasmic reticulum calcium with the SERCA pump inhibitor thapsigargin (TG) there was a significant increase in levels of free intracellular calcium only when calcium was re-introduced to the extracellular medium. With this positive control for the functional presence of endoplasmic reticulum-based SOCE in our cultured neurons, we conducted similar experiments with agents that de-acidify endolysosomes and release calcium from endolysosome stores. Even after depleting ER pools of calcium with TG, application of NAADP-AM, BAF and GPN still caused increased influx of extracellular calcium and still induced increased levels of intracellular calcium. Thus, in these neurons there appeared to be at least two separate and functional store-operated calcium mechanisms; one governed by endoplasmic reticulum and the other by endolysosomes.

In testing the distinctive nature of the two store-operated calcium mechanisms governed by endoplasmic reticulum or endolysosomes we used pharmacological and molecular/genetic strategies. Using siRNA to knock-down protein expression levels of STIM1, a protein that is central to SOCE, and the SOCE blockers SKF-96365 and 2-APB we were able to block significantly TG-induced release of calcium from endoplasmic reticulum, but we were unable to block significantly NAADP-AM-, BAF- and GPN-induced calcium influx. However, we were able to block significantly NAADP-AM-, BAF- and GPN-induced calcium influx with the selective N-type calcium channel (NTCC) blocker (ω -conotoxin). The selective and specific nature of this inhibition by ω -conotoxin was confirmed further by showing that NAADP-AM-, BAF- and GPN-induced calcium influx was not blocked by inhibitors of L-type (nimodipine, verapamil) and P/Q-type (ω -agatoxin) calcium channels. Moreover, we confirmed these pharmacological findings by showing that siRNA knockdown of NTCCs attenuated significantly NAADP-AM-, BAF- and GPN-induced calcium influx, but did not affect TG-induced SOCE. Together, the above results demonstrated that calcium released from endolysosomes can be distinct from calcium released from endoplasmic reticulum through SOCE mechanisms and that the calcium released from endolysosomes is capable of activating cell surface calcium channels to stimulate calcium influx. These findings support and extend earlier findings that calcium released from endolysosomes did not stimulate endoplasmic reticulum-dependent SOCE in MDCK epithelial cells²³. Accordingly, this new mechanism was termed by us as “acidic store-operated calcium entry” (aSOCE)³⁴.

Role of lysosome exocytosis in acidic store-operated calcium entry (aSOCE)

Multiple mechanisms might control aSOCE involving NTCCs. One such mechanism might involve lysosome exocytosis because we have shown using a quantitative biotinylation of surface proteins assay that NAADP-AM, BAF and GPN all increased cell surface protein expression levels of NTCCs and lysosome-associated glycoprotein 1 (LAMP1). Next, we addressed the possibility that lysosome exocytosis and NTCCs were linked directly by conducting co-immunoprecipitation studies and found a physical interaction between

NTCCs and LAMP1. Because LAMP1 is critical for lysosome exocytosis⁴⁰, those observations suggested to us that lysosome exocytosis might be a functional partner in aSOCE especially because aSOCE was inhibited following siRNA knockdown of protein expression levels of LAMP1. Thus, de-acidification of endolysosomes might be of central importance because NAADP-AM, BAF and GPN through very different initial mechanisms all appeared to enhance lysosome exocytosis and the recycling of NTCCs to the plasma membrane where they participated in calcium influx generally and aSOCE more specifically. Physically, this makes sense as well because of findings that de-acidification of endolysosomes changes cellular distribution patterns of these organelles from a mostly perinuclear pattern to one where the endolysosomes migrate close to the plasma membrane⁴¹. Thus, functionally and physically there is evidence favoring endolysosomes and endolysosome exocytosis in calcium entry.

Physical interactions and functional relevance of inter-organelle signalling

In addition to physical interactions between endolysosomes and plasma membranes, it is becoming increasingly clear that endolysosomes physically and functionally interact as well with other intracellular organelles including mitochondria and endoplasmic reticulum. Such recognition has led to an appreciation for dynamic physical and chemical communications between intracellular organelles including those regulated by pH and calcium.

Physical interactions between mitochondria and endoplasmic reticulum were first described about 60 years ago and the functional significance of mitochondria-associated membranes was first characterized about 30 years ago⁴². Even today, there continues to be work focused on the physical and functional interactions between organelles⁴¹ as well as the role that organellar interactions plays in the pathogenesis of neurodegenerative diseases^{44,45}. As it relates to endolysosomes, it is now known that there are extensive physical interactions between endolysosomes and mitochondria and that these inter-organelle communications participate in lipid and metabolite exchange as well as mitochondrial quality control⁴⁶. Conversely, mitochondrial dysfunction has been found to negatively affect lysosome structure and function through reactive oxygen species-dependent mechanisms⁴⁷. Extensive membrane contact sites have been described between lysosomes and endoplasmic reticulum, that these contact sites were evolutionarily conserved, and that calcium released from lysosomes was sufficient to stimulate endoplasmic reticulum-dependent calcium-induced calcium release^{48,49,50}. However, only recently was it shown that endolysosomes maintain their thousand-fold calcium concentration gradient in cells in part by refilling endolysosome stores of calcium from IP₃-regulated stores of calcium in endoplasmic reticulum⁵¹. Some of the differences in findings as to calcium movements between organelles might be because of cell-specific mechanisms. In addition, the difficult nature of understanding inter-organelle calcium dynamics is highlighted by work showing that STIM1 and STIM2 are expressed in endolysosomes, at least in platelets, and that depletion of acidic organellar stores of calcium can increase protein-protein interactions between STIM proteins with Orai1 and TRPC channels to induce SOCE⁵². It is further complicated by findings that calcium released through endolysosome-resident TRML1 channels can cause calcium release from endoplasmic reticulum and calcium influx⁵³ and that NAADP has been implicated in this “cross-talk”⁵⁴.

Possible role of endolysosomes and aSOCE in pathogenesis of Alzheimer's disease and HIV-1 associated neurocognitive disorder (HAND)

Disturbance in endolysosome structure and/or function have been noted in a variety of neurodegenerative disorders including Alzheimer's disease (AD) and HIV-1 neurocognitive disorder (HAND)^{55,56,57,58,59}. AD is a devastating age-related neurodegenerative disease that is the commonest cause of dementia in people over the age of 65. People with HAND, on the other hand, exhibit neurological complications ranging from mild (mild cognitive impairment) to severe (dementia). In the current era of anti-retroviral therapeutics HIV-1 infected individuals are living almost full life-spans, but are now experiencing a prevalence rate of over 50% for HAND^{60,61}. Clinically and pathologically people living with neuroHIV-1 are exhibiting AD-like symptoms including learning and memory deficits as well as increased amyloidogenesis. Although the pathogenesis of HAND is not fully understood, HIV-1 proteins including the HIV-1 transactivator of transcription protein Tat have been implicated by others and us to be causative virotoxins in HAND^{62,63,64,65,66,67,68,69}. Among the HIV-1 viral proteins, HIV-1 Tat is present in brains of HIV-1 infected individuals and its levels stay elevated in CSF even when HIV-1 viral levels are immeasurable⁷⁰. Others and we have shown that HIV-1 Tat directly excites neurons^{65,71,72}, disturbs neuronal calcium homeostasis^{64,73}, disrupts synaptic integrity^{74,75}, and induces neurotoxicity^{68,76}.

Endolysosome dysfunction has been implicated in the development of at least two pathological hallmarks of AD and HAND; A β accumulation and neurofibrillary tangle formation. Endolysosomes are very important for amyloidogenic processing of A β PP to A β because amyloid β precursor protein (A β PP) is first endocytosed, the amyloidogenic enzymes BACE-1 and γ -secretase are almost exclusively located in endosomes and lysosomes, the acidic environment of endolysosomes is favorable for amyloidogenic metabolism of A β PP, and A β can be either accumulated in or released by exocytosis from endolysosomes^{77,78,79,80,81,82,83}. Tau is a microtubule-associated protein, and when hyperphosphorylated it aggregates and contributes to the formation of neurofibrillary tangles. Tau aggregates can be degraded by cathepsin D in autophagosomes-lysosomes^{84,85}, and endolysosome dysfunction contributes to tau aggregation and neurofibrillary tangle formation^{86,87}. On the other hand, transcriptional activation of lysosome biogenesis can clear aggregated tau⁸⁸. Thus, endolysosomes are important sites for development these neurological disorders.

Dysregulation of intracellular calcium has also been implicated in the neuropathogenesis of these same neurological disorders. And it is clear (see above) that de-acidification of endolysosomes releases calcium from these acidic stores^{28,89,90}. We found that HIV-1 Tat protein elevated endolysosome pH and disturbed the structure and function of endolysosomes⁷⁴, a prominent and early pathological feature of HAND^{57,58}. Clearly, endolysosome calcium stores contribute to neuronal calcium signaling and function^{91,92,93} and calcium release from endolysosomes triggers calcium release from endoplasmic reticulum^{11,17} and through plasma membranes via aSOCE (see above).

HIV-1, HIV-1 proteins including HIV-1 Tat, and anti-retroviral therapeutic drugs contribute to the development of AD-like pathology including increases in A β levels^{94,95,96,97,98,99}. HIV-1 Tat enters neurons via receptor-mediated endocytosis^{100,101,102}. The Tat-induced deacidification of endolysosomes and resulting effects on calcium dyshomeostasis likely results from the ability of HIV-1 Tat to decrease the levels and activity of vacuolar-ATPase as well as compensatory increases in cathepsin D and LAMP-1¹⁰¹. The consequences of such alterations in calcium dynamics and homeostasis are synaptic disruption and neurotoxicity^{104,105,106}.

Endolysosomes contain physiologically important levels of calcium that is readily releasable by a number of stimuli and insults. The calcium can exit through a number of channels including TRPML and two pore channels. Once released the calcium can signal other organelles to release calcium and for greater influx of calcium through plasma membrane-resident calcium channels especially N-type calcium channels. These effects on endolysosome structure and function have clear implications to the pathogenesis of AD and HAND; neurological disorders that show overlap in terms of clinical and pathological features. We are excited to be part of this emerging area of cell biology focused on inter-organellar signaling and look forward to further studies elucidating physiological and pathological consequences of calcium release from endolysosome stores.

Acknowledgements

The authors gratefully acknowledge the funding provided by the NIH for our work; P30GM103329, R01MH100972, R21DA040519, and R01MH105329

Reference List

1. Berridge MJ, Bootman MD, Roderick HL. Calcium signalling: Dynamics, homeostasis and remodelling. Vol. 4, Nature Reviews Molecular Cell Biology. 2003 p. 517–29. [PubMed: 12838335]
2. Jaiswal JK. Calcium – how and why? J Biosci. 2001;26(3):357–63. [PubMed: 11568481]
3. Case RM, Eisner D, Gurney A, Jones O, Muallem S, Verkhatsky A. Evolution of calcium homeostasis: From birth of the first cell to an omnipresent signalling system. Cell Calcium. 2007;42(4–5):345–50. [PubMed: 17574670]
4. Wideman JG, Leung KF, Field MC, Dacks JB. The Cell Biology of the Endocytic System from an Evolutionary Perspective. Cold Spring Harb Perspect Biol [Internet]. 2014 4 1 [cited 2018 Jul 16];6(4):a016998–a016998. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24478384> [PubMed: 24478384]
5. Carafoli E The fateful encounter of mitochondria with calcium: How did it happen? Biochim Biophys Acta - Bioenerg [Internet]. 2010 6 1 [cited 2018 Jul 16];1797(6–7):595–606. Available from: <https://www.sciencedirect.com/science/article/pii/S0005272810001301#fig1>
6. Patel S, Cai X. Evolution of acidic Ca²⁺-stores and their resident Ca²⁺-permeable channels [Internet]. Vol. 57, Cell Calcium. Churchill Livingstone; 2015 [cited 2018 Jul 14]. p. 222–30. Available from: <https://www.sciencedirect.com/science/article/pii/S0143416014002012> [PubMed: 25591931]
7. Raffaello A, Mammucari C, Gherardi G, Rizzuto R. Calcium at the Center of Cell Signaling: Interplay between Endoplasmic Reticulum, Mitochondria, and Lysosomes. 2016 [cited 2017 May 17]; Available from: [http://www.cell.com/trends/biochemical-sciences/pdf/S0968-0004\(16\)30147-5.pdf](http://www.cell.com/trends/biochemical-sciences/pdf/S0968-0004(16)30147-5.pdf)
8. Christensen K a, Myers JT, Swanson J a. pH-dependent regulation of lysosomal calcium in macrophages. J Cell Sci. 2002;115(Pt 3):599–607. [PubMed: 11861766]

9. Morgan AJ, Platt FM, Lloyd-Evans E, Galione A. Molecular mechanisms of endolysosomal Ca²⁺ signalling in health and disease. *Biochem J* [Internet]. 2011;439(3):349–74. Available from: <http://www.biochemj.org/content/439/3/349.abstract> [PubMed: 21992097]
10. Moreno SNJ, Docampo R. The role of acidocalcisomes in parasitic protists. In: *Journal of Eukaryotic Microbiology*. 2009 p. 208–13.
11. Patel S, Docampo R. Acidic calcium stores open for business: Expanding the potential for intracellular Ca²⁺-signaling. Vol. 20, *Trends in Cell Biology*. 2010 p. 277–86. [PubMed: 20303271]
12. Brailoiu E, Churamani D, Cai X, Schrlau MG, Brailoiu GC, Gao X, et al. Essential requirement for two-pore channel 1 in NAADP-mediated calcium signaling (a). *J Cell Biol*. 2009;186(2):201–9. [PubMed: 19620632]
13. Calcraft PJ, Ruas M, Pan Z, Cheng X, Arredouani A, Hao X, et al. NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature*. 2009;459(7246):596–600. [PubMed: 19387438]
14. Zong X, Schieder M, Cuny H, Fenske S, Gruner C, Rötzer K, et al. The two-pore channel TPCN2 mediates NAADP-dependent Ca²⁺-release from lysosomal stores. *Pflügers Arch Eur J Physiol* [Internet]. 2009 9 26 [cited 2018 Jul 14];458(5):891–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19557428> [PubMed: 19557428]
15. Ruas M, Rietdorf K, Arredouani A, Davis LC, Lloyd-Evans E, Koegel H, et al. Purified TPC Isoforms Form NAADP Receptors with Distinct Roles for Ca²⁺-Signaling and Endolysosomal Trafficking. *Curr Biol* [Internet]. 2010 4 27 [cited 2018 Jul 27];20(8):703–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20346675> [PubMed: 20346675]
16. Schieder M, Rötzer K, Brüggemann A, Biel M, Wahl-Schott CA. Characterization of two-pore channel 2 (TPCN2)-mediated Ca²⁺ currents in isolated lysosomes. *J Biol Chem* [Internet]. 2010 7 9 [cited 2018 Jul 14];285(28):21219–22. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20495006> [PubMed: 20495006]
17. Zhu MX, Ma J, Parrington J, Calcraft PJ, Galione A, Evans AM. Calcium signaling via two-pore channels: local or global, that is the question. *AJP Cell Physiol* [Internet]. 2010;298(3):C430–41. Available from: <http://ajpcell.physiology.org/cgi/doi/10.1152/ajpcell.00475.2009>
18. Camacho M, Machado JD, Alvarez J, Borges R. Intravesicular calcium release mediates the motion and exocytosis of secretory organelles: A study with adrenal chromaffin cells. *J Biol Chem*. 2008;283(33):22383–9. [PubMed: 18562320]
19. Machado JD, Camacho M, Alvarez J, Borges R. On the role of intravesicular calcium in the motion and exocytosis of secretory organelles. *Commun Integr Biol*. 2009;2(2):71–3. [PubMed: 19704891]
20. Starkus JG, Fleig A, Penner R. The calcium-permeable non-selective cation channel TRPM2 is modulated by cellular acidification. *J Physiol*. 2010;588(8):1227–40. [PubMed: 20194125]
21. Kiselyov K, Colletti GA, Terwilliger A, Ketchum K, Lyons CWP, Quinn J, et al. TRPML: Transporters of metals in lysosomes essential for cell survival? Vol. 50, *Cell Calcium*. 2011 p. 288–94. [PubMed: 21621258]
22. Cao Q, Zhong XZ, Zou Y, Murrell-Lagnado R, Zhu MX, Dong XP. Calcium release through P2X4 activates calmodulin to promote endolysosomal membrane fusion. *J Cell Biol*. 2015;209(6):879–94. [PubMed: 26101220]
23. Haller T, Dietl P, Deetjen P, Völkl H. The lysosomal compartment as intracellular calcium store in MDCK cells: A possible involvement in InsP3-mediated Ca²⁺-release. *Cell Calcium*. 1996;19(2):157–65. [PubMed: 8689673]
24. McGuinness L, Bardo SJ, Emptage NJ. The lysosome or lysosome-related organelle may serve as a Ca²⁺-store in the boutons of hippocampal pyramidal cells. *Neuropharmacology*. 2007;52(1):126–35. [PubMed: 16930634]
25. Repnik U, esen MH, Turk B. The endolysosomal system in cell death and survival. *Cold Spring Harb Perspect Biol* [Internet]. 2013 1 1 [cited 2017 Oct 16];5(1):a008755 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23284043> [PubMed: 23284043]
26. Ferguson SM. Neuronal lysosomes. *Neurosci Lett* [Internet]. 2018; Available from: <http://linkinghub.elsevier.com/retrieve/pii/S030439401830260X>

27. Goo MS, Sancho L, Slepak N, Boassa D, Deerinck TJ, Ellisman MH, et al. Activity-dependent trafficking of lysosomes in dendrites and dendritic spines. *J Cell Biol*. 2017;216(8):2499–513. [PubMed: 28630145]
28. Galione A, Morgan AJ, Arredouani A, Davis LC, Rietdorf K, Ruas M, et al. NAADP as an intracellular messenger regulating lysosomal calcium-release channels. *Biochem Soc Trans* [Internet]. 2010;38(6):1424–31. Available from: <http://biochemsoctrans.org/lookup/doi/10.1042/BST0381424> [PubMed: 21118101]
29. Shen D, Wang X, Li X, Zhang X, Yao Z, Dibble S, et al. Lipid storage disorders block lysosomal trafficking by inhibiting a TRP channel and lysosomal calcium release. *Nat Commun*. 2012;3.
30. Macgregor A, Yamasaki M, Rakovic S, Sanders L, Parkesh R, Churchill GC, et al. NAADP Controls Cross-talk between Distinct Ca^{2+} Stores in the Heart. *J Biol Chem* [Internet]. 2007 5 18 [cited 2018 Jul 16];282(20):15302–11. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17387177> [PubMed: 17387177]
31. Galione A, Parrington J, Funnell T. Physiological roles of NAADP-mediated Ca^{2+} signaling. *Sci China Life Sci* [Internet]. 2011;54(8):725–32. Available from: <http://link.springer.com/10.1007/s11427-011-4207-5> [PubMed: 21786195]
32. Putney JW. A model for receptor-regulated calcium entry. *Cell Calcium*. 1986;7(1):1–12. [PubMed: 2420465]
33. Putney JW. Capacitative calcium entry: From concept to molecules. Vol. 231, *Immunological Reviews*. 2009 p. 10–22. [PubMed: 19754887]
34. Hui L, Geiger NH, Bloor-Young D, Churchill GC, Geiger JD, Chen X. Release of calcium from endolysosomes increases calcium influx through N-type calcium channels: Evidence for acidic store-operated calcium entry in neurons. *Cell Calcium* [Internet]. 2015 [cited 2017 May 31];58:617–27. Available from: http://ac.els-cdn.com/S0143416015001529/1-s2.0-S0143416015001529-main.pdf?_tid=d155fae0-460a-11e7-a743-00000aabb0f02&acdnat=1496239973_5d092efe55666657b3cd6c5dc4881cf0 [PubMed: 26475051]
35. Arredouani A, Ruas M, Collins SC, Parkesh R, Clough F, Pillinger T, et al. Nicotinic acid adenine dinucleotide phosphate (NAADP) and endolysosomal two-pore channels modulate membrane excitability and stimulus-secretion coupling in mouse pancreatic β cells. *J Biol Chem* [Internet]. 2015 8 28 [cited 2018 Jul 14];290(35):21376–92. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26152717> [PubMed: 26152717]
36. Moccia F, Lim D, Nusco GA, Ercolano E, Santella L. NAADP activates a Ca^{2+} current that is dependent on F-actin cytoskeleton. *FASEB J*. 2003;17(13):1907–9. [PubMed: 12923070]
37. Moccia F, Billington RA, Santella L. Pharmacological characterization of NAADP-induced Ca^{2+} signals in starfish oocytes. *Biochem Biophys Res Commun*. 2006;348(2):329–36. [PubMed: 16890912]
38. Naylor E, Arredouani A, Vasudevan SR, Lewis AM, Parkesh R, Mizote A, et al. Identification of a chemical probe for NAADP by virtual screening. *Nat Chem Biol*. 2009;5(4):220–6. [PubMed: 19234453]
39. Churchill GC, O'Neill JS, Masgrau R, Patel S, Thomas JM, Genazzani AA, et al. Sperm deliver a new second messenger: NAADP. *Curr Biol*. 2003;13(2):125–8. [PubMed: 12546785]
40. Yogalingam G, Bonten EJ, van de Vlekkert D, Hu H, Moshiah S, Connell SA, et al. Neuraminidase 1 Is a Negative Regulator of Lysosomal Exocytosis. *Dev Cell*. 2008;15(1):74–86. [PubMed: 18606142]
41. Li X, Rydzewski N, Hider A, Zhang X, Yang J, Wang W, et al. A molecular mechanism to regulate lysosome motility for lysosome positioning and tubulation. *Nat Cell Biol*. 2016;18(4):404–17. [PubMed: 26950892]
42. Herrera-Cruz MS, Simmen T. Over six decades of discovery and characterization of the architecture at mitochondria-associated membranes (MAMs). In: *Advances in Experimental Medicine and Biology*. 2017 p. 13–31.
43. Wu Y, Whiteus C, Xu CS, Hayworth KJ, Weinberg RJ, Hess HF, et al. Contacts between the endoplasmic reticulum and other membranes in neurons. *Proc Natl Acad Sci* [Internet].

- 2017;114(24):E4859–67. Available from: <http://www.pnas.org/lookup/doi/10.1073/pnas.1701078114> [PubMed: 28559323]
44. Schon EA, Area-Gomez E. Mitochondria-associated ER membranes in Alzheimer disease. *Mol Cell Neurosci* [Internet]. 2013 7 [cited 2018 Jul 27];55:26–36. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22922446> [PubMed: 22922446]
 45. Joshi AU, Kornfeld OS, Mochly-Rosen D. The entangled ER-mitochondrial axis as a potential therapeutic strategy in neurodegeneration: A tangled duo unchained. Vol. 60, *Cell Calcium*. 2016 p. 218–34. [PubMed: 27212603]
 46. Soto-Herederó G, Baixauli F, Mittelbrunn M. Interorganelle Communication between Mitochondria and the Endolysosomal System. *Front Cell Dev Biol* [Internet]. 2017;5 Available from: <http://journal.frontiersin.org/article/10.3389/fcell.2017.00095/full> [PubMed: 28271061]
 47. Demers-Lamarche J, Guillebaud G, Tlili M, Todkar K, Bélanger N, Grondin M, et al. Loss of mitochondrial function impairs Lysosomes*. *J Biol Chem* [Internet]. 2016 5 6 [cited 2017 Dec 8];291(19):10263–76. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26987902> [PubMed: 26987902]
 48. Kilpatrick BS, Eden ER, Schapira AH, Futter CE, Patel S. Direct mobilisation of lysosomal Ca²⁺ triggers complex Ca²⁺ signals. *J Cell Sci* [Internet]. 2013;126(Pt 1):60–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23108667> <http://jcs.biologists.org/content/joces/126/1/60.full.pdf> [PubMed: 23108667]
 49. Penny CJ, Kilpatrick BS, Eden ER, Patel S. Coupling acidic organelles with the ER through Ca²⁺-microdomains at membrane contact sites. Vol. 58, *Cell Calcium*. 2015 p. 387–96. [PubMed: 25866010]
 50. Hariri H, Ugrankar R, Liu Y, Henne WM. Inter-organelle ER-endolysosomal contact sites in metabolism and disease across evolution. Vol. 9, *Communicative and Integrative Biology*. 2016.
 51. Garrity AG, Wang W, Collier CM, Levey SA, Gao Q, Xu H. The endoplasmic reticulum, not the pH gradient, drives calcium refilling of lysosomes. *Elife* [Internet]. 2016 5 23 [cited 2017 May 11];5 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27213518>
 52. Zbidi H, Jardin I, Woodard GE, Lopez JJ, Berna-Erro A, Salido GM, et al. STIM1 and STIM2 are located in the acidic Ca²⁺ stores and associates with Orai1 upon depletion of the acidic stores in human platelets. *J Biol Chem*. 2011;286(14):12257–70. [PubMed: 21321120]
 53. Kilpatrick BS, Yates E, Grimm C, Schapira AH, Patel S. Endo-lysosomal TRP mucolipin-1 channels trigger global ER Ca²⁺ release and Ca²⁺ influx. *J Cell Sci* [Internet]. 2016 [cited 2018 Jul 27];129(20):3859–67. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27577094> [PubMed: 27577094]
 54. Ronco V, Potenza DM, Denti F, Vullo S, Gagliano G, Tognolina M, et al. A novel Ca²⁺-mediated cross-talk between endoplasmic reticulum and acidic organelles: Implications for NAADP-dependent Ca²⁺-signalling. *Cell Calcium* [Internet]. 2015 2 1 [cited 2018 Jul 14];57(2):89–100. Available from: <https://www.sciencedirect.com/science/article/pii/S0143416015000020> [PubMed: 25655285]
 55. Tate BA, Mathews PM. Targeting the Role of the Endosome in the Pathophysiology of Alzheimer's Disease: A Strategy for Treatment. *Sci Aging Knowl Environ* [Internet]. 2006 6 28 [cited 2018 Jul 27];2006(10):re2–re2. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16807486>
 56. Boland B, Kumar A, Lee S, Platt FM, Wegiel J, Yu WH, et al. Autophagy Induction and Autophagosome Clearance in Neurons: Relationship to Autophagic Pathology in Alzheimer's Disease. *J Neurosci* [Internet]. 2008 7 2 [cited 2018 Jul 27];28(27):6926–37. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18596167> [PubMed: 18596167]
 57. Gelman BB, Soukup VM, Holzer CE 3rd, Fabian RH, Schuenke KW, Keherly MJ, et al. Potential role for white matter lysosome expansion in HIV-associated dementia. *J Acquir Immune Defic Syndr*. 2005;39(4):422–5. [PubMed: 16010164]
 58. Spector SA, Zhou D. Autophagy: an overlooked mechanism of HIV-1 pathogenesis and neuroAIDS? *Autophagy* [Internet]. 2008 7 [cited 2018 Jul 27];4(5):704–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18424919> [PubMed: 18424919]
 59. Cysique LA, Hewitt T, Croitoru-Lamoury J, Taddei K, Martins RN, Chew CS, et al. APOE ε4 moderates abnormal CSF- β levels, while neurocognitive impairment is associated with

- abnormal CSF tau levels in HIV+ individuals – a cross-sectional observational study. *BMC Neurol* [Internet]. 2015 12 1 [cited 2018 Jul 27];15(1):51 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25880550> [PubMed: 25880550]
60. Ellis RJ, Rosario D, Clifford DB, McArthur JC, Simpson D, Alexander T, et al. Continued High Prevalence and Adverse Clinical Impact of Human Immunodeficiency Virus–Associated Sensory Neuropathy in the Era of Combination Antiretroviral Therapy. *Arch Neurol* [Internet]. 2010 5 1 [cited 2018 Jul 27];67(5):552 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20457954> [PubMed: 20457954]
 61. Heaton RK, Clifford DB, Franklin DR, Woods SP, Ake C, Vaida F, et al. HIV-associated neurocognitive disorders persist in the era of potent antiretroviral therapy: Charter Study. *Neurology* [Internet]. 2010 12 7 [cited 2017 Sep 13];75(23):2087–96. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21135382> [PubMed: 21135382]
 62. Sabatier JM, Vives E, Mabrouk K, Benjouad A, Rochat H, Duval A, et al. Evidence for neurotoxic activity of tat from human immunodeficiency virus type 1. *J Virol* [Internet]. 1991 2 [cited 2018 Jul 27];65(2):961–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/1898974> [PubMed: 1898974]
 63. Weeks BS, Lieberman DM, Johnson B, Roque E, Green M, Loewenstein P, et al. Neurotoxicity of the human immunodeficiency virus type 1 tat transactivator to PC12 cells requires the Tat amino acid 49–58 basic domain. *J Neurosci Res* [Internet]. 1995 9 1 [cited 2018 Jul 27];42(1):34–40. Available from: <http://doi.wiley.com/10.1002/jnr.490420105> [PubMed: 8531224]
 64. Haughey NJ, Holden CP, Nath A, Geiger JD. Involvement of inositol 1,4,5-trisphosphate-regulated stores of intracellular calcium in calcium dysregulation and neuron cell death caused by HIV-1 protein Tat. *J Neurochem* [Internet]. 1999 10 [cited 2018 Jul 27];73(4):1363–74. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10501179> [PubMed: 10501179]
 65. Nath A, Haughey NJ, Jones M, Anderson C, Bell JE, Geiger JD. Synergistic neurotoxicity by human immunodeficiency virus proteins Tat and gp120: Protection by memantine. *Ann Neurol* [Internet]. 2000 2 [cited 2018 Jul 27];47(2):186–94. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10665489> [PubMed: 10665489]
 66. Pérez A, Probert AW, Wang KK, Sharmeen L. Evaluation of HIV-1 Tat induced neurotoxicity in rat cortical cell culture. *J Neurovirol* [Internet]. 2001 2 1 [cited 2018 Jul 27];7(1):1–10. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11519477>
 67. King JE, Eugenin EA, Buckner CM, Berman JW. HIV tat and neurotoxicity. *Microbes Infect* [Internet]. 2006 4 [cited 2018 Jul 27];8(5):1347–57. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16697675> [PubMed: 16697675]
 68. Buscemi L, Ramonet D, Geiger JD. Human immunodeficiency virus type-1 protein Tat induces tumor necrosis factor- α -mediated neurotoxicity. *Neurobiol Dis* [Internet]. 2007 6 [cited 2018 Jun 1];26(3):661–70. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17451964> [PubMed: 17451964]
 69. Agrawal L, Louboutin J-P, Reyes BAS, Van Bockstaele EJ, Strayer DS. HIV-1 Tat neurotoxicity: A model of acute and chronic exposure, and neuroprotection by gene delivery of antioxidant enzymes. *Neurobiol Dis* [Internet]. 2012 2 [cited 2018 Jul 27];45(2):657–70. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22036626> [PubMed: 22036626]
 70. Johnson TP, Patel K, Johnson KR, Maric D, Calabresi PA, Hasbun R, et al. Induction of IL-17 and nonclassical T-cell activation by HIV-Tat protein. *Proc Natl Acad Sci U S A* [Internet]. 2013 8 13 [cited 2017 Jul 20];110(33):13588–93. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23898208> [PubMed: 23898208]
 71. Magnuson DSK, Jsnudsen BE, Geiger JD, Brownstone RM, Nath A. Human Immunodeficiency Virus type 1 Tat Activates Non-N-Methyl-D-Aspartate Excitatory Amino Acid Receptors and Causes Neurotoxicity. *Ann Neurol* [Internet]. 1995 [cited 2017 Nov 20];37(3):373–80. Available from: <https://medund.illiad.oclc.org/illiad/illiad.dll?Action=10&Form=75&Value=116096> [PubMed: 7695237]
 72. Nath A, Psooy K, Martin C, Knudsen B, Magnuson DS, Haughey N, et al. Identification of a human immunodeficiency virus type 1 Tat epitope that is neuroexcitatory and neurotoxic. *J Virol* [Internet]. 1996 [cited 2018 Jun 13];70(3):1475–80. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC189968/pdf/701475.pdf> [PubMed: 8627665]

73. Haughey NJ, Mattson MP. Calcium dysregulation and neuronal apoptosis by the HIV-1 proteins Tat and gp120. *J Acquir Immune Defic Syndr* [Internet]. 2002 10 1 [cited 2018 Jul 27];31 Suppl 2:S55–61. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12394783> [PubMed: 12394783]
74. Kim HJ, Martemyanov KA, Thayer SA. Human immunodeficiency virus protein Tat induces synapse loss via a reversible process that is distinct from cell death. *J Neurosci* [Internet]. 2008 11 26 [cited 2018 Jul 27];28(48):12604–13. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19036954> [PubMed: 19036954]
75. Fitting S, Xu R, Bull C, Buch SK, El-Hage N, Nath A, et al. Interactive comorbidity between opioid drug abuse and HIV-1 Tat: chronic exposure augments spine loss and sublethal dendritic pathology in striatal neurons. *Am J Pathol* [Internet]. 2010 9 [cited 2018 Jul 27];177(3):1397–410. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20651230> [PubMed: 20651230]
76. Hui L, Chen X, Haughey NJ, Geiger JD. Role of Endolysosomes in HIV-1 Tat-Induced Neurotoxicity. *ASN Neuro* [Internet]. 2012 4 3 [cited 2018 Jul 27];4(4):AN20120017 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22591512>
77. Rajendran L, Annaert W. Membrane Trafficking Pathways in Alzheimer's Disease. *Traffic* [Internet]. 2012 6 [cited 2018 Jul 27];13(6):759–70. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/22269004> [PubMed: 22269004]
78. Morel E, Chamoun Z, Lasiecka ZM, Chan RB, Williamson RL, Vetanovetz C, et al. Phosphatidylinositol-3-phosphate regulates sorting and processing of amyloid precursor protein through the endosomal system. *Nat Commun* [Internet]. 2013 12 2 [cited 2018 Jul 27];4(1):2250 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23907271> [PubMed: 23907271]
79. Jiang S, Li Y, Zhang X, Bu G, Xu H, Zhang Y. Trafficking regulation of proteins in Alzheimer's disease. *Mol Neurodegener* [Internet]. 2014 1 11 [cited 2018 Jul 27];9:6 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24410826> [PubMed: 24410826]
80. Nixon RA. Endosome function and dysfunction in Alzheimer's disease and other neurodegenerative diseases. *Neurobiol Aging* [Internet]. 2005 3 [cited 2018 Jul 27];26(3):373–82. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15639316> [PubMed: 15639316]
81. Rajendran L, Schneider A, Schlechtingen G, Weidlich S, Ries J, Braxmeier T, et al. Efficient Inhibition of the Alzheimer's Disease -Secretase by Membrane Targeting. *Science* (80–) [Internet]. 2008 4 25 [cited 2018 Jul 27];320(5875):520–3. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18436784>
82. Shimizu H, Tosaki A, Kaneko K, Hisano T, Sakurai T, Nukina N. Crystal Structure of an Active Form of BACE1, an Enzyme Responsible for Amyloid Protein Production. *Mol Cell Biol* [Internet]. 2008 6 1 [cited 2018 Jul 27];28(11):3663–71. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18378702> [PubMed: 18378702]
83. Sannerud R, Declerck I, Peric A, Raemaekers T, Menendez G, Zhou L, et al. ADP ribosylation factor 6 (ARF6) controls amyloid precursor protein (APP) processing by mediating the endosomal sorting of BACE1. *Proc Natl Acad Sci* [Internet]. 2011 8 23 [cited 2018 Jul 27];108(34):E559–68. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21825135> [PubMed: 21825135]
84. Hamano T, Gendron TF, Causevic E, Yen S-H, Lin W-L, Isidoro C, et al. Autophagic-lysosomal perturbation enhances tau aggregation in transfectants with induced wild-type tau expression. *Eur J Neurosci* [Internet]. 2008 3 [cited 2018 Jul 27];27(5):1119–30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18294209> [PubMed: 18294209]
85. Chesser AS, Pritchard SM, Johnson GVW. Tau Clearance Mechanisms and Their Possible Role in the Pathogenesis of Alzheimer Disease. *Front Neurol* [Internet]. 2013 9 3 [cited 2018 Jul 27];4:122 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24027553> [PubMed: 24027553]
86. Jo C, Gundemir S, Pritchard S, Jin YN, Rahman I, Johnson GVW. Nrf2 reduces levels of phosphorylated tau protein by inducing autophagy adaptor protein NDP52. *Nat Commun* [Internet]. 2014 12 25 [cited 2018 Jul 27];5(1):3496 Available from: <http://www.nature.com/articles/ncomms4496> [PubMed: 24667209]
87. Bi X, Liao G. Erratum: Autophagic-lysosomal dysfunction and neurodegeneration in Niemann-Pick type C mice: Lipid starvation or indigestion? (Autophagy) [Internet]. Vol. 3, *Autophagy*. 2007 [cited 2018 Jul 27]. p. 646–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17921694> [PubMed: 17921694]

88. Polito VA, Li H, Martini-Stoica H, Wang B, Yang L, Xu Y, et al. Selective clearance of aberrant tau proteins and rescue of neurotoxicity by transcription factor EB. *EMBO Mol Med* [Internet]. 2014 9 1 [cited 2018 Jul 27];6(9):1142–60. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25069841> [PubMed: 25069841]
89. Masgrau R, Churchill GC, Morgan AJ, Ashcroft SJH, Galione A. NAADP: a new second messenger for glucose-induced Ca²⁺ responses in clonal pancreatic beta cells. *Curr Biol* [Internet]. 2003 2 4 [cited 2018 Jul 27];13(3):247–51. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12573222> [PubMed: 12573222]
90. Mitchell KJ, Lai FA, Rutter GA. Ryanodine Receptor Type I and Nicotinic Acid Adenine Dinucleotide Phosphate Receptors Mediate Ca²⁺ Release from Insulin-containing Vesicles in Living Pancreatic β -Cells (MIN6). *J Biol Chem* [Internet]. 2003 3 28 [cited 2018 Jul 27];278(13):11057–64. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12538591> [PubMed: 12538591]
91. Haas E, Bhattacharya I, Brailoiu E, Damjanovic M, Brailoiu GC, Gao X, et al. Regulatory Role of G Protein-Coupled Estrogen Receptor for Vascular Function and Obesity. *Circ Res* [Internet]. 2009 2 13 [cited 2018 Jul 27];104(3):288–91. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19179659> [PubMed: 19179659]
92. Pandey V, Chuang C-C, Lewis AM, Aley PK, Brailoiu E, Dun NJ, et al. Recruitment of NAADP-sensitive acidic Ca²⁺ stores by glutamate. *Biochem J* [Internet]. 2009 9 15 [cited 2018 Jul 27];422(3):503–12. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19548879> [PubMed: 19548879]
93. Dickinson GD, Churchill GC, Brailoiu E, Patel S. Deviant nicotinic acid adenine dinucleotide phosphate (NAADP)-mediated Ca²⁺ signaling upon lysosome proliferation. *J Biol Chem* [Internet]. 2010 4 30 [cited 2018 Jul 27];285(18):13321–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20231291> [PubMed: 20231291]
94. Rempel HC, Pulliam L. HIV-1 Tat inhibits neprilysin and elevates amyloid β . *AIDS* [Internet]. 2005 1 28 [cited 2018 Jul 27];19(2):127–35. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15668537> [PubMed: 15668537]
95. Giunta B, Hou H, Zhu Y, Rrapo E, Tian J, Takashi M, et al. HIV-1 Tat contributes to Alzheimer's disease-like pathology in PSAPP mice. *Int J Clin Exp Pathol* [Internet]. 2009 [cited 2018 Jul 27];2(5):433–43. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19294002> [PubMed: 19294002]
96. Aksenov MY, Aksenova MV, Mactutus CF, Booze RM. HIV-1 protein-mediated amyloidogenesis in rat hippocampal cell cultures. *Neurosci Lett* [Internet]. 2010 5 21 [cited 2018 Jul 27];475(3):174–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20363291> [PubMed: 20363291]
97. Chen X, Hui L, Geiger NH, Haughey NJ, Geiger JD. Endolysosome involvement in HIV-1 transactivator protein-induced neuronal amyloid beta production. *Neurobiol Aging* [Internet]. 2013 10 [cited 2017 Aug 8];34(10):2370–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23673310> [PubMed: 23673310]
98. Kim J, Yoon JH, Kim YS. HIV-1 tat interacts with and regulates the localization and processing of amyloid precursor protein. Chauhan A, editor. *PLoS One* [Internet]. 2013 11 29 [cited 2018 Jul 27];8(11):e77972 Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24312169> [PubMed: 24312169]
99. Fields JA, Dumaop W, Crews L, Adame A, Spencer B, Metcalf J, et al. Mechanisms of HIV-1 Tat neurotoxicity via CDK5 translocation and hyper-activation: role in HIV-associated neurocognitive disorders. *Curr HIV Res* [Internet]. 2015 [cited 2018 Jul 27];13(1):43–54. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25760044> [PubMed: 25760044]
100. Liu Y, Jones M, Hingtgen CM, Bu G, Larabee N, Tanzi RE, et al. Uptake of HIV-1 Tat protein mediated by low-density lipoprotein receptor-related protein disrupts the neuronal metabolic balance of the receptor ligands. *Nat Med* [Internet]. 2000 12 1 [cited 2018 Jul 27];6(12):1380–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11100124> [PubMed: 11100124]
101. Deshmane SL, Mukerjee R, Fan S, Sawaya BE. High-Performance Capillary Electrophoresis for Determining HIV-1 Tat Protein in Neurons. Kashanchi F, editor. *PLoS One* [Internet]. 2011 1 7

[cited 2018 Jul 27];6(1):e16148 Available from: <http://dx.plos.org/10.1371/journal.pone.0016148> [PubMed: 21249135]

102. Vendeville A, Rayne F, Bonhoure A, Bettache N, Montcourrier P, Beaumelle B. HIV-1 Tat enters T cells using coated pits before translocating from acidified endosomes and eliciting biological responses. *Mol Biol Cell* [Internet]. 2004 5 [cited 2018 Jul 27];15(5):2347–60. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15020715> [PubMed: 15020715]
103. Mangieri LR, Mader BJ, Thomas CE, Taylor CA, Luker AM, Tse TE, et al. ATP6V0C Knockdown in Neuroblastoma Cells Alters Autophagy-Lysosome Pathway Function and Metabolism of Proteins that Accumulate in Neurodegenerative Disease. Srinivasula SM, editor. *PLoS One* [Internet]. 2014 4 2 [cited 2018 Jul 27];9(4):e93257 Available from: <http://dx.plos.org/10.1371/journal.pone.0093257> [PubMed: 24695574]
104. Bendiske J, Caba E, Brown QB, Bahr BA. Intracellular Deposition, Microtubule Destabilization, and Transport Failure: An “Early” Pathogenic Cascade Leading to Synaptic Decline. *J Neuropathol Exp Neurol* [Internet]. 2002 7 1 [cited 2018 Jul 27];61(7):640–50. Available from: <https://academic.oup.com/jnen/article-lookup/doi/10.1093/jnen/61.7.640> [PubMed: 12125743]
105. Bendiske J, Bahr BA. Lysosomal activation is a compensatory response against protein accumulation and associated synaptopathogenesis--an approach for slowing Alzheimer disease? *J Neuropathol Exp Neurol* [Internet]. 2003 5 [cited 2018 Jul 27];62(5):451–63. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12769185> [PubMed: 12769185]
106. Kanju PM, Parameshwaran K, Vaithianathan T, Sims CM, Huggins K, Bendiske J, et al. Lysosomal dysfunction produces distinct alterations in synaptic alphaamino-3-hydroxy-5-methylisoxazolepropionic acid and N-methyl-D-aspartate receptor currents in hippocampus. *J Neuropathol Exp Neurol* [Internet]. 2007 9 [cited 2018 Jul 27];66(9):779–88. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17805008> [PubMed: 17805008]

Figure 1:

HIV-1 proteins and other neurotoxic insults can cause deacidification of endolysosomes. Increasing endolysosome pH can release calcium and other cations from endolysosomes. Calcium released from readily releasable stores in endolysosomes can increase the release of calcium from other intracellular stores and can increase the influx of extracellular calcium. Such increases in pH and calcium levels can cause endoplasmic reticulum (ER) and mitochondrial dysfunction, Alzheimer's disease (AD)-like pathology, and synaptodendritic damage.

Figure 2:

HIV-1 Tat de-acidifies endolysosomes, increases amyloidogenesis, and releases calcium from readily releasable stores in endolysosomes. Calcium released from endolysosomes can affect mitochondrial and endoplasmic reticulum (ER) calcium stores, and increase store operated calcium (SOCE) mechanisms. Mechanistically, following de-acidification endolysosome calcium is released through TRPML1 and two pore channels (TPCs). The calcium signals can be amplified by releasing calcium from other organelles including mitochondria and ER, and by activating ER-based SOCE involving STIM1 and Orai channels as well as acidic store operated calcium entry involving N-type calcium channels (NTCCs).