Parachute mitral valve: Morphology and surgical management

Paraşüt mitral kapak: Morfoloji ve cerrahi tedavi

Shi-Min Yuan

Department of Cardiothoracic Surgery, The First Hospital of Putian, Teaching Hospital, Fujian Medical University, Putian, China

ABSTRACT

Background: This review aims to discuss morphology and surgical management of parachute mitral valve.

Methods: A total of 62 articles in the English language with 330 parachute mitral valve patients were retrieved from the PubMed, HighWire Press, and Cochrane Library databases using specific MeSH terms and keywords between January 2000 and December 2018. In these articles, morphology of parachute mitral valve and surgical treatment options were investigated.

Results: A non-syndromic parachute mitral valve was present in 287 patients (87.0%) and a syndromic parachute mitral valve was present in 43 patients (13.0%). A higher number of patients with a non-syndromic parachute mitral valve presented with congestive heart failure compared to syndromic ones. The patients with a non-syndromic parachute mitral valve often had mitral regurgitation, while syndromic parachute mitral valve patients often had mitral stenosis.

Conclusion: Parachute mitral valves are usually not an isolated lesion and are often characterized by a constellation of pathological changes of the mitral valve leaflets, annulus, commissures, subvalvular apparatus, and supravulvular mitral ring. Therefore, the majority of the patients need one or more surgical operations. The incidence of adverse events such as reintervention, postoperative complete heart block, and mortality is high in these patients.

Keywords: Cardiac surgery; mitral valve annuloplasty; mitral valve stenosis.

ÖZ

Amaç: Bu derlemede paraşüt mitral kapak morfolojisi ve cerrahi tedavisi tartışıldı.

Çalışma planı: Ocak 2000 - Aralık 2018 tarihleri arasında spesifik MeSH terimleri ve anahtar sözcükleri kullanılarak PubMed, HighWire Press ve Cochrane Kütüphanesi veri tabanlarından İngilizce dilinde 330 paraşüt mitral kapak hastasını kapsayan 62 makaleye ulaşıldı. Bu çalışmalarda paraşüt mitral kapak morfolojisi ve cerrahi tedavi seçenekleri araştırıldı.

Bulgular: Hastaların 287'sinde (%87.0) sendromik olmayan paraşüt mitral kapak ve 43'ünde (%13.0) sendromik paraşüt mitral kapak mevcuttu. Sendromik olanlara kıyasla, sendromik olmayan paraşüt mitral kapaklı daha fazla sayıda hasta konjestif kalp yetmezliği ile başvurdu. Sendromik olmayan paraşüt mitral kapaklı hastalarda sıklıkla mitral yetmezlik varken, sendromik paraşüt mitral kapaklı hastalarda sıklıkla mitral darlık vardı.

Sonuç: Paraşüt mitral kapaklar genellikle izole bir lezyon değildir ve sıklıkla mitral kapak yaprakçıkları, anülüs, komissürler, subvalvüler aparat ve supravulvüler mitral halkanın patolojik değişikliklerinin birlikteliği ile karakterizedir. Bu nedenle, hastaların birçoğu bir veya daha fazla cerrahi işleme ihtiyaç duyar. Bu hastalarda yeniden girişim, ameliyat sonrası tam kalp bloku ve mortalite gibi advers olayların insidansı yüksektir.

Anahtar sözcükler: Kalp ameliyatı; mitral kapak anüloplastisi; mitral kapak darlığı.

In 1963, Shone et al.^[1] firstly reported the mitral valve pathology of “an insertion of the chordae into a single papillary muscle, producing a funnel-shaped valve”, and they defined this lesion as a parachute mitral valve (PMV). Subsequently, Bett and Stovin^[2]

reported a patient with PMV and bicuspid aortic valve. In PMV, all chords are typically shortened and thickened, and attached to the posteromedial papillary muscle, while the anterolateral papillary muscle is absent.^[3-5]

Received: March 14, 2019 Accepted: August 02, 2019 Published online: January 23, 2020

Correspondence: Shi-Min Yuan, MD. Department of Cardiothoracic Surgery, The First Hospital of Putian, Teaching Hospital, Fujian Medical University, 351100 Putian, China. Tel: 86 594 6923117 e-mail: shiminyuan@126.com

Cite this article as:

Yuan SM. Parachute mitral valve: Morphology and surgical management. Turk Gogus Kalp Dama 2020;28(1):219-226

©2020 All right reserved by the Turkish Society of Cardiovascular Surgery.

Parachute mitral valve can be an isolated lesion, or one of the constellations of Shone syndrome. McElhinney et al.^[6] reported that 25.9% patients with severe congenital mitral stenosis had a PMV. Aslam et al.^[7] also reported Shone syndrome in 1.17% of all congenital heart lesions. Shone syndrome mainly consists of four defects: supralvalvular mitral membrane, PMV, subaortic stenosis (membranous or muscular), and coarctation of the aorta.^[1,8] The solitary papillary muscle and orientation of a severely affected PMV contributes to subaortic stenosis.^[9] Although surgical management of PMV is constantly reported, the morphological features of PMV and surgical indications of PMV have been described in limited cases.

In this review, pertinent morphological aspects and surgical management of PMV are discussed.

MATERIALS AND METHODS

Publications were systematically searched in the PubMed, HighWire Press, and Cochrane Library databases between January 2000 and December 2018. The MeSH terms and keywords were used to identify articles including “parachute mitral valve”, “supralvalvular mitral ring”, “single papillary muscle”, “Shone syndrome”, “congenital mitral stenosis”, “mitral valve repair”, and “mitral valve replacement”, in the English language. The screening of the bibliographic references helped in completing the literature retrieval. A total of 62 articles including 17 retrospective studies, one case series, and 44 case reports which met the inclusion criteria during initial screening were included in this review. Double-blind, randomized-controlled clinical studies were excluded.

The data independently extracted from each publication were the patient demographics, clinical presentations, mitral valve morphology, cardiac surgical procedures, and patient outcomes.

Statistical analysis

Statistical analysis was performed using the IBM SPSS version 22.0 software (IBM Corp., Armonk, NY, USA). Descriptive data were expressed in mean $\pm$ standard deviation (SD), median (min-max) or number and frequency. Independent samples t-test was used to compare quantitative variables. The categorical variables were compared using the chi-square (χ^2) or Fisher's exact test with continuity correction. A *p* value of <0.05 was considered statistically significant.

RESULTS

In total, 62 articles^[3,5-7,9-66] with 330 patients were included. A non-syndromic PMV was present in 287

(87.0%) patients and a syndromic PMV (as constellation of pathology of Shone syndrome) was present in 43 (13.0%) patients.

Gender was described for 236 (236/330 patients, 71.5%) and 135 (57.2%) were males and 101 (42.8%) were females. The male-to-female ratio did not differ between patients with non-syndromic and syndromic PMV (128/95 vs. 7/6, $\chi^2=0.1$, *p*=1.000). The mean age of the patients was 21.1 ± 22.0 (range, 0 to 85) years. Age was not specified in 224 patients. A total of 166 (74.1%) patients were pediatric and 58 (25.9%) were adult patients ($\chi^2=104.1$, *p*<0.001). In the non-syndromic PMV group, 145 (145/177, 81.9%) patients were pediatrics and 32 (32/177, 18.1%) were adults, while in the syndromic PMV group, 21 (21/47, 44.7%) patients were pediatrics and 26 (26/47, 55.3%) patients were adults.

Most PMV patients were symptomatic, while a few patients were asymptomatic in either non-syndromic or syndromic PMV patients ($\chi^2=42.6$, *p*<0.001 for non-syndromic patients, and $\chi^2=16.7$, *p*<0.001 for syndromic patients). Among pediatrics, 17 patients (21.3%) were asymptomatic and 63 patients (78.8%) were symptomatic, while among adults, nine patients (36%) were asymptomatic and 16 patients (64%) were symptomatic ($\chi^2=2.2$, *p*=0.184). A higher number of patients with a non-syndromic PMV presented with congestive heart failure compared to syndromic ones. The patients with a non-syndromic PMV were mostly associated with an atrioventricular septal defect and hypoplastic left ventricle than those with a syndromic PMV. However, the patients with a syndromic PMV had a higher incidence of coarctation of the aorta, bicuspid aortic valve, and subaortic obstruction (Table 1).

Hemodynamic studies showed that the peak and mean mitral pressure gradients did not significantly differ between the groups (Table 1). All patients had a single papillary muscle. Most patients had a thickened mitral valve leaflet, shortened chords, and mitral stenosis or regurgitation. The patients with a non-syndromic PMV often had mitral regurgitation, while those with a syndromic PMV had mitral stenosis (Table 2).

Management was described for 216 patients. Accordingly, 20 patients (9.3%) were not operated due to conservative treatment/on a follow-up/waiting for surgical operation/operation refusal/sudden death, and loss of surgical opportunity,^[12,13,15,21,22,25,26,29,41,42,45,48-52,57,66] while 196 patients (90.7%) received a total of 198 surgical operations as follows: non-mitral valve operations (*n*=70, 35.4%)^[3,11,14,26,30,33,36,39,46,61] and

Table 1. Comparisons between non-syndromic and syndromic PMV patients

Clinical presentation	Non-syndromic PMV		Syndromic PMV		Test	χ^2	<i>p</i>
	n	%	n	%			
Case number	287	87.0	43	13.0	Chi-square	360.8	<0.001
Gender						0.1	1.000
Male (n=135)	128	94.8	7	5.2	Chi-square		
Female (n=101)	95	94.1	6	5.9	Chi-square		
Child/adult						26.8	<0.001
Child (n=166)	145	87.3	21	12.7	Chi-square		
Adult (n=58)	32	55.2	26	44.8	Chi-square		
Asymptomatic (n=26)	25*	26.3	1	8.3	Fisher exact test with continuity correction	1.9	0.286
Symptomatic (n=81)	70	73.7	11	91.7	Chi-square	1.9	
Heart failure	38	54.3	1	9.1	Fisher exact test with continuity correction	7.8	0.007
Dyspnea/tachypnea	14	20.0	8	72.7	Chi-square	13.4	0.001
Associated cardiac anomaly							
Coarctation of the aorta	82	28.6	33	76.7	Chi-square	38.2	<0.001
Bicuspid aortic valve	63	22.0	30	69.8	Chi-square	68.3	<0.001
Hypoplastic left ventricle	36	12.5	0	0	Fisher exact test with continuity correction	6.1	0.008
Atrioventricular septal defect	26	9.1	0	0	Fisher exact test with continuity correction	4.2	0.034
Subaortic obstruction	25	8.7	12	27.9	Chi-square	13.8	0.001
Tricuspid regurgitation	3	1.0	4	9.3	Fisher exact test with continuity correction	12.3	0.007
Anomalous origin of coronary artery	1	0.3	2	4.7	Fisher exact test with continuity correction	7.7	0.046
Left superior vena cava	1	0.3	7	16.3	Fisher exact test with continuity correction	40.1	<0.001
Supravalvular aortic stenosis	1	0.3	3	7.0	Fisher exact test with continuity correction	13.7	0.008

PMV: parachute mitral valve; * One of the asymptomatic patients developed dyspnea later.^[41]

Table 2. Hemodynamic, metrology, and morphology of PMV

Clinical presentation	Non-syndromic PMV (n=287)			Syndromic PMV (n=43)			Test	χ^2 /t-value	p
	n	%	Mean±SD	n	%	Mean±SD			
Mean mitral pressure gradient (mmHg)			12.7±7.9			9.9±1.9	Independent samples t test	0.763	0.458
Peak mitral pressure gradient (mmHg)			26.0±2.6			21.5±11.0	Independent samples t test	0.673	0.520
Annulus diameter (mm)			11.6±4.4						
z-value			-1.6±2.0						
Mitral valve area (cm ²)			1.8±1.0			2.0±0.6	Independent samples t test	0.352	0.732
Dominant papillary muscle	<ul style="list-style-type: none"> • Posteromedial (n=41)^[39,43,45,48,53,62] • Anterolateral (n=14)^[39,65] • Inferolateral^[56] • Left (n=1)^[32] • Central^[54] 			<ul style="list-style-type: none"> • Apical^[63] 					
Supravalvular mitral ring	10 ^[9,10,38,39]	3.5		11 ^[4,7,12,15,20,21,28,37,49,63,66]	25.6		Chi-square	23.0	<0.001
Mitral valve	99	100		36	100				
MR	50	50.5		6	16.7		Chi-square	12.5	<0.001
MS	36	36.4		29	80.6		Chi-square	20.7	<0.001
MS + MR	7	7.1		1	2.8		Fisher exact test with continuity correction	0.9	0.681
Normal	6	6.1		0	0		Fisher exact test with continuity correction	2.3	0.342

PMV: parachute mitral valve; MR: Mitral regurgitation; MS: Mitral stenosis.

Table 3. Morphology of PMV^[17]

Mitral structure	Morphology
Valve	Deformed, restricted leaflet mobility
Mitral valve orifice	Stenosis, regurgitant, or normal
Commissures	Barely distinguishable, frequently underdeveloped
Papillary muscle	Single, or one prominent and one absent/hypoplastic
Chord	Short and thick, convergent papillary insertion
The functional communication between the left atrium and the left ventricle	Interchordal spaces

PMV: Parachute mitral valve.

mitral valve operations (n=128, 64.6%). Mitral valve operations were prevailed by double patch repair for atrioventricular septal defects (n=20, 15.6%),^[44] surgical mitral valvuloplasty (n=17, 13.3%),^[5,66] mitral valve replacement (MVR) (n=16, 12.5%),^[3,5,9,26,34,36,59,62,66] and zone of apposition closure (n=16, 12.5%).^[58] Reintervention was required in 19 patients (19/196, 9.7%).^[3,9,16,17,26,58,60]

The mean follow-up of the patients was 150.7±145.6 months (range, 1 month to 20 years) (n=21). The outcomes of 231 patients were reported as follows: 191 (82.7%) recovered, six (2.6%) were complicated, and 34 (14.7%) died.

DISCUSSION

Morphology

Parachute mitral valve is formed by specific malformations of the mitral leaflets *per se*, as well as subvalvular structures (Table 3). The major morphology of PMV is a single papillary muscle, or one papillary muscle is severely hypoplastic. Chauvaud^[67] proposed that, in PMV patients, mitral regurgitation might be caused by hypoplasia of one papillary muscle, commissural enlargement, valve leaflet defects, and shortened chords. The authors reported that the hypoplastic papillary muscle was usually the posterior one, while the other papillary muscle was medially displaced. However, some others^[39,43,45,48,53,62] reported the dominant papillary muscle was the posterior one. The results of this study supported the posterior papillary muscle was the dominant one.

The combination of lesions can give rise to a funnel configuration of the mitral valve. Three-dimensional echocardiography can visualize all characteristic findings of PMV including the absence of one papillary muscle, a funnel-shaped mitral valve, a doming-shaped elongated chordae tendineae, and a pear-shaped left

atrium.^[18] A supramitral ring in a form of membranous or fibrous shelf is often an integral part of the PMV, thereby, significantly reducing the effective orifice area of the mitral valve.^[68]

The characterized single papillary muscle which receives all chords confirms true PMV. Conversely, two papillary muscles with all chordae inserting into one muscle and the other being hypoplastic indicate a parachute-like mitral valve.

According to the literature review, the mean mitral valve annulus diameter was 8.2 (range, 7 to 10) mm, which corresponds to a z-score of -0.665 on the basis of the normal range for newborns (10±2.6 mm).^[66]

Surgical treatment and outcomes

Serraf et al.^[69] reported that PMV-related mitral stenosis often caused failure of biventricular repair in newborns with borderline small left ventricles, thus strengthening the importance of the left ventricular inflow status in decision making for either a uni- or a biventricular treatment strategy.

Balloon mitral valvuloplasty decreased the peak and the mean mitral valve gradients by a median of 33% and 38%, respectively; however, 54.5% (6/11) patients with a supravulvar mitral ring developed significant mitral regurgitation following balloon mitral valvuloplasty.^[6]

Mitral repair has been a preferred procedure as opposed to MVR. In some patients, repair of a stenotic PMV was performed through a papillary muscle incision and leaflet fenestration.^[70] In children, MVR shows several drawbacks, such as high operative mortality, significant incidence of complete heart block and pacemaker implantation, lack of prosthetic valves with sizes and with growth potential that are suitable for small children, difficulties in postoperative anticoagulant therapy, and rapid bioprosthetic valve deterioration.^[71]

Shone et al.^[1] reported that mitral valve obstruction was the most critical problem of this lesion. The severity of the mitral valve obstruction was found to be inversely correlated with long-term outcomes, and the operative mortality of patients with Shone syndrome was eventually adversely affected.^[66] However, Marino et al.^[39] found no significant association between progressive mitral stenosis and PMV type, dominant papillary muscle, sex, or any surgical or interventional therapies.

As PMVs are usually not isolated lesions and are characterized by a constellation of pathological changes of the mitral valve leaflets, annulus, commissures, subvalvular apparatus, and supralvalvular mitral ring, the majority of patients need one or more surgical operations and the reintervention rate is high.^[39]

In conclusion, about two-thirds of parachute mitral valve patients require surgical treatment of the mitral valve lesions. Parachute mitral valves are curable by mitral valve repair in most cases, and mitral valve replacement is indicated only for patients with severe mitral valve lesions.

Declaration of conflicting interests

The author declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding

The author received no financial support for the research and/or authorship of this article.

REFERENCES

- Shone JD, Sellers RD, Anderson RC, Adams P Jr, Lillehei CW, Edwards JE. The developmental complex of "parachute mitral valve," supralvalvular ring of left atrium, subaortic stenosis, and coarctation of aorta. *Am J Cardiol* 1963;11:714-25.
- Bett JH, Stovin PG. Parachute deformity of the mitral valve. *Thorax* 1969;24:632-6.
- Brown JW, Ruzmetov M, Vijay P, Hoyer MH, Girod D, Rodefeld MD, et al. Operative results and outcomes in children with Shone's anomaly. *Ann Thorac Surg* 2005;79:1358-65.
- Séguéla PE, Houyel L, Acar P. Congenital malformations of the mitral valve. *Arch Cardiovasc Dis* 2011;104:465-79.
- Turkoz R, Ayabakan C, Vuran C, Omay O, Yildirim SV, Tokel NK. Duplicate mitral valve in an infant with Shone's anomaly. *Ann Thorac Surg* 2009;88:1683-5.
- McElhinney DB, Sherwood MC, Keane JF, del Nido PJ, Almond CS, Lock JE. Current management of severe congenital mitral stenosis: outcomes of transcatheter and surgical therapy in 108 infants and children. *Circulation* 2005;112:707-14.
- Aslam S, Khairy P, Shohoudi A, Mercier LA, Dore A, Marcotte F, et al. Shone complex: an under-recognized congenital heart disease with substantial morbidity in adulthood. *Can J Cardiol* 2017;33:253-9.
- Carey LS, Sellers RD, Shone JD. Radiologic findings in the developmental complex of parachute mitral valve, supralvalvular ring of left atrium, subaortic stenosis, and coarctation of aorta. *Radiology* 1964;82:1-10.
- Schaverien MV, Freedom RM, McCrindle BW. Independent factors associated with outcomes of parachute mitral valve in 84 patients. *Circulation* 2004;109:2309-13.
- Abelson M. Parachute mitral valve and a large ventricular septal defect in an asymptomatic adult. *Cardiovasc J S Afr* 2001;12:212-4.
- Agnoletti G, Raisky O, Boudjemline Y, Ou P, Bonnet D, Sidi D, et al. Neonatal surgical aortic commissurotomy: predictors of outcome and long-term results. *Ann Thorac Surg* 2006;82:1585-92.
- Akgüllü C, Erdoğan E. Adult patient with Shone's syndrome and patent ductus arteriosus: a case report. *Türk Kardiyol Dern Ars* 2012;40:350-3.
- Almeida S, Cotrim C, Miranda R, Lopes L, Almeida AR, Loureiro MJ, et al. The role of echocardiography in assessing parachute mitral valve. *Rev Port Cardiol* 2009;28:335-9.
- Bobylev D, Meschenmoser L, Boethig D, Horke A. Surgical repair of Shone's complex with anomalous origin of the left coronary artery arising from the right pulmonary artery. *Interact Cardiovasc Thorac Surg* 2015;20:439-42.
- Chandra S, Kharwar RB, Mundhekar A. Shone complex. *J Cardiovasc Ultrasound* 2015;23:54-5.
- Cho S, Kim WH, Kwak JG, Lee JR, Kim YJ. Surgical results of mitral valve repair for congenital mitral valve stenosis in paediatric patients. *Interact Cardiovasc Thorac Surg* 2017;25:877-82.
- Delmo Walter EM, Hetzer R. Repair for congenital mitral valve stenosis. *Semin Thorac Cardiovasc Surg Pediatr Card Surg Annu* 2018;46-57.
- Demirkol S, Baysan O, Celik T, Kurt IH, Kucuk U, Balta S. Incremental value of three-dimensional transesophageal echocardiography in a patient with parachute mitral valve. *Echocardiography* 2012;29:E24-5.
- Demirkol S, Arslan Z, Balta S, Küçük U. A parachute mitral valve accompanying persistent left superior vena cava: assessment by three-dimensional transthoracic echocardiography. *Anadolu Kardiyol Derg* 2012;12:E23-4.
- Dibardino DJ, Fraser CD Jr, Dickerson HA, Heinle JS, McKenzie ED, Kung G. Left ventricular inflow obstruction associated with persistent left superior vena cava and dilated coronary sinus. *J Thorac Cardiovasc Surg* 2004;127:959-62.
- Ekici B, Erkan AF, Sökmen Y, Tüfekçioğlu O. Shone's complex with dextrocardia and situs inversus totalis: a case report. *Türk Kardiyol Dern Ars* 2012;40:612-4.
- Espinola-Zavaleta N, Chugh R, Ramírez GM. Parachute mitral valve with severe mitral regurgitation in an adult patient. *Echocardiography* 2012;29:E122-5.
- Fisher D, Dipietro A, Murdison KA, Lemieux CA. Full monosomy 21: echocardiographic findings in the third molecularly confirmed case. *Pediatr Cardiol* 2013;34:733-5.
- Gokhroo RK, Bisht DS, Padmanabhan D, Gupta S. Association of large muscular ventricular septal defect with

- parachute mitral valve: A rare case report. *Indian J Cardiol* 2013;16:64-6.
25. Gori T, Salerno D, Donati G. 'Parachute' accessory mitral leaflet and pulmonary valve stenosis in an asymptomatic 85-year-old man. *Eur Heart J* 2008;29:223.
26. Grimaldi A, Vermi AC, Maisano F, Sacco F, Castiglioni A, Zangrillo A, et al. Echocardiographic patterns of incomplete Shone's syndrome in adults. *J Heart Valve Dis* 2011;20:552-6.
27. Henaine R, Nloga J, Wautot F, Yoshimura N, Rabilloud M, Obadia JF, et al. Long-term outcome after annular mechanical mitral valve replacement in children aged less than five years. *Ann Thorac Surg* 2010;90:1570-6.
28. Hosseinpour AR, Amanullah M, Ramnarine IR, Stümper O, Barron DJ, Brawn WJ. Combined atrial arterial switch operation (double switch) for hearts with Shone syndrome and pulmonary hypertension. *J Thorac Cardiovasc Surg* 2006;131:471-3.
29. Isılak Z, Deveci OS, Yalcın M, Uz O, Dogan M, Uzun M. A unique case of parachute mitral valve in an adult: an abnormally long chorda tendinea. *Echocardiography* 2012;29:E245-6.
30. Jain D, Martel M, Reyes-Múgica M, Parkash V. Heterotopic nephrogenic rests in the colon and multiple congenital anomalies: possibly related association. *Pediatr Dev Pathol* 2002;5:587-91.
31. Jiang ZY, Pircova A, Sekarski N, Hack I, Laurini R, Janzer R, et al. Transposition of the great arteries, pulmonary atresia, and multiple ventricular septal defects associated with multiple cardiac rhabdomyomas in a case of tuberous sclerosis. *Pediatr Cardiol* 2000;21:165-9.
32. Kim WH, Lee TY, Kim SC, Kim SJ, Lee YT. Unbalanced atrioventricular septal defect with parachute valve. *Ann Thorac Surg* 2000;70:1711-2.
33. Kurtul A, Duran M, Akkaya E, Ornek E. Parachute mitral and tricuspid valves together with ventricular septal defect. *Türk Kardiyol Dern Ars* 2013;41:264.
34. López-Pardo F, Urbano-Moral JA, González-Calle A, Laviana-Martinez F, Esteve-Ruiz I, Lagos-Degrande O, et al. Three-Dimensional Transesophageal Echocardiography in the Anatomical Assessment of Isolated Parachute Mitral Valve in an Adult Patient. *Echocardiography* 2015;32:1732-5.
35. Lorenzo N, Claver E, Alió J, Aguilar R. Incomplete Shone's complex in the sixth decade of life: echo and cardiac magnetic resonance imaging assessment. *Eur Heart J* 2017;38:459.
36. Lu M, Tang H. 3166: Echocardiographic in the diagnosis of parachute mitral valve. *Ultrasound Med Biol* 2006;32:P219.
37. Ma XJ, Huang GY, Liang XC, Liu XQ, Jia B. Atypical Shone's complex diagnosed by echocardiography. *Pediatr Cardiol* 2011;32:442-8.
38. Mądry W, Karolczak MA, Grabowski K. Supravalvar mitral ring with a parachute mitral valve and subcoarctation of the aorta in a child with hemodynamically significant VSD. A study of the morphology, echocardiographic diagnostics and surgical therapy. *J Ultrason* 2017;17:206-11.
39. Marino BS, Kruger LE, Cho CJ, Tomlinson RS, Shera D, Weinberg PM, et al. Parachute mitral valve: morphologic descriptors, associated lesions, and outcomes after biventricular repair. *J Thorac Cardiovasc Surg* 2009;137:385-93.e4.
40. Meenakshi K, Chidambaram S, Dhandapani VE, Rameshwar R. A rare association of parachute mitral valve with double outlet right ventricle and severe pulmonary hypertension in an adult. *J Assoc Physicians India* 2014;62:50-2.
41. Mohan JC, Shukla M, Mohan V, Sethi A. Parachute mitral valve and Pacman deformity of the ventricular septum in a middle-aged male. *Indian Heart J* 2016;68:S126-S30.
42. Mohan JC, Shukla M, Sethi A. Parachute deformity of both atrioventricular valves with congenitally corrected transposition in an adult. *Indian Heart J* 2015;67:565-9.
43. Motoda H, Murata M, Iwanaga S, Matsushita K, Nakamizo H, Wakino S, et al. Parachute mitral valve incidentally diagnosed in an adult patient with hypertension. *J Echocardiogr* 2010;8:28-9.
44. Myers PO, del Nido PJ, Bautista-Hernandez V, Marx GR, Emani SM, Pigula FA, et al. Biventricular repair for common atrioventricular canal defect with parachute left atrioventricular valve. *Eur J Cardiothorac Surg* 2016;49:546-51.
45. Nikolic A, Joksimovic Z, Jovovic L. Exuberant accessory mitral valve tissue with possible true parachute mitral valve: a case report. *J Med Case Rep* 2012;6:292.
46. Nkoke C, Lekoubou A, Yonta EW, Dzudie A, Kengne AP. Shone's anomaly: a report of one case in sub-Saharan Africa. *Cardiovasc Diagn Ther* 2014;4:495-8.
47. Park SJ, Kwak CH, Hwang JY. Long-term survival in double inlet left ventricle combined with pulmonary stenosis and parachute mitral valve: a rare case. *Int Heart J* 2007;48:261-7.
48. Patsouras D, Korantzopoulos P, Kountouris E, Siogas K. Isolated parachute mitral valve as an incidental finding in an asymptomatic hypertensive adult. *Clin Res Cardiol* 2007;96:38-41.
49. Popescu BA, Jurcut R, Serban M, Parascan L, Ginghina C. Shone's syndrome diagnosed with echocardiography and confirmed at pathology. *Eur J Echocardiogr* 2008;9:865-7.
50. Prunier F, Furber AP, Laporte J, Geslin P. Discovery of a parachute mitral valve complex (Shone's anomaly) in an adult. *Echocardiography* 2001;18:179-82.
51. Purvis J, Sharma D. Adult parachute mitral valve detected in pregnancy. *Heart* 2011;97:1192.
52. Purvis JA, Smyth S, Barr SH. Multi-modality imaging of an adult parachute mitral valve. *J Am Soc Echocardiogr* 2011;24:351.e1-3.
53. Rybicka J, Dobrowolski P, Kuśmierczyk M, Rózański J, Kowalski M, Hoffman P. Parachute mitral valve in a young adult with recurrent pulmonary oedema. *Acta Cardiol* 2011;66:401-3.
54. Sadeghian H, Savand-Roomi Z. Parachute mitral valve. In: Sadeghian H, Savand-Roomi Z, editors. *Echocardiographic Atlas of Adult Congenital Heart Disease*. Cham: Springer; 2015. p. 349-50.
55. Saroli T, Gelehrter S, Gomez-Fifer CA, van der Velde ME, Bove EL, Ensing GJ. Anomalies of left coronary artery origin affecting surgical repair of hypoplastic left heart syndrome and Shone complex. *Echocardiography* 2008;25:727-31.
56. Saura D, Campos JV, Villegas M, Picó F, de la Morena G, Valdés-Chávarri M. Heart-hand syndrome. *Int J Cardiol* 2008;129:e7-9.

57. Sethi R, Kharwar RB, Sharma A, Kumar V, Narain VS. Multimodality imaging of holmes heart with parachute mitral valve. *Echocardiography* 2014;31:E132-5.
58. Sharma V, Burkhart HM, Schaff HV, Cetta F, Cabalka A, Dearani JA. Management of zone of apposition in parachute left atrioventricular valve in atrioventricular septal defect. *Ann Thorac Surg* 2013;95:1665-9.
59. Shehatha JS, Taha AY, Mirza AJ. Late Shone complex: A case report and literature review. *J Egypt Soc Cardio-Thorac Surg* 2018;26:133-5
60. Stellin G, Padalino MA, Vida VL, Boccuzzo G, Orrù E, Biffanti R, et al. Surgical repair of congenital mitral valve malformations in infancy and childhood: a single-center 36-year experience. *J Thorac Cardiovasc Surg* 2010;140:1238-44.
61. Takawira FF, Sinyangwe G, Mwangi MN, Mathivha TM. Case report and images in cardiology: Shone's complex variant associated with a patent ductus arteriosus: simultaneous treatment of coarctation and patent ductus arteriosus using a covered stent. *SAHeart* 2010;7:282-6.
62. Vistarini N, Belaidi M, Desjardins G, Pellerin M. Parachute Mitral Valve. *Can J Cardiol* 2016;32:1261.e5-6.
63. Vivas MF, Politi MT, Riznyk LM, Castro MF, Avegliano G, Ronderos R. Shone's syndrome: Insights from three-dimensional echocardiography. *Echocardiography* 2018;35:417-9.
64. Wagner F, Gebauer RA, Paech C. Newborn in cardiogenic shock. *Heart* 2018;104:467.
65. Yamamoto T, Onishi T, Omar AM, Norisada K, Tatsumi K, Matsumoto K, et al. Isolated true parachute mitral valve in an asymptomatic elderly patient. *J Echocardiogr* 2010;8:131-2.
66. Zucker N, Levitas A, Zalstein E. Prenatal diagnosis of Shone's syndrome: parental counseling and clinical outcome. *Ultrasound Obstet Gynecol* 2004;24:629-32.
67. Chauvaud S. Surgery of congenital mitral valve disease. *J Cardiovasc Surg (Torino)* 2004;45:465-76.
68. Remenyi B, Gentles TL. Congenital mitral valve lesions: correlation between morphology and imaging. *Ann Pediatr Cardiol* 2012;5:3-12.
69. Serraf A, Piot JD, Bonnet N, Lacour-Gayet F, Touchot A, Bruniaux J, et al. Biventricular repair approach in ducto-dependent neonates with hypoplastic but morphologically normal left ventricle. *J Am Coll Cardiol* 1999;33:827-34.
70. Zias EA, Mavroudis C, Backer CL, Kohr LM, Gotteiner NL, Rocchini AP. Surgical repair of the congenitally malformed mitral valve in infants and children. *Ann Thorac Surg* 1998;66:1551-9.
71. Jaquiss RD. Excellent Early; Not-So-Excellent Late. *Semin Thorac Cardiovasc Surg* 2016;28:461-2.