Published in final edited form as:

J Am Chem Soc. 2019 January 30; 141(4): 1435–1440. doi:10.1021/jacs.8b11465.

Sydnone-Based Approach to Heterohelices through 1,3-Dipolar-Cycloadditions

Expédite Yen-Pon[†], Pier Alexandre Champagne[§], Lucie Plougastel[†], Sandra Gabillet[†], Pierre Thuéry[‡], Mizuki Johnson^{||}, Gilles Muller^{||}, Grégory Pieters[†], Frédéric Taran[†], K. N. Houk[§], Davide Audisio^{*,†}

[†]Service de Chimie Bio-organique et Marquage (SCBM), JOLIOT, CEA, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

[‡]NIMBE, CEA, CNRS, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

[§]Department of Chemistry and Biochemistry, University of California, Los Angeles, California 90095, United States

^{||}Department of Chemistry, San José State University, San José, California 95192-0101, United States

Abstract

The first approach to pyrazole-containing helices via sydnone-aryne [3 + 2]-cycloaddition is described. An unprecedented regioselectivity in the cycloaddition step toward the more sterically constrained product was observed in the presence of extended aromatic scaffolds. DFT calculations enabled understanding the origin of this unexpected selectivity.

Ortho-fused aromatic rings belong to a class of helical-shaped molecules named helices.¹ Since their discovery, these variegated aromatics have fascinated chemist practitioners due to their elegant architecture, inherent chirality, and structural complexity. Helical structures are particularly interesting, as they are found in many biomacromolecules inspiring chemists involved in the field of asymmetric catalysis,² and their enhanced chiroptical properties have attracted much attention in material science.³ The presence of a heteroatom in the fused polycyclic system considerably alters the electronic structure and helps tune various optoelectrical properties.⁴ In particular, nitrogen-containing helices, including pyridine,⁵ pyrrole,⁶ pyrazine,⁷ and imidazolium^{4a,8} have attracted much attention. Despite the broad interest in this family of compounds, synthetic access still remains challenging and often requires cumbersome multistep approaches.¹ Ideally, the desired helical motif would be assembled in the last step of the sequence, thus enabling divergent opportunities for structure diversity. Sydnone-arynes are azomethine imines, well-known for their 1,3-dipolar-cycloadditions

*Corresponding Author: davide.audisio@cea.fr.

Supporting Information

The Supporting Information is available free of charge on the [ACS Publications website](https://pubs.acs.org) at DOI: 10.1021/jacs.8b11465.

Experimental procedures and computational details NMR spectra for obtained compounds (PDF)

Crystallographic data for compounds **10a** (CCDC 1544556), **10b** (CCDC 1872990), and **11b** (CCDC 1544557) (CIF)

The authors declare no competing financial interest.

with linear and strained alkynes, generating pyrazoles, a pharmaceutically and agrochemically relevant heterocyclic scaffold.⁹ We reasoned that properly designed prohelical sydnones **3**, bearing *ortho*-extended aromatic substitutions, would be suitable partners with arynes bearing an extended aromatic core.¹⁰ After cycloaddition, the subsequent loss of carbon dioxide would deliver the desired pyrazole-containing helicenes, a family of heterohelicenes so far unreported (Figure 1).

We now describe a novel disconnection allowing a direct access to a range of unreported helical pyrazoles, based on a key sydnone-aryne cycloaddition. In the process, we discovered a unique example of selective cycloaddition involving these mesoionic betaines in favor of the sterically hindered helical product and determined the reasons behind such selectivity with analysis of DFT calculations.

The implementation of this strategy relies on the prerequisite preparation of *ortho*-fused aromatic azomethine imine dipoles. *N*-Methyl sydnone **1** was identified as a key component to build up such *ortho*-aromatic structures. Readily available in two steps from commercial sarcosine, **1** provides a versatile handle for further derivatization. It was envisioned that metal catalyzed functionalization at the C₄ position of the sydnone with 2-halobenzaldehyde would provide a key intermediate that might undergo intramolecular Knoevenagel condensation under basic conditions to give the desired product **3**. Initial attempts showcased that the whole cascade could be performed in one single operation. Product **3a** was first isolated in moderate yield together with the intermediate uncyclized aldehyde **S3a**.¹¹ After some experimentation, it was found that, in the presence of catalytic amounts of Pd(OAc)₂ and triphenylphosphine, with an excess of K₂CO₃ (4 equiv), the desired mesoionic compound **3a** could be obtained in 88% yield. Similarly, the tetracyclic *ortho*-sydnone **3b** could be isolated in 66% yield starting from the corresponding 1-bromo-2-naphthaldehyde (Figure 2A).

With a reliable access to the prerequisite *ortho*-fused aromatic azomethine imines **3a** and **3b** secured, we turned our attention to the key 1,3-dipolar-cycloaddition between polycyclic sydnones and aryne precursors. The sydnone-aryne cycloaddition is a 50 year old transformation pioneered by Gotthardt, Huisgen, and Knorr and later by Kato and Tsuge in 1974,¹² but only recently the synthetic potential of this transformation has been studied in detail.¹³ Initially, we reacted sydnones **3a** and **3b** in the presence of silyl triflate **4a** (1.5 equiv) and TBAF (1.5 equiv) at room temperature. As expected, the desired products **5** and **6** were isolated in good yields, 77% and 70%, respectively (Figure 2C). It is worth mentioning that **5** and **6** have been synthesized in only two steps from readily available *N*-methyl sydnone.¹⁴ The sequence could be extended to the *ortho*-fused 1,2-naphthyne precursor **4b** (Figure 2D). In presence of tricyclic sydnone **3a** the formation of the two possible cycloadducts, the desired hetero-[5]-helicene **7a** and the S-shaped product **7b**, was observed. As expected, no degree of selectivity was achieved and the cycloadducts were isolated in a 1:1 ratio and an overall 74% yield. The tetracyclic sydnone **3b** gave a low degree of selectivity in favor of the sterically hindered hetero-[6]-helicene **8a** (**8a/8b**, ratio 63/37). The ¹H NMR analysis of the two regioisomers showed well-resolved signals, which could be clearly assigned with COSY and NOESY measurements.¹¹ Intrigued by this unexpected result, we investigated the reactivity of *ortho*-sydnones **3a** and **3b** in the presence of 3,4-

phenanthryne precursor **4c** (Figure 2D). This *ortho*-annulated aryne is particularly interesting because it should allow the formation of [6]- and [7]-helicenes. The reaction of **3a** with **4c** in the presence of TBAF in THF delivered a crude mixture with useful selectivity in favor of the helical product (crude ^1H NMR ratio 90:10). After purification, the two components of the reaction could be isolated in 79% and 6% yield. The 2D-NMR measurements determined the identity of the major compound as the desired [6]-helical pyrazole **9a**.¹⁵ When **4c** was reacted in the presence of **3b**, [7]-helicene **10a** was isolated in 62% yield with high selectivity (**10a/10b** 96/4). Crystals of both isomers **10a** and **10b** were grown by slowly evaporating dichloromethane solutions, and the structures were determined by single-crystal X-ray diffraction (Figure 3A).

The chiroptical properties of [7]-helicene **10a** were subjected to a preliminary evaluation. Enantiomers of **10a** were resolved from a racemic mixture using HPLC with a chiral-phase column.¹⁶ In Figure 3B, the circular dichrograms of the enantiopure samples are shown, which exhibit both positive and negative Cotton effects at 235 and 357 nm. The spectra of the enantiomers are mirror images of each other. The measurements conducted in degassed dichloromethane for the set of enantiomers led to the observation of a mirror-image CPL signal (Figure 3C). The g_{lum} values are -0.001 and $+0.001$ at about the maximum emission wavelength for **10a**.¹⁷ These values are of the same order of magnitude as those for other examples of organic CPL-active helicenes.¹⁸ These results confirm that the solution of [7]-helicene **10a** in degassed dichloromethane exhibits active CPL signals and also that the emitted light is polarized in opposite directions for the two enantiomeric forms for this helicene-like structure.

The selectivities observed for these sydnone-aryne cycloadditions were unexpected, given precedents in the literature. First, sydnones are known to be poorly regioselective in their cycloadditions with asymmetrical alkyne dipolarophiles.^{13,19–23} Moreover, the Houk–Garg distortion-based model to explain the preferred regioselectivity of attack of nucleophiles on strained alkynes,²⁴ which is based on the difference in internal angles of the alkyne carbons, predicts low selectivity with 1,2-naphthyne or 3,4-phenanthryne, albeit in the observed direction. DFT optimizations of the two structures (M06–2X/6–31+G(d,p))¹¹ reveal that the two aryne carbons have similar internal bond angles (Figure 4A).

In order to highlight whether the peculiar structure of sydnones **3a** and **3b** is responsible for the unusual selectivity, sydnone **3c** was synthesized. As shown in Figure 4B, when *N*-phenyl sydnone **3c** was reacted with **4c** the opposite selectivity was observed (ratio 33:66 in favor of **11b**).²⁵ This result suggests that the origin of the selectivity might be related to the structure of the polycyclic sydnone itself. To understand the origins of such a dichotomy, we calculated the free energy profiles for the reactions of **3a–3c** with both 1,2-naphthyne and 3,4-phenanthryne, using the same DFT method described above. Profiles for the reaction of **3a** with 3,4-phenanthryne are shown in Figure 4C.¹¹ In all cases examined, the regio-determining cycloaddition step is also rate-determining and fully irreversible, as the formation of the intermediate cycloadduct (**int**) is highly exergonic ($G = -33$ to -58 kcal/mol). The CO_2 extrusion step (**TS 2**) is then extremely facile and once again very exergonic. All cycloaddition transition states (**TS 1**) have very low activation barriers ($G^\ddagger$ between 9 and 12 kcal/mol). Our calculations also quantitatively reproduce the experimental

selectivities, that is formation of the helical regioisomer is predicted to be favored for sydnones **3a** and **3b**, but unfavored for **3c**.

The TSs leading to the two regioisomers for the representative reaction of **3a** with 3,4-phenanthryne are shown in Figure 4D. The TSs for the other pairs of reactants can be found in the Supporting Information (SI) and display similar arrangements. First, the TS leading to the major helical isomer is more asynchronous, with the shorter forming bond being between the more nucleophilic terminus of the azomethine imine of the sydnone and the more electrophilic (linear) carbon of the aryne.²⁶ Indeed, for the fused sydnones **3a** and **3b**, the nitrogen atom bears more HOMO character than the carbon, while, for *N*-phenyl sydnone **3c**, the opposite is found. Second, the forming bonds are fairly long, indicative of very early TSs, which are in accord with the high exergonicity of the cycloaddition steps and the Hammond postulate. Third, the major TS seems to benefit from stabilizing C–H $\cdots\pi$ interactions (also called face-to-edge π – π interactions) between the polycyclic backbones of the reactants, while the minor TS does not.²⁷

Distortion/interaction analysis²⁸ confirms this behavior: as the TSs are so early, the reactants are barely distorted from their ground-state geometries, and total distortion energies are, at most, 4.0 kcal/mol. As such, even though the helical regioisomer of the cycloadducts and pyrazoles is more sterically encumbered and always higher in energy than the S-shape isomer, this effect is not important in **TS 1** since the reactants are still far from each other. Conversely, interaction energies range from –7.4 to –12.4 kcal/mol, and are greater for the helical vs the S-shape regioisomer. In fact, for the six systems studied, the activation energies correlate with interaction energies, but have no correlation with distortion energies.

¹¹ To confirm the stabilization offered by the dispersive aromatic–aromatic interactions, we computed the binding energies of aromatic dimers in the same geometries as the helical TSs.

¹¹ For the four combinations evaluated, the binding energies were between –0.4 to –1.9 kcal/mol, demonstrating the stabilization offered by the C–H $\cdots\pi$ interactions for the helical TSs, in addition to the more favorable primary orbital interactions. Thus, the TSs that benefit from the most interactions are also earlier, further lowering the cost to distort the reactants. These results indicate that, with other very reactive partners, regioselective cycloadditions might be possible when interactions with polycyclic backbones are present. Indeed, when substituted sydnones **3d–3l** were reacted with **4c**, similar values of regioselectivity (>85/15) were observed in favor of the corresponding [6]- and [7]-helicenes **12a–20a** (Figure 5). While the presence of electron-neutral and -withdrawing substituents on the sydnone does not affect the transformation, disubstituted electron-rich dipole **3h** was poorly reactive and the desired [6]-helicene **16a** was isolated in 18% yield.²⁹ Derivative **20a** with the presence of a chloride substituent offered a useful handle for further functionalization. Under catalytic conditions, the products of Suzuki cross-coupling reaction **21a** and **22a** were isolated in 96% and 56% yield. This preliminary proof-of-concept showed the possibility to further monofunctionalize this helical scaffold by means of metal catalysis which could be of interest for potential applications.

In summary, we have developed a method to access [4]-, [5]-, [6]-, and [7]-helicenes containing pyrazoles through sydnone 1,3-dipolar cycloadditions. This process involves the design and synthesis of *ortho*-substituted polyaromatic sydnones, which are more

nucleophilic than conventional ones, and highlights the first example of regioselective cycloaddition of such mesoionic dipoles with aryne derivatives. Calculations showed that primary orbital interactions and C–H $\cdots\pi$ dispersive interactions control the regioselectivity of this transformation. This reaction will ultimately provide a modular access to substituted derivatives and could be amenable to the synthesis of other helicenes families.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

This work was supported by CEA and ANR (ANR-17-CE07-0035-01) and the U.S. National Science Foundation (CHE-1361104). P.A.C. gratefully acknowledges the Fonds de recherche du Québec - Nature et Technologies for a postdoctoral fellowship. Computations were performed on the Hoffman2 cluster at UCLA. G.M. acknowledges the NIH Minority Biomedical Research Support (Grant 1 SC3 GM089589-08) and the Henry Dreyfus Teacher-Scholar Award for financial support. The authors thank C. Chollet, E. Marcon, A. Goudet, S. Lebrequier, and D.-A. Buisson (DRF-JOLIOT-SCBM, CEA) for excellent analytical support. We wish to acknowledge Dr. P. Dauban (ICSN, CNRS, France) for helpful discussions.

REFERENCES

- (1). (a)Shen Y; Chen C-F Helicenes: synthesis and applications. *Chem. Rev* 2012, 112, 1463–1535. [PubMed: 22017405] (b)Gingras M One hundred years of helicene chemistry. Part 1: non-stereoselective syntheses of carbohelicenes. *Chem. Soc. Rev* 2013, 42, 968–1006. [PubMed: 23151799] (c)Gingras M; Félix G; Peresutti R One hundred years of helicene chemistry. Part 2: stereoselective syntheses and chiral separations of carbohelicenes. *Chem. Soc. Rev* 2013, 42, 1007–1050. [PubMed: 23151610] (d)Gingras M One hundred years of helicene chemistry. Part 3: applications and properties of carbohelicenes. *Chem. Soc. Rev* 2013, 42, 1051–1095. [PubMed: 23151680] (e)Chen C-F; Shen Y Helicene Chemistry, From Synthesis to Applications, Springer-Verlag: Berlin, Heidelberg, Germany, 2017.
- (2). (a)Takenaka N; Chen J; Captain B; Sarangthem RS; Chandrakumar A Helical Chiral 2-aminopyridinium ions: a new class of hydrogen bond donor Catalysts. *J. Am. Chem. Soc* 2010, 132, 4536–4537. [PubMed: 20232867] (b)Aillard P; Voituriez A; Dova D; Cauteuruccio S; Licandro E; Marinetti A Phosphathiahelicenes: synthesis and uses in enantioselective gold catalysis. *Chem. - Eur. J* 2014, 20, 12373–12376 and references cited therein. [PubMed: 25113927] (c)González-Fernández E; Nicholls LDM; Schaaf LD; Farés C; Lehmann CW; Alcarazo M Enantioselective synthesis of [6]carbohelicenes. *J. Am. Chem. Soc* 2017, 139, 1428–1431. [PubMed: 28112916] (d)Ferreira M; Naulet G; Gallardo H; Dechambenoit P; Bock H; Durola F A Naphtho-fused double [7]helicene from a maleate-bridged Chrysene Trimer. *Angew. Chem., Int. Ed* 2017, 56, 3379–3382.
- (3). (a)Hatakeyama T; Hashimoto S; Oba T; Nakamura M Azaboradibenzo[6]helicene: carrier inversion induced by helical homochirality. *J. Am. Chem. Soc* 2012, 134, 19600–19603. [PubMed: 23167918] (b)Yang Y; da Costa RC; Fuchter MJ; Campbell AJ Circularly polarized light detection by a chiral organic semiconductor transistor. *Nat. Photonics* 2013, 7, 634–638.
- (4). (a)Otani T; Tsuyuki A; Iwachi T; Someya S; Tateno K; Kawai H; Saito T; Kanyiva KS; Shibata T Facile two-step synthesis of 1,10-phenanthroline-derived polyaza[7]helicenes with high fluorescence and CPL efficiency. *Angew. Chem, Int. Ed.* 2017, 56, 3906–3910. [PubMed: 28267252] (b)Collins SK; Vachon MP Unlocking the potential of thiaheterohelicenes: chemical synthesis as the key. *Org. Biomol. Chem* 2006, 4, 2518–2524. [PubMed: 16791312]
- (5). (a)Míšek J; Teplý F; Stará IG; Tichý M; Šaman D; Císa ová I; Vojtíšek P; Starý IA Straightforward route to helically chiral N-heteroaromatic compounds: practical synthesis of racemic 1,14-diaza[5]helicene and optically pure 1- and 2-aza[6]helicenes. *Angew. Chem, Int. Ed.* 2008, 47, 3188–3191. [PubMed: 18338415] (b)Takenaka N; Sarangthem RS; Captain B Helical chiral pyridine N-oxides: a new family of asymmetric catalysts. *Angew. Chem, Int. Ed.*

- 2008, 47, 9708–9710. [PubMed: 18988214] (c) Kaneko E; Matsumoto Y; Kamikawa K Synthesis of azahelicene N-oxide by palladium-catalyzed direct C–H annulation of a pendant (Z)-bromovinyl side chain. *Chem. - Eur. J* 2013, 19, 11837–11841. [PubMed: 23873825] (d) Bosson J; Labrador GM; Pascal S; Miammay F-A; Yushchenko O; Li H; Bauffier L; Sojic N; Tovar RC; Muller G; Jacquemin D; Laurent AD; Guennic BL; Vauthey E; Lacour J Physicochemical and electronic properties of cationic [6]helicenes: from chemical and electrochemical stabilities to far-red (polarized) luminescence. *Chem. - Eur. J* 2016, 22, 18394–18403. [PubMed: 27885721] (e) Nakamura K; Furumi S; Takeuchi M; Shibuya T; Tanaka K Enantioselective synthesis and enhanced circularly polarized luminescence of S-shaped double azahelicenes. *J. Am. Chem. Soc* 2014, 136, 5555–5558. [PubMed: 24670158] (f) Nejedlý J; Šámal M; Rybáček J; Tobrmanová M; Szydło F; Coudret C; Neumeier M; Vacek J; Chocholoušová JV; Budšínský M; Šaman D; Bednářová L; Sieger L; Stará IG; Starý I Synthesis of long oxahelicenes by polycyclization in a flow reactor. *Angew. Chem, Int. Ed.* 2017, 56, 5839–5843. [PubMed: 28417538]
- (6). (a) Goto K; Yamaguchi R; Hiroto S; Ueno H; Kawai T; Shinokubo H Intermolecular oxidative annulation of 2-aminoanthracenes to diazaacenes and aza[7]helicenes. *Angew. Chem, Int. Ed.* 2012, 51, 10333–10336. [PubMed: 22965455] (b) Shi L; Liu Z; Dong G; Duan L; Qiu Y; Jia J; Guo W; Zhao D; Cui D; Tao X Synthesis, Structure, Properties, and application of a carbazole-based diaza[7]helicene in a deep-blue-emitting OLED. *Chem. - Eur. J* 2012, 18, 8092–8099. [PubMed: 22592951] (c) Upadhyay GM; Talele HR; Bedekar AV Synthesis and photophysical properties of aza[n]helicenes. *J. Org. Chem* 2016, 81, 7751–7759. [PubMed: 27439443] (d) Krzeszewski M; Kodama T; Espinoza EM; Vullev VI; Kubo T; Gryko DT Nonplanar butterfly-shaped π -expanded pyrrolopyrroles. *Chem. - Eur. J* 2016, 22, 16478–16488. [PubMed: 27659591]
- (7). Sakamaki D; Kumano D; Yashima E; Seki S A Facile and versatile approach to double N-heterohelicenes: tandem oxidative CN couplings of N-heteroacenes via cruciform dimers. *Angew. Chem, Int. Ed.* 2015, 54, 5404–5407. [PubMed: 25772706]
- (8). Jiřková M; Šaman D; Koval D; Kašík V; Klepetář B; Čísarová I; Teplý F Modular Synthesis of Helicene-Like Compounds Based on the Imidazolium Motif. *Eur. J. Org. Chem* 2014, 2014, 5681–5685.
- (9). (a) Browne DL; Harrity JPA Recent developments in the chemistry of sydnones. *Tetrahedron* 2010, 66, 553–568. (b) Decuy-père E; Plougastel L; Audisio D; Taran F Sydnone–lkyne cycloaddition: applications in synthesis and bioconjugation. *Chem. Commun* 2017, 53, 11515–11527.
- (10). For important pioneering work on helicenes formation by [4 + 2] cycloaddition of benzyne, see: (a) del Mar Real M; Pérez Sestelo J; Sarandeses LA Inner–outer ring 1,3-bis(trimethylsilyloxy)-1,3-dienes as useful intermediates in the synthesis of helicenes. *Tetrahedron Lett.* 2002, 43, 9111–9114. (b) Truong T; Daugulis O Divergent reaction pathways for phenol arylation by arynes: synthesis of helicenes and 2-arylphenols. *Chem. Sci* 2013, 4, 531–535. [PubMed: 24077102]
- (11). See SI for additional details.
- (12). For the very first report, see: (a) Gotthardt H; Huisgen R; Knorr R 1.3-Dipolare cycloadditionen, XXXVIII. Reaktionen der sydnone mit benz-in und mit einigen heteromehrfachbindungen. *Chem. Ber* 1968, 101, 1056–1058. For other pioneering reports, see: (b) Nakazawa S; Kiyosawa T; Hirakawa K; Kato H Selectivity in the thermal and photochemical fragmentation of the cycloadduct from benzyne and a mesoionic thiazol-4-one. *J. Chem. Soc., Chem. Commun* 1974, 621a. (c) Kato H; Nakazawa S; Kiyosawa T; Hirakawa K Heterocycles by cycloaddition. Part II. Cycloaddition–extrusion reactions of five-membered mesoionic compounds with benzyne: preparation of benz[c]azole and benzo[c]thiophen derivatives. *J. Chem. Soc., Perkin Trans 1* 1976, 672–675. (d) Tsuge O; Samura H The formation of 1H-dibenzo[b,g][1,4,5]triazapentalene from 2-(o-nitrophenyl)- and 2-(o-azidophenyl)-2H-indazole. *Org. Prep. Proced. Int* 1974, 6, 161–167.
- (13). (a) Wu C; Fang Y; Larock RC; Shi F Synthesis of 2H-indazoles by the [3 + 2] cycloaddition of arynes and sydnones. *Org. Lett* 2010, 12, 2234–2237. [PubMed: 20394430] (b) Fang Y; Wu C; Larock RC; Shi F Synthesis of 2H-indazoles by the [3 + 2] dipolar cycloaddition of sydnones with arynes. *J. Org. Chem* 2011, 76, 8840–8851. [PubMed: 21970468] (c) Soares MIL; Nunes CM; Gomes CSB; Pinho e Melo TMVD Thiazolo[3,4-b]indazole-2,2-dioxides as masked extended dipoles: pericyclic reactions of benzodiazafulvenium methides. *J. Org. Chem* 2013, 78, 628–637. [PubMed: 23215255]

- (14). For a previous synthesis of compound 5, see: (a) Zhao J; Wu C; Li P; Ai W; Chen H; Wang C; Larock RC; Shi F Synthesis of pyrido[1,2-b]indazoles via aryne [3 + 2] cycloaddition with Ntosylpyridinium Imides. *J. Org. Chem* 2011, 76, 6837–6843. [PubMed: 21755977] (b) Zhao J; Li P; Wu C; Chen H; Ai W; Sun R; Ren H; Larock RC; Shi F Aryne [3 + 2] cycloaddition with Nsulfonylpyridinium imides and in situ generated N-sulfonylisoquinolinium imides: a potential route to pyrido[1,2-b]indazoles and indazolo[3,2-a]isoquinolines. *Org. Biomol. Chem* 2012, 10, 1922–1930. [PubMed: 22278679] (c) Zheng Q-Z; Feng P; Liang Y-F; Jiao N Pd-Catalyzed tandem C–H azidation and N–N bond formation of arylpyridines: a direct approach to pyrido[1,2-b]indazoles. *Org. Lett* 2013, 15, 4262–4265. [PubMed: 23915282]
- (15). In contrast to [6]-carbohelicenes, [6]-helicene 9a is not configurationally stable at room temperature. Computed racemization energy barrier is 22.5 kcal/mol (B3LYP/6–31G(d)). Enantiomers of 9a were resolved by chiral-phase HPLC, and a rapid racemization was observed. For additional information regarding the racemization barriers of other pyrazolohelicenes herein reported, see ref 11. For a nice overview on carbohelicenes racemization, see: Barroso J; Cabellos JL; Pan S; Murillo F; Zarate Z; Fernandez-Herrera MA; Merino G Revisiting the racemization mechanism of helicenes. *Chem. Commun* 2018, 54, 188–191.
- (16). The measured racemization barrier for pyrazolo[7]helicene **10a** was found to be 32.7 kcal/mol. This value is in agreement with the calculated racemization energy barrier for **10a**: 31.6 kcal/mol (B3LYP/6–31G(d)). See ref 11 for details **10a10a**.
- (17). The degree of CPL is given by the luminescence dissymmetry ratio, $g_{lum}(\lambda) = 2 \ I/I = 2(I_L - I_R)/(I_L + I_R)$, where I_L and I_R refer, respectively, to the intensity of left and right circularly polarized emissions s_{lum} LRLRL.
- (18). (a) Kaseyama T; Furumi S; Zhang X; Tanaka K; Takeuchi M Hierarchical assembly of a phthalhydrazide-functionalized helicene. *Angew. Chem, Int. Ed.* 2011, 50, 3684–3687. [PubMed: 21416571] (b) Sawada Y; Furumi S; Takai A; Takeuchi M; Noguchi K; Tanaka K Rhodium-catalyzed enantioselective synthesis, crystal structures, and photophysical properties of helically chiral 1,1'-bitriphenylenes. *J. Am. Chem. Soc* 2012, 134, 4080–4083. [PubMed: 22335235] (c) Shen C; Anger E; Srebro M; Vanthuyne N; Deol KK; Jefferson TD; Muller G; Williams JAG; Toupet L; Roussel C; Autschbach J; Réau R; Crassous J Straightforward access to mono- and bis-cycloplatinated helicenes displaying circularly polarized phosphorescence by using crystallization resolution methods. *Chem. Sci* 2014, 5, 1915–1927. [PubMed: 24855556]
- (19). We are aware of only two single examples of regioselective sydnone-aryne cycloaddition (>85/15), in the presence of 3-silylbenzyne: (a) Ikawa T; Masuda S; Takagi A; Akai S 1,3- and 1,4-Benzdiyne equivalents for regioselective synthesis of polycyclic heterocycles. *Chem. Sci* 2016, 7, 5206–5211. [PubMed: 30155171] (b) Ikawa T; Masuda S; Nakajima H; Akai S 2-(Trimethylsilyl)phenyl trimethylsilyl ethers as stable and readily accessible benzyne precursors. *J. Org. Chem* 2017, 82, 4242–4253. [PubMed: 28247765]
- (20). (a) Houk KN; Sims J; Duke RE; Strozier RW; George JK Frontier molecular orbitals of 1,3 dipoles and dipolarophiles. *J. Am. Chem. Soc* 1973, 95, 7287–7301. (b) Houk KN; Sims J; Watts CR; Luskus LJ Origin of reactivity, regioselectivity, and periselectivity in 1,3-dipolar cycloadditions. *J. Am. Chem. Soc* 1973, 95, 7301–7315. (c) Padwa A; Burgess EM; Gingrich HL; Roush DM On the problem of regioselectivity in the 1,3-dipolar cycloaddition reaction of munchedones and sydnones with acetylenic dipolarophiles. *J. Org. Chem* 1982, 47, 786–791.
- (21). McMahon TC; Medina JM; Yang Y-F; Simmons BJ; Houk KN; Garg NK Generation and regioselective trapping of a 3,4-piperidyne for the synthesis of functionalized heterocycles. *J. Am. Chem. Soc* 2015, 137, 4082–4085. [PubMed: 25768436]
- (22). Shah TK; Medina JM; Garg NK Expanding the strained alkyne toolbox: generation and utility of oxygen-containing strained alkynes. *J. Am. Chem. Soc* 2016, 138, 4948–4954. [PubMed: 26987257]
- (23). Noteworthy, in the presence of unsymmetric cyclooctynes, sydnones have been reported to deliver a 1:1 mixture of regioisomers: Narayanam MK; Liang Y; Houk KN; Murphy JM Discovery of new mutually *orthogonal* bio*orthogonal* cycloaddition pairs through computational screening. *Chem. Sci* 2016, 7, 1257–1261. [PubMed: 29910881]
- (24). (a) Cheong PH-Y; Paton RS; Bronner SM; Im G-YJ; Garg NK; Houk KN Indolyne and aryne distortions and nucleophilic regioselectivities. *J. Am. Chem. Soc* 2010, 132, 1267–1269. [PubMed: 20058924] (b) Medina JM; Mackey JL; Garg NK; Houk KN The role of aryne

distortions, steric effects, and charges in regioselectivities of aryne reactions. *J. Am. Chem. Soc.* 2014, 136, 15798–15805. [PubMed: 25303232]

- (25). The nature of pyrazoles **11a** and **11b** was unambiguously established by NOE and X-ray crystallography (see SI for details) **11a11b**.
- (26). In all cases, the reactions are HOMO (sydnone) – LUMO (aryne) controlled. See SI for details.
- (27). For a review on aromatic interactions as control elements in stereoselective reactions, see: Krenske EH; Houk KN Aromatic interactions as control elements in stereoselective organic reactions. *Acc. Chem. Res.* 2013, 46, 979–989. [PubMed: 22827883]
- (28). Bickelhaupt FM; Houk KN Analyzing reaction rates with the distortion/interaction-activation strain model. *Angew. Chem, Int. Ed.* 2017, 56, 10070–10086. [PubMed: 28447369]
- (29). Unreacted sydnone **3h** was present in the reaction crude even after a prolonged reaction time **3h**.

Helical pyrazoles

- Novel scaffold
- Straightforward synthesis
- Divergent approach
- Unprecedented regioselectivity

Figure 1.
Design of pyrazole-based helicenes.

Figure 2.

(A) One-pot synthesis of polycyclic sydnones **3a** and **3b**; (B) aryne precursor; (C) synthesis of helical pyrazoles **5** and **6**. (D) Synthesis of helical pyrazoles **7a**, **8a**, **9a**, and **10a**. * Isomer ratio was measured by ^1H NMR of the crude mixture.

Figure 3.

(A) Molecular structure of **10a**: (I) top and (II) side views. Molecular structure of **10b**: (III) top and (IV) side views. Ellipsoids are set at 40% probability; hydrogen atoms are omitted for clarity. (B) UV (gray curve) and CD spectra of (+)-**10a** (black curve) and (–)-**10a** (red curve); (C) CPL (upper curve) and total luminescence (lower curves) spectra of (+)-**10a** (black curve) and (–)-**10a** (red curve) in degassed dichloromethane at 295 K, upon excitation at 430 nm. See Supporting Information for details.

Figure 4.

(A) DFT-optimized structures of polycyclic arynes derived from **4b** and **4c**. (B) Reaction between nonplanar *N*-phenyl sydnone **3c** and 3,4-phenanthryne precursor **4c**. * Isomer ratio was measured by ^1H NMR of the crude mixture. (C) Energy profile for the reaction of **3a** with 3,4-phenanthryne to form **9a** or **9b**. Free energies (*enthalpies*) are in kcal/mol and were obtained at the M06–2X/6–311+G(2d,2p)/SMD(THF)/M06–2X/6–31+G(d,p) level of theory. (D) Cycloaddition TSs leading to regioisomers **9a** and **9b**. M06–2X/6–311+G(2d,2p)/SMD(THF)/M06–2X/6–31+G(d,p). Free energies in kcal/mol.

Figure 5. Synthesis of helical pyrazoles **12a–22a**. * Isomer ratio was measured by ^1H NMR of the crude mixture. n.i. = not isolated. ^a See Supporting Information for detailed conditions.