Published in final edited form as:

Aerosp Med Hum Perform. 2019 May 01; 90(5): 475–479. doi:10.3357/AMHP.5284.2019.

Cooling to facilitate metabolic suppression in healthy individuals

Jon C. Rittenberger, MD, MS, Kathryn L. Flickinger, MS, Alexandra Weissman, MD, Melissa Repine, BS, Jonathan Elmer, MD, Francis X. Guyette, MD, MPH, Clifton W. Callaway, MD, PhD

University of Pittsburgh, Applied Physiology Laboratory, Department of Emergency Medicine

Abstract

Background—Carbon dioxide (CO₂) toxicity could be catastrophic for astronauts. Suppressing metabolism by lowering body temperature decreases CO₂ production and may facilitate rescue in the event of a crippled ship. Lowering body temperature requires shivering suppression. We evaluated dexmedetomidine to facilitate cooling of healthy individuals.

Methods—Following consent, we administered a 1 mcg/kg bolus of dexmedetomidine followed by continuous infusion (0.5–1.4 mcg/kg/hr) for 3 hours of cooling. We cooled subjects using a bolus of 30cc/kg of 4°C saline followed by surface cooling. We measured vital signs, thermal and comfort scales, sedation and shivering for 3 hours and during recovery. ANOVA evaluated changes in measures over time.

Results—Nine subjects completed the study. Mean age was 31 (SD 8) years, mean mass was 71 (SD 14) kg, height of 168 (SD 9) cm and body mass index of 25 (SD 3). Median time to 1°C drop in core temperature was 16 (IQR 15, 32) minutes. Temperature changed over time with median lowest temperature was 33.1°C (IQR 32.8°C, 34.1°C). Neither heart rate nor diastolic blood pressures changed over time. Systolic blood pressure decreased over time. Subjects responded to verbal stimuli and completed tasks throughout the protocol. During cooling and maintenance, subjects reported discomfort and the sensation of being cold.

Conclusion—Dexmedetomidine facilitates shivering suppression during prolonged cooling in healthy individuals. Subjects are easily roused, have mild decreases in systolic blood pressure and note sensations of discomfort and cold. Cooling to suppress metabolism is a feasible countermeasure to prolong astronaut endurance.

Keywords

Hypothermia; cognition; metabolism; dexmedetomidine

Introduction

One potential threat to astronaut safety is a carbon dioxide (CO₂) scrubbing module failure. At baseline, astronauts in the International Space Station breathe atmosphere containing

approximately 4 mmHg CO₂, compared to terrestrial CO₂ of approximately 0.3 mmHg. CO₂ levels greater than 5 mmHg cause poor concentration, fatigue, dizziness, headaches, and difficulty sleeping.⁷ Thus, a malfunctioning CO₂ scrubber could rapidly result in impairment and eventually threaten survival. Suppressing astronaut metabolism and CO₂ production could slow CO₂ accumulation and prolong survival time while correcting or rescuing from such a malfunction. Similarly, metabolic suppression could extend time available for rescue, facilitating additional options in situations like the Apollo 13 or STS-107 missions. Preclinical data suggest that metabolic rate decreases 7-8% for each 1°C decrease in core temperature.⁸ Thus, a small change in core temperature could significantly reduce energy needs and CO₂ production by the crew.

Forced reductions of core body temperature below about 36.5°C in normal humans triggers shivering, which increases metabolism and prevents further reductions in core temperature unless extraordinary measures are applied. Lowering core body temperature as a strategy to reduce metabolism requires reliable and safe shivering suppression. A number of anesthetic medications can suppress shivering, but these medications also cause sedation or respiratory depression.

We have tested the ability of several drugs to suppress shivering during forced cooling of healthy humans.^{6,9} Dexmedetomidine is an alpha-2-adrenergic agonist that reduces shivering without impairing breathing, and that also allows rapid awakening.⁴ Dexmedetomidine induces a drowsy state that resembles physiological sleep rather than the anesthetic or intoxicated state created by other drugs.¹ Whether dexmedetomidine is safe and effective for maintaining lower body temperature in spontaneously breathing humans over prolonged periods is unknown. We hypothesized that a bolus of dexmedetomidine followed by infusion will suppress shivering during forced cooling of healthy individuals for 3 hours with mild side effects.

Methods

Subjects

The University of Pittsburgh Human Research Protection Organization (formerly IRB) approved this study. All subjects provided written informed consent prior to participation. We recruited healthy individuals between the ages of 18-49 years old. Subject screening included a physical examination to confirm and an electrocardiogram to exclude the presence of conduction abnormalities. Each subject self-reported they were free from heart disease, vascular disease, renal or hepatic impairment, or any allergy to dexmedetomidine. Women could not be pregnant or currently breastfeeding and provided a negative urine pregnancy test on each protocol day.

Procedure

Experiments took place in the Applied Physiology Laboratory at the University of Pittsburgh.⁴ Intervention days were separated by at least one week. Subjects refrained from food, alcohol, and tobacco for 6 hours prior to the study protocol. We allowed subjects to drink water. After obtaining baseline post-void weight and body anthropomorphic

measurements, we placed 18g or 20g intravenous catheters bilaterally in veins proximal to the wrist. We monitored subjects using three-lead electrocardiogram, continuous pulse oximetry, nasal cannula continuous waveform capnography, and non-invasive blood pressure. We placed a naso-esophageal temperature probe was placed to monitor core (esophageal) body temperature. Surface cooling consisted of water circulating pads with a gel adherent to the skin (Arctic Sun, Medivance, Inc. Louisville, CO), which were placed on torso and legs. All experiments occurred between 8am and 2pm with subjects lightly clothed in a lab with ambient temperature between 20-22°C.

We recorded thermal sensation, comfort, sedation, and shivering scales every five minutes. Subjects reported comfort and thermal sensation scores verbally or by pointing to a score on a placard with each scale. The general (overall) comfort scale (1-4) had anchors of 1, “comfortable,” and 4 “very uncomfortable”. Thermal sensation scale (1-5) had anchors by 1, “comfortable,” and 5 “very cold.” An investigator rated sedation using the Richmond Agitation and Sedation Scale (RASS).^{5,11} The RASS is a 10-point scale ranging from -5 (unresponsive to voice or painful stimulation) to 4 (combative and danger to self or staff). An investigator visually and manually assessed intensity of shivering using the Bedside Shivering Assessment Scale (BSAS).² The BSAS (0-3) ranges from 0 (no shivering) to 3 (severe shivering- generalized or sustained upper/lower extremity shivering).

After placement of all monitors and baseline measurements, we administered a 1 mcg/kg bolus of dexmedetomidine (Hospira, Lake Forest, IL) over 10 minutes followed by an infusion starting at 0.5 mg/kg/hr. The infusion was increased by 0.2 mg/kg/hr every 5 minutes is the patient was shivering (BSAS = 1) or had an increase in core temperature to a maximum dose of 1.4mg/kg/hr.

To induce cooling, we infused 30 ml/kg cold (4°C) normal saline intravenously via a pressure bag (approximately 10 min/L) through an 18g or 20g IV in the antecubital fossa. Maximum bolus volume was 2 L. Following the cold saline bolus, we activated the surface cooling pads at their maximal cooling rate (water circulating at 4°C).

In the first two experiments, the cold saline bolus started contemporaneously with the bolus of dexmedetomidine. One subject experienced vagal symptoms of malaise and nausea, with profound bradycardia during the infusions, which rapidly resolved with stopping both infusions. This subject did not complete the protocol and is not included in the final data. After this subject, we changed the protocol to administer the cold saline bolus after completion of the initial dexmedetomidine bolus.

Investigators stopped drug administration or cooling if heart rate declined to less than 45 bpm for 2 consecutive minutes, SBP declined to <90 mmHg for 5 consecutive minutes, tissue pulse oximetry (SpO₂) declined to <90% for one minute, or the subject became deeply sedated (RASS-4). Dexmedetomidine infusion continued for up to 3 hours. After 3 hours, surface cooling stopped, and subjects were allowed blankets to promote passive warming. After stopping dexmedetomidine, we monitored subjects for a minimum of 120 minutes or until RASS was 0 and they were able to eat, void and ambulate without difficulty.

Statistical Analysis

Data were analyzed using descriptive statistics. Time-course analyses were completed using Analysis of variance (ANOVA) STATA 14.0 (College Station, TX).

Results

Ten subjects (9 men) started the protocol, and one subject failed to complete the experiment after having severe bradycardia during simultaneous infusion of dexmedetomidine and cold saline. After amending the protocol to separate these infusions, nine subjects (8 men) completed 20 experiments which comprise the final data for analysis. The mean age was 31 (SD 8) years. Mean mass was 71 (SD 14) kg, height of 168 (SD 9) cm and body mass index of 25 (SD 3). One subject experienced asymptomatic atrial fibrillation, which resolved when core temperature became $>36^{\circ}\text{C}$.

Median time to 1°C drop in core temperature was 16 (IQR 15, 32) minutes. Temperature changed over time with median lowest temperature of 33.1°C (IQR 32.8°C , 34.1°C ; $F(11)=14.4$, $p<0.0001$). [Figure 1]

The median temperature at the end of protocol was 33.8°C (IQR 33.1°C , 34.3°C). Neither heart rate [baseline 68 (SD 8) bpm, nadir 47 (SD 15) bpm; $F(11)=0.73$, $p=0.71$] nor diastolic blood pressures [baseline 75 (SD 13) mmHg, nadir 59 (SD 8) mmHg; $F(11)=1.76$, $p=0.007$] changed over time. Systolic blood pressure decreased from a pre-intervention measurement of 129 (SD 11) mmHg to a nadir of 105 (SD 9) mmHg over time ($F(11)=2.79$, $p=0.0035$). [Figure 2]

Shivering was mild (BASS -1-2) and increased during induction (BSAS 1; SD 1.3 at 30 minutes) and maintenance of cooling and decreased after the cooling pads were removed (BSAS 0; SD 0.3 at 210 minutes; $F(11)=3.57$, $p=0.0003$). [Figure 3a]

Most subjects experienced mild (RASS -1 to -2) sedation during induction and maintenance of cooling ($F(11)=4.38$, $p=0.0001$). [Figure 3b] Sedation resolved after dexmedetomidine infusion stopped at minute 180. During cooling and maintenance, subjects experienced discomfort and a sensation of being cold (Thermal: $F(11)=10.01$, $p=0.0001$; Comfort: $F(11)=14.35$, $p=0.0001$). [Figure 4]

These symptoms improved with the cessation of cooling but required time to abate entirely.

Discussion

Dexmedetomidine facilitates shivering suppression, as measured by BSAS, during prolonged cooling in healthy individuals. A bolus and infusion of dexmedetomidine suppresses shivering adequately to force core body temperature to $33\text{--}34^{\circ}\text{C}$. Interestingly, the reduction in core body temperature is long-lasting. Core temperature rarely returned to normal before the end of monitoring time. Despite this, subjects were able to eat, ambulate and void without difficulty.

Subjects are easily roused and have mild decreases in systolic blood pressure. These data suggest that individuals could participate in self-rescue during hypothermia. While subjects appeared grossly returned to baseline within hours after cooling, we did not complete detailed cognitive or psychomotor testing, such as the Cognition Test Battery.¹⁰ Long-lasting effects or hangover from metabolic suppression regimens will be important areas to investigate in the future.

The timing of interventions is important. Simultaneous bolus of cold saline and dexmedetomidine caused profound bradycardia in one subject. Our impression from all of these experiments is that cold saline infusion causes a significant vagal response, as most patients report a deep visceral sensation during the infusion. Dexmedetomidine has well-documented direct inhibition on cardiac AV nodal conduction.³ When administered together, the increased reflex vagal tone and direct AV block profoundly reduce atrial-ventricular conduction, and subjects with slow ventricular escape rhythms will not tolerate the combined infusion. The simplest solution was to separate these infusions in time, and we saw no recurrence after doing so. Future experiments could test whether pharmacological interventions might also reduce this risk.

We determined potential side effects to address for practical application of forced cooling, particularly if this is used in remote locations. For example, asymptomatic atrial fibrillation occurred in one subject but resolved during rewarming. This provoked dysrhythmia was likely a result of hypothermia rather than drug infusion, and it has little consequence over a few hours. However, a cardiac rhythm change would not be acceptable over days or longer. The decrease in systolic blood pressure and bradycardia did not cause symptoms. These changes in vital signs are consistent with the decreased metabolic rate at lower core body temperature. These predictable effects define some minimum monitoring that should be in place for safe application.

During induction and maintenance of cooling, subjects experienced the sensation of being cold and rated it as moderate to severe in the level of discomfort. In fact, one subject experienced significant shivering, resulting in tachycardia during cooling. This illustrates the need for adequate shivering suppression to facilitate cooling. Future studies should evaluate methods to mitigate this perception as it may help facilitate temperature modulation. Medications that affect the transient receptor potential cation channel subfamily M member 8 (TrpM8) and transient receptor potential cation channel subfamily V member 1 (TrpV1) are associated with skin perception of cold and are thus potential targets. For longer cooling durations or frequent cooling bouts to be pursued, this will need to be addressed.

Our study is limited in its duration of cooling. Any application to a mission to space or potential rescue operations would require hours to days of cooling to derive maximal benefit.

Prolonged cooling of non-anesthetized healthy individuals is feasible using intravascular and surface cooling methods. Dexmedetomidine facilitates prolonged cooling by suppressing shivering with tolerable side effects in most subjects. Cooling to suppress metabolism is a feasible countermeasure to prolong astronaut endurance in the event of a failed CO₂

scrubber. It may also be considered in other life support or vehicle failures necessitating prolonged exposure to CO₂.

References

1. Akeju O, Hobbs LE, Gao L, Burns SM, Pavone KJ, Plummer GS, Walsh EC, Houle TT, Kim S, Bianchi MT, Ellenbogen JM, Brown EN. Dexmedetomidine promotes biomimetic nonrapid eye movement stage 3 sleep in humans: a pilot study. *Clinical Neurophysiology*. 2018;129:69–78. PMID: 29154132. [PubMed: 29154132]
2. Badjatia N, Strongilis E, Gordon E, Prescutti M, Fernandez L, Fernandez A, et al. Metabolic impact of shivering during therapeutic temperature modulation: The bedside shivering assessment scale. *Stroke; a journal of cerebral circulation*. 2008;39:3242–3247
3. Bloor BC, Ward DS, Belleville JP, Maze M. Effects of intravenous dexmedetomidine in humans. *Anesthesiology*. 1992;77:1134–1142. [PubMed: 1361311]
4. Callaway CW, Elmer J, Guyette FX, Molyneaux BJ, Anderson KB, Empey PE, Gerstel SJ, Holquis K, Repine MJ, Rittenberger JC. Dexmedetomidine reduces shivering during mild hypothermia in waking subjects. *PLoS One* 2015 8 3;10(8):e0129709. [PubMed: 26237219]
5. Ely EW, Truman B, Shintani A, Thomason JW, Wheeler AP, Gordon S, et al. Monitoring sedation status over time in ICU patients: Reliability and validity of the Richmond agitation-sedation scale (RASS). *JAMA*. 2003;289:2983–2991 [PubMed: 12799407]
6. Hostler D, Northington WE, Callaway CW. High-dose diazepam facilitates core cooling during cold saline infusion in healthy volunteers. *Appl Physiol Nutr Metab*. 2009 Aug;34(4):582–6. [PubMed: 19767791]
7. Law J, Young M, Alexander D, Mason SS, Wear ML, Mendez CM, Stanley D, Ryder VM, Van Baaler M. Carbon dioxide physiological training at NASA. *Aerosp Med Hum Perform*. 2017;88:897–902. [PubMed: 28923137]
8. Michenfelder JD, Theye RA. Hypothermia: effect on canine brain and whole-body metabolism. *Anesthesiology*. 1968;29(6):1107–12. PMID:5726746. [PubMed: 5726746]
9. Moore TM, Callaway CW, Hostler D. Core temperature cooling in healthy volunteers after rapid intravenous infusion of cold and room temperature saline solution. *Ann Emerg Med*. 2008 2;51(2):153–9. [PubMed: 18045737]
10. Moore TM, Basner M, Nasrini J, Hermosillo E, Kabadi S, Roalf DR, McGuire S, Ecker AJ, Ruparel K, Port AM, Jackson CT, Dinges DF, Gur RC. Validation of the Cognition Test Battery for spaceflight in a sample of highly educated adults. *Aerosp Med Hum Perform*. 2017;88:937–946. [PubMed: 28923143]
11. Sessler CN, Gosnell MS, Grap MJ, Brophy GM, O'Neal PV, Keane KA, et al. The Richmond agitation-sedation scale: Validity and reliability in adult intensive care unit patients. *American journal of respiratory and critical care medicine*. 2002;166:1338–1344 [PubMed: 12421743]

Figure 1.
Core temperature over time. Error bars represent one standard deviation.

Figure 2.

A. Heart rate over time. B. Systolic and diastolic blood pressure over time. Error bars represent one standard deviation.

Figure 3.

A. Bedside Shivering Assessment Scale (BSAS) over time. A higher score delineates more shivering. **B.** Richmond Agitation Sedation Scale (RASS) over time. A lower score indicates deeper sedation. Error bars represent one standard deviation.

Figure 4.

A. Comfort scale over time. A higher score delineates more discomfort. **B.** Thermal sensation over time. A higher score indicates a sensation of more intense cold by the subject. Error bars represent one standard deviation.