Published in final edited form as:

J Am Chem Soc. 2019 April 03; 141(13): 5444–5460. doi:10.1021/jacs.9b00328.

Pseudoephedrine-Derived Myers Enolates: Structures and Influence of Lithium Chloride on Reactivity and Mechanism

Yuhui Zhou, Janis Jermaks, Ivan Keresztes, Samantha N. MacMillan, David B. Collum*

Department of Chemistry and Chemical Biology, Baker Laboratory, Cornell University, Ithaca, New York 14853–1301

Abstract

The structures and reactivities of pseudoephedrine-derived dianionic Myers enolates are examined. A combination of NMR and IR spectroscopic, crystallographic, and computational data reveal that the homoaggregated dianions form octalithiated tetramers displaying S_4 -symmetric Li_8O_8 cores and overall C_2 symmetry. Computational and isotopic labeling studies reveal strong N–Li contacts in the carboxamide enolate moiety. The method of continuous variations proves deceptive, as octalithiated tetrameric homoaggregates afford hexalithiated trimeric heteroaggregates. A lithium diisopropylamide–lithium enolate mixed aggregate is found to be a C_2 -symmetric hexalithiated species incorporating two enolate dianions and two lithium diisopropylamide (LDA) subunits. Structural and rate studies show that lithium chloride has little effect on the dynamics of the enolate homoaggregates but forms adducts of unknown structure. Rate studies of alkylations indicate that the aging of the aggregates can have effects spanning orders of magnitude. The LiCl –enolate adduct dramatically accelerates the reaction but requires superstoichiometric quantities owing to putative autoinhibition. Efforts and progress toward eliminating the requisite large excess of LiCl are discussed.

Graphical Abstract

Introduction

In 1994, Myers and co-workers reported that dianions of type **2** derived from acylated pseudoephedrines (**1**) undergo functionalizations with a range of electrophiles in high yields and selectivities (eq 1).¹ The products are deprotected by facile hydrolysis that appears to benefit from anchimeric assistance from the pendant hydroxy moiety noted previously by Evans using pyrrolidinol-derived amides.² A key observation was that 5.0 equiv of LiCl

*David B. Collum, dbc6@cornell.edu.

Supporting Information: Spectroscopic, crystallographic, computational, and rate data as well as additional Job plots. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interests.

was essential for optimal yields. Although no direct evidence attests to the role of LiCl, an artist's rendition (**3**) was offered to illustrate the opposite selectivities observed for alkyl halide and epoxides. Despite the need for large quantities of LiCl, which has the effect of imposing concentration limits, the alkylations have found applications in a number of pharmaceutical-scale drug syntheses.³ They are now part of the pantheon of methods used for asymmetric synthesis, and they display special value in generating quaternary centers in exceptional yields and selectivities.¹

(1)

Understanding reactivity and selectivity in solutions teeming with observable and fleeting species demands a sound understanding of solution structure. Challenges to determining the structures of Myers enolates promised to be especially daunting: (1) possible aggregate topologies illustrated in Chart 1 are *representative* but are by no means a comprehensive compilation of possible structural motifs⁴; (2) positional isomerism represented generically in eq 2 would amplify spectral complexity; and (3) the difficulty of characterizing dianions⁵ is exacerbated by the absence of observable scalar coupling of Li–O bonds.

(2)

We describe herein studies of the structure and reactivity of Myers enolates. We take unusual care to illustrate some of the trials and tribulations—the often unstated pain and suffering en route to the answer—while trying not to make the reader suffer proportionately. The story is one of tactics and strategies. Although we characterized several heteroaggregates and a lithium diisopropylamide (LDA)–enolate mixed aggregate spectroscopically (*vide infra*), our best efforts to understand the all-important homoaggregate of **2** left us with too many possible structures and too many niggling details that simply did not afford an unapologetic

assignment. A low-resolution crystal structure that would be unpublishable in isolation provided us the answer key—the Rosetta stone. The crystallographic and spectroscopic data proved fully self-consistent. Which structure of the many under consideration was correct? In a word, none. We had many of the details right but missed the topology of the Li_8O_8 core that was, ironically, lurking in the Cambridge Structural Database.⁶

Dropping any pretense of presenting the data to the reader and *then* making the assignment, we provide the plot-spoiling solution structure of Myers enolate **2** as C_2 -symmetric octalithiated prism **4** constituted from four dianionic subunits manifesting an S_4 -symmetric core and overall C_2 symmetry. The four N–Li contacts were foreshadowed computationally. Although four coordinated THF ligands are omitted for clarity, they were the subject of considerable experimental focus. As part of our efforts to use the method of continuous variations (MCV)—a favorite of ours for studying solution structures of enolates⁷—we generated and characterized heteroaggregates containing subunits of opposite absolute configuration (heterochiral) that afforded *hexalithiated* prisms of type **5**. In short, MCV failed to provide heteroaggregates that attest to the structures of the homoaggregates. Control experiments to understand the consequences of excess LDA revealed C_2 -symmetric hexalithiated mixed aggregate **6** (comprising two LDA subunits and two dianionic subunits), which bears little relationship to the enolate homoaggregates but shares several unusual features with heteroaggregate **5**. Rate studies of alkylations showed profound aging effects spanning orders of magnitude.⁸ As to the influence of LiCl, we observed nondescript lithium enolate–LiCl mixed aggregates.

Results

We confront the age-old problem of presenting data from a variety of sources that, taken in total, provide the assignments yet must be presented serially. We try to assist the reader by prefacing the results with a detailed survey of the methods, as well as the structural and stereochemical consequences of the polygonal prisms. The nonspecialist will find an adequate survey of the results in the Discussion.

Substrates.

Acylated pseudoephedrine-based enolates **2** and **7–22** (Chart 3) were prepared using standard protocols.¹ [^{15}N](*R,R*)-**2** and [^{15}N](*S,S*)-**2** to probe N–Li contacts originated from [^{15}N] NH_4Cl , as shown in Scheme 1. [^6Li]LDA and [$^6\text{Li},^{15}\text{N}$]LDA were isolated as LiCl-free solids.⁹ Solvation was probed in several ways, including by using pyridine as a ^6Li chemical shift reagent¹⁰ and [^{15}N]isoquinoline (eq 3) as an inexpensive,¹¹ relatively nonvolatile pyridine surrogate.

(3)

Myers enolates were generated at $-78\text{ }^{\circ}\text{C}$ with subsequent aging for 10 min at $25\text{ }^{\circ}\text{C}$ owing to *slow aggregate equilibration*.⁸ Although we focused on aggregates **4–6** prepared from propionate enolate **2**, we examined the behavior of 17 enolates that fall into several distinct categories: (1) enolates **8** and **11** bearing small aliphatic enolate substituents share common properties with enolate **2** that we comfortably assigned by analogy to homoaggregate **4**; (2) **7, 9, 10, 12, 19**, and **20**, which manifest complex ensembles of otherwise sharp resonances suggesting isomerism is at play yet form well-defined heterochiral heteroaggregates sharing spectroscopic properties with **5**; (3) arylated enolates **13–18**, which are similar to one another but quite distinct from their aliphatic counterparts, yet again appear to form distinct type **5** heteroaggregates; and (4) enolate **21**, which is insoluble, and enolate **22**, which displays hopelessly broad resonances (possibly owing to instability mentioned by Myers).¹

MCV.

We have developed general protocols for using MCV in conjunction with NMR spectroscopy to characterize lithium enolates.⁷ In its simplest form, MCV involves mixing binary pairs of lithium enolate homoaggregates, $\mathbf{A}_n$ and $\mathbf{B}_n$, of unknown aggregation state, n , to generate an ensemble of homo- and heteroaggregates (eq 4). In principle, the homo- and heteroaggregates can be monitored using ^1H , ^6Li , ^{13}C , ^{15}N , or ^{19}F NMR spectroscopies. In practice, only ^6Li and ^{19}F NMR spectroscopies played roles in this study. (Caveat: Large T_1 's for ^6Li mandated up to 50 s pulse delays to attain accurate ^6Li integrations.) ^{15}N NMR spectroscopy has been useful in MCV when labeled substrates are available.¹² Despite the requisite resolution indicating that it would work well, $^{15}\text{N}[(R,R)\text{-1}]$ is simply too precious. Plotting the relative concentrations of different aggregates versus measured mole fraction of $\mathbf{A}_n$ (X_{A}) or $\mathbf{B}_n$ (X_{B}) affords a Job plot. (See Figure 7 for an example.)

Choosing the right pairing partners to ensure sufficient spectroscopic resolution of all homo- and heteroaggregates is critical and largely empirical and was particularly challenging in this case. The presumption is that the number and symmetries of the heteroaggregates attest to the aggregation number of homoaggregates $\mathbf{A}_n$ and $\mathbf{B}_n$. For the first time in what is now hundreds of applications, the homo- and heteroaggregates did *not* share a common aggregation number. Therefore, we defer discussing MCV-based studies until after presenting a more conventional approach to assigning of the structure of homoaggregate **4**.

Two-Dimensional NMR Spectroscopy.

Critical spatial relationships stem from a composite of NMR spectroscopies (COSY, TOCSY, HSQC, HMBC, and ROESY).¹³ Detailed discussions of the cross-correlations without a computer interface are poorly suited for print media. We only briefly allude to important correlations herein. In a pedagogically more tractable approach, a video tutorial in the supporting information walks the viewer through some of the assignments, as done in a previous paper.¹¹ Pulsed field-gradient spin echo NMR spectroscopy (PFGSE)—or diffusion NMR spectroscopy—was used semi-quantitatively to determine relative aggregate sizes.¹⁴

Computations.

Density functional theory (DFT) calculations were carried out at the B3LYP/6–31G(d) level with energies from single-point calculations at the MP2 level of theory.¹⁵ They largely showed plausibility of structural motifs, provided support to the correlation spectroscopies, and allowed us to probe experimentally elusive three-dimensional details. The energies proved *unhelpful* in distinguishing structural and stereoisomers owing to resulting narrow energetic ranges. The computations were *crucial*, however, in *predicting* strong N–Li contacts that were eventually confirmed spectroscopically and crystallographically. They also played a central role in developing mechanistic and stereochemical models.

Isomerism.

Three general classes of isomerism alluded to throughout the case studies required constant attention because of their implications for symmetry and spectral complexity. We also occasionally saw evidence of minor, lower symmetry isomers. The three major forms of isomerism are as follows.

1. Helical isomerism is exemplified by mirror-image fragments **23a** and **23b**, which become diastereomers when stereogenic centers are included. Asymmetry within the prismatic structure of **4** rendered concerns about helical isomerism moot, but the possibility of such isomerism lingered within our models for months.

2. Isomerism resulting from N–Li bond scission and reassociation to form different N–Li contacts (Scheme 2) is represented by isomeric pairs **24/25** and **26/27**. Chelates in **24** and **26** have phenyl moieties on the concave chelate face, whereas **25** and **27** have them on the convex face. Two-dimensional spectroscopic probes of **4** and **5** show both types (*vide infra*). Should only one of the multiple subunits reverse chelate orientation, however, any existing symmetry would be lost and the resonance counts would spike as each ⁶Li becomes magnetically distinct.
3. Positional isomerism such as that manifested in **24/26** or **25/27** pairs in Scheme 2 results from reversal of the enolate and alkoxide roles. Such a transposition would break the high symmetry.

Case 1: Homoaggregate of (*R,R*)-**2**.

Structural studies of the homoaggregate of propionate enolate (*R,R*)-**2** are the focus of this paper. Applications of MCV are deferred to subsequent case studies.

^6Li NMR spectra of (*R,R*)-**2** generated at $-78\text{ }^\circ\text{C}$ show complex mixtures of resonances (Figure 1A). Full aggregate equilibration required aging at $25\text{ }^\circ\text{C}$ for 10 min, affording two groups of ^6Li resonances at chemical shift extremes in carefully measured 1:1:1:1 proportions (Figure 1B). (The details of this aging process are delineated below.) Varying the THF concentration between 2.0 equiv per Li and 12.3 M (neat THF) and the enolate concentration from 0.020 M to 0.20 M resulted in no change in relative integrations, indicating that the resonances arose from either a single aggregate with four magnetically inequivalent lithiums or two isomeric aggregates in a precise 1:1 proportion. Failure to achieve coalescence at $>40\text{ }^\circ\text{C}$ owing to slow inter- and *intra*aggregate exchange,¹⁶ in conjunction with requisite aging at ambient temperature, attested to the remarkable robustness of the aggregate core.

Early in the study, DFT computations of dianion subunits in isolation or as part of larger aggregates evidenced strong N–Li contacts, prompting us to insert an ^{15}N label as shown in Scheme 1. [^{15}N](*R,R*)-**2** showed ^6Li – ^{15}N coupling in the downfield resonances at 2.32 and 2.35 ppm ($J_{\text{N-Li}} = 0.90$ and 1.40 Hz , respectively), whereas the upfield resonances remained as sharp singlets (Figure 1C). The corresponding ^{15}N NMR spectrum shows two 1:1:1 ^{15}N triplets correlated by single-frequency decoupling (Figure 1D).

Studies of solvation showed that only two of the four lithiums—logically those lacking N–Li contacts—contain one ligated THF each. For example, titrations of toluene solutions of **2** with THF showed that only 0.5 equiv THF per lithium was required to convert broad mounds to the four characteristic sharp resonances; additional THF caused minimal further change. More surprisingly, free and bound THF were in slow exchange on the NMR timescale *at room temperature* (Figure 2A). Such slow ethereal solvent exchange on lithium is rarely observed¹⁷ and is without precedent at ambient temperature. Single-frequency decoupling of the β protons on the bound THF revealed a single AB quartet for the α protons (Figure 2B). Given that two magnetically inequivalent THF ligands would show as many as *four* distinct AB quartets, we presumed that a combination of free rotation about the Li–O bond and near magnetic equivalence is at play. The decoupling was mimicked using 3,3,4,4-tetradeuteriotetrahydrofuran (THF-*d*₄),¹⁸ affording the same AB quartet.

Pyridine functions as a ^6Li NMR chemical shift reagent, imparting large ($>1\text{ ppm}$) downfield shifts for pyridine-ligated ^6Li nuclei and muted downfield shifts for more distal ^6Li resonances.¹⁰ Indeed, serial additions of pyridine to **2** showed large (1.1–1.4 ppm) shifts of the formerly THF-bearing lithiums and small shifts ($\approx 0.4\text{ ppm}$) of the ^6Li resonances coordinated by the carboxamide nitrogens. Readily available and relatively inexpensive [^{15}N]isoquinoline¹¹ caused analogous shifts and showed two of the four ^6Li resonances as doublets ($J_{\text{N-Li}} = 3.9$ and 3.6 Hz) and the remaining two as sharp singlets (Figure 3). ^{15}N NMR spectra showed broad mounds rather than sharp resonances. Broad-band ^{15}N decoupling caused the two ^6Li doublets to collapse to singlets. The coupling constants are large compared with those of other ^6Li – ^{15}N dative bonds.^{17a}

The resonance count of the homoaggregate of **2** excludes a single hexalithiated form by symmetry. The remaining options shown in Chart 4 include: (1) a 1:1 mixture of isomeric dilithiated forms, such as isomers **28** and **29**, displaying different convexity, each with two

magnetically inequivalent lithium nuclei; (2) a 1:1 mixture of two high-symmetry tetralithio isomers, such as **30** and **31**; (3) a single low-symmetry tetralithio form, such as **32** or **33**; (4) either **30** or **31** paired with its helical isomer (not drawn but akin to **23a,b**); (5) two high-symmetry hexalithio aggregates **34** and **35** (see below) or analogous pairs of octalithio or higher polygonal prismatic helical isomers; and (6) any one of a host of octalithio derivatives with a symmetry element affording a reduced resonance count (*vide infra*).

The full gamut of two-dimensional NMR spectroscopies led to the assignment of all protons and carbons within the two magnetically inequivalent subunits of (*R,R*)-**2** (Figure 4). The critical spatial correlations are as follows:

1. Multiple correlations between the two magnetically distinct dilithio subunits excluded *all* models based on pairs of isomers such as dilithio isomers **28** and **29**, high-symmetry tetralithio isomers **30** and **31**, and helical isomers **34** and **35**.
2. Strong inter-subunit correlations resulting from Ph–Ph contacts (*a*) and Me–H contacts (*b*) exclude tetralithio isomers **30** or **31**, as well as unsymmetrical tetralithio isomers **32** and **33**.
3. Apparent *intra*-subunit correlations suggest opposite convexities, as drawn in Figure 4. The Me–Ph correlations (*c*) and Me–H correlation (*d*) place the phenyl moieties on the convex and concave faces, respectively. We hasten to add, however, that deception by *inter*-subunit interactions between *two magnetically equivalent subunits* was a persistent concern. Moreover, the isomer with the phenyl on the convex face, as in subunit 1, is also calculated using simple (dilithio) models to be >2.0 kcal/mol more stable.
4. Correlations *a* and *b* seem to exclude a number of plausible aggregates by showing that *both* faces of *both* subunits are proximate.

By eliminating all di-, tetra-, and hexalithio homoaggregates, we were left with octalithio derivatives. Additional support came from two independent measurements. A diffusion NMR spectrum¹⁴ showed that the homoaggregate is larger than a demonstrably hexalithio heteroaggregate (*vide infra*). We tried using mass spectrometry (MS) on a DART MS, but the crude effort suffered from considerable protonation problems and left us leery of potential for user bias.¹⁹

Considering all connectivities of octagonal prisms and stacked cubes consistent with two inequivalent subunits and four magnetically inequivalent lithium nuclei left eight prisms and 18 stacked cubes as possible structures. Including the restriction that subunits 1 and 2 have opposite convexities (Figure 4), the number of possibilities reduces to six prisms and 12 stacked cubes (Chart 5). The DFT-computed relative energies showed plausibility but would not make the assignment for us. The nastier problem was that none accounted for the concurrent correlations without some overtly optimistic interpretation.

We *needed* an X-ray crystal structure. Early efforts failed, but we eventually obtained a low-resolution crystal structure that provided the needed insight (Figure 5). The structure of **4** shows an S_4 -symmetric core,⁶ which becomes a C_2 -symmetric octalithiated aggregate with the chiral appendages. The lithium counts, N–Li contacts, and most importantly, through-

space interactions match the resonance correlations without exception. Differential ^6Li - ^{15}N coupling noted above is reflected in markedly different N-Li bond lengths. Even with a graphic interface, one has to look carefully to see two pairs of subunits of opposite convexity. The pairs of magnetically inequivalent THF ligands reside in environments that appear so similar that the spectroscopic detection of a single observable THF subunit is not surprising.

Case 2: (*R,R*)-2-(*R,R*)-8 Homochiral Heteroaggregates.

The use of MCV to characterize homoaggregated enolates is based on the notion that mixtures of two homoaggregates will produce a highly characteristic ensemble of heteroaggregates (eq 4). Taking a cue from previous studies of chiral β -amino ester enolates,²⁰ we examined three pairing strategies. The first involved mixing combinations of enolates within a homochiral series—mixtures of (*R,R*)-2 and (*R,R*)-8, for example—to generate type-**i** heteroaggregates. Heteroaggregation was evidenced by intractable spectral complexity. This strategy was doomed from the start given that two structurally distinct Myers enolates can form seven distinct heteroaggregated analogs of homoaggregate **4** manifesting 48 ^6Li resonances.

Case 3: (*S,S*)-2-(*R,R*)-2 Heterochiral Heteroaggregates.

Mixing enantiomers of a single enolate—systematically generating scalemic mixtures—affords type-**ii** heteroaggregates with heterochiral subunits. A substantial advantage of this strategy is that $x:y$ and $y:x$ heteroaggregates are mirror images, which makes them appear spectroscopically as a single species, thereby markedly reducing the resonance count and minimizing resolution problems. By chance, heterochiral enolates also optimize resolution *and* favor heteroaggregates relative to homoaggregates nonstatistically. The fundamental presumption of MCV for determining aggregation state fails, however. Scalemic mixtures of octalithiated homoaggregates of enolates afforded *hexalithiated* heteroaggregates without exception (eq 5).

For example, scalemic mixtures formed from (*R,R*)-**2** and (*S,S*)-**2** with requisite aging at 25 °C display six new ⁶Li resonances in equal proportions (Figure 6A). The resonance count precludes an octalithiated aggregate. Resonances with mid-range chemical shifts (1.0–2.0 ppm) proved to be hallmarks of such heterochiral aggregates.

Monitoring the proportions of homo- and heteroaggregates as a function of mole fraction $X_{(R,R)\text{-}2}$ afforded the Job plot in Figure 7. Although the data adequately fit to an all-hexalithio (trimer) model, the fit in Figure 7 corresponds to that derived for octalithio homoaggregates (tetramers) and hexalithio heteroaggregates (trimers).²⁰ (Recall that only two spectroscopically discrete species exist.) The requisite fitting protocols are in the supporting information. The maximum at 0.50 derives from the superposition of enantiomeric 2:1 and 1:2 heteroaggregates. One could easily imagine confusing the maximum as evidence of a tetralithio heteroaggregate (dimer) if not for the resonance count and poor fit.

The relative and absolute aggregation state of heteroaggregate **5** was supported by diffusion NMR spectroscopy showing that the hexalithio appeared to be smaller than the octalithio (Figure 8).¹⁴ In fact, showing the reverse—the octalithio is larger than the more easily characterized hexalithio—was more important.

The gamut of two-dimensional NMR spectroscopies provided the key resonance correlations for the three discrete subunits of **5** shown in Figure 9. The convexities as drawn derive from the correlations labeled *a*, *d*, and *e*. Notably, subunit 3 is isolated from subunits 1 and 2 and displays no intersubunit correlation.

Labeling studies showed that the hexalithio heteroaggregates represented an aggregate that was altogether different from that of octalithio homoaggregate **4**. In mixtures of [¹⁵N](*R,R*)-**2** and [¹⁵N](*S,S*)-**2**, the resonance at 1.95 ppm appeared as a doublet ($J_{\text{N-Li}} = 1.25$ Hz), whereas the resonance at 1.83 ppm appeared as a doublet of doublets ($J_{\text{N-Li}} = 1.30$ and 2.17 Hz). This result indicates that *two nitrogens are bound to a common lithium* (Figure 6B). The remaining four upfield resonances are singlets. The ¹⁵N NMR spectra showed three ¹⁵N triplets with correlations confirmed by single-frequency decoupling. Mixtures of (*S,S*)-**2** and [¹⁵N](*R,R*)-**2** in which only *one* substrate is ¹⁵N labeled caused the lithium central to the N–Li–N fragment to appear as a doublet at all proportions of the two enolates. Thus, the subunits forming the N–Li–N contact—subunits 1 and 2 in Figure 9—are necessarily heterochiral as drawn. From connectivities alone, we could draw six isomeric cores with a central N–Li–N contact. Including additional restrictions imposed by the heterochiral core and established convexities, we were left with two pseudo-octahedral partial structures, **36** and **37**, displaying cores that would be delta isomers.²² There are eight possible isomers obtained by the placement of either (*R,R*)-**2** or (*S,S*)-**2** as the third subunit (with positional isomerism considered). The gross structure of hexalithiated aggregate **5** relied on ¹H and ¹³C correlation spectroscopy (supporting information). From the eight possible isomers, DFT computations suggested that the isomer shown as aggregate **5** with the **36** core is the most stable by >4 kcal/mol (Figure 10).

Case 4: (*R,R*)-**8**–(*S,S*)-**11** Heterochiral Heteroaggregates.

Pairing enolates of opposite absolute configuration *and* containing different enolate substituents affords type-iii heteroaggregates displaying many of the properties of the heterochirality observed in case 3 but with magnetically distinct $R'S_2$ and R'_2S heteroaggregates (eq 6). Heterochiral binary combinations of **2**, **8**, and **11** were all tractable. Emblematically, mixtures of (*S,S*)-**8** and (*R,R*)-**11** displayed fully desymmetrized 2:1 and 1:2 heteroaggregates each with affiliated six equal-intensity resonances (Figure 11). Unlabeled debris in the spectra of both homo- and heteroaggregates is reproducible, implicating minor stereoisomers. Although detailed correlating spectroscopic analyses would offer marginal gain at great effort, monitoring the aggregates versus mole fraction of (*R,R*)-**11**, $X_{(R,R)\text{-}11}$, affords the Job plot shown in Figure 12. The use of measured rather than intended mole fraction²¹ is especially suited for such mixtures.

Case 5: (*R,R*)-**14**–(*S,S*)-**16** Heterochiral Heteroaggregates.

Myers enolates with sterically demanding yet anion-stabilizing aryl substituents show behaviors that are distinctly different from those of their aliphatic counterparts (see Figure 1A). Each displays a major upfield resonance accompanied by a minor resonance in ratios that vary from enolate to enolate (Figure 13). The intensities are invariant to changes in enolate concentration, suggesting that they are isomers.²³ The absence of downfield resonances shows that the aryls cause deep-seated structural changes. High steric demands and anion stabilization of the phenyl moiety would be expected to promote lower aggregates.²⁴ Diffusion NMR spectroscopy confirmed that the homoaggregate is smaller than the hexalithio heteroaggregate (*vide infra*). Attempts to obtain X-ray-quality crystals of the homoaggregates were unsuccessful.

Heterochiral mixtures of arylated enolates (*S,S*)-**14** and (*R,R*)-**16** showed clear evidence of hexalithiated heteroaggregates, but spectral overlap precluded a Job plot using ^6Li NMR spectroscopy. The single fluorine nucleus per subunit (as opposed to two ^6Li nuclei) enables ^{19}F NMR spectroscopy while cutting the resonance counts in half. The ^{19}F NMR spectrum shows a single resonance (Figure 14) despite normally high-resolution of ^{19}F NMR spectroscopy. Mixtures showed that the 1:2 and 2:1 ^{19}F pairs are consistent with a hexalithiated species. Minor isomers of the 2:1 and 1:2 heteroaggregates also were observable. Monitoring the homo- and heteroaggregates versus $X_{(R,R)\text{-}14}$ and arbitrarily assigning the homoaggregates as hexalithio, tetralithio, or even dilithio afforded reasonable Job plots. The plot in Figure 15 arbitrarily assumes a tetralithiated homoaggregate.

Case 6: (*R,R*)-2-LDA Mixed Aggregate.

As a standard control experiment, lithium enolates were generated from [⁶Li,¹⁵N]LDA⁹ to identify any resonances attributable to LDA–enolate mixed aggregates. Although resulting LDA-derived mixed aggregate **6** bore little relationship to homoaggregate **4**, it foreshadowed the unusual N–Li–N connectivity in heteroaggregate **5**.

Mixtures of enolate (*R,R*)-**2** with varying amounts of excess [⁶Li,¹⁵N]LDA included LDA homodimer, homoaggregate **4**, and mixed dimer **6**. The key characteristics of **6** were as follows: (1) four ⁶Li resonances appeared in a 1:2:2:1 ratio (Figure 16A), indicating a hexalithiated species with a symmetry element; (2) the two major ⁶Li resonances appeared as doublets (*J* = 5.45 and 5.45 Hz) owing to splitting by a single LDA-derived ¹⁵N nucleus (Figure 16B); (3) the mixed aggregate prepared from [¹⁵N](*R,R*)-**2** and [⁶Li]LDA caused one of the minor resonances *not* connected to the LDA fragment to appear as a triplet (*J* = 2.4 Hz; Figure 16C), which correlated with *one* ¹⁵N triplet (1:1:1). The gamut of ¹H and ¹³C NMR spectroscopies showed a single subunit with the phenyl moiety in the convex face, as drawn for **6**. The stoichiometry, symmetry, and coupling demanded a hexalithio mixed aggregate with a *C*₂ symmetry axis. The symmetry demanded that the dianions reside on opposite faces of **6**. Although pyridine caused marked chemical shifts, attempts to probe solvation using [¹⁵N]-isoquinoline failed owing to decomposition. DFT computations showed that **6** is viable, revealed favorable N–Li contacts, and supported tetrasolvation, as displayed in Figure 17.

Lithium Chloride: Influence on Structure.

Addition of LiCl to a *pre-aged* solution of aggregate **4** had no measurable effect (Figure 18A): aggregate **4** was observed along with free LiCl at the anticipated relative proportions. Warming the probe in 20 °C increments caused no broadening of the resonances of **4** or appearance of new resonances. Careful integration of the resonances, however, showed that the broad mound corresponding to LiCl increased at the expense of **4** (Figure 18, B–F). The mound appeared to include free LiCl in rapid exchange with a LiCl–enolate adduct(s) denoted nondescriptly as **2**_{LiCl}. Assuming conservation of mass, the proportions of homoaggregate **4** and **2**_{LiCl} could be teased out. Plotting the percent of enolate **2** that existed as homoaggregate **4** versus temperature (Figure 19) showed limited adduct formation at low temperatures and a marked change (a discontinuity) at ≈0 °C. Reversing the protocol—cooling the probe in 20 °C increments—did *not* reverse the changes; **2**_{LiCl} remained dominant.²⁵ The spectrum re-recorded at –80 °C *may* include additional debris that corresponds to species in slow exchange at –80 °C, but it appears insignificant.¹ Adduct formation was confirmed by LiCl-dependent chemical shifts observed in the ¹H and ¹³C NMR spectra.²⁶ The time-averaged resonances may be concealing multiple LiCl adducts of varying LiCl stoichiometries as discussed below.¹ Although adduct **2**_{LiCl} cannot be characterized, it is decidedly important (*vide infra*).

Lithium Chloride: Influence on Subunit Exchange.

In a fairly crude experiment, kinetically formed (pre-aged) ensembles of **2** (see Figure 1A) were warmed in 20 °C increments with cooling to –80 °C at each temperature level to record

the ^6Li spectra. The spectra indicate that aging to form **4** becomes consequential at $>0^\circ\text{C}$. Repeating the experiment with added LiCl *showed little or no effect on the rate of aging*.

In a cleaner experiment, samples of (*S,S*)-**2** and (*R,R*)-**2** were *separately and fully aged* before mixing to form a racemic mixture of aggregate **4**. Warming in 20°C increments revealed hexalithiated heteroaggregate **5**, evidencing that *interaggregate subunit exchange on laboratory timescales* emerges only at 40°C . (Subunit exchange in the octalithiated homoaggregate is slower than exchange in unaged ensembles.) The same experiment with added LiCl showed that rate of conversion of the two antipodes of **4** to give **5** occurs at $>0^\circ\text{C}$. Thus, *LiCl influences the aging rate on laboratory timescales but only marginally and, as we discuss below, much more slowly than the alkylation*.

Aging Effects on Enolate Reactivity.

We now present some curious observations on how aggregate aging and the stabilities of the aggregates influence their reactivities. Alkylations using allyl bromide (eq 7) were monitored using in situ IR spectroscopy following the formation of the product alkoxide at 1610 cm^{-1} . (The enolate absorbance at 1640 cm^{-1} was poorly resolved from an absorbance of the product alkoxide.) The apparent percent conversions were confirmed with workup and ^1H NMR spectroscopic analysis of the crude products. Fully aged homoaggregate **4** was marginally reactive toward allyl bromide at 0°C for 1.0 h (Figure 20). Myers also noted such low conversions (10–13%). Varying the THF molarity offered no improvement. Elevated temperatures promoted O-alkylation byproducts. Reasoning that unaged samples contain structures that are less stable and, thus, more reactive, we found that addition of allyl bromide 1.0 min after enolization at -20°C improved the conversion to 40%. Completely unaged samples kinetically formed at -78°C (see Figure 1A) *alkylated at -78°C to 67% conversion before stalling*. Surmising that the ensemble of unstable aggregates generated at -78°C might depend on the source, we showed that metalation with lithium 2,2,6,6-tetramethylpiperide increased the conversion to 78%, which proved to be the upper limit. Attempts to push the conversion of unaged samples through gentle warming failed, presumably because of competing aging to less reactive forms.

(7)

Influence of LiCl on Reactivity.

It is not surprising that homoaggregate **4** is inert; it *looks* inert. Addition of 5.0 equiv of LiCl as part of Myers's optimized protocol is imperative. A more granular view obtained by adding various concentrations of LiCl to the allylation in eq 7 reveals a burst of reactivity followed by stalling to the rate that approximates the LiCl-free rate (Figure 20). Both the initial rates and the percent conversions were promoted by elevated LiCl concentrations. We

attribute the acceleration to the nondescript adduct(s) denoted as $\mathbf{2}_{\text{LiCl}}$ and the stalling to an equally nondescript autoinhibition resulting from a lithium alkoxide–LiCl mixed aggregate that occludes the LiCl.²⁸ Monitoring the alkylation with ^6Li NMR spectroscopy revealed only homoaggregate $\mathbf{4}$. The ^6Li resonances of the product alkoxide were absorbed into the broad LiCl-derived mound in rapid time-averaged exchange.

Monitoring the initial alkylation rates proved instructive. The initial rates indicate a clean first-order dependence on allyl bromide (Figure 21) attesting to the general quality of the data and a rate-limiting alkylation.²⁹ Curvatures in the LiCl dependence (Figure 22) and enolate dependence (Figure 23) could be construed as fractional-order dependencies emblematic of dimer–monomer and tetramer–monomer dissociations.²⁸ Detection and quantitation of $\mathbf{2}_{\text{LiCl}}$, however, clearly supported the curvatures in Figures 22 and 23 as saturation kinetics.

The red curve in Figure 22 represents first-order saturation behavior required if the equilibrium in eq 8 affords $\mathbf{2}_{\text{LiCl}}$ as a monomer-based tetralithiated mixed aggregate, (enolate)(LiCl)₂ (eq 8, $1/n = 1$ and $1/m = 1/4$). The blue curve shows a fit with n as an adjustable parameter and fits to a second-order LiCl saturation ($1/n = 2$ and $1/m = 1$) consistent with $\mathbf{2}_{\text{LiCl}}$ as a tetramer-based mixed aggregate, (enolate)₄(LiCl)₄. We hasten to add that the quality of the fits distinguishes the various possible structures for $\mathbf{2}_{\text{LiCl}}$ in principle but, in our opinion, not in practice. The enolate dependence in Figure 23 is fit to a first-order saturation behavior (eq 8, $1/m = 1$), but a fit to a fractional-order saturation behavior ($1/m = 1/4$) is, practically speaking, indistinguishable. Plotting the initial rates versus THF concentration in hexane (Figure 24) reveals an upward curvature suggesting intermediate first or second order.

The saturation behaviors are consistent with the generic rate law in eq 9. Two critical pieces of information remained outstanding: (1) the structure of $\mathbf{2}_{\text{LiCl}}$ as described in eq 8 is unknown, leaving the values of m and n in eq 9 undefined; and (2) the reaction order in $\mathbf{2}_{\text{LiCl}}$ could not be measured owing to solubility constraints; knowing this would provide an important clue about the structure of $\mathbf{2}_{\text{LiCl}}$. We consider three mechanisms in the Discussion.

$$d[\text{product}]/dt = k[\text{RBr}][\text{LiCl}]^{1/n}[\text{enolate}]^{1/m}/(1 + k'[\text{LiCl}]^{1/n}[\text{enolate}]^{1/m}) \quad (9)$$

Discussion

Myers's central insight, which made pseudoephedrine-derived enolates popular reagents for asymmetric alkylations, was that excess LiCl is critical to obtaining high yields. From a structural and mechanistic perspective, the dianionic enolates present a host of challenges outlined in the introduction that are unlike any we have confronted to date.

Structural Studies: Protocols.

The aggregate structures are summarized in Scheme 3, but they tell only a small part of the story. The number of analytical methods and tactics is unusual. These include a bevy of two-dimensional correlation spectroscopies, ^6Li – ^{15}N double-labeling studies, diffusion NMR spectroscopy, mass spectrometry, IR spectroscopy, MCV, X-ray crystallography, and DFT computations. This breadth of effort was demanded by the elusiveness of the answers rather than by some predilection toward thoroughness.

Standard ^1H and ^{13}C NMR two-dimensional correlation spectroscopies provided critical information but did not lead us to a definitive structural assignment. Attempts to use MS to determine aggregation¹⁹ presented potentially resolvable technical challenges that rendered the data from our first effort at risk of user-dependent interpretation. DFT computations often help resolve structural issues, but there were simply too many possible isomers in a narrow energetic range. After several years of study, a low-quality crystal structure provided the answer key: the solution structural studies came together into a fully coherent structural model. While we had obsessed over octagonal prisms and stacked cubes, the octalithiated aggregate proved to be **4**, the core of which was known.⁶

MCV consumed considerable attention. By intent, mixtures of structurally similar homoaggregates afford ensembles of heteroaggregates with numbers and symmetries that attest to the structures of the homoaggregates. However, mixtures of octalithiated homoaggregates—dozens of combinations in total—afforded *hexalithiated* heteroaggregates. To the best of our knowledge, this failure of MCV is a first for us. A detailed study of two antipodes of enolate **2** revealed the *heterochiral* heteroaggregate of type **5**. Aside from the different aggregation numbers of **4** and **5**, the hexalithiated heteroaggregates include strikingly different structural details.

Structures.

Let us take a granular peek at structural details of the homo- and heteroaggregates and mixed aggregates³⁰ summarized in Scheme 3.

- i. *Aging Effects.* Implicitly embedded in Myers's alkylation protocols are aging effects—requisite warming cycles—that prove to have a firm structural basis and became a major focus of our studies. Enolization at $-78\text{ }^\circ\text{C}$ affords a complex mixture that converges to form octalithiated aggregate **4** only on warming to ambient temperatures. The same aging effects are observed during formation of heteroaggregates of type **5**. Although aging to form **4** could be construed as the structural change that Myers addressed by warming the samples, this conclusion is incorrect. The aging was needed to generate LiCl adducts (below).
- ii. *N–Li contacts.* The N–Li contacts of carboxamide-derived enolates appear to have gone undocumented until now. We discovered them computationally and then took them seriously enough only to confirm them using ^{15}N -labeled substrate. Contemporaneously, Williard and co-workers crystallographically detected a number of such N–Li contacts in structurally simple carboxamide enolates.³¹ As to how general they are in solution remains unknown; a simple

¹⁵N-labeled propionamide enolate affords no N–Li coupling.³² From the outset, the importance of the N–Li contacts and the resulting convexity in the Myers enolates seemed certain to underlie the stereocontrol.

- iii. *Slow THF exchange.* Slow exchange of lithium-bound THF on **4** with free THF at ambient temperatures is surprising. It is rare that THF can be observed in the slow exchange limit even at low temperatures.¹⁷ We suspect that the slow exchange on aggregate **4** derives, at least in part, from suppression of the associative pathway.^{17a} Although slow alkylations of **4** probably render rates of THF exchange irrelevant, highly reactive electrophiles could suffer consequentially (stereochemically) from restricted access to coordination sites.
- iv. *Stereochemically sensitive aggregation.* The formation of octalithiated homochiral aggregates of type **4** and hexalithiated heterochiral heteroaggregates of type **5** are, in one sense, an idiosyncratic artifact of using MCV to study aggregation. (We are likely the only group to intentionally study racemates of chiral lithium enolates used in asymmetric synthesis.)²⁰ Nonetheless, the hexalithiated heteroaggregates display a fundamentally different doubly chelated N–Li–N core (see **36** and **37**). This structural feature is shared by LDA mixed aggregate **6** (Figure 17). The hexalithiated aggregates are also formed nonstatistically. The good news is that DFT studies suggest that the *homochiral* hexalithio derivatives prefer heterochirality (eq 10) and the octalithiated forms prefer homochirality (eq 11). The dubious result is that, when combined, DFT computations completely fail to predict hexalithiated aggregates of type **5** to be competitive with homoaggregate **4** (eq 12).

Aging Effects on Reactivity.

If one supposes that all aggregated forms converge on a fleeting dilithiated (monomeric) intermediate, then aggregate stability and reactivity should correlate inversely (Scheme 4). Fully aged **4** reacts slowly at 0 °C and stalls at <10% conversion, whereas unaged aggregates—the complex ensemble of species formed kinetically at –78 °C—are orders of magnitude more reactive, eventually stalling at 67–78% conversion. Presumably, the kinetically formed ensemble contains aggregates of varying reactivities. One cannot help but worry that the depiction of **5** may be susceptible to correction in its detail.

The Role of LiCl: Enolate Structure and Reactivity.

Adding excess LiCl is imperative, as noted by Myers,¹ and it serves the dual function of accelerating the alkylation and allowing the reaction to proceed to full conversion (Figure 20). We considered several possible roles for LiCl.

- i. *Catalyzed Deaggregation.* One could imagine that LiCl *catalyzes* tetramer–monomer exchange, shifting the rate-limiting step from deaggregation to alkylation of the dissociated monomer.^{28,34} LiCl-catalyzed deaggregation³³ was studied and documented in excruciating detail for LiCl-catalyzed deaggregation of LDA.²⁸ Presuming that a catalyzed deaggregation would affiliate with a catalyzed exchange of subunits, we probed the influence of LiCl on rates of subunit exchange from a number of angles. Minor accelerations of subunit exchanges occurring *on laboratory timescales* near ambient temperatures were observed, but they could not explain LiCl-mediated alkylations at –60 °C.
- ii. *Mixed Aggregation.* Lithium chloride forms a nondescript mixed aggregate denoted simply as **2_{LiCl}** (eq 8). Formation of **2_{LiCl}** is slow at –80 °C and occurs in earnest only near ambient temperatures on laboratory timescales (eq 13). *It is the slow formation of 2_{LiCl}* that Myers seems to have overcome by warming the enolate–LiCl mixtures before alkylating the enolates. An odd temperature dependence (Figure 19) shows that the formation of **2_{LiCl}** is slow and irreversible. It also suggests, however, that other partial adducts at intermediate temperatures may exhibit limited reactivity. Once **2_{LiCl}** forms, the time averaging of free and bound LiCl in the ⁶Li NMR spectra (eq 14) made it difficult to detect—we almost missed it—and difficult to characterize to date.³⁵ The potentially critical observation is that exchange of free and bound LiCl, as well as *intraaggregate* exchanges of the ⁶Li nuclei within **2_{LiCl}**, are exceedingly facile on *NMR timescales* at –80 °C.

(13)

(14)

- iii. *Autoinhibition.* Alkylation of enolate **2** with allyl bromide stalls at percent conversions that qualitatively correlate with the added LiCl. In theory, this result could stem from incomplete formation of $\mathbf{2}_{\text{LiCl}}$, but we cannot push the percent conversions effectively even under conditions in which LiCl could be recycled. We surmise that the product of the reaction—an alkoxide (R'OLi)—occludes LiCl in a mixed aggregate (eq 15). Once again, strikingly fast LiCl exchange did not refute the existence of such an adduct but did preclude structural studies.

(15)

It is instructive to ponder the structure of $\mathbf{2}_{\text{LiCl}}$ (Chart 6). The deep-seated structural changes in LiCl adducts **38** and **39** are appealing in light of remarkably low rates of LiCl incorporation, although **39** contains one or two LiCl fragments depending on whether the counterion is a simple or a cationic triple ion.³⁶ Chloride-derived “ate” complexes analogous to **39** and **40**, in which one could think of chloride as a hexamethylphosphoramide surrogate, have been observed³⁷ and are computationally viable (supporting information). Retention of the intact octalithio framework, as depicted in **40**, cannot be rigorously excluded—it is a computational minimum—but the spectroscopic properties of $\mathbf{2}_{\text{LiCl}}$ seem to argue against the intact Li₈O₈ core.

Rate Studies and Mechanism.

Detailed rate studies reveal curvatures consistent with saturation kinetics reflecting the lithium enolate- and LiCl-dependent formation of $\mathbf{2}_{\text{LiCl}}$ (Figures 22 and 23). The accompanying THF dependence (Figure 24) is compatible with almost any model with some imagination. We have considered a number of mechanisms for the alkylation that hinge on the structure of $\mathbf{2}_{\text{LiCl}}$ with a focus on mixed aggregates **38** or **39**. We present a working hypothesis based on **38** as described by eq 16 and delineated in Scheme 5.

Computational studies suggest a preference for **38a** relative to **38b**, displaying the opposite convexity in the resting state that superficially follows through to the most stable transition structures **41a** and **41b**. The mechanism also includes no provisions for an additional THF suggested by the rate studies, but that does not trouble us too much. The more systemic flaw is that the preference for transition structure **41b** predicts the *wrong (minor) isomer* compared with that observed experimentally.

An alternative mechanism that could account for THF-dependent rates (maybe too much so depending on the resting state of **2_{LiCl}**) is based on LiCl ate complexes such as **39**. Such ate complexes have been observed³⁷ as have simple ⁺Li(THF)_{*n*} and more complex triple-ion-based counterions.³⁶ The model reveals a strong (2.5 kcal/mol) preference for transition structure **42** relative to the second most stable of three other isomeric transition (supporting information). This model is consistent with the experimental results.

The third and simplest of the three mechanisms is based on direct alkylation via dilithiated monomer-based transition structure **43** stripped free of any LiCl influence. Although the model could readily account for the THF dependence, it also predicts a net inverse LiCl dependence of some form owing to requisite dissociation of LiCl from **2_{LiCl}**. We can easily imagine such details evading our detection by being embedded in the saturation kinetics required to form **2_{LiCl}** (Figure 22). This monomer-based transition structure **43** is computationally preferred relative to the three other choices (supporting information) and is consistent with experiment.

On first inspection, the notion that forming a LiCl adduct to simply dissociate it before alkylation seems to defy the Principle of Detailed Balance, which suggests that in such an ensemble the LiCl should impart a positive influence on rates only if it is included in the rate-limiting transition state.³⁸ Yet the Principle of Detailed Balance applies only to ensembles of species *in equilibrium*. The Myers enolate and its LiCl adduct (eq 17) are *not* at equilibrium; the reluctant formation of **2_{LiCl}** requires ambient temperatures, affording **2_{LiCl}**, which seems to be in *rapid* exchange with monomer but *slow* exchange with homoaggregate **4**. The role of LiCl is to overcome the otherwise insurmountable barrier to dissociating octalithiated prismatic tetramer **4**. An analogous example of aggregate ensembles in which not all species are at equilibrium was recently observed for Evans enolates in which facile dimer–monomer equilibration occurs concurrently with slow formation of the more stable and unreactive tetramer.^{8d}

(17)

Inspection of the transition structures (12 in total) with the aid of a graphical interface reveals no self-evident insight into the stereochemical preferences. There is no visibly discernible preference for a specific convexity of the chelated dianion or for an approach of the allyl bromide to the enolate that can be generalized. Notably, however, one can see a lengthening of the N–Li bond at the transition structure as the loss of the enolate character at the transition states that attenuates the Lewis basicity of the nitrogen ligand. IRC calculations indicate that the alkylation causes complete scission of the N–Li bond during the alkylation.

Omitting Lithium Chloride?

A potential impediment to using Myers enolates on pilot-plant or plant scales stems from the relatively high dilutions (0.10 M) imposed by the limited solubility (and cost) of the 5-fold LiCl excess. Evidence that precluding enolate aging markedly enhanced reactivities and percent conversions in halide free solutions offered reason for optimism, but we could not push past 78% conversion. Investigators at Merck were able to exclude LiCl in the alkylation of a highly stabilized enolate bearing an aryl group akin to enolates **13–18** using THF/TMEDA,^{3e} but we showed those are structurally quite different than the more standard Myers enolates. It would be tempting to add ligands to selectively extract LiCl from the putative product mixed aggregate by forming $^+\text{Li}(\text{ligand})/\text{Cl}^-$ (eq 18), but according to the Principle of Detailed Balance, such a side equilibrium would *decrease* the concentration of 2_{LiCl} as well. Nonetheless, we tried a few chelating ligands to no avail. We have, however, been making progress using a radically different strategy, but that story will have to wait.

(18)

Conclusions

Studies of dianionic Myers enolates certainly presented challenges and surprises. The influence of aging on aggregate structure and reactivity is profound and reinforces the notion that enolates age remarkably slowly compared with other organolithiums. The pseudoephedrine-derived Myers enolates are slower to equilibrate than most, but they are by no means unique.⁸ Indeed, limited solution structural studies of enolates to date suggest that *most* enolates—particularly the structurally complex variants—undergo aggregate exchanges so slowly that the unsuspecting user may be getting whipsawed by slow aggregate equilibrations without realizing it. Functionalizing an enolate under conditions in which aggregates have not equilibrated may be constructive or destructive. Worse, however, is when functionalizations of *partially* equilibrated aggregates wreak havoc on attempts to develop rational models to control and understand reactivity and selectivity.

In some ways, the study of Myers enolates is about refining protocols for characterizing complex organolithiums in solution. We are positioning to move our gaze to Myers's second-generation enolates, derived from the 1,2-diphenyl-substituted amino ethanol. The

new perspective might fill in some details that eluded us in the current investigation. Moreover, there are many dianionic enolates in which ancillary lithium alkoxides, lithiated sulfonamides, or lithiated Boc-amines are used in lieu of standard protecting groups. We are optimistic that we now have the tools to probe structure–reactivity relationships.

Experimental

Reagents and Solvents.

THF and toluene were distilled from solutions containing sodium benzophenone ketyl. Pyridine was distilled from solutions sodium benzophenone ketyl. LDA, [^6Li]LDA, and [^6Li , ^{15}N]LDA were prepared as described previously.⁹ Air- and moisture-sensitive materials were manipulated under argon using standard glove box, vacuum line, and syringe techniques. Myers enolate precursors were either purchased or prepared as described previously.¹ Several previously unreported precursors were prepared by acylating the pseudoephedrine as described in supporting information.

NMR Spectroscopy.

Individual stock solutions of substrates and LDA were prepared at room temperature. An NMR tube under vacuum was flame-dried on a Schlenk line and allowed to return to room temperature. It was then backfilled with argon and placed in a dry ice/acetone bath at -78°C . The appropriate amounts of acylated pseudoephedrine and LDA (1.1 equiv) were added sequentially via syringe. The tube was sealed under partial vacuum and vortexed three times on a vortex mixer for 5 s. Samples could be stored indefinitely in a freezer -86°C . Each sample routinely contained 0.10 M total enolate with a 0.005 M excess of LDA. Standard ^1H , ^6Li , ^{13}C , ^{15}N , and ^{19}F NMR spectra were recorded on a 500 MHz spectrometer at 499.92, 73.57, 125.72, 50.67, and 470.33 MHz, respectively. Quantitated ^6Li integrations required the relaxation delay set to seven times the decay half-life. Integration of NMR signal was carried out using the peak area generated by a line-fitting method. The ^6Li and ^{13}C resonances are referenced to 0.30 M [^6Li]LiCl/MeOH at -90°C (0.0 ppm) and the CH_2O resonance of THF at -80°C (67.57 ppm). Two-dimensional NMR spectroscopies (COSY, TOCSY, HSQC, HMBC, and ROESY) were recorded using standard pulse sequences.

IR Spectroscopic Analyses.

IR spectra were recorded using an in situ IR spectrometer fitted with a 30-bounce, silicon-tipped probe. The spectra were acquired in 16 scans at a gain of 1 and a resolution of 4 cm^{-1} . A representative reaction was carried out as follows: The IR probe was inserted through a nylon adapter and O-ring seal into an oven-dried, cylindrical flask fitted with a magnetic stir bar and a T-joint. The T-joint was capped with a septum for injections and a nitrogen line. After evacuation under full vacuum, heating, and flushing with nitrogen, the flask was charged with a stock solution of enolate **2** in THF or THF/hexane and LiCl in THF. The vessel was cooled to -60°C in a constant-temperature bath controlled with a cold finger and prepared with fresh acetone. After a background spectrum was recorded, neat allyl bromide was added with stirring. IR spectra were recorded every 6 s with monitoring of the absorbance at 1610 cm^{-1} over the course of the reaction.

MCV.

The mathematical treatment for creating Job plots using Mathematica has been described previously.²⁰ The specific case in which octalithiated homoaggregates (formally tetramers) afford hexalithiated heteroaggregates (trimers) required additional coding as described in the supporting information.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments.

We thank the National Institutes of Health (GM077167) for support.

References and Footnotes

1. (a) Yang BH; Chen H; Gleason JL; Myers AG Use of Pseudoephedrine as a Practical Chiral Auxiliary for Asymmetric Synthesis. *J. Am. Chem. Soc* 1994, 116, 9361. (b) McKinstry L; Myers AG Practical Syntheses of Enantiomerically Enriched γ -Lactones and γ -Hydroxy Ketones by the Alkylation of Pseudoephedrine Amides with Epoxides and their Derivatives. *J. Org. Chem* 1996, 61, 2428. (c) Yang BH; Chen H; McKinstry L; Kopecky DJ; Gleason JL; Myers AG Pseudoephedrine as a Practical Chiral Auxiliary for the Synthesis of Highly Enantiomerically Enriched Carboxylic Acids, Alcohols, Aldehydes, and Ketones. *J. Am. Chem. Soc* 1997, 119, 6496.
2. Evans DA; Takacs JM Enantioselective Alkylation of Chiral Enolates. *Tetrahedron Lett.* 1980, 21, 4233.
3. (a) Sandham DA; Taylor RJ; Carey JS; Fässler A A convergent synthesis of the renin inhibitor CGP60536B. *Tetrahedron Letters*, 2000, 41, 10091. (b) Dragovich PS; Prins TJ; Zhou R Formal, stereoselective synthesis of hydroxyethylene dipeptide isosteres utilizing pseudoephedrine amides. *J. Org. Chem* 1997, 62, 7872. (c) Sinz C; Bittner A; Brady E; Candelore M; Dallas-Yang Q; Ding V; Jiang G; Lin Z; Qureshi S; Salituro G; Saperstein R Discovery of *N*-Aryl-2-acylindole human glucagon receptor antagonists. *Bioorganic & medicinal chemistry letters* 2011, 21, 7124. [PubMed: 22030028] (d) Narasimhulu CP; Das P Stereoselective Preparation of C1–C10 and C11–O14 Fragments of Narbonolide: Exploiting the Versatility of Thiazolidinethione Chiral Auxiliary. *Synthesis* 2009, 474. (e) Smitrovich JH; DiMichele L; Qu C; Boice GN; Nelson TD; Huffman MA; Murry J Michael Reactions of Pseudoephedrine Amide Enolates: Effect of LiCl on Syn/Anti Selectivity. *J. Org. Chem* 2004, 69, 1903. [PubMed: 15058935]
4. (a) Seebach D Structure and Reactivity of Lithium Enolates. From Pinacolone to Selective C-Alkylations of Peptides. Difficulties and Opportunities Afforded by Complex Structures. *Angew. Chem., Int. Ed. Engl* 1988, 27, 1624. (b) Braun M Lithium Enolates: 'Capricious' Structures - Reliable Reagents for Synthesis. *Helv. Chim. Acta* 2015, 98, 1.
5. (a) Gruver JM; West SP; Collum DB; Sarpong R Experimental Characterization and Computational Study of Unique C,N-Chelated Lithium Dianions. *J. Am. Chem. Soc* 2010, 132, 13212. [PubMed: 20822143] (b) Li D; Keresztes I; Hopson R; Williard PG Characterization of Reactive Intermediates by Multinuclear Diffusion-Ordered NMR Spectroscopy (DOSY). *Acc. Chem. Res* 2009, 42, 270. [PubMed: 19105594] (c) Gareyev R; Ciula JC; Streitwieser A Lithium and Cesium Ion-Pair Acidities of Dibenzyl Ketone. Aggregation of Lithium and Cesium Ion Pairs of the Enolate Ion and Dianion. *J. Org. Chem* 1996, 61, 4589. [PubMed: 11667384] (d) Brand H; Capriotti JA; Arnold J New Lithium Porphyrin Derivatives: Synthesis of Li2(P)(Et2O)2 (P = TTP, TBPP) and Solution Structure of Li2(TTP)(Et2O)2 by 7Li and 15N NMR *Inorg. Chem* 1994, 33, 4334. (e) Günther H Selected Topics from Recent NMR Studies of Organolithium Compounds. *J. Braz. Chem. Soc* 1999, 10, 241. (f) Jacobson MA; Keresztes I; Williard PG On the Mechanism of THF Catalyzed Vinylic Lithiation of Allylamine Derivatives: Structural Studies Using 2-D and Diffusion-Ordered NMR Spectroscopy. *J. Am. Chem. Soc* 2005, 127, 4965. [PubMed: 15796563] (g) Cohen Y; Roelofs NH; Reinhardt G; Scott LT; Rabinovitz M Novel carbocyclic dianions: NMR study of charge

- delocalization, paratropicity, and structure in the dianions of acephenanthrylene and aceanthrylene. *J. Org. Chem* 1987, 52, 4207.(h)Matsuo T; Mizue T; Sekiguchi A Synthesis and Molecular Structure of a Dilithium Salt of the cis-Diphenylcyclobutadiene Dianion. *Chem. Lett* 2000, 896.
- (i)Henderson KW; Dorigo AE; MacEwan GJ; Williard PG Structure of an unsymmetrical heptalithium cage complex containing aldolate and enolized aldolate dianion. *Tetrahedron* 2011, 67, 10291.
6. (a)MacLellan JG; Mulvey RE; Nichols PJ; Andrews PC New homo- and hetero-alkali metal alkoxide cages; crystal structures of $[\text{Me}_2\text{N}(\text{CH}_2)_2\text{OLi}]_8$ and $[\{\text{Me}_2\text{N}(\text{CH}_2)_2\text{O}\}_2\text{Li}_8\text{K}_6]\text{O}$. *J. Chem. Soc., Dalton Trans*, 2002, 1651.(b)Allan JF; Nassar R; Specht E; Beatty A; Calin N; Henderson KW Characterization of a kinetically stable, highly ordered, octameric form of lithium tert-butoxide and its implications regarding aggregate formation. *J. Am. Chem. Soc* 2004, 126, 484. [PubMed: 14719943] (c)Boyle TJ; Alam TM; Peters KP; Rodriguez MA Structural diversity of lithium neopentoxide compounds. *Inorg. Chem* 2001, 40, 6281. [PubMed: 11703131]
7. Renny JS; Tomasevich LL; Tallmadge EH; Collum DB Method of Continuous Variations: Applications of Job Plots to the Study of Molecular Associations in Organometallic Chemistry. *Angew. Chem., Int. Ed* 2013, 52, 11998.
8. (a)Casy BM; Flowers RA On the Nature of the Oxidative Heterocoupling of Lithium Enolates. *J. Am. Chem. Soc* 2011, 133, 11492. [PubMed: 21721524] (b)Jin KJ; Collum DB Solid-State and Solution Structures of Glycinimine-Derived Lithium Enolates *J. Am. Chem. Soc* 2015, 137, 14446. [PubMed: 26554898] (c)For a marked influence of aggregate aging on the reactivity and selectivity of organolithiums, see: Xu F; Reamer RA; Tillyer R; Cummins JM; Grabowski EJJ; Reider PJ; Collum DB; Huffman JC Highly Enantioselective 1,2-Addition of Lithium Acetylide-Ephedrate Complexes: Spectroscopic Evidence for Reaction Proceeding via a 2:2 Tetramer, and X-ray Characterization of Related Complexes. *J. Am. Chem. Soc* 2000, 122, 11212.(d)Tallmadge EH; Jermaks J; Collum DB Structure-Reactivity Relationships in Lithiated Evans Enolates: Influence of Aggregation and Solvation on the Stereochemistry and Mechanism of Aldol Additions. *J. Am. Chem. Soc* 2016, 138, 345. [PubMed: 26639525]
9. Ma Y; Hoepker AC; Gupta L; Faggin MF; Collum DB 1,4-Addition of Lithium Diisopropylamide to Unsaturated Esters: Role of Rate-Limiting Deaggregation, Autocatalysis, Lithium Chloride Catalysis and Other Mixed Aggregation Effects. *J. Am. Chem. Soc* 2010, 132, 15610. [PubMed: 20961095]
10. Jermaks J; Tallmadge EH; Keresztes I; Collum DB Lithium Amino Alkoxide-Evans Enolate Mixed Aggregates: Aldol Addition with Matched and Mismatched Stereocontrol. *J. Am. Chem. Soc* 2018, 140, 3077. [PubMed: 29457718]
11. (a)Garratt PJ; Vollhardt KPC Homophthalaldehyde. *Synthesis*, 1971, 8, 423.(b)Huang X; Tanaka KS; Bennet AJ Glucosidase-catalyzed hydrolysis of α -D-glucopyranosyl pyridinium salts: kinetic evidence for nucleophilic involvement at the glucosidation transition state. *J. Am. Chem. Soc* 1997, 119, 11147.
12. Algera RF; Ma Y; Collum DB Sodium Diisopropylamide: Aggregation, Solvation, and Stability. *J. Am. Chem. Soc* 2017, 139, 7921. [PubMed: 28557426]
- 13.(a). Friebolin H Basic One- and Two-Dimensional NMR Spectroscopy; Wiley VCH: Weinheim, 2010.(b)Claridge TDW High-Resolution NMR Techniques in Organic Chemistry, 2nd ed; Elsevier: Amsterdam, 2009.
- 14.(a). Sørland GH Dynamic Pulsed-Field-Gradient NMR, Springer: Berlin, 2014.(b)Pöppler A-C; Granitzka M; Herbst-Irmer R; Chen Y-S; Iversen BB; John M; Mata RA; Stalke D Characterization of a Multicomponent Lithium Lithiate from a Combined X-Ray Diffraction, NMR Spectroscopy, and Computational Approach. *Angew. Chem. Int. Ed* 2014, 53, 13282. (c)Neufeld R; Teuteberg TL; Herbst-Irmer R; Mata RA; Stalke D Solution Structures of Hauser Base i-Pr₂NMgCl and Turbo- Hauser Base i-Pr₂NMgCl·LiCl in THF and the Influence of LiCl on the Schlenk Equilibrium. *J. Am. Chem. Soc* 2016, 138, 4796. [PubMed: 27011251]
15. Gaussian 09, Revision A.02, Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Scalmani G, Barone V, Petersson GA, Nakatsuji H, Li X, Caricato M, Marenich A, Bloino J, Janesko BG, Gomperts R, Mennucci B, Hratchian HP, Ortiz JV, Izmaylov AF, Sonnenberg JL, Williams-Young D, Ding F, Lipparini F, Egidi F, Goings J, Peng B, Petrone A, Henderson T, Ranasinghe D, Zakrzewski VG, Gao J, Rega N, Zheng G, Liang W, Hada M, Ehara M, Toyota K, Fukuda R, Hasegawa J, Ishida M, Nakajima T, Honda Y, Kitao O, Nakai H, Vreven

- T, Throssell K, Montgomery JA Jr., Peralta JE, Ogliaro F, Bearpark M, Heyd JJ, Brothers E, Kudin KN, Staroverov VN, Keith T, Kobayashi R, Normand J, Raghavachari K, Rendell A, Burant JC, Iyengar SS, Tomasi J, Cossi M, Millam JM, Klene M, Adamo C, Cammi R, Ochterski JW, Martin RL, Morokuma K, Farkas O, Foresman JB, and Fox DJ, Gaussian, Inc., Wallingford CT, 2016.
16. It is often the case that exchange of magnetically inequivalent ^6Li nuclei within aggregates can, on warming, undergo rapid intraaggregate exchange, causing each aggregate to display a single ^6Li resonance. See reference 20.
 17. (a)Lucht BL; Collum DB Lithium Hexamethyldisilazide: A View of Lithium Ion Solvation Through a Glass-Bottom Boat. *Acc. Chem. Res* 1999, 32, 1035.(b)Boche G; Fraenkel G; Cabral J; Harms K; van Eikema Hommes NJR; Lohrenz J; Marsch M; Schleyer P. v. R. *exo,exo*-[1,3-Bis(trimethylsilyl)allyl]lithium-*N,N,N',N'*-tetramethylethylenediamine Complex: Crystal Structure and Dynamics in Solution. *J. Am. Chem. Soc* 1992, 114, 1562.(c)Hilmersson G; Davidsson OA Multinuclear NMR Study of a Chiral Lithium Amide with an Intramolecular Chelating Methoxy Group in Coordinating Solvents at the Slow Ligand Exchange Limit. *J. Org. Chem* 1995, 60, 7660.(d)Reich HJ; Kulicke KJ Dynamics of Solvent Exchange in Organolithium Reagents. Lithium as a Center of Chirality. *J. Am. Chem. Soc* 1996, 118, 273.
 18. (a)Bissell ER; Finger M Organic Deuterium Compounds. II. Some Deuterated Tetrahydrofurans. *J. Org. Chem* 1959, 24, 1259.(b)Yang YK; Bergman RG Synthesis, molecular structure, and thermal chemistry of (η -5-cyclopentadienyl) dicarbonylrhenacyclopentane. *Organometallics* 1985, 4, 129.
 19. (a)Putau A; Koszinowski K Probing Cyanocuprates by Electrospray Ionization Mass Spectrometry. *Organometallics* 2010, 29, 3593.(b)Cva ka J; Svatoš A Matrix-assisted laser desorption/ionization analysis of lipids and high molecular weight hydrocarbons with lithium 2,5-dihydroxybenzoate matrix. *Rapid Commun. Mass Spectrom* 2003, 17, 2203. [PubMed: 14515318]
 20. Liou LR; McNeil AJ; Toombes GES; Collum DB Structures of β -Amino Ester Enolates: New Strategies Using the Method of Continuous Variation. *J. Am. Chem. Soc* 2008, 130, 17334. [PubMed: 19090750]
 21. The intended mole fraction refers to the mole fraction based on what was added to the samples. The measured mole fraction—the mole fraction within only the ensemble of interest—eliminates the distorting effects of impurities.
 22. Haq I; Lincoln P; Suh D; Norden B; Chowdhry BZ; Chaires JB Interaction of .DELTA.- and .LAMBDA.-[Ru(phen)2DPPZ]2+ with DNA: A Calorimetric and Equilibrium Binding Study. *J. Am. Chem. Soc* 1995, 117, 4788.
 23. Additional resonances appear at low THF concentrations.
 24. Reich HJ Role of Organolithium Aggregates and Mixed Aggregates in Organolithium Mechanisms. *Chem. Rev* 2013, 113, 7130. [PubMed: 23941648]
 25. Allowing samples to stand at low temperature for weeks causes no detectable increase in free enolate concentration, suggesting that the failure to observe appreciable 4 is not the result of a suppressed rate of exchange.
 26. In some cases, adding LiCl to hexalithiated heteroaggregates causes considerable additional debris to appear, attesting to significant formation of structurally complicated LiCl-enolate mixed aggregates.
 27. Monitoring the influence of LiCl on the IR absorbance of enolate 4 at 1640 cm^{-1} revealed broadening of the peak with no chemical shift changes.⁻¹
 28. Algera RF; Gupta L; Hoepker AC; Liang J; Ma Y; Singh KJ; Collum DB Lithium Diisopropylamide: Non-Equilibrium Kinetics and Lessons Learned about Rate Limitation. *J. Org. Chem* 2017, 82, 4513. [PubMed: 28368117]
 29. LiBr is approximately 15-fold less effective at mediating the alkylation. The low concentrations generated during the alkylation are inconsequential.
 30. We refer to (LiX)_n and (LiX)_m(LiX')_n as a “homoaggregate” and “heteroaggregate,” respectively, and reserve the term “mixed aggregate” for (LiX)_m(LiY)_n.
 31. Williard PG, unpublished.
 32. For a discussion of the relationship of Li-N coupling and Li-N bonding, see:Engelhardt F; Maaß C; Andrada DM; Herbst-Irmer R; Stalke D Benchmarking lithium mide versus amine bonding by

charge density and energy decomposition analysis arguments. Chem. Sci 2018, 9, 3111. [PubMed: 29732094]

33. In some literature reports, mixed aggregates containing a single subunit of substrate and inorganic salts are referred to as “deaggregated.” We find this usage to be unproductive.
34. Lithium sulfonate in catalytic quantities influenced the stereochemistry of acetylide additions without influencing the structure of the observable mixed aggregate: Parsons RL Jr. Development of Commercially Viable, Enantioselective Processes for HIV-1 NNRTIs. Curr. Opin. Drug Discovery Dev 2000, 3, 783.
35. Attempts to grow crystals of 2LiCl afforded only homoaggregate 4.
36. Yakimansky AV; Müller AHE; Van Beylen M Density Functional Theory Study on the Aggregation and Dissociation Behavior of Lithium Chloride in THF and Its Interaction with the Active Centers of the Anionic Polymerization of Methyl Methacrylate and Styrene. Macromolecules 2000, 33, 5686.
37. Romesberg FE; Collum DB Lithium Dialkylamide Mixed Aggregation: An NMR Spectroscopic Study of the Influence of Hexamethylphosphoramide (HMPA). J. Am. Chem. Soc 1994, 116, 9198.
38. Alberty RA Principle of Detailed Balance in Kinetics. J. Chem. Educ 2004, 81, 1206 Hammes, G. G. Principles of Chemical Kinetics; Academic Press: New York, 1978; pp 14–15.

Figure 1. ^6Li NMR spectra of a solution of $0.10\text{ M } [^6\text{Li}](R,R)\text{-2}$ in neat (12.3 M) tetrahydrofuran (THF) recorded at $-80\text{ }^\circ\text{C}$. (A) Unaged; (B) aged at $25\text{ }^\circ\text{C}$ for 10 min. Inset in (B) shows the resolution at $-40\text{ }^\circ\text{C}$. (C) and (D) NMR spectra of an aged solution of $0.10\text{ M } [^6\text{Li}, ^{15}\text{N}](R,R)\text{-2}$. (C) ^6Li NMR spectrum recorded at $-20\text{ }^\circ\text{C}$ (to improve resolution). (D) ^{15}N NMR spectrum recorded at $-80\text{ }^\circ\text{C}$.

Figure 2. Partial ¹H NMR spectra of an aged solution of 0.10 M (*R,R*)-**2** in 0.44 M THF and 9.1 M toluene-*d*₈ at 20 °C. (A) Coupled; (B) single-frequency irradiated at 1.40 ppm.

Figure 3. ^6Li NMR spectra of aged solutions of 0.10 M $[^6\text{Li}](R,R)\text{-2}$ at $-95\text{ }^\circ\text{C}$. (A) 0.20 M THF; (B) 0.40 M $[^{15}\text{N}]$ isoquinoline and 0.20 M THF. The asterisk (*) flags THF-isoquinoline mixed-solvated **4**.

Figure 4.
Critical resonance correlations of the *(R,R)*-2-derived homoaggregate.

Figure 5. Partial X-ray crystal structure of octalithiated enolate **4** showing the core with one subunit and the total structure missing four THF ligands.

Figure 6.

^6Li NMR spectra of aged solutions of enolate mixtures in neat THF at $-80\text{ }^\circ\text{C}$: (A) 0.050 M $[\text{}^6\text{Li}](S,S)\text{-2}$ and 0.050 M $[\text{}^6\text{Li}](R,R)\text{-2}$; and (B) 0.050 M $[\text{}^6\text{Li}, \text{}^{15}\text{N}](S,S)\text{-2}$ and 0.050 M $[\text{}^6\text{Li}, \text{}^{15}\text{N}](R,R)\text{-2}$. The resonance labeled “alkoxides” corresponds to monolithiated substrate.

Figure 7.

Job plot showing the relative integrations of octalithio homoaggregates (R_4/S_4) and hexalithio heteroaggregates (RS_2/R_2S) versus intended mole fractions²¹ of $[^6\text{Li}](R,R)\text{-2}$ (X_R) for 0.10 M mixtures of enantiomeric lithium enolates $[^6\text{Li}](R,R)\text{-2}$ (**R**) and $[^6\text{Li}](S,S)\text{-2}$ (**S**) in neat THF at -80°C monitored with ^6Li NMR spectroscopy. The curves result from a parametric fit to a tetramer–trimer model.²⁰

Figure 8.

Plot of diffusion coefficient (D) from diffusion NMR spectroscopy versus molecular weight of an aged solution of 0.040 M (S,S)-**2** and 0.060 M (R,R)-**2** in neat THF at $-60\text{ }^\circ\text{C}$; $y = a + bx$, $a = -4.622$, $b = 0.443$. This experiment assumes four coordinated THF ligands per homotetramer and four per heterotrimer.

Figure 9.

Critical resonance correlations of subunits in the hexalithiated heterochiral aggregate formed from (*R,R*)-**2** and (*S,S*)-**2**.

Figure 10. DFT-computed structure of the most stable hexalithiated enolate aggregate, **5**, showing the Li_6O_6 core (left) and distorted octahedral core described by **36** (right).

Figure 11.

^6Li NMR spectra of 0.10 M solutions of $[\text{}^6\text{Li}](S,S)\text{-8}$ (S) and $[\text{}^6\text{Li}](R,R)\text{-11}$ (R') in neat THF at -80°C . The measured mole fractions of $[\text{}^6\text{Li}](R,R)\text{-11}$ ($X_{R'}$) are 0.00, 0.40, 0.57, and 1.00 for A–D, respectively.

Figure 12.

Job plot showing the relative integrations of octalithio homoaggregates and hexalithio heteroaggregates versus measured mole fractions of $[\text{}^6\text{Li}](R,R)\text{-}\mathbf{11}$ ($X_{R'}$) for 0.10 M mixtures of lithium enolates $[\text{}^6\text{Li}](S,S)\text{-}\mathbf{8}$ (S) and $[\text{}^6\text{Li}](R,R)\text{-}\mathbf{11}$ (R') in neat THF at $-80\text{ }^{\circ}\text{C}$ monitored with ^6Li NMR spectroscopy (Figure 11). The curves result from a parametric fit to a tetramer-trimer model.²⁰

Figure 13.

^6Li NMR spectra of 0.10 M solutions of $[\text{}^6\text{Li}](S,S)\text{-16}$ (**S**) and $[\text{}^6\text{Li}](R,R)\text{-14}$ (**R'**) in neat THF at -80°C at the ratios indicated on the spectra.

Figure 14.

^{19}F NMR spectra of 0.10 M solutions of $[\text{}^6\text{Li}](S,S)\text{-16}$ (S) and $[\text{}^6\text{Li}](R,R)\text{-14}$ (R') in neat THF at -80°C . The measured mole fractions of $[\text{}^6\text{Li}](R,R)\text{-14}$ ($X_{R'}$) are 0.00, 0.34, 0.67, and 1.00 for A–D, respectively.

Figure 15.

Job plot showing the relative integrations of tetralithio homo- and hexalithio heteroaggregates versus measured mole fractions of $[^6\text{Li}](R,R)\text{-14}$ ($X_{R'}$) for 0.10 M mixtures of lithium enolates $[^6\text{Li}](S,S)\text{-16}$ (**S**) and $[^6\text{Li}](R,R)\text{-14}$ (**R'**) in neat THF at $-80\text{ }^{\circ}\text{C}$ monitored with ^{19}F NMR spectroscopy (see Figure 14). The curves result from a parametric fit to a dimer–trimer model.²⁰

Figure 16. ^6Li NMR spectra of 0.10 M solutions of $[\text{Li}](R,R)\text{-2}$ and 0.20 M LDA in neat THF at -80°C using (A) $[\text{Li}]\text{LDA}$; (B) $[\text{Li},^{15}\text{N}]\text{LDA}$; and (C) $[\text{Li},^{15}\text{N}]\text{LDA}$ and $[\text{Li},^{15}\text{N}](R,R)\text{-2}$.

Figure 17.

Structure of enolate-LDA mixed aggregate **6**. The four solvents supported by DFT computations have been omitted in the computed rendition for clarity.

Figure 18.

^6Li NMR spectra of a pre-aged 0.10 M solution of $[\text{}^6\text{Li}](R,R)\text{-2}$ with subsequent addition of 0.20 M LiCl in neat $\text{THF-}d_8$ recorded at incrementally rising temperatures.

Figure 19.

Plot of percent enolate **4** in solutions of a pre-aged 0.10 M solution of $[\text{}^6\text{Li}](R,R)\text{-2}$ with subsequent addition of 0.20 M LiCl in neat THF- d_8 versus temperature (Figure 18). Incremental temperature increases (black) were followed by incremental temperature decreases (blue).

Figure 20.

Rates and percent conversions of alkylation of lithium enolate (*R,R*)-**2** (0.040 M) in neat THF at $-60\text{ }^{\circ}\text{C}$ by allyl bromide (0.092 M) with various amounts of LiCl.

Figure 21.

Plot of initial rate constants (k_{obsd}) versus allyl bromide concentration for alkylation of 0.040 M (*R,R*)-**2** and 0.040 M LiCl in neat THF at $-60\text{ }^{\circ}\text{C}$. The curve depicts an unweighted least-squares fit to the function $f(x) = ax^n$ such that $a = (1.2 \pm 0.26) \times 10^{-2}$ and $n = 1.1 \pm 0.19$.

Figure 22.

Plot of initial rate constants (k_{obsd}) versus LiCl equivalence (per enolate) for alkylation of 0.040 M (*R,R*)-**2** and 0.092 M allyl bromide in neat THF at $-60\text{ }^{\circ}\text{C}$. The red curve depicts an unweighted least-squares fit to the function $f(x) = ax^n/(bx^n + 1)$ such that $a = (1.6 \pm 0.34) \times 10^{-4}$, $b = 0.57 \pm 0.25$, and $n = 1$ (set value). The blue curve depicts an unweighted least-squares fit with n as an adjustable parameter such that $a = (2.3 \pm 0.86) \times 10^{-4}$, $b = 1.2 \pm 0.52$, and $n = 2.1 \pm 0.77$.

Figure 23.

Plot of initial rate constants (k_{obsd}) versus the concentration of (*R,R*)-**2** for allylation in eq 7 with 0.080 M LiCl and 0.18 M allyl bromide in neat THF at -60°C . The asterisk (*) denotes a point at 0.010 M enolate; the absorbance at 0.010 M enolate is too low to be considered fully reliable. The curve depicts an unweighted least-squares fit to the function $f(x) = ax^m/(bx^m + 1)$ such that $a = 0.014 \pm 0.0043$, $b = 48 \pm 18$, and $m = 1$ (set value).

Figure 24.

Plot of initial rate constants (k_{obsd}) versus THF concentration for alkylation of 0.040 M (*R,R*)-**2** with 0.040 M LiCl by 0.092 M allyl bromide in hexanes cosolvent at -60°C . The curve depicts an unweighted least-squares fit to the function $f(x) = ax^n$ such that $a = (0.042 \pm 0.020) \times 10^{-4}$ and $n = 1.5 \pm 0.21$.

Scheme 1. Synthesis of ^{15}N -labeled enolate precursors.

(a) PhC(=O)H , MgCl_2 , TMSCl , NEt_3 , EtOAc , 24 h, rt; (b) TFA , MeOH , 10 min, rt; (c) TBDMSCl , imidazole, DMF , 18 h, rt; (d) LiOH , H_2O_2 , THF , 6 h, 0°C to rt; (e) $(\text{COCl})_2$, CH_2Cl_2 , 2 h, rt; (f) $^{15}\text{NH}_4\text{Cl}$, NaOH , 30 min, 0°C to rt; (g) [hydroxyl(tosyloxy)iodo]benzene, CH_3CN , 2 h, reflux; (h) acetic formic anhydride, CH_2Cl_2 , 1 h, -30°C ; (i) LiAlH_4 , THF , overnight, reflux; (j) propionyl chloride, NEt_3 , THF , 10 min, 0°C .

Scheme 2.

Isomerism arising from reversal of chelate orientation or enolate and alkoxide positioning.

Scheme 3.
Structural summary

Scheme 4.

Reactivity of Myers enolate **2** without added LiCl as measured by percent conversion after 1 h.

Scheme 5.
Mechanism of alkylation of cubic mixed aggregates.

Author Manuscript

Chart 2.

Structurally characterized homo- and heteroaggregates and mixed aggregates (**4–6**) of Myers enolates.

Chart 3.
Myers enolates.

Chart 4.
Possible di-, tetra-, and hexalithio aggregates of enolate (*R,R*)-2.

Chart 5.
Renditions of DFT-computed structures showing a representative stacked cube and octagonal prism of octalithio homoaggregates of *(R,R)*-2.

Chart 6.
Possible structures of enolate-LiCl adduct 2LiCl .