Published in final edited form as:

ACS Infect Dis. 2019 July 12; 5(7): 1139–1149. doi:10.1021/acsinfecdis.9b00010.

Discovery and Mechanistic Study of a Novel Human STING Agonist

Xiaohui Zhang^{1,2}, Bowei Liu^{1,3}, Liudi Tang⁴, Qing Su¹, Nicky Hwang¹, Mohit Sehgal¹, Junjun Cheng¹, Julia Ma¹, Xuexiang Zhang¹, Yinfei Tan⁵, Yan Zhou⁶, Zhongping Duan², Victor R. DeFilippis⁷, Usha Viswanathan¹, John Kulp¹, Yanming Du¹, Ju-Tao Guo^{1,*}, Jinhong Chang^{1,*}

¹Department of Experimental Medicine, Baruch S. Blumberg Institute, 3805 Old Easton Rd., Doylestown, Pennsylvania 18902, USA.

²Artificial Liver Center, Beijing Youan Hospital, Capital Medical University, 8 Xitoutiao, Fengtai, Beijing 100069, China.

³Department of Gastroenterology, Henan Provincial People's Hospital, People's Hospital of Zhengzhou University, 7 Weiwu Rd., Jinshui, Zhengzhou, Henan 450016, China.

⁴Microbiology and Immunology graduate program, Drexel University College of Medicine, 2900 W Queen Ln, Philadelphia, Philadelphia, Pennsylvania 19129, USA.

⁵Genomics Facilities, Fox Chase Cancer Center, 333 Cottman Ave., Philadelphia 19111, Pennsylvania, USA.

⁶Bioinformatics and Biostatistics Facility, Fox Chase Cancer Center, 333 Cottman Ave., Philadelphia 19111, USA.

⁷Vaccine and Gene Therapy Institute, Oregon Health and Science University, 505 NW 185th Avenue, Beaverton, Oregon 97006, USA

Abstract

Stimulator of interferon genes (STING) is an integral ER membrane protein that can be activated by 2'3'-cGAMP synthesized by cyclic guanosine monophosphate-adenosine monophosphate synthase (cGAS) upon binding of double-stranded DNA and activates interferon (IFN) and inflammatory cytokine response to defend against microorganism infection. Pharmacologic activation of STING has been demonstrated to induce an antiviral state and boost antitumor immunity. We previously reported a cell-based high throughput screening assay that allowed for identification of small molecule cGAS-STING pathway agonists. We report herein a compound, 6-bromo-N-(naphthalen-1-yl)benzo[d][1,3] dioxole-5-carboxamide (BNBC), that induces proinflammatory cytokine response in a human STING-dependent manner. Specifically, we showed that BNBC induced type I and III IFN dominant cytokine responses in primary human fibroblasts and peripheral blood mononuclear cells (PBMCs). BNBC also induced cytokine response in PBMC-derived myeloid dendritic cells and promoted their maturation, suggesting that

*Corresponding authors: Jinhong Chang, jinhong.chang@bblumberg.org, Ju-Tao Guo, ju-tao.guo@bblumberg.org.

Supporting Information. Methods for the synthesis and characterization of BNBC and its analogs.

STING agonist treatment could potentially regulate the activation of CD4⁺ and CD8⁺ T lymphocytes. As anticipated, treatment of primary human fibroblast cells with BNBC induced an antiviral state that inhibited the infection of several members of flaviviruses. Taken together, our results indicate that BNBC is a human STING agonist that not only induces innate antiviral immunity against a broad spectrum of viruses but may also stimulate the activation of adaptive immune response, which is important for treatment of chronic viral infections and tumors.

Graphical Abstract

Keywords

high throughput assay; STING; Innate immune; antiviral

Pattern recognition receptors (PRRs) are host cellular proteins that recognize pathogen associated molecular patterns, such as viral nucleic acids, capsids, bacterial peptidoglycans, lipopolysaccharides or other metabolites¹, and subsequently activate a proinflammatory cytokine response and cell death pathways²⁻³. These innate immune responses not only limit the proliferation and spread of microorganisms in the early phase of infection, but also facilitate the induction of a more powerfully adaptive immune response that can ultimately resolve the infections⁴⁻⁵. In addition, PRR-mediated innate immune response also plays an important role in surveillance of tumor genesis and invasion⁶⁻⁸. Stimulator of interferon genes (STING) is an endoplasmic reticulum (ER) membrane protein that plays a central role in the activation of innate immune response by multiple cellular DNA sensors⁹⁻¹⁰. Particularly, recognition of cytoplasmic double-stranded DNA by cyclic GMP-AMP synthase (cGAS), the primary cytoplasmic DNA sensor¹¹⁻¹³, catalyzes the synthesis of a cyclic dinucleotide c[G(2',5')pA(3',5')p], or 2'3'-cGAMP for short¹⁴, which binds to STING and induces its dimerization and translocation from the ER membrane to perinuclear vesicles and activates downstream signaling components, such as NF κ B and TBK-1/IRF3¹⁵⁻¹⁷, leading to the expression of interferons and other proinflammatory cytokines and chemokines¹⁷⁻¹⁸. STING is expressed in professional innate and adaptive immune cells and plays important roles in innate and adaptive immune response to various pathogens and tumors^{6, 19-21}. It is also well documented that STING responds to mitochondrial

damages^{22–25}, DNA double-strand breaks^{26–27}, and its over-activation may contribute to the onset of autoimmune diseases such as systemic lupus erythematosus^{28–29}.

Due to its critical role in host immune responses, pharmacological modulation of STING activity has been considered as a viable immunotherapeutic approach for treatment of pathogen infections, tumors and autoimmune disorders. Indeed, recent studies showed that intra-tumor administration of 2'3'-cGAMP induced profound regression of established tumors in mice and generated substantial systemic immune responses capable of rejecting distant metastases and providing long-lived immunologic memory^{30–31}. STING agonists had also been demonstrated to potentiate the efficacy of immune checkpoint blockade therapy^{32–33} and enhance the immunogenicity of vaccines^{34–35}. In addition, we and others have demonstrated that STING agonist therapy was able to induce a host immune response to control the infection of influenza A virus³⁶, hepatitis B virus^{18, 37}, herpes simplex virus³⁸ and human immunodeficiency virus³⁹. These studies prove the concept that pharmacologic activation of STING is an attractive immunotherapeutic approach to treat viral infection and cancers. Moreover, recent discovery of a small molecular inhibitor of STING paves the way for treatment of systemic lupus erythematosus and other autoimmune diseases⁴⁰.

Although several analogs of cyclic dinucleotide (CDN)-based STING agonists have been tested in tumor models in mice and demonstrated antitumor activity through activation of anticipated immune response³⁴, their medical application has been significantly limited by the low cell membrane permeability⁴¹. Accordingly, extensive efforts have been made to discover non-cyclic dinucleotide small molecule human STING agonists. One approach is to modify the mouse-specific STING agonist 5',6'-dimethylxanthene-4-acetic acid (DMXAA), which has not yet resulted in identification of novel derivatives capable of binding human STING⁴². However, α -mangostin, a dietary xanthone with antitumor and antiviral activities^{43–44}, was demonstrated to bind and activate both mouse and human STING⁴⁵. Another approach is through biochemistry- or cell-based high throughput screening (HTS). Screening of small molecules that can compete the binding of a radio-labeled cyclic dinucleotide STING agonist to the C-terminal domain of human STING identified a aminobenzimidazole compound and its dimer derivatives as potent human STING agonists⁴⁶. Alternatively, taking cell-based HTS approaches, three chemotypes of compounds have been identified to activate a cytokine response in a human STING-dependent manner^{47–49}. We report herein another novel human STING specific agonist from a cell-based HTS campaign and its pharmacological properties. Our preliminary medicinal chemistry studies indicate that this carboxamide STING agonist is a good candidate for further lead optimization and development toward a therapeutic agent for treatment of chronic viral infections and tumors.

Results

Discovery of a carboxamide compound that activates cGAS-STING pathway

We previously established a HepG2 cell-derived cell line with reconstituted human cGAS-STING pathway and ISG54 promotor-driven luciferase for high throughput screening of human cGAS-STING pathway agonists⁴⁹. Screening of the NCI Diversity Set V collection of 1,594 small molecule compounds identified a compound, 6-bromo-N-(naphthalen-1-

yl)benzo[d][1,3]dioxole-5-carboxamide (BNBC) (Fig. 1A), that significantly induced ISG54-driven luciferase expression in HepAD38/cGAS-STING/ISG54 cells. As shown in Fig 1B, treatment of HepAD38/cGAS-STING/ISG54Luc cells with BNBC for 4 h induced luciferase expression in a concentration-dependent manner. On the contrary, BNBC did not induce luciferase expression in HepAD38/cGAS-STING⁻/ISG54Luc cells that express human STING with truncation of carboxyl-terminal domain (CTD). Because the CTD of STING is essential for the activation of IRF3 and subsequent induction of cytokines, the cGAS-STING⁻ reporter cell line serves as a negative control¹⁸. While significant ISG54 promoter-driven luciferase activity was detected at low micromolar concentration in cells expressing human STING, no cytotoxicity was detected at up to 200 μ M concentration. (Fig. 1C). Furthermore, BNBC also concentration-dependently induced ISG54 driven luciferase expression in HepG2/STING/ISG54 reporter cells lacking cGAS (Fig. 1B). These results thus suggest that BNBC is a cGAS/STING pathway activator and most likely targets a cellular component down-stream of cGAS, but at or up-stream of STING.

BNBC is a human STING specific agonist

In order to precisely determine the cellular target of BNBC, we first examined the effects of BNBC on proinflammatory cytokine gene expression in HepG2 cells reconstituted with human or mouse STING, respectively, using mouse STING specific agonist DMXAA as a control. As shown in Fig. 2A and B, consistent with results obtained with ISG54 reporter assays, BNBC concentration-dependently induced IFN- β , IL29 (IFN- λ 1) and TNF- α mRNA expression in HepG2/STING cells (Fig. 2A), but not in parental HepG2 cells, HepG2/STING⁻ cells and HepG2-derived cell line expressing mouse STING (HepG2/mSTING) (Fig. 2B). As expected, DMXAA only induced IFN- β mRNA expression in HepG2/mSTING cells, but not in HepG2/STING cells expressing human STING. Because the only difference between HepG2/hSTING and HepG2/mSTING cell lines is the expression of STING from human or mice, the results thus indicate that as DMXAA is a specific agonist of mouse STING, BNBC should be a specific agonist of human STING. In agreement with this notion, while DMXAA treatment only induced the peri-nuclear translocation of mSTING and nuclear translocation of IRF3, BNBC treatment induced the peri-nuclear translocation of HepG2 cells expressing human STING, but not mouse STING (Fig. 2C). Interestingly, although BNBC treatment induced peri-nuclear translocation of both full-length and C-terminally truncated hSTING, it only induced IRF3 phosphorylation and nuclear translocation in HepG2/STING cells, but not in HepG2/STING⁻ cells (Fig. 2D). These results imply that although BNBC might specifically bind to human STING and induce its subcellular translocation in a CTD-independent manner, its activation of IRF3 phosphorylation and nuclear translocation is CTD-dependent, which is consistent with the essential role of STING CTD in recruitment of TBK1 and phosphorylation of IRF3¹⁷. Taken together, our results presented above indicate that BNBC is a human STING specific agonist.

BNBC induces a proinflammatory cytokine response and establishment of antiviral state in human foreskin fibroblast cells (THF) in a STING-dependent manner.

THF is a cell line derived from primary human diploid foreskin fibroblasts (HFF)⁵⁰ and expresses physiologically relevant levels of STING^{48–49}. As observed in HepG2-derived cell

line ectopically expressing functional human STING, BNBC concentration-dependently induced IFN- β , IL-28A and IL-6 mRNA expression as well as secretion of IFN- β into culture media in THF cells (Fig. 3A and B) in the absence of cytotoxicity (Fig. 3C). Also consistent with its role as a STING agonist, knocking out the expression of STING, but not TRIF or IPS-1 (MAVS), in THF cells completely abolished BNBC-induced IFN- β mRNA expression (Fig. 3D).

To determine the BNBC-induced gene expression profile, RNAseq analysis was performed with RNA extracted from THF cells harvested at 4, 6 and 8 h post BNBC treatment. As illustrated in Fig. 3E and F, BNBC altered the expression of 132, 211 and 310 genes at 4, 6 and 8 h of treatment, respectively. Ingenuity canonical pathway analysis indicated that many components of PRR and interferon signaling pathways, including downstream factors of IRF3 and IRF7, STAT1 and STAT3, were up-regulated (Table 1). In addition, several top regulator effect networks being altered after BNBC treatment are related to the activation of antigen presenting cells, myeloid cells and leukocytes. The RNAseq analyses thus revealed that BNBC treatment induced a typical innate proinflammatory cytokine response gene expression profile consistent with the activation of STING pathway.

In agreement with its induced gene expression profile that indicates the induction of antiviral state, BNBC treatment efficiently protected Dengue virus (DENV), Yellow fever virus (YFV) and Zika virus (ZIKV) infection of THF cells. As shown in Fig. 4, pre-treatment of THF cells with BNBC for 8 h concentration-dependently reduced the intracellular DENV, YFV and ZIKV RNA levels by up to 90, 150 and 800-fold, respectively. Similarly, pre-treatment of THF cells with another small molecule human STING agonist, G10⁴⁸, reduced viral RNAs in similar potencies against the three flaviviruses.

BNBC induced inflammatory cytokine response in human peripheral blood mononuclear cells (PBMCs) and promoted PBMC-derived dendritic cell maturation

STING is expressed in professional innate and adaptive immune cells and plays important roles in innate and adaptive immune response to various pathogens and tumors^{6, 51}. In order to investigate the effects of BNBC on the function of professional immune cells, we first examined BNBC-induced cytokine response in human PBMCs and demonstrated that BNBC efficiently induced expression of type I and type III IFN, but to a lesser extent, TNF- α (Fig. 5A). Interestingly, while cGAMP significantly induced the expression of IFN- β , IL-28A and IL-29 in PBMCs derived from all three donors (Fig. 5B), BNBC and G10 treatment only significantly enhanced type I and III IFN mRNA expression in PBMCs derived from 2 out of 3 donors (Fig. 5B). It remains to be further investigated whether the unresponsiveness of PBMC-derived from donor 1 is associated with STING gene variation that may impede the binding and activation by BNBC and G10 or other physiological factors that affects the metabolism of the non-cyclic dinucleotide STING agonists.

Dendritic cells are professional antigen presentation cells that function at the interface of innate and adaptive immune responses and play a critical role in immune control of virus infection. It has been shown that activation of pattern recognition receptors (PRRs) by pathogen-associated molecular patterns (PAMPs) in dendritic cells initiates essential innate immune signaling cascades resulting in maturation of dendritic cells and up-regulation of

their antigen presenting machinery as well as costimulatory molecules such as CD80. As shown in Fig. 6A, treatment of PBMC-derived dendritic cells with BNBC (100 μ M) or LPS (10 μ g/ml) for 6 h efficiently induced the expression of IFN- β and IL-29 mRNAs and to a lesser extent, TNF- α and IL-6 mRNAs. Moreover, treatment of PBMC-derived dendritic cells with BNBC (50 μ M) or LPS (10 μ g/ml) for 48 h significantly induced expression of CD80 (Fig. 6B and C). These results suggest that activation of STING is indeed able to induce the maturation of dendritic cells.

Preliminary structure-activity-relationship studies identified active pharmacophores of BNBC

BNBC is an amide with two fused rings on each side of the acid and amine with a MW of 370 and calculated logP of 4.0. The necessary pharmacophores of BNBC were determined through evaluating the analogs with structural alterations around B ring (57093Z, 57057Z), C ring (57071Z, 57076, and 59011Z), or both B and C rings (59016Z) (Table 2). BNBC analogs were prepared in house and tested for their activities to induce ISG54 promoter-driven luciferase expression in HepG2/hSTING cells and ranked by the Minimum Effective Concentration to induce 5-fold luciferase activity (MinEC_{5X}) relative to the mock-treated controls. As shown in Table 2, introduction of a more electronegative nitrogen atom to either B ring (57057Z) or D ring (57076) will make the resulting products more polar and thus more likely improve water solubility. However, both compounds lost activities to induce the ISG54 promoter-driven luciferase expression. Similar inactivation was also observed when a polar and water solubilizing group, morpholine, was added to the 2-position of the C ring (59011Z). These results suggested that the binding site of BNBC might be highly hydrophobic. In contrast, two analogs of BNBC with either opened A ring (57093Z) or D ring (57071Z) maintained low MinEC_{5X} values of 2.5 and 2.6 μ M, respectively, similar to that of BNBC (2.2 μ M). However, compound 59016Z, with both opened A and D rings, has a much higher MinEC_{5X} of 50 μ M, suggesting a reduced activity in activating STING. Interestingly, the compound 57093Z has significantly increased maximum fold of induction of ISG54 promoter activity indicating that it might be a more potent STING activator. These results suggested that either A ring or D ring can be opened and substituted with alkyl, halogen, or other groups to maintain or improve the STING activation activity and thus established a base for further chemical modification.

To evaluate the pharmacological properties of BNBC and its active analogs, 57071Z and 57093Z, an *in vitro* ADME profiling studies were performed. As summarized in Table 3, BNBC has good caco-2 permeability and acceptable plasma protein bindings, which predict a favorable oral absorption rate if the molecule can be made in solution. A panel of 9 cytochrome P450 (Cyp) enzymes (CYP1A2, 2A6, 2B6, 2C8, 2C9, 2C19, 2D6, CYP2E1 and 3A4 with two different substrates) were analyzed in 10 reactions. While BNBC and 57071Z each inhibits 5 out of 10 Cyps/reactions, including relatively more common enzymes CYP3A4 (determined with Midazolam and Testosterone) and/or CYP2D6, 57093Z, a more potent BNBC analog with an opened A ring, exhibits an improved Cyp inhibition profile and inhibits 2 out of 10 Cyp enzymes, CYP1A2 and CYP2C9, which are not considered to be as critical as CYP3A4 and 2D6. One major issue of BNBC and its analogs is the poor aqueous solubility in both pH 4 and pH 7.4 at sub- or low-micromolar. Further chemical modification

is necessary to improve the solubility, since an extremely low aqueous solubility is hard to overcome even through proper formulation and dosing as a suspension. As shown in Table 3, BNBC and its two analogs apparently have very rapid metabolic rate in both human and mouse liver microsomes, predicting a low metabolic stability *in vivo* and raising concern for insufficient overall drug exposure. However, since high systematic exposure is the major concern for STING agonist to cause systematic inflammatory side effect, the metabolic property of BNBC predicts a low systematic exposure after oral dosing and favors intrahepatic STING activation, which may reduce systematic toxicity and is thus beneficial for treatment of chronic HBV infection.

Discussion

BNBC is a carboxamide compound from NCI Diversity Set V compound library and was identified to induce the expression of interferon and other proinflammatory cytokines in HepG2 cells expressing functional human, but not mouse STING, suggesting that BNBC is a human STING specific agonist (Fig. 2). In agreement with this notion, BNBC induced STING peri-nuclear translocation as well as phosphorylation and nuclear translocation of IRF3 (Fig. 2). Moreover, BNBC induced cytokine response in THF and PBMCs. Its activity to induce cytokine response in THF cells was dependent on the expression of STING, but not MAVS and TRIF, adaptor for RIG-I-like receptors and TLR3/TLR4, respectively (Figs. 3 and 5). Importantly, BNBC was able to induce a gene expression profile that is consistent with STING activation (Fig. 3) and established an antiviral state in treated cells to restrict the infection of three different flaviviruses (Fig. 4). Although our biological data presented herein suggest that STING is likely the target of BNBC, it remains to be determined whether BNBC can directly bind to STING.

Due to its critical role in innate and adaptive immunity, STING is an important therapeutic target for diseases with dysfunctional immune response, particularly, cancers and chronic viral infections. Preclinical and clinical studies with different formulations of CDN-based STING agonists demonstrated promising efficacy to various tumors and adjuvant activity to induce more robust cellular and humoral immune responses. However, due to its poor cell permeability and metabolic stability, the CNDs were administrated *via* intra-tumor injection, which limited their application, particularly for treatment of viral infections and non-solid tumors. In fact, studies with DMXAA in mice clearly demonstrated that systematic administration of STING agonists is efficacious to tumors and viral infections^{52–53}. However, all the non-CDN STING agonists discovered through cell-based HTS thus far only specifically activate human, but not mouse STING^{48–49}. This species specificity impeded the *in vivo* evaluation of their therapeutic efficacy in mice models. Interestingly, α -mangostin and amidobenzimidazole derivative appear to activate both human and mouse STING, indicating that the discovery and development of human and mouse dual active non-CDN STING agonists is possible^{45–46}. Further investigation into the mechanism of different agonists to activate STING should provide clues for the development of dual active STING agonists.

Considering the mode of action on STING agonist therapy of tumors, the current experimental evidence indicates that the activation of STING in tumor cells can induce a

cytokine response and cell death^{31, 34}, which will enhance cross-presentation of tumor antigens by dendritic cells and activate T cell immune response against tumors. In addition, activation of STING in tumor stromal cells and infiltrated immune cells may also contribute to the induction of efficient anti-tumor immune response. In the case of chronic viral infection, such as chronic hepatitis B, while activation of the cytokine response in either virus infected cells or non-infected cells may inhibit viral replication, restoration of host adaptive antiviral response is essential to control the persistent viral infection and achieve clinical cure^{54–56}. Therefore, in the course of developing STING agonist, it is important to not only evaluate its activity on cytokine induction but should also assess its ability to enhance adaptive antiviral immunity in relevant cell culture and *in vivo* models. Our studies reported herein showed that BNBC not only activates an inflammatory cytokine response, but also induces dendritic cell maturation (Fig. 6). With this encouraging observation, we will further investigate immune modulating activity of BNBC and its derivatives (Table 2) *in vivo* in humanized mice models with human hepatocytes and an adaptive immune system^{57–58}.

Conclusion

BNBC represents a new chemotype human STING agonist that can activate innate and adaptive immune response for treatment of viral infections and tumors.

Materials and Methods

Cells and viruses

HepG2 is a human hepatoblastoma cell line obtained from ATCC. HepG2-derived cell lines with reconstituted human cGAS and/or STING, with/without firefly luciferase under the control of an ISG54 promoter (HepAD38/cGAS-STING/ISG54Luc, HepG2/STING/ISG54Luc and HepG2/STING), with reconstituted human STING with 39 amino acid deletion on CTD required for downstream signaling, but not ligand binding (HepAD38/cGAS-STING C/ISG54Luc, HepG2/STING C), and with reconstituted mouse STING (HepG2/mSTING) were described previously^{18, 49}. THF cells are primary human diploid foreskin fibroblasts (HFF) with reconstitution of the catalytic subunit of human telomerase⁵⁰. THF with STING knockout (THF- STING), IPS-1 knockout (THF- IPS-1) or TRIF knockout (THF- TRIF) were described and characterized previously⁵⁹. Human peripheral blood mononuclear cells (PBMCs) were isolated from the whole blood of four healthy donors (Biological Specialty) using Ficoll-Paque density gradient centrifugation (Miltenyi Biotech). Freshly isolated PBMCs were cultured in RPMI-1640 medium (Corning) containing L-Glutamine and 10% FBS. To isolate PBMC-derived dendritic cells, the suspended cells were removed from freshly isolated PBMCs after overnight culture. The remaining adherent cells were stimulated with human recombination GM-CSF (50 ng/ml, Gibco) and IL-4 (50 ng/ml, Gibco) for 7 days to induce the production of dendritic cells.

Zika virus (PRVABC59 strain) was purchased from ATCC. Yellow fever virus (17D strain) and dengue virus (serotype 2, New Guinea C strain) stocks were produced by electroporation of Huh7.5 cells with *in vitro* transcribed RNAs from the corresponding cDNA constructs, as described previously^{49, 60–61}.

Chemicals

Diversity Set V compound library containing 1,594 structure diversified small molecule compounds were acquired from the National Cancer Institute (NCI). 6-bromo-N-(naphthalen-1-yl)benzo[d][1,3]dioxole-5-carboxamide (BNBC) and its analogs were synthesized in house with > 95% purity. Synthesis and characterization data of BNBC and its analogs are shown in Supplemental Materials. BNBC and its analogs were dissolved in DMSO at a stock concentration of 50 mM. 2'3'-cGAMP and LPS were purchased from InvivoGen. DMXAA was purchased from AdooQ BioScience. G10, a previously reported human STING agonist, was purchased from Aobious⁴⁸.

ISG54 luciferase reporter assay

HepAD38/cGAS-STING/ISG54Luc cells, HepAD38/cGAS STING C/ISG54Luc cells or HepG2/STING/ISG54Luc cells were seeded in black wall/clear bottom 96-well plates at a density of 4×10^4 /well in 0.1 mL of medium for overnight. HTS and hit identification were as described previously⁴⁹. To determine the activity of BNBC, the cells were treated with 1% DMSO (mock treated control) or indicated concentrations of BNBC (or its analogs) for 4 h. The firefly luciferase activities were measured by using Steady-Glo substrate (Promega) with TopCount (Perkin Elmer)⁴⁹.

Cytotoxicity assay

Cell viability was measured using MTT assay (Sigma).

Analysis of cytokine expression

IFN- β mRNA, TNF- α mRNA, IL-28A, IL-28B, IL-29 mRNA, IL-1 β and IL-6 mRNA were measured using total cellular RNA, by quantitative RT-PCR (qRT-PCR) assays^{18, 49}. β -actin mRNA served as internal. Cytokine secreted into culture medium was detected by ELISA assay (RD Systems).

Western blot assay

Cells were lysed with NuPAGE® LDS sample buffer (Thermo Fisher Scientific) supplemented with 2.5% 2-Mercaptoethanol (Sigma) followed by electrophoresis in NuPAGE 4–12% Bis-Tris Gel (Thermo Fisher Scientific) and transfer onto a PVDF membrane (Thermo Fisher Scientific). Primary antibodies against total and phosphorylated IRF3 or β -actin were from Cell Signaling. Secondary antibodies were from LI-COR. Bolts were imaged with LI-COR Odyssey system (LI-COR Biotechnology).

Immunofluorescent assay

HepG2/STING, HepG2/STING C and HepG2/mSTING cells were treated with either BNBC or DMXAA for 2 h. The cells were fixed with PBS containing 4% paraformaldehyde followed by incubation with 0.1% Triton X-100 for 20 min. Antibodies against IRF3 or STING were from Cell Signaling. Alexa Fluor 488-conjugated secondary antibody was from Invitrogen⁴⁹. Cell nuclei were stained with DAPI.

RNAseq and data analysis

mRNA sequencing (mRNA-Seq) assays were performed in the Genomics Facility of Fox Chase Cancer Center. RNA sample quantity was measured by UV absorbance at 260, 280, and 230 nm with a NanoDrop 1000 spectrophotometer (ThermoScientific). RNA integrity was determined by bioAnalyzer using an RNA Nano chip instrument (Agilent Technologies). 1000 ng total RNAs from each sample were used to make mRNA-seq library using the Truseq stranded mRNA library kit. Specifically, mRNAs were enriched twice via poly-T based RNA purification beads, and fragmented at 94 °C for 8 min. The first strand cDNA was synthesized by Superscript II and random primers at 42 °C for 15 min, followed by second strand synthesis at 16 °C for 1 h. Adapters with illumine P5, P7 sequences as well as indices were ligated to the cDNA fragment at 30 °C for 10 min. After Ampure bead (BD) purification, a 15-cycle of PCR reaction was used to enrich the fragments. Sample libraries were subsequently pooled and loaded to the Illumina Hiseq2500. Raw sequence reads were aligned to the mouse genome (mm10) using the Tophat algorithm⁶². Cufflinks algorithm⁶³ was implemented to assemble transcripts and estimate their abundance. Cuffdiff⁶⁴ was used to statistically assess expression changes in quantified genes in different conditions. Genes with false discovery rate <0.2 and fold change >2 were considered significant.

YFV, DENV and ZIKV infection and antiviral assays

THF cells were treated with indicated concentrations of STING agonists for 8 h followed by infection with DENV, YFV or ZIKV at a multiplicity of infection (MOI) of 0.1 for 1 h. Two days post infection, the viral RNAs were quantified by qRT-PCR assays using LightCycler 480II (Roche). β -actin mRNA was quantified to normalize the levels of viral RNA. Primers used for amplification of viral RNA were reported previously^{60, 65}.

Flowcytometry analysis of PBMC-derived dendritic cells

PBMC-derived dendritic cells were washed twice with PBS containing 2% FBS and 1 mM EDTA, and stained using anti-CD11c FITC and anti-CD80 PE antibodies (eBioscience) for 30 min at room temperature. Data were acquired by GUAVA (Millipore) and analyzed with FlowJo.

In vitro adsorption, distribution, metabolism and elimination (ADME) profiling of BNBC and its analogs

In vitro ADME profiling studies were performed by Pharmaron. Aqueous solubility in PBS at pH 4.0 and 7.4 was determined at up to 300 μ M. Metabolic stability in human and mouse liver microsomes was determined at 0, 15, 30, 45 and 60 min post incubation and expressed as time of 50% reduction compared to 0 min (T_{1/2}). Permeability in human epithelial colorectal adenocarcinoma cells Caco-2 was assessed and expressed as transportation rate in both directions (apical to basolateral (A-B) and basolateral to apical (B-A)) across the cell monolayer, as well as efflux ratio, an indicator of a compound's active efflux. Plasma protein binding (PPB) in samples of human or mouse origins was determined using equilibrium dialysis method and expressed as percentage of binding. Inhibition of each of the 10 cytochrome P450 (CYP) isozymes (CYP1A2, CYP2A6, CYP2B6, CYP2C8,

CYP2C9, CYP2C19, CYP2D6, CYP2E1 and CYP3A4) was determined as greater than 50% inhibition at 10 μ M concentration.

Statistics

P values of unpaired *t* test were calculated using GraphPad QuickCalcs.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank Wade Bresnahan at University of Minnesota for providing THF cells. A cDNA clone of serotype 2 dengue virus, pACYC177-NGC-DENV-2, was a gift from Pei-Yong Shi at University of Texas Medical Branch. A cDNA clone of Yellow fever virus 17D strain, pACNR/FLYF-17Dx, was a gift from Charles Rice at Rockefeller University.

Funding

This work was supported by grants from the National Institutes of Health, USA (AI134732 and AI113267), Arbutus Biopharma Inc. and the Commonwealth of Pennsylvania through the Hepatitis B Foundation.

Abbreviations

BNBC	6-bromo-N-(naphthalen-1-yl)benzo[d][1,3]dioxole-5-carboxamide
cGAS	cyclic guanosine monophosphate-adenosine monophosphate synthase
DMXAA	5,6-Dimethylxanthenone-4-acetic acid
HBV	hepatitis B virus
HFF	human foreskin fibroblast
IFN	interferon
LPS	Lipopolysaccharides
mDCs	Myeloid dendritic cells
PAMPs	pathogen-associated molecular patterns
PBMCs	peripheral blood mononuclear cells
PRRs	pattern recognition receptors
STING	stimulator of interferon genes
TLR	toll-like receptor
TNF-α	tumor necrosis factor-alpha

REFERENCES

1. Zhou P; She Y; Dong N; Li P; He H; Borio A; Wu Q; Lu S; Ding X; Cao Y; Xu Y; Gao W; Dong M; Ding J; Wang DC; Zamyatina A; Shao F, Alpha-kinase 1 is a cytosolic innate immune receptor for bacterial ADP-heptose. *Nature* 2018, 561 (7721), 122–126. [PubMed: 30111836]
2. Akira S; Uematsu S; Takeuchi O, Pathogen recognition and innate immunity. *Cell* 2006, 124 (4), 783–801 DOI: 10.1038/s41586-018-0433-3. [PubMed: 16497588]
3. Shi J; Gao W; Shao F, Pyroptosis: Gasdermin-Mediated Programmed Necrotic Cell Death. *Trends Biochem Sci* 2017, 42 (4), 245–254 DOI: 10.1016/j.tibs.2016.10.004. [PubMed: 27932073]
4. Iwasaki A; Medzhitov R, Control of adaptive immunity by the innate immune system. *Nat Immunol* 2015, 16 (4), 343–53 DOI: 10.1038/ni.3123. [PubMed: 25789684]
5. Chang J; Block TM; Guo JT, The innate immune response to hepatitis B virus infection: implications for pathogenesis and therapy. *Antiviral Res* 2012, 96 (3), 405–13 DOI: 10.1016/j.antiviral.2012.10.001. [PubMed: 23072881]
6. Corrales L; McWhirter SM; Dubensky TW Jr.; Gajewski TF, The host STING pathway at the interface of cancer and immunity. *J Clin Invest* 2016, 126 (7), 2404–11 DOI: 10.1172/JCI86892. [PubMed: 27367184]
7. Chiba S; Baghdadi M; Akiba H; Yoshiyama H; Kinoshita I; Dosaka-Akita H; Fujioka Y; Ohba Y; Gorman JV; Colgan JD; Hirashima M; Uede T; Takaoka A; Yagita H; Jinushi M, Tumor-infiltrating DCs suppress nucleic acid-mediated innate immune responses through interactions between the receptor TIM-3 and the alarmin HMGB1. *Nat Immunol* 2012, 13 (9), 832–42 DOI: 10.1038/ni.2376. [PubMed: 22842346]
8. Cen X; Liu S; Cheng K, The Role of Toll-Like Receptor in Inflammation and Tumor Immunity. *Frontiers in pharmacology* 2018, 9, 878 DOI: 10.3389/fphar.2018.00878. [PubMed: 30127747]
9. Kondo T; Kobayashi J; Saitoh T; Maruyama K; Ishii KJ; Barber GN; Komatsu K; Akira S; Kawai T, DNA damage sensor MRE11 recognizes cytosolic double-stranded DNA and induces type I interferon by regulating STING trafficking. *Proc Natl Acad Sci U S A* 2013, 110 (8), 2969–74 DOI: 10.1073/pnas.1222694110. [PubMed: 23388631]
10. Chen Q; Sun L; Chen ZJ, Regulation and function of the cGAS-STING pathway of cytosolic DNA sensing. *Nat Immunol* 2016, 17 (10), 1142–9 DOI: 10.1038/ni.3558. [PubMed: 27648547]
11. Wu J; Sun L; Chen X; Du F; Shi H; Chen C; Chen ZJ, Cyclic GMP-AMP is an endogenous second messenger in innate immune signaling by cytosolic DNA. *Science* 2013, 339 (6121), 826–30 DOI: 10.1126/science.1229963. [PubMed: 23258412]
12. Sun L; Wu J; Du F; Chen X; Chen ZJ, Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* 2013, 339 (6121), 786–91. [PubMed: 23258413]
13. Cai X; Chiu YH; Chen ZJ, The cGAS-cGAMP-STING pathway of cytosolic DNA sensing and signaling. *Mol Cell* 2014, 54 (2), 289–96 DOI: 10.1126/science.1232458. [PubMed: 24766893]
14. Zhang X; Shi H; Wu J; Zhang X; Sun L; Chen C; Chen ZJ, Cyclic GMP-AMP containing mixed phosphodiester linkages is an endogenous high-affinity ligand for STING. *Mol Cell* 2013, 51 (2), 226–35 DOI: 10.1016/j.molcel.2013.05.022. [PubMed: 23747010]
15. Burdette DL; Monroe KM; Sotelo-Troha K; Iwig JS; Eckert B; Hyodo M; Hayakawa Y; Vance RE, STING is a direct innate immune sensor of cyclic di-GMP. *Nature* 2011, 478 (7370), 515–8 DOI: 10.1038/nature10429. [PubMed: 21947006]
16. Yin Q; Tian Y; Kabaleeswaran V; Jiang X; Tu D; Eck MJ; Chen ZJ; Wu H, Cyclic di-GMP sensing via the innate immune signaling protein STING. *Mol Cell* 2012, 46 (6), 735–45 DOI: 10.1016/j.molcel.2012.05.029. [PubMed: 22705373]
17. Tanaka Y; Chen ZJ, STING specifies IRF3 phosphorylation by TBK1 in the cytosolic DNA signaling pathway. *Science signaling* 2012, 5 (214), ra20 DOI: 10.1126/scisignal.2002521. [PubMed: 22394562]
18. Guo F; Tang L; Shu S; Sehgal M; Sheraz M; Liu B; Zhao Q; Cheng J; Zhao X; Zhou T; Chang J; Guo JT, Activation of STING in hepatocytes suppresses the replication of hepatitis B virus. *Antimicrob Agents Chemother* 2017, 61 (10), e00771–17 DOI: 10.1128/AAC.00771-17. [PubMed: 28717041]

19. Li XD; Wu J; Gao D; Wang H; Sun L; Chen ZJ, Pivotal roles of cGAS-cGAMP signaling in antiviral defense and immune adjuvant effects. *Science* 2013, 341 (6152), 1390–4 DOI: 10.1126/science.1244040. [PubMed: 23989956]
20. Ma Z; Damania B, The cGAS-STING Defense Pathway and Its Counteraction by Viruses. *Cell Host Microbe* 2016, 19 (2), 150–8 DOI: 10.1016/j.chom.2016.01.010. [PubMed: 26867174]
21. Gao D; Wu J; Wu YT; Du F; Aroh C; Yan N; Sun L; Chen ZJ, Cyclic GMP-AMP synthase is an innate immune sensor of HIV and other retroviruses. *Science* 2013, 341 (6148), 903–6 DOI: 10.1126/science.1240933. [PubMed: 23929945]
22. Rongvaux A; Jackson R; Harman CC; Li T; West AP; de Zoete MR; Wu Y; Yordy B; Lakhani SA; Kuan CY; Taniguchi T; Shadel GS; Chen ZJ; Iwasaki A; Flavell RA, Apoptotic caspases prevent the induction of type I interferons by mitochondrial DNA. *Cell* 2014, 159 (7), 1563–77 DOI: 10.1016/j.cell.2014.11.037. [PubMed: 25525875]
23. Aguirre S; Fernandez-Sesma A, Collateral Damage during Dengue Virus Infection: Making Sense of DNA by cGAS. *J Virol* 2017, 91 (14) DOI: 10.1128/JVI.01081-16.
24. Andreeva L; Hiller B; Kostrewa D; Lassig C; de Oliveira Mann CC; Jan Drexler D; Maiser A; Gaidt M; Leonhardt H; Hornung V; Hopfner KP, cGAS senses long and HMGB/TFAM-bound U-turn DNA by forming protein-DNA ladders. *Nature* 2017, 549 (7672), 394–398 DOI: 10.1038/nature23890. [PubMed: 28902841]
25. McArthur K; Whitehead LW; Heddleston JM; Li L; Padman BS; Oorschot V; Geoghegan ND; Chappaz S; Davidson S; San Chin H; Lane RM; Dramicanin M; Saunders TL; Sugiana C; Lessene R; Osellame LD; Chew TL; Dewson G; Lazarou M; Ramm G; Lessene G; Ryan MT; Rogers KL; van Delft MF; Kile BT, BAK/BAX macropores facilitate mitochondrial herniation and mtDNA efflux during apoptosis. *Science* 2018, 359 (6378) DOI: 10.1126/science.aao6047.
26. Deng L; Liang H; Xu M; Yang X; Burnette B; Arina A; Li XD; Mauceri H; Beckett M; Darga T; Huang X; Gajewski TF; Chen ZJ; Fu YX; Weichselbaum RR, STING-Dependent Cytosolic DNA Sensing Promotes Radiation-Induced Type I Interferon-Dependent Antitumor Immunity in Immunogenic Tumors. *Immunity* 2014, 41 (5), 843–52 DOI: 10.1016/j.immuni.2014.10.019. [PubMed: 25517616]
27. Harding SM; Benci JL; Irianto J; Discher DE; Minn AJ; Greenberg RA, Mitotic progression following DNA damage enables pattern recognition within micronuclei. *Nature* 2017, 548 (7668), 466–470 DOI: 10.1038/nature23470. [PubMed: 28759889]
28. Yan N, Immune Diseases Associated with TREX1 and STING Dysfunction. *J Interferon Cytokine Res* 2017, 37 (5), 198–206 DOI: 10.1089/jir.2016.0086. [PubMed: 28475463]
29. Crowl JT; Gray EE; Pestal K; Volkman HE; Stetson DB, Intracellular Nucleic Acid Detection in Autoimmunity. *Annu Rev Immunol* 2017, 35, 313–336 DOI: 10.1146/annurev-immunol-051116-052331. [PubMed: 28142323]
30. Corrales L; Glickman LH; McWhirter SM; Kanne DB; Sivick KE; Katibah GE; Woo SR; Lemmens E; Banda T; Leong JJ; Metchette K; Dubensky TW Jr.; Gajewski TF, Direct Activation of STING in the Tumor Microenvironment Leads to Potent and Systemic Tumor Regression and Immunity. *Cell reports* 2015, 11 (7), 1018–30 DOI: 10.1016/j.celrep.2015.04.031. [PubMed: 25959818]
31. Iurescia S; Fioretti D; Rinaldi M, Targeting Cytosolic Nucleic Acid-Sensing Pathways for Cancer Immunotherapies. *Frontiers in immunology* 2018, 9, 711 DOI: 10.3389/fimmu.2018.00711. [PubMed: 29686682]
32. Wang H; Hu S; Chen X; Shi H; Chen C; Sun L; Chen ZJ, cGAS is essential for the antitumor effect of immune checkpoint blockade. *Proc Natl Acad Sci U S A* 2017, 114 (7), 1637–1642 DOI: 10.1073/pnas.1621363114. [PubMed: 28137885]
33. Ghaffari A; Peterson N; Khalaj K; Vitkin N; Robinson A; Francis JA; Koti M, STING agonist therapy in combination with PD-1 immune checkpoint blockade enhances response to carboplatin chemotherapy in high-grade serous ovarian cancer. *British journal of cancer* 2018 DOI: 10.1038/s41416-018-0188-5.
34. Fu J; Kanne DB; Leong M; Glickman LH; McWhirter SM; Lemmens E; Metchette K; Leong JJ; Lauer P; Liu W; Sivick KE; Zeng Q; Soares KC; Zheng L; Portnoy DA; Woodward JJ; Pardoll DM; Dubensky TW Jr.; Kim Y, STING agonist formulated cancer vaccines can cure established

tumors resistant to PD-1 blockade. *Sci Transl Med* 2015, 7 (283), 283ra52 DOI: 10.1126/scitranslmed.aaa4306.

35. Hanson MC; Crespo MP; Abraham W; Moynihan KD; Szeto GL; Chen SH; Melo MB; Mueller S; Irvine DJ, Nanoparticulate STING agonists are potent lymph node-targeted vaccine adjuvants. *J Clin Invest* 2015, 125 (6), 2532–46 DOI: 10.1172/JCI79915. [PubMed: 25938786]
36. Shirey KA; Nhu QM; Yim KC; Roberts ZJ; Teijaro JR; Farber DL; Blanco JC; Vogel SN, The anti-tumor agent, 5,6-dimethylxanthenone-4-acetic acid (DMXAA), induces IFN-beta-mediated antiviral activity in vitro and in vivo. *J Leukoc Biol* 2011, 89 (3), 351–7 DOI: 10.1189/jlb.0410216. [PubMed: 21084628]
37. Guo F; Han Y; Zhao X; Wang J; Liu F; Xu C; Wei L; Jiang JD; Block TM; Guo JT; Chang J, STING agonists induce an innate antiviral immune response against hepatitis B virus. *Antimicrob Agents Chemother* 2015, 59 (2), 1273–81 DOI: 10.1128/AAC.04321-14. [PubMed: 25512416]
38. Skouboe MK; Knudsen A; Reinert LS; Boullaran C; Lioux T; Perouzel E; Thomsen MK; Paludan SR, STING agonists enable antiviral cross-talk between human cells and confer protection against genital herpes in mice. *PLoS Pathog* 2018, 14 (4), e1006976 DOI: 10.1371/journal.ppat.1006976. [PubMed: 29608601]
39. Aroh C; Wang Z; Dobbs N; Luo M; Chen Z; Gao J; Yan N, Innate Immune Activation by cGMP-AMP Nanoparticles Leads to Potent and Long-Acting Antiretroviral Response against HIV-1. *J Immunol* 2017, 199 (11), 3840–3848 DOI: 10.4049/jimmunol.1700972. [PubMed: 29084836]
40. Haag SM; Gulen MF; Reymond L; Gibelin A; Abrami L; Decout A; Heymann M; van der Goot FG; Turcatti G; Behrendt R; Ablasser A, Targeting STING with covalent small-molecule inhibitors. *Nature* 2018, 559 (7713), 269–273 DOI: 10.1038/s41586-018-0287-8. [PubMed: 29973723]
41. Li L; Yin Q; Kuss P; Maliga Z; Millan JL; Wu H; Mitchison TJ, Hydrolysis of 2'3'-cGAMP by ENPP1 and design of nonhydrolyzable analogs. *Nat Chem Biol* 2014, 10 (12), 1043–8 DOI: 10.1038/nchembio.1661. [PubMed: 25344812]
42. Hwang J; Kang T; Lee J; Choi BS; Han S, Design, synthesis, and biological evaluation of C7-functionalized DMXAA derivatives as potential human-STING agonists. *Organic & biomolecular chemistry* 2018 DOI: 10.1039/c8ob01798k.
43. Zheng S; Liu J; Faried A; Richard SA; Gao X, Novel Chemically Synthesized, Alpha-Mangostin-Loaded Nano-Particles, Enhanced Cell Death Through Multiple Pathways Against Malignant Glioma. *Journal of biomedical nanotechnology* 2018, 14 (11), 1866–1882 DOI: 10.1166/jbn.2018.2627. [PubMed: 30165924]
44. Phan TKT; Shahbazzadeh F; Pham TTH; Kihara T, Alpha-mangostin inhibits the migration and invasion of A549 lung cancer cells. *PeerJ* 2018, 6, e5027 DOI: 10.7717/peerj.5027. [PubMed: 29967723]
45. Zhang Y; Sun Z; Pei J; Luo Q; Zeng X; Li Q; Yang Z; Quan J, Identification of alpha-Mangostin as an Agonist of Human STING. *ChemMedChem* 2018 DOI: 10.1002/cmdc.201800481.
46. Ramanjulu JM; Pesiridis GS; Yang J; Concha N; Singhaus R; Zhang SY; Tran JL; Moore P; Lehmann S; Eberl HC; Muelbauer M; Schneck JL; Clemens J; Adam M; Mehlmann J; Romano J; Morales A; Kang J; Leister L; Graybill TL; Charnley AK; Ye G; Nevins N; Behnia K; Wolf AI; Kasparcova V; Nurse K; Wang L; Li Y; Klein M; Hopson CB; Guss J; Bantscheff M; Bergamini G; Reilly MA; Lian Y; Duffy KJ; Adams J; Foley KP; Gough PJ; Marquis RW; Smothers J; Hoos A; Bertin J, Design of amidobenzimidazole STING receptor agonists with systemic activity. *Nature* 2018, 564 (7736), 439–443 DOI: 10.1038/s41586-018-0705-y. [PubMed: 30405246]
47. Pryke KM; Abraham J; Sali TM; Gall BJ; Archer I; Liu A; Bambina S; Baird J; Gough M; Chakhtoura M; Haddad EK; Kirby IT; Nilsen A; Streblow DN; Hirsch AJ; Smith JL; DeFilippis VR, A Novel Agonist of the TRIF Pathway Induces a Cellular State Refractory to Replication of Zika, Chikungunya, and Dengue Viruses. *mBio* 2017, 8 (3) DOI: 10.1128/mBio.00452-17.
48. Sali TM; Pryke KM; Abraham J; Liu A; Archer I; Broeckel R; Staverosky JA; Smith JL; Al-Shammari A; Amsler L; Sheridan K; Nilsen A; Streblow DN; DeFilippis VR, Characterization of a Novel Human-Specific STING Agonist that Elicits Antiviral Activity Against Emerging Alphaviruses. *PLoS Pathog* 2015, 11 (12), e1005324 DOI: 10.1371/journal.ppat.1005324. [PubMed: 26646986]

49. Liu B; Tang L; Zhang X; Ma J; Sehgal M; Cheng J; Zhang X; Zhou Y; Du Y; Kulp J; Guo JT; Chang J, A cell-based high throughput screening assay for the discovery of cGAS-STING pathway agonists. *Antiviral Res* 2017, 147, 37–46 DOI: 10.1016/j.antiviral.2017.10.001. [PubMed: 28982551]
50. Bresnahan WA; Hultman GE; Shenk T, Replication of wild-type and mutant human cytomegalovirus in life-extended human diploid fibroblasts. *J Virol* 2000, 74 (22), 10816–8. [PubMed: 11044129]
51. Larkin B; Ilyukha V; Sorokin M; Buzdin A; Vannier E; Poltorak A, Cutting Edge: Activation of STING in T Cells Induces Type I IFN Responses and Cell Death. *J Immunol* 2017, 199 (2), 397–402 DOI: 10.4049/jimmunol.1601999. [PubMed: 28615418]
52. Baguley BC, Small-molecule cytokine inducers causing tumor necrosis. *Curr Opin Investig Drugs* 2001, 2 (7), 967–75.
53. Ching LM; Young HA; Eberly K; Yu CR, Induction of STAT and NFkappaB activation by the antitumor agents 5,6-dimethylxanthene-4-acetic acid and flavone acetic acid in a murine macrophage cell line. *Biochem Pharmacol* 1999, 58 (7), 1173–81. [PubMed: 10484075]
54. Guo JT; Guo H, Metabolism and function of hepatitis B virus cccDNA: Implications for the development of cccDNA-targeting antiviral therapeutics. *Antiviral Res* 2015, 122, 91–100 DOI: 10.1016/j.antiviral.2015.08.005. [PubMed: 26272257]
55. Tang L; Zhao Q; Wu S; Cheng J; Chang J; Guo JT, The current status and future directions of hepatitis B antiviral drug discovery. *Expert opinion on drug discovery* 2017, 12 (1), 5–15 DOI: 10.1080/17460441.2017.1255195. [PubMed: 27797587]
56. Penna A; Artini M; Cavalli A; Levrero M; Bertoletti A; Pilli M; Chisari FV; Rehmann B; Del Prete G; Fiaccadori F; Ferrari C, Long-lasting memory T cell responses following self-limited acute hepatitis B. *J Clin Invest* 1996, 98 (5), 1185–94. [PubMed: 8787682]
57. Zhu D; Liu L; Yang D; Fu S; Bian Y; Sun Z; He J; Su L; Zhang L; Peng H; Fu YX, Clearing Persistent Extracellular Antigen of Hepatitis B Virus: An Immunomodulatory Strategy To Reverse Tolerance for an Effective Therapeutic Vaccination. *J Immunol* 2016, 196 (7), 3079–87 DOI: 10.4049/jimmunol.1502061. [PubMed: 26936879]
58. Bility MT; Cheng L; Zhang Z; Luan Y; Li F; Chi L; Zhang L; Tu Z; Gao Y; Fu Y; Niu J; Wang F; Su L, Hepatitis B virus infection and immunopathogenesis in a humanized mouse model: induction of human-specific liver fibrosis and M2-like macrophages. *PLoS Pathog* 2014, 10 (3), e1004032 DOI: 10.1371/journal.ppat.1004032. [PubMed: 24651854]
59. Gall B; Pryke K; Abraham J; Mizuno N; Botto S; Sali TM; Broeckel R; Haese N; Nilsen A; Placzek A; Morrison T; Heise M; Streblow D; DeFilippis V, Emerging Alphaviruses Are Sensitive to Cellular States Induced by a Novel Small-Molecule Agonist of the STING Pathway. *J Virol* 2018, 92 (6) DOI: 10.1128/JVI.01913-17.
60. Guo F; Wu S; Julander J; Ma J; Zhang X; Kulp J; Cuconati A; Block TM; Du Y; Guo JT; Chang J, A Novel Benzodiazepine Compound Inhibits Yellow Fever Virus Infection by Specifically Targeting NS4B Protein. *J Virol* 2016 DOI: 10.1128/JVI.01253-16.
61. Ma J; Zhang X; Soloveva V; Warren T; Guo F; Wu S; Lu H; Guo J; Su Q; Shen H; Solon E; Comunale MA; Mehta A; Guo JT; Bavari S; Du Y; Block TM; Chang J, Enhancing the antiviral potency of ER alpha-glucosidase inhibitor IHVR-19029 against hemorrhagic fever viruses in vitro and in vivo. *Antiviral Res* 2017 DOI: 10.1016/j.antiviral.2017.12.008.
62. Trapnell C; Pachter L; Salzberg SL, TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* 2009, 25 (9), 1105–11 DOI: 10.1093/bioinformatics/btp120. [PubMed: 19289445]
63. Trapnell C; Williams BA; Pertea G; Mortazavi A; Kwan G; van Baren MJ; Salzberg SL; Wold BJ; Pachter L, Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat Biotechnol* 2010, 28 (5), 511–5 DOI: 10.1038/nbt.1621. [PubMed: 20436464]
64. Trapnell C; Hendrickson DG; Sauvageau M; Goff L; Rinn JL; Pachter L, Differential analysis of gene regulation at transcript resolution with RNA-seq. *Nat Biotechnol* 2013, 31 (1), 46–53 DOI: 10.1038/nbt.2450. [PubMed: 23222703]
65. Ma J; Zhang X; Soloveva V; Warren T; Guo F; Wu S; Lu H; Guo J; Su Q; Shen H; Solon E; Comunale MA; Mehta A; Guo JT; Bavari S; Du Y; Block TM; Chang J, Enhancing the antiviral

potency of ER alpha-glucosidase inhibitor IHVR-19029 against hemorrhagic fever viruses in vitro and in vivo. *Antiviral Res* 2018, 150, 112–122 DOI: 10.1016/j.antiviral.2017.12.008. [PubMed: 29253498]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1. Carboxamide compound BNBC activates cGAS-STING pathway in a functional human STING-dependent manner.

(A) Chemical structure of BNBC. (B) Activation of ISG54 promoter activities by BNBC were determined in HepAD38/cGAS-STING/ISG54Luc, HepAD38/cGAS-STING C/ISG54Luc cells and HepG2/STING/ISG54Luc cells. Luciferase activity was measured at 4 h post treatment and expressed as fold of induction (mean $\pm$ standard deviation, $n=3$). * indicates $p < 0.05$ compared to mock treated control. (C) Cytotoxicity was determined in HepAD38/cGAS-STING/ISG54Luc and HepAD38/cGAS-STING C/ISG54Luc cells after 4 h treatment and expressed as percent of mock treated control (mean $\pm$ standard deviation, $n=3$).

Figure 2. BNBC is a human STING specific agonist.

(A) HepG2/STING cells were treated with indicated concentrations of BNBC for 6 h. The cytokine mRNAs were measured by qRT-PCR and expressed as fold of induction (mean $\pm$ standard deviation, n=4). (B) HepG2, HepG2/STING Δ C and HepG2/mSTING cells were treated with indicated concentrations of BNBC or DMXAA for 6 h. Induction of IFN- β mRNAs were determined by qRT-PCR and expressed as fold of induction (mean $\pm$ standard deviation, n=4). * indicates $p < 0.05$ compared to mock treated control. (C) HepG2/STING, HepG2/STING Δ C and HepG2/mSTING were mock treated or treated with 100 μ M of BNBC or 50 μ M of DMXAA for 2 h. The expression and subcellular localization of STING and IRF3 were determined by immunofluorescent staining (green). Cell nuclei were stained with DAPI (blue). (D) HepG2/STING and HepG2/STING Δ C cells were treated with 100 μ M BNBC for 2 h. Total and phosphorylated IRF3 were determined by Western blot assay. β -actin served as a loading control.

Figure 3. BNBC activation of inflammatory cytokine response in THF cells is STING-dependent.

(A) THF cells were treated with indicated concentrations of BNBC for 6 h, followed by quantification of cytokine mRNAs by qRT-PCR assays. The levels of cytokine mRNA were expressed as fold of induction (mean ± standard deviation, n=4). (B) THF cells were treated with indicated concentrations of BNBC or cGAMP for 18 h, followed by detection of IFN-β in culture medium with ELISA assay (mean ± standard deviation, n=4). (C) Cytotoxicity was determined in THF cells and expressed as percent of mock treated control (mean ± standard deviation, n=3). (D) Parental THF cells as well as THF cells with knockout of STING, IPS-1 or TRIF were treated with 50 μM of BNBC for 6 h. IFN-β mRNA was detected by qRT-PCR and expressed as fold of induction (mean ± standard deviation, n=4). * indicates $p < 0.05$ relative to mock treated control. (E and F) RNAseq analysis was performed using RNA samples extracted from THF cells treated with 50 μM of BNBC for indicated time periods. Venn diagram (E) illustration of numbers of up- or downregulated genes with fold change greater than 2 and FDR < 20%, relative to mock treated control. The heat map (F) is expressed as the log₂ fold difference.

Figure 4. Pretreatment of THF cells with BNBC inhibited the replication of Dengue, Yellow fever, and Zika viruses.

THF cells were treated with indicated concentrations of BNBC or control compound G10 for 8 h. Cells were then infected with DENV (A), YFV (B), or ZIKV (C) at MOI of 0.1 for 1 h and cultured for additional 48 h. Viral RNAs were determined by qRT-PCR, and expressed as values relative to mock infected control (mean $\pm$ standard deviations, $n=3$). * indicates $p < 0.05$ compared to mock treated control.

Figure 5. Activity of BNBC in human PBMCs.

(A) PBMCs freshly isolated from human blood were treated with indicated concentration of BNBC for 6 h and followed by detection of cytokine mRNAs by qRT-PCR. The levels of cytokine mRNA were expressed as fold of induction (mean $\pm$ standard deviation, $n=4$). (B) PBMCs freshly isolated from human blood of additional three healthy donors were treated with 15 μ M of 2'3'-cGAMP, 60 μ M of BNBC or G10 for 6 h. The mRNAs of IFN- β , IL-28A, IL-29 and TNF α were detected by qRT-PCR and expressed as fold of induction (mean $\pm$ standard deviation, $n=3$). * indicates $p < 0.05$ compared to mock treated control.

Figure 6. BNBC activated type I and III interferon dominated inflammatory cytokine response in PBMC-derived dendritic cells, and promoted dendritic cell maturation.

(A) Human PBMC-derived dendritic cells were treated with 10 μ g/ml LPS or 100 μ M of BNBC for 6 h. Cytokine mRNAs were detected by qRT-PCR and expressed as fold of induction relative to mock treated control (mean $\pm$ standard deviation, n=3). (B and C) Human PBMC-derived dendritic cells were treated with 10 μ g/ml LPS or 50 μ M of BNBC for 48 h. CD80 expression was gated from CD11c+ cells (B). Fluorescent intensity of CD80 from 4 independent experiments using PBMCs from 2 donors were expressed as mean $\pm$ standard deviation (MFI) (C). * indicates $p < 0.05$ compared to mock treated control.

Table 1.

Analysis of gene alterations using Ingenuity analysis

Ingenuity canonical pathway	<i>p</i> value	Overlap
Interferon signaling	4.8E-13	27.8% (10/36)
Activation of IRF by cytosolic PRR	2.0-E10	15.9% (10/63)
Role of PRR in recognition of bacteria and viruses	3.9E-08	8.0% (11/137)
Upstream regulator	<i>p</i> value	Predicted activation
IRF7	2.5E-57	Activated
STAT1	6.1-E46	Activated
IRF3	5.7E-43	Activated
STAT3	8.4E-36	Activated
NKX2-3	4.6E-34	Inhibited
Top regulator effect networks	Consistency score	Diseases & functions
CNOT7, EF1, IFI16, IRF1,3,5,7, NFATC2, NKX2-3...	76.42	Activation of APCs
CNOT7, EBF1, ESR1, HIF1A, IFI16, IRF1,3,4,5...	48.91	Accumulation of myeloid cells
HIF1A, IFI16, IRF5,7, KLF2, STAT4...	28.74	Activation of leukocytes

Table 2.

Structure-activity relationship study of BNBC

Cmpd*	BNBC	57093Z	57057Z	57071Z	57076	59011Z	59016Z
Structure
$I_{\text{MinEC}_{50}}$ (μM)	2.2	2.5	> 200	2.6	> 200	> 200	50
$I_{\text{Maximum induction}}$ (fold)	20.5	28.0	-	17.3	-	-	7.8
$^2\text{CC}_{50}$ (μM)	> 200	> 200	> 200	> 200	> 200	> 200	> 200

* compound.

$I_{\text{Luciferase}}$ activity was determined in HepG2/STING/ISG54Luc cells and expressed as Minimum Effective Concentration to induce 5-fold luciferase activity or maximum fold of luciferase induction.

2 Cytotoxicity was determined in HepG2/STING/ISG54Luc cells by MTT assay.

Table 3.

ADME profile of BNBC and its analogs

Cmpd [*]	Aqueous solubility (μM)		Metabolic stability T _{1/2} (min)		Caco-2			Cyp [#]	PPB (%)	
	PH = 4	PH = 7.4	Human	Mouse	P _{app} (A-B) (10 ⁻⁶ , cm/s)	P _{app} (B-A) (10 ⁻⁶ , cm/s)	Efflux Ratio	IC ₅₀ < 10 μM	Human	Mouse
BNBC	3.57	2.62	19.96	2.21	14.46	5.98	0.41	5/10	97.34	97.88
57093Z	0.28	0.26	9.50	2.70	ND	ND	ND	2/10	ND	ND
57071Z	4.14	3.94	15.39	1.81	ND	ND	ND	5/10	ND	ND

* compound.

[#] Cyps inhibited by BNBC are 1A2, 2C19, 2D6, 3A4 (Midazolam) and 3A4 (Testosterone); Cyps inhibited by 57093Z are 1A2 and 2C19; Cyps inhibited by 57071Z are 1A2, 2B6, 2C19, 3A4 (Midazolam) and 3A4 (Testosterone);