Therapeutic Potential of Lipoxin A₄ in Chronic Inflammation: Focus on Cardiometabolic Disease

Ting Fu, Muthukumar Mohan, Eoin P. Brennan, Owen L. Woodman, Catherine Godson, Phillip Kantharidis, Rebecca H. Ritchie,[#] and Cheng Xue Qin^{*,#}

Cite This: *ACS Pharmacol. Transl. Sci.* 2020, 3, 43–55

Read Online

ACCESS |

Metrics & More

Article Recommendations

ABSTRACT: Several studies have shown that failure to resolve inflammation may contribute to the progression of many chronic inflammatory disorders. It has been suggested targeting the resolution of inflammation might be a novel therapeutic approach for chronic inflammatory diseases, including inflammatory bowel disease, diabetic complications, and cardiometabolic disease. Lipoxins [LXs] are a class of endogenously generated mediators that promote the resolution of inflammation. Biological actions of LXs include inhibition of neutrophil infiltration, promotion of macrophage polarization, increase of macrophage efferocytosis, and restoration of tissue homeostasis. Recently, several studies have demonstrated that LXs and synthetic analogues protect tissues from acute and chronic inflammation. The mechanism includes down-regulation of pro-inflammatory cytokines and chemokines (e.g., interleukin-1 β and tumor necrosis factor- α), inhibition of the activation of the master pro-inflammatory pathway (e.g., nuclear factor κ -light-chain-enhancer of activated B cells pathway) and increased release of the pro-resolving cytokines (e.g., interleukin-10). Three generations of LXs analogues are well described in the literature, and more recently a fourth generation has been generated that appears to show enhanced potency. In this review, we will briefly discuss the potential therapeutic opportunity provided by lipoxin A₄ as a novel approach to treat chronic inflammatory disorders, focusing on cardiometabolic disease and the current drug development in this area.

KEYWORDS: Lipoxin A₄, resolution of inflammation, chronic inflammatory disorders, therapeutic target

INFLAMMATION AND RESOLUTION OF INFLAMMATION

Overview of Inflammation and Resolution of Inflammation. Inflammation acts as a self-protective response to remove harmful stimuli and begin the repair process. The inflammatory response is characterized by distinct phases including initiation and resolution.^{1,2} The initiation phase reflects leukocyte recruitment to injured tissue in response to locally derived protein peptides and lipid mediators. This process is outlined in Figure 1.³ Prostaglandins (PGs) and leukotrienes (LTs) are among the earliest pro-inflammatory lipid mediators derived from arachidonic acid (AA), which stimulate vessel permeability to evoke neutrophil infiltration and promote leukocyte chemotactic responses, ultimately

amplifying the inflammatory response.⁴ During this phase, pro-inflammatory cytokines [Interleukin (IL)-1 β , tumor necrosis factor (TNF)- α , and IL-8] are released by immune cells (such as neutrophil, B-cell, and T-cell) following the activation of specific pattern recognition receptors (PRRs)^{5,6} in response to damage-associated molecular patterns (DAMPs) and pathogen-associated molecular patterns (PAMPs).

Special Issue: Emerging Leaders in Pharmacology and Translational Science

Received: November 5, 2019

Published: January 17, 2020

Figure 1. An overview of immune cell infiltration and pro-inflammatory/pro-resolving mediator release in the setting of inflammation. The initiation phase of inflammation is characterized by the release of pro-inflammatory mediators and the accumulation of neutrophils. During the resolution phase, anti-inflammatory and pro-resolving mediators (e.g., LXs, Rvs, MaRs, and protectins) are released. Pro-resolving mediators limit neutrophil infiltration and promote M1-like to M2-like macrophage phenotype switching. The M2-like macrophages exert an anti-inflammatory effect through macrophage efferocytosis and the release of anti-inflammatory cytokines such as TGF- β_1 and IL-10. IL = interleukin; TNF- α = tumor necrosis factor- α ; LTs = leukotrienes; GM-CSF = granulocyte-macrophage colony-stimulating factor; VCAM = vascular cell adhesion molecule; ICAM = intercellular adhesion molecule; Tx = thromboxane; PGs = prostaglandins; LXs = lipoxins; Rvs = resolvins; MaRs = maresins; TGF- β = transforming growth factor- β ; VEGF = vascular endothelial growth factor; PAF = platelet-activating factor; PGE₂ = prostaglandin E₂. The figure was adapted from ref 3, copyright 2011, American Chemical Society. This schema was created using images provided by Inkscape and Servier Medical Art by Servier (<https://smart.servier.com/image-set-download/>), licensed under a Creative Commons Attribution 3.0 Unported License (<https://creativecommons.org/licenses/by/3.0/>).

Resolution of inflammation is dynamically regulated and coupled to the generation of bioactive mediators including lipids and peptides. It does not merely reflect the diminution in chemotactic gradients. One of the crucial aspects of a self-limiting inflammatory response is the activation of a switch in lipid mediator class production, that initiates the resolution phase.⁷ The eicosanoids lipoxins (LXs) [Lipoxin A₄ (5S,6R,15S-trihydroxy-7,9,13-trans-11-ciseicosatetraenoic acid, LXA₄) and lipoxin B₄ (5S,14R,15S-trihydroxy-6,10,12-trans-8-ciseicosatetraenoic acid, LXB₄)] are the first class of specialized pro-resolving mediators (SPMs) to be released and are derived from AA.⁸ It has previously been demonstrated that the biosynthesis of LXs through 15-lipoxygenase (LOX) can reverse the production of pro-inflammatory mediators leukotrienes B₄ (LTB₄).⁹ This phase of lipid mediator class switching marks the cessation of the inflammatory response and the initiation of the resolution of inflammation. Thus, the balance between the formulation of LXs and LTs has served as crucial evidence of the active resolution phase of inflammation.¹⁰ Following the identification of LXs,¹¹ several ω -3 polyunsaturated fatty acid (PUFA) derived SPMs have been identified, including resolvins (Rvs),¹² protectins,¹³ and maresins (MaRs).¹⁴

It is essential to switch to the resolution phases of inflammation to restore tissue homeostasis. Failure to resolve inflammation leads to excessive tissue damage and ultimately progression to chronic low-grade inflammation, which underlies many common disorders.¹⁵ Several recent studies have demonstrated that chronic inflammatory diseases may be associated with unresolved inflammation, including cancer, asthma, inflammatory bowel disease (IBD), and diabetic complications.^{16–19} Here we review the role of SPMs, with a focus on LXA₄ in the resolution phase of inflammation, highlighting their novel therapeutic potential of this approach in many common acute and chronic inflammatory conditions.

Consequences of Failure of Resolution of Inflammation. Resolution of inflammation is essential to restore tissue homeostasis and to prevent the exaggerated accumulation of apoptotic cells, to inhibit further neutrophil infiltration, and to activate various anti-inflammatory cytokines, as detailed below.¹⁵ Failure to resolve acute inflammatory response could lead to chronic inflammation in the cardiometabolic disease. Any impairment of these processes may lead to chronic inflammatory disorders, tissue fibrosis, and eventually end-organ failure.^{24,25} For example, severe acute inflammatory insult in the heart, such as myocardial infarction (MI) with unresolved inflammation, leads to subsequent HF post-

Table 1. Markers and Functions of M1- and M2-like Macrophages⁴⁴

	M1-like macrophages	M2- like macrophages
Surface markers	MHC-II, CD86, CD80	CD206, CD163, Arginase-1/2
Extracellular mediators	Release of superoxide and NO	Fibronectin 1, IGF, PDGF-C, TGF- β , FXIIIa, collagen IV
Cytokines	IL-1, IL-12, IL-23, TNF- α , CCL2, CCL5, CCL-10, CCL-11, MCP-1, iNOS	IL-1Ra, IL-10, IL-4, CCL22
Function

MI.^{20–23} DAMPs, key mediators that activate the innate immune pathways via increased neutrophil and monocyte migration.^{26,27} As a result, excessive neutrophil levels, in parallel with increased local production of inflammatory mediators (e.g., TNF- α , IL-1 β , IL-6) and proteinases can further damage the heart.^{28,29} While macrophages can phagocytose dead cells, the excessive accumulation of macrophages can increase the action of matrix metalloproteinases (MMP)-1, MMP-7, MMP-8, and MMP-12, leading to proteolytic degradation of extracellular matrix (ECM) and cardiac remodelling.³⁰ Furthermore, macrophages may also produce pro-fibrotic mediators, such as transforming growth factor- β (TGF- β) directly.³¹ The role of TGF- β in heart failure (HF) is pleiotropic, and dependent on the duration and dose.³² Excessive levels of TGF- β can activate the pro-inflammatory nuclear factor κ -light-chain-enhancer of activated B cells (NF- κ B) pathway in cardiomyocytes, ultimately leading to the development of diabetic cardiomyopathy and progression to HF.³³ However, during a self-limited inflammation, TGF- β can inhibit the release of pro-inflammatory mediators by macrophages to promote the resolution of inflammation.³⁴ By advancing our understanding of the relationship between the resolution of inflammation and chronic inflammatory disorders, this may lead to the development of a novel class of pro-resolving medicine, targeting many common inflammatory diseases.³⁵

Role of Macrophage Phenotype in the Resolution of Inflammation. Macrophages are one of the essential immune cells in different microtissue environments (Table 1). They have a high degree of plasticity and adaptability.³⁶ Several studies have suggested that macrophage differentiation plays a key role in various phases of inflammation. In chronic low-

grade inflammatory states, such as atherosclerosis, monocytes are recruited to the vessel wall in response to chemotactic gradients and upregulation of adhesion molecule expression.³⁷ Two classically described monocyte populations contribute to the inflammatory process, namely Ly6C^{high} (M1-like) and Ly6C^{low} (M2-like) monocytes. Ly6C^{high} monocytes accumulate in atherosclerotic plaques in response to local production of pro-inflammatory mediators, such as monocyte chemoattractant protein (MCP)-1.³⁷ The main role of the M1-like macrophages is to drive a pro-inflammatory program by the release of pro-inflammatory cytokines (e.g., IL-6, TNF- α).³⁸ Another subset of monocyte is Ly6C^{low}. They also accumulate in inflammatory areas and differentiate into M2-like macrophages. During the tissue healing phase, M2-like macrophages are thought to play a predominant role in promoting tissue repair.³⁹ Thus, an imbalance of the M1-/M2-like macrophages ratio can contribute to the unresolved chronic inflammatory state.^{40,41} In murine models of diabetic-induced-atherosclerosis, increased plaque area is seen concomitant with the increased expression of M1-like macrophages markers (e.g., MMP-2, IL-1) and the decreased expression of M2-like macrophages (e.g., Arg1, IL-10).⁴² This finding has consisted of bone marrow-derived macrophages (BMDMs).⁴² Furthermore, Rahman et al. have reported that continued accumulation of Ly6C^{high} monocytes and increased polarization of the M2-like macrophages are beneficial for the resolution of inflammation of atherosclerosis.⁴³ This finding has suggested that continued new monocytes recruitment is also critical for resolution.

Clinical Evidence of the Beneficial Effects of Specialized Pro-resolving Mediators. In 1984, Serhan's laboratory reported that PUFAs produce a family of pro-

Figure 2. Biosynthesis of lipoxins.⁶⁶ Three major pathways involved in the synthesis of LXs involving LOX, including 5-LOX, 15-LOX, and 12-LOX. They all contribute to the synthesis of LXs from arachidonic acid. AA = arachidonic acid; LOX = lipoxygenase; 15-HPETE = 15-hydroxy-peroxyeicosatetraenoic acid; COX-2 = cyclooxygenase-2; LTA₄ = leukotriene A₄. This schema was created using images provided by Inkscape.

resolving lipid mediators derived from ω -6.¹¹ With more sophisticated structural analysis and LC–MS/MS approaches, it is clear that LXs temporally appear in the resolution phase along with the production of resolvins, protectins, and maresins, thus forming the superfamily of mediators produced during the resolution phase of inflammation.⁴⁵ The term of “specialized pro-resolving mediators” was introduced to cover the superfamily.⁴⁶ SPMs can reduce the magnitude and duration of the inflammatory response and stimulate re-epithelialization, injury healing, and tissue regeneration.⁴⁷

The potential therapeutic action of ω -3/6 PUFA supplementation as a source of SPMs has been demonstrated *in vivo*⁴⁸ and also in humans.⁴⁹ In a large clinical trial, 11324 patients with MI (trial identifier: NCT01422317) randomly received ω -3 PUFAs 1 g/day, vitamin E 300 mg/day, or both for 3.5 years. At the end point, these data indicated that using ω -3 PUFAs can lower the risk of a fatal event.⁴⁹ Another randomized clinical trial also demonstrated that the use of ω -3 PUFAs-enriched emulsion results in improved liver function and limits the inflammatory response in patients with gastrointestinal disease ($n = 64$, trial identifier: NCT00292279).⁵⁰ In cancer studies, dietary ω -3 PUFAs limited the inflammatory response and decreased the level of IL-6 in colorectal cancer patients.⁵¹ In a Japanese breast cancer study ($n = 38234$; approval number, 13–021), there was no overall significant relationship between treatment (ω -3/6 PUFAs) and the risk of breast cancer; however, positive interaction between ω -6 PUFAs and certain subsets (on estrogen and progesterone receptor status) of breast cancer was demonstrated.⁵² Mori’s laboratory demonstrated the production and identification of resolvins that were supplemented with ω -3 PUFAs for 5 days in 21 healthy volunteers

(registration number: ACTRN12610000740099).⁵³ In addition, a clinical trial (trial identifier: NCT03434236) about using SPMs (Lipinova) as a daily nutritional supplement to improve the pro-resolving capability in patients with knee arthroplasty is currently recruiting patients, the SPMs cumulative levels in plasma and serum will be measured. To further support the therapeutic potential of LXA₄ mimetics, this class of chemicals has been shown to relieve symptoms in eczematous infants, with minimal adverse events.⁵⁴ This pro-resolution activity of LXA₄ is elicited through binding and activating a G-protein-coupled receptor (GPCR), N-formyl peptide receptor 2 (FPR2),⁵⁵ also known as ALX/FPR2.

■ INTRODUCTION TO LIPOXINS

An Overview of Lipoxins. The term “lipoxin” is derived from lipoxygenase interaction products. LXA₄ and LXB₄ are the two main lipid mediators formed in mammals.¹¹ Four LXs have been identified: LXA₄, LXB₄, aspirin-triggered lipoxin A₄ (15-epi-LXA₄), and aspirin-triggered lipoxin B₄ (15-epi-LXB₄).⁵⁶ Aspirin-induced acetylation of COX2 promotes the generation of 15-epi LXs which demonstrate similar bioactivity to native LXs.⁵⁷ As the host defense and regulation of inflammation are important in human immune disorders, LXs play a key role during the resolution phase and thus may provide a new approach for the treatment of chronic inflammatory diseases.⁵⁸ Currently, most studies are focusing on LXA₄ and 15-epi-LXA₄, as the receptors for LXB₄ and 15-epi-LXB₄ have not been identified.⁵⁹

The Biosynthesis of Lipoxins. Serhan and colleagues first isolated LXs from human leukocytes, reporting a structure consisting of four conjugated double bonds (Figure 2).¹¹ Hamberg and Samuelsson demonstrated that LXs are a class of

Figure 3. Function and distribution of FPRs in different cell types. FPR1 and ALX/FPR2 are widely distributed on the membrane of monocytes, eosinophils, neutrophils, and epithelial cells. Pathogen-associated molecular patterns (PAMPs), DAMPs, or pro-resolving mediators can activate FPRs. The common function of FPRs is regulating the resolution of inflammation. Image adapted from refs 88 and 89. Copyright 2015, 2018, Elsevier. This schema was created using images provided by Inkscape.

metabolites produced from AA. In mammalian tissues, LXs are produced by different biosynthetic pathways, involving the interaction of activated neutrophils within the epithelium, endothelium, and platelets.⁶⁰ The first pathway is the major source of LXs in the vasculature and involves the action of 5-LOX and 12-LOX (in platelets).¹¹ The second is via 15-LOX (in epithelium) and 5-LOX (in neutrophils).⁶¹ Both of these routes ultimately result in the production of LXA₄ and LXB₄. As mentioned before, the second route not only produces LXs but also pro-inflammatory LTs. One study demonstrated that the synthesis of LXs can decrease the production of LTs via the activation of 12-LOX in platelets.^{62,63} The last route is triggered by aspirin, forming lipoxin epimer structures (15-epi-LXA₄ and 15-epi-LXB₄), through the cyclooxygenase-2 (COX-2) pathway in epithelial cells.⁵⁶ Interestingly, several pro-inflammatory cytokines contribute to these biosynthetic routes. For instance, one study showed that increased levels of IL-1 β and TNF- α induce the release of COX-2, as a result of the synthesis of 15-epi-LXA₄ and 15-epi-LXB₄.⁶⁴ Another study demonstrated that granulocyte-macrophage colony-stimulating factor (GM-CSF) can increase the expression of 5-LOX in neutrophils.⁶⁵ The pathways from AA to LXs are shown in Figure 2.⁶⁶ LXs are rapidly metabolized via dehydrogenation.⁶⁷ As a result, there is a great deal of interest in developing synthetic lipoxin mimetics (sLXMs) which offer enhanced stability *in vivo* and may represent a novel therapeutic approach.^{68,69}

Lipoxin A₄ Receptors. Formyl peptide receptors (FPRs) are members of the seven-transmembrane G protein-coupled receptor (GPCR) family and are PRRs.^{70,71} The FPR family is involved in the regulation and the resolution of inflammation, highlighting their potential to modulate inflammatory responses in many disease conditions, such as infection, cancer, Alzheimer's disease, obesity, and diabetes.^{20,72,73} A number of studies have demonstrated that the immune response to FPRs involves different cell types (hepatocytes,

fibroblasts, endothelial cells, and cardiomyocytes) and tissues (lung, heart, the tunica media of coronary arteries, kidney, stomach, and colon).^{22,74–76} The typical roles of FPRs are to contribute to the immune response and the clearance of damaged cells.⁷⁷ FPRs not only respond to bacterial and mitochondrial formylated peptides but also recognize endogenous nonformylated peptides and some lipids. The human FPRs are classified into three isoforms: hFPR1, hFPR2, and hFPR3.²¹ The hFPR1 and hFPR2 are present on several host cell types, including inflammatory cells, monocytes, epithelial, endothelial, and fibroblasts.^{78,79} It has also been reported that FPRs play a key role in the pro-inflammatory response, as a result of chemotaxis, degranulation, and the oxidative burst.⁸⁰ hFPR3 is commonly expressed in dendritic cells, but its role is not well understood.²⁰ LXA₄ and their metabolic stable analogues elicit cellular responses by acting as the specific receptor A lipoxin/formyl peptide receptor 2 (ALX/FPR2), now renamed to FPR2.⁸¹ Interestingly, during the initial phase, the production of serum-amyloid-A (SAA) is increased and FPR2 is activated. By the end of the initial inflammatory stage, pro-resolving mediators (RvD1, AnxA1, and LXA₄) are produced to activate FPR2,¹⁵ and LXA₄ inhibits SAA from binding to FPR2.⁸²

FPR2 and FPR1 exhibit similar sequence homology and functions.⁸³ hFPR1 can be activated by bacterial and mitochondrial proteins and N-formylated peptide.⁸⁴ The release of N-formylmethionyl-leucyl-phenylalanine (fMLF) from the bacterial wall activates polymorphonuclear leukocytes (PMNs) bactericidal response. As a result, neutrophils degranulate, intracellular mediators are released, and excess superoxide is generated.⁸⁵ In contrast, FPR2 is a low-affinity receptor for many fMLFs.⁸⁶ It can also be activated by endogenous nonformylated ligands, such as LXA₄, synthetic lipoxin mimetics, annexin-A1, and SAA.^{21,73,87} FPR1 is predominantly involved in the pro-inflammatory immune defense against bacteria, whereas FPR2 is mainly associated

Figure 4. The bioactions of LXA₄ in a self-limiting inflammatory process. Dashed red lines indicate inhibition of LXA₄, and green solid lines indicate the pro-resolving functions of LXA₄. Data adapted from refs 20, 92, and 45. The schema was created using images provided by Inkscape and Servier Medical Art by Servier (<https://smart.servier.com/image-set-download/>), licensed under a Creative Commons Attribution 3.0 Unported License (<https://creativecommons.org/licenses/by/3.0/>).

with inflammatory disease. Overall, the FPR family plays a key role during the host immune response and resolution of inflammation (Figure 3).^{88,89} It has been suggested that LXA₄ can also activate other receptors. For instance, a study has shown that LXA₄ acts as a GPR32 agonist to increase the Ca²⁺ metabolism and promote mucin secretion in human goblet cells.⁹⁰ In addition, one study has demonstrated that the FPR2 might not be the primary receptor of LXA₄ in the liver, for which the data indicated that LXA₄ can lead to hepatic steatosis due to activation of the aryl hydrocarbon receptor.⁹¹

The Role of Lipoxin A₄ during the Resolution of Inflammation. The bioactivity of the pro-resolution lipid mediator LXA₄ includes the inhibition of neutrophil infiltration, stimulation of macrophage-mediated phagocytosis of apoptotic cells (efferocytosis), prevention of fibroblast activation, and inhibition of cellular proliferation (Figure 4).^{20,92,45} Studies show that LXs can stimulate the production of nitric oxide in endothelial cells⁹³ and up-regulate monocyte

chemotaxis and adherence, without causing any degranulation or elevation of ROS.⁹⁴ In addition, LXs can decrease the activity of NF-κB and up-regulate the expression of anti-inflammatory miRNA.^{95,96} When NF-κB action in human leukocytes is inhibited, the release of several pro-inflammatory cytokines is reduced (e.g., inhibit the production of IL-8 about 50–65%), which is associated with the accumulation and activation of neutrophils.⁹⁷

At the initial inflammatory phase, LTs, PGs, and other eicosanoids enhance the pro-inflammatory response, stimulating transendothelial neutrophil infiltration which may lead to tissue injury.⁹⁴ At the conclusion of the initial inflammatory phase, there is a temporal switch from LTB₄ to LXA₄ production.⁹⁸ As described earlier, LXs can also stimulate macrophages to change their phenotype (from M1-like to M2-like), enhance the clearance of apoptotic neutrophils,⁴⁷ and prolong the lifespan of macrophages.⁹⁹ M2-like macrophages can produce anti-inflammatory cytokines (e.g., IL-10) and

Table 2. Level of Lipoxins in Human Disease^{103,109}

SPMs	category	disease	result summary
LXA ₄ and ATL	cardiometabolic disease	metabolic syndrome	level of LXA ₄ has decreased in human blood from obese patients ¹¹⁰
		Type 2 diabetes mellitus	decreased the levels of serum LXA ₄ , may be associated with the progression of diabetic complications ¹¹¹
		peripheral artery disease	lower plasma levels of ATL in patients with symptomatic peripheral artery disease compared to healthy volunteers ¹¹²
	others	renal disease	lower levels of ATL in plasma in patients with chronic kidney disease ¹¹³
		COPD	excess expression of serum-amyloid-A and less LXA ₄ in blood ⁸²
		IBD	decreased the urinary levels of LXA ₄ and increased levels of LTs in IBD patients; the levels reflect the severity of inflammation ¹¹⁴
		localized aggressive periodontitis	decreased levels of LXA ₄ in patients with the disease compared to healthy volunteers ¹¹⁵
		rheumatoid arthritis	low levels of LXA ₄ in synovial fluid from rheumatoid arthritis patients, higher levels of pro-inflammatory mediator LTB ₄ and PGE ₂ ¹¹⁶
		colon and pancreatic cancer	insufficient production of LXA ₄ ¹⁰⁷

bioactive SPMs (e.g., LXs) to promote the resolution of inflammation, see Table 1.¹⁰⁰ One study has indicated that LXs up-regulated phosphoinositide 3-kinases/protein kinase B and extracellular signal-regulated kinases/nuclear factor erythroid 2-related factor 2 in murine and human macrophages, which are related to inhibit the apoptosis of macrophages.⁹⁹ Therefore, LXs serve as one of the earliest classes of endogenous lipid mediators generated to inhibit the accumulation of neutrophils, arrest inflammation, and promote the resolution of inflammation.¹⁰¹

Evidence for the Therapeutic Opportunity of LX-Based Approaches. An excessive immune response may lead to chronic inflammation, further tissue damage, and even to autoimmune disease.¹⁶ As a consequence, it is essential to limit both the initial inflammatory response, for the resolution of inflammation.¹⁰² The relationship between circulating LX levels and common diseases is highlighted below in Table 2.¹⁰³ Given the growing evidence of the protective role of LXs in inflammation, there is a great interest in developing novel “resolution-targeted” therapy, which could effectively treat inflammatory diseases, such as colitis, rheumatoid arthritis, asthma, and others. For instance, in renal disease, one study showed that LX treatment resulted in a higher level of mRNA suppressors of cytokine signaling (such as cytokines, growth factors, adhesion molecules, etc.), indicating that LXs exert a positive influence against acute renal failure in mice.¹⁰⁴ Another study has demonstrated that LXA₄ attenuates the accumulation of neutrophils in ischemic renal failure.¹⁰⁵

LXs may also offer a potential treatment for solid tumors (e.g., colon and pancreatic cancers), with studies showing that LXs can decrease the expression of Nrf2 and NF-κB.^{106,107} Excessive production of SAA is a key mediator in chronic obstructive pulmonary disease (COPD), especially when a large quantity of VEGF and TNF-α are present. Interestingly, the administration of LXA₄ in this setting has been shown to inhibit the expression of SAA in mice.⁸² Furthermore, the ATL analogue (15-epi-16-p-fluorophenoxy-lipoxin A₄ methyl ester) has been shown to decrease the level of inflammation in different skin conditions, including psoriasis and dermatitis, hinting at the therapeutic potential for LXs in this disease setting.¹⁰⁸

Therapeutic Use of Lipoxin A₄ in Cardiometabolic Disease. The cardiometabolic disease is a metabolic disorder influenced by multiple factors (e.g., diabetes, hypertension, etc.).¹¹⁷ Diabetes mellitus (DM) is a chronic condition that occurs when the level of glucose in the blood is elevated as a

result of poor pancreatic function.¹¹⁸ As a chronic metabolic disease, DM leads to long-term complications in several major organs, which can be divided into macrovascular and microvascular complications.¹¹⁹ Major microvascular complications include retinopathy (damage to eyes, leading to blindness), neuropathy (damage to neurons, leading to polyneuropathy), and nephropathy (damage to the kidney, leading to renal failure).^{120,121} Chronic low-grade inflammation is believed to play a key role in diabetes-associated complications, as evidenced by the elevated level of circulating inflammatory cytokines in patients with such complications.^{122–124} Thus, using immunomodulators and immunosolvents such as LXA₄ may be a therapeutic strategy.¹²⁵

Some evidence shows that LXA₄ and the more stable mimetics can limit inflammation and symptoms in animal models of diabetic complications. Borgesón et al. reported that LXA₄ and a benzo-LXA₄ analogue reduce adipose inflammation, with protection also observed in the kidney and liver in high-fat diet-induced obese mice.⁷² Interestingly, they found LXs decreased the percentage of M1-like macrophages and increased the percentage of M2-like macrophages during adipose inflammation, without a significant impact on macrophage infiltration. Further, LXA₄ reduced the mouse weight as well as levels of hepatic triglycerides and renal injury.⁷² Using the same benzo-LXA₄ analogue, Brennan and colleagues also demonstrated that LXs protects against diabetes-associated complications, including accelerated atherosclerosis⁶⁹ and kidney disease.⁶⁸ By using an ApoE^{−/−} murine model of streptozotocin-induced diabetes, they showed that metabolic parameters were not changed by LXs, suggesting that LXA₄-mediated protection is independent of glycaemic control. The total atherosclerotic plaque burden, as well as aortic expression of pro-inflammatory genes (such as IL-1β, IL-6), was reduced in the LXA₄ treatment group. Here, LXs suppressed the activation of vascular smooth muscle cells, improving endothelial cell function.

Current State of Drug Development for Lipoxin A₄. There are four separate generations of LXA₄ analogues, first introduced in Serhan et al., to prolong the half-life and bioactivation of native LXA₄.⁶⁷ The second generation, 3-oxa-LXA₄ analogues were applied in different disease models (e.g., asthma and colitis) to evaluate the PK parameters of LXA₄.^{126–128} The third generation was the pyridine/benzo-LXA₄ analogues, used extensively by this group and others.^{129,130} The benzo analogues were shown to activate the LXA₄ receptor and demonstrated pro-resolving actions,

Table 3. Overview of the Development of LXs-Based Therapy

including the enhancement of macrophage phagocytic activity.^{72,131,132} More recently, a fourth-generation LXA₄ analogue has been described, which incorporates imidazole into the structure, and initial data from this analogue indicates potent anti-inflammatory responses *in vitro*.¹³³

It has previously been shown that circulating levels of LXs are reduced in several inflammatory disorders, suggesting that LX-replacement therapy may offer a novel therapeutic approach.^{82,110,111} As a result, there is also great interest in determining the potential of using LXA₄ levels as a biomarker of disease progressions, such as COPD (trial identifier: NCT03609541) and asthma (trial identifier: NCT01898767). Currently, the ASTHMIRINE trial is investigating aspirin as a monotherapy for the treatment of uncontrolled asthma, with LXA₄ levels being measured throughout the study (trial identifier: NCT02906761, phase 3 trial, *n* = 180). The second-generation lipoxin analogue (3-oxa-lipoxin) has been developed with Berlex-Schering and has successfully moved to Phase I trial for treating IBD in 2007. The final data showed the safety of 3-oxa-lipoxin in healthy individuals. A phase I trial is also ongoing to evaluate the safety and efficacy of the LX analogue BLXA₄-ME as an oral rinse for gingivitis (trial identifier: NCT02342691). This double-blind intervention trial, has 125 participants enrolled. While the development of LXA₄ and its analogues for treating the chronic inflammatory disorder is at an early stage, it is anticipated that these clinical trials will further highlight the important role of LXA₄ in many common inflammatory diseases (Table 3).

CONCLUSION

Traditional anti-inflammatory approaches aim to block the inflammatory response and these include steroids (inhibit the activation of phospholipase A₂),¹³⁴ nonsteroidal anti-inflammatory drugs (NSAIDs) (inhibit the activation of COX), and biological drugs (e.g., TNF- α inhibitor).¹³⁵ However, these approaches are only effective in about 50% of patients with many undesirable side effects (e.g., stomach bleeding, kidney toxicity).¹³⁶ Those drugs lead to delay of the resolution phase and tissue repairing, leading to further tissue damage. It is crucial to promote the resolution phase to terminate the initial phase of inflammation. Mimicking the action of endogenous pro-resolving mediators may be a novel approach to treat chronic inflammatory diseases with fewer side effects and better outcomes for patients.¹³⁷ Since the discovery of LXA₄ several decades ago, there is now a large body of evidence in the literature demonstrating potent protective responses. It promotes the resolution of inflammation and homeostasis in many chronic inflammatory disorders, such as IBD, COPD, and diabetic complications. LXA₄ produced during the resolution phase limits neutrophil infiltration and enhances macrophage phagocytosis of apoptotic cell debris as well as microbes, to promote resolution of inflammation. The LXA₄ receptor FPR2 has been identified; the properties and distribution of the receptor are known, and we are now starting to appreciate the complex intracellular signal transduction events post-LXA₄ receptor activation, which appear to be cell type-dependent. A recent study demonstrated that LXA₄ regulates the MAPK signaling pathway via inhibition of

the phosphorylation of p38 and JNK in the airway.¹³⁸ Tabas and colleagues have provided the first evidence that MerTK signaling pathways are mobilized to produce LXA₄ in macrophages.¹³⁹ In addition, 15-epi-LXA₄ suppresses PI3K/AKT signaling pathway to regulate TNF- α -induced tissue factor in vascular endothelial cells.¹⁴⁰ From a drug development perspective, the short half-life, rapid inactivation, and costly synthesis of LXA₄ limit its drug development. However, the recent development of imidazole- and oxazole mimetics¹³³ and the development of other stable mimetics,¹⁴¹ are opening the way to the use of LXA₄-based therapy for the treatment of chronic inflammatory diseases.

AUTHOR INFORMATION

Corresponding Author

Cheng Xue Qin – Heart Failure Pharmacology, Baker Heart and Diabetes Institute, Melbourne 3004, Australia; Department of Pharmacology and Therapeutics, University of Melbourne, Parkville 3010, Australia; Department of Diabetes, Central Clinical School, Monash University, Clayton 3800, Australia; orcid.org/0000-0003-2169-2686; Email: ChengxueHelena.Qin@baker.edu.au

Authors

Ting Fu – Heart Failure Pharmacology, Baker Heart and Diabetes Institute, Melbourne 3004, Australia; Department of Pharmacology and Therapeutics, University of Melbourne, Parkville 3010, Australia

Muthukumar Mohan – Department of Diabetes, Central Clinical School, Monash University, Clayton 3800, Australia

Eoin P. Brennan – UCD Diabetes Complications Research Centre, UCD Conway Institute, UCD School of Medicine, University College Dublin, Dublin 4, Ireland

Owen L. Woodman – Heart Failure Pharmacology, Baker Heart and Diabetes Institute, Melbourne 3004, Australia

Catherine Godson – UCD Diabetes Complications Research Centre, UCD Conway Institute, UCD School of Medicine, University College Dublin, Dublin 4, Ireland

Phillip Kantharidis – Department of Diabetes, Central Clinical School, Monash University, Clayton 3800, Australia

Rebecca H. Ritchie – Department of Diabetes, Central Clinical School and Drug Discovery Biology, Monash Institute of Pharmaceutical Sciences, Monash University, Clayton 3800, Australia; Heart Failure Pharmacology, Baker Heart and Diabetes Institute, Melbourne 3004, Australia; Department of Pharmacology and Therapeutics, University of Melbourne, Parkville 3010, Australia

Complete contact information is available at: <https://pubs.acs.org/10.1021/acspsci.9b00097>

Author Contributions

#Contributed equally to this work.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported in part by both the National Health and Medical Research Council (NHMRC) of Australia, including NHMRC APP1159460, APP1122176, and the Victorian Government's Operational Infrastructure Support Program. Rebecca H. Ritchie is an NHMRC Senior Research

Fellow (APP1059960), Chengxue Qin is a NHF Future Fellow and Baker Fellow.

ABBREVIATIONS

AA, arachidonic acid; ATL, aspirin-triggered lipoxin; COPD, chronic obstructive pulmonary disease; COX-2, cyclooxygenase-2; CTGF, connective tissue growth factor; DAMPS, damage-associated molecular patterns; DM, diabetes mellitus; ECM, extracellular matrix; fMLF, *N*-formylmethionyl-leucyl-phenylalanine; FPRs, formyl peptide receptors; GM-CSF, granulocyte-macrophage colony-stimulating factor; GPCR, G protein-coupled receptor; HF, heart failure; IBD, inflammatory bowel disease; IL, interleukin; LOX, lipoxygenase; LTs, leukotrienes; LTB₄, leukotrienes B₄; LXA₄, lipoxin A₄; LXB₄, lipoxin B₄; LXs, lipoxins; Ly-6c, lymphocyte antigen 6c; MaRs, maresins; MCP-1, monocyte chemoattractant protein-1; MI, myocardial infarction; MMP, matrix metalloproteinases; NF- κ B, nuclear factor κ -light-chain-enhancer of activated B cells; PAF, platelet-activating factor; PAMPs, pathogen associated molecular patterns; PGs, prostaglandins; PRRs, pattern recognition receptors; PUFAs, polyunsaturated fatty acids; ROS, reactive oxygen species; Rvs, resolvins; SAA, serum-amyloid-A; SPMs, specialized pro-resolving mediators; TGF- β , transforming growth factor- β ; TNF- α , tumour necrosis factor- α

REFERENCES

- (1) Lawrence, T., Willoughby, D. A., and Gilroy, D. W. (2002) Anti-Inflammatory Lipid Mediators and Insights into the Resolution of Inflammation. *Nat. Rev. Immunol.* 2 (10), 787–795.
- (2) Feehan, K. T., and Gilroy, D. W. (2019) Is Resolution the End of Inflammation? *Trends Mol. Med.* 25 (3), 198–214.
- (3) Serhan, C. N. (2007) Resolution Phase of Inflammation: Novel Endogenous Anti-Inflammatory and Proresolving Lipid Mediators and Pathways. *Annu. Rev. Immunol.* 25, 101–137.
- (4) Harris, S. G., Padilla, J., Koumas, L., Ray, D., and Phipps, R. P. (2002) Prostaglandins as Modulators of Immunity. *Trends Immunol.* 23 (3), 144–150.
- (5) Zhang, J.-M., and An, J. (2007) Cytokines, Inflammation, and Pain. *Int. Anesthesiol. Clin.* 45 (2), 27–37.
- (6) Takeuchi, O., and Akira, S. (2010) Pattern Recognition Receptors and Inflammation. *Cell* 140 (6), 805–820.
- (7) Serhan, C. N. (2017) Treating Inflammation and Infection in the 21st Century: New Hints from Decoding Resolution Mediators and Mechanisms. *FASEB J.* 31 (4), 1273–1288.
- (8) Serhan, C. N., and Chiang, N. (2008) Endogenous Pro-Resolving and Anti-Inflammatory Lipid Mediators: A New Pharmacologic Genus. *Br. J. Pharmacol.* 153 (Suppl.1), S200–15.
- (9) Wu, S.-H., Liao, P.-Y., Yin, P.-L., Zhang, Y.-M., and Dong, L. (2009) Elevated Expressions of 15-Lipoxygenase and Lipoxin A4 in Children with Acute Poststreptococcal Glomerulonephritis. *Am. J. Pathol.* 174 (1), 115–122.
- (10) Kieran, N. E., Maderna, P., and Godson, C. (2004) Lipoxins: Potential Anti-Inflammatory, Proresolution, and Antifibrotic Mediators in Renal Disease. *Kidney Int.* 65 (4), 1145–1154.
- (11) Serhan, C. N., Hamberg, M., and Samuelsson, B. (1984) Lipoxins: Novel Series of Biologically Active Compounds Formed from Arachidonic Acid in Human Leukocytes. *Proc. Natl. Acad. Sci. U. S. A.* 81 (17), 5335–5339.
- (12) Serhan, C. N., Clish, C. B., Brannon, J., Colgan, S. P., Chiang, N., and Gronert, K. (2000) Novel Functional Sets of Lipid-Derived Mediators with Antiinflammatory Actions Generated from Omega-3 Fatty Acids via Cyclooxygenase 2-Nonsteroidal Antiinflammatory Drugs and Transcellular Processing. *J. Exp. Med.* 192 (8), 1197–1204.

- (13) Bazan, N. G. (2005) Neuroprotectin D1 (NPD1): A DHA-Derived Mediator That Protects Brain and Retina against Cell Injury-Induced Oxidative Stress. *Brain Pathol.* 15 (2), 159–166.
- (14) Serhan, C. N., Yang, R., Martinod, K., Kasuga, K., Pillai, P. S., Porter, T. F., Oh, S. F., and Spite, M. (2009) Maresins: Novel Macrophage Mediators with Potent Antiinflammatory and Proresolving Actions. *J. Exp. Med.* 206 (1), 15–23.
- (15) Serhan, C. N., Brain, S. D., Buckley, C. D., Gilroy, D. W., Haslett, C., O'Neill, L. A. J., Perretti, M., Rossi, A. G., and Wallace, J. L. (2007) Resolution of Inflammation: State of the Art, Definitions and Terms. *FASEB J.* 21 (2), 325–332.
- (16) Khatami, M. (2011) Unresolved Inflammation: “immune Tsunami” or Erosion of Integrity in Immune-Privileged and Immune-Responsive Tissues and Acute and Chronic Inflammatory Diseases or Cancer. *Expert Opin. Biol. Ther.* 11 (11), 1419–1432.
- (17) Vachier, I., Bonnans, C., Chavis, C., Farce, M., Godard, P., Bousquet, J., and Chanez, P. (2005) Severe Asthma Is Associated with a Loss of LX4, an Endogenous Anti-Inflammatory Compound. *J. Allergy Clin. Immunol.* 115 (1), 55–60.
- (18) Purvis, G. S. D., Chiazza, F., Chen, J., Azevedo-Loiola, R., Martin, L., Kusters, D. H. M., Reutelingsperger, C., Fountoulakis, N., Gnudi, L., Yaqoob, M. M., et al. (2018) Annexin A1 Attenuates Microvascular Complications through Restoration of Akt Signalling in a Murine Model of Type 1 Diabetes. *Diabetologia* 61 (2), 482–495.
- (19) Purvis, G. S. D., Collino, M., Loiola, R. A., Baragetti, A., Chiazza, F., Brovelli, M., Sheikh, M. H., Collotta, D., Cento, A., Mastrocola, R., et al. (2019) Identification of AnnexinA1 as an Endogenous Regulator of RhoA, and Its Role in the Pathophysiology and Experimental Therapy of Type-2 Diabetes. *Front. Immunol.* 10 (MAR), 571.
- (20) Qin, C., Yang, Y. H., May, L., Gao, X., Stewart, A. G., Tu, Y., Woodman, O. L., and Ritchie, R. H. (2015) Cardioprotective Potential of Annexin-A1 Mimetics in Myocardial Infarction. *Pharmacol. Ther.* 148, 47–65.
- (21) Qin, C. X., Finlayson, S. B., Al-Sharea, A., Tate, M., De Blasio, M. J., Deo, M., Rosli, S., Prakoso, D., Thomas, C. J., Kiriazis, H., et al. (2017) Endogenous Annexin-A1 Regulates Haematopoietic Stem Cell Mobilisation and Inflammatory Response Post Myocardial Infarction in Mice In Vivo. *Sci. Rep.* 7 (1), 16615.
- (22) Qin, C. X., Rosli, S., Deo, M., Cao, N., Walsh, J., Tate, M., Alexander, A. E., Donner, D., Horlock, D., Li, R., et al. (2019) Cardioprotective Actions of the Annexin-A1 N-Terminal Peptide, Ac2–26, Against Myocardial Infarction. *Front. Pharmacol.* 10 (APR), 269.
- (23) Ong, S.-B., Hernandez-Resendiz, S., Crespo-Avilan, G. E., Mukhametshina, R. T., Kwek, X.-Y., Cabrera-Fuentes, H. A., and Hausenloy, D. J. (2018) Inflammation Following Acute Myocardial Infarction: Multiple Players, Dynamic Roles, and Novel Therapeutic Opportunities. *Pharmacol. Ther.* 186, 73–87.
- (24) Lawrance, I. C., Wu, F., Leite, A. Z. A., Willis, J., West, G. A., Fiocchi, C., and Chakravarti, S. (2003) A Murine Model of Chronic Inflammation-Induced Intestinal Fibrosis down-Regulated by Antisense NF-Kappa B. *Gastroenterology* 125 (6), 1750–1761.
- (25) Deswal, A., Petersen, N. J., Feldman, A. M., Young, J. B., White, B. G., and Mann, D. L. (2001) Cytokines and Cytokine Receptors in Advanced Heart Failure: An Analysis of the Cytokine Database from the Vesnarinone Trial (VEST). *Circulation* 103 (16), 2055–2059.
- (26) Venereau, E., Ceriotti, C., and Bianchi, M. E. (2015) DAMPs from Cell Death to New Life. *Front. Immunol.* 6, 422.
- (27) Liu, L., Wang, Y., Cao, Z.-Y., Wang, M.-M., Liu, X.-M., Gao, T., Hu, Q.-K., Yuan, W.-J., and Lin, L. (2015) Up-Regulated TLR4 in Cardiomyocytes Exacerbates Heart Failure after Long-Term Myocardial Infarction. *J. Cell. Mol. Med.* 19 (12), 2728–2740.
- (28) Bratton, D. L., and Henson, P. M. (2011) Neutrophil Clearance: When the Party Is over, Clean-up Begins. *Trends Immunol.* 32 (8), 350–357.
- (29) Torre-Amione, G., Kapadia, S., Benedict, C., Oral, H., Young, J. B., and Mann, D. L. (1996) Proinflammatory Cytokine Levels in Patients with Depressed Left Ventricular Ejection Fraction: A Report from the Studies of Left Ventricular Dysfunction (SOLVD). *J. Am. Coll. Cardiol.* 27 (5), 1201–1206.
- (30) Van Linthout, S., Miteva, K., and Tschope, C. (2014) Crosstalk between Fibroblasts and Inflammatory Cells. *Cardiovasc. Res.* 102 (2), 258–269.
- (31) Kuwahara, F., Kai, H., Tokuda, K., Kai, M., Takeshita, A., Egashira, K., and Imaizumi, T. (2002) Transforming Growth Factor-Beta Function Blocking Prevents Myocardial Fibrosis and Diastolic Dysfunction in Pressure-Overloaded Rats. *Circulation* 106 (1), 130–135.
- (32) Dobaczewski, M., Chen, W., and Frangogiannis, N. G. (2011) Transforming Growth Factor (TGF)- β Signaling in Cardiac Remodeling. *J. Mol. Cell. Cardiol.* 51 (4), 600–606.
- (33) Gupta, S., Young, D., Maitra, R. K., Gupta, A., Popovic, Z. B., Yong, S. L., Mahajan, A., Wang, Q., and Sen, S. (2008) Prevention of Cardiac Hypertrophy and Heart Failure by Silencing of NF-KappaB. *J. Mol. Biol.* 375 (3), 637–649.
- (34) Huynh, M.-L. N., Fadok, V. A., and Henson, P. M. (2002) Phosphatidylserine-Dependent Ingestion of Apoptotic Cells Promotes TGF-Beta1 Secretion and the Resolution of Inflammation. *J. Clin. Invest.* 109 (1), 41–50.
- (35) Lawrence, T., Bebie, M., Liu, G. Y., Nizet, V., and Karin, M. (2005) IKKalpha Limits Macrophage NF-KappaB Activation and Contributes to the Resolution of Inflammation. *Nature* 434 (7037), 1138–1143.
- (36) Hirayama, D., Iida, T., and Nakase, H. (2018) The Phagocytic Function of Macrophage-Enforcing Innate Immunity and Tissue Homeostasis. *Int. J. Mol. Sci.* 19 (1), 92.
- (37) Tacke, F., Alvarez, D., Kaplan, T. J., Jakubzick, C., Spanbroek, R., Llodra, J., Garin, A., Liu, J., Mack, M., van Rooijen, N., et al. (2007) Monocyte Subsets Differentially Employ CCR2, CCR5, and CX3CR1 to Accumulate within Atherosclerotic Plaques. *J. Clin. Invest.* 117 (1), 185–194.
- (38) Tarique, A. A., Logan, J., Thomas, E., Holt, P. G., Sly, P. D., and Fantino, E. (2015) Phenotypic, Functional, and Plasticity Features of Classical and Alternatively Activated Human Macrophages. *Am. J. Respir. Cell Mol. Biol.* 53 (5), 676–688.
- (39) Das, A., Sinha, M., Datta, S., Abas, M., Chaffee, S., Sen, C. K., and Roy, S. (2015) Monocyte and Macrophage Plasticity in Tissue Repair and Regeneration. *Am. J. Pathol.* 185 (10), 2596–2606.
- (40) Zhu, W., Yu, J., Nie, Y., Shi, X., Liu, Y., Li, F., and Zhang, X. (2014) Disequilibrium of M1 and M2 Macrophages Correlates with the Development of Experimental Inflammatory Bowel Diseases. *Immunol. Invest.* 43 (7), 638–652.
- (41) Khanna, S., Biswas, S., Shang, Y., Collard, E., Azad, A., Kauh, C., Bhasker, V., Gordillo, G. M., Sen, C. K., and Roy, S. (2010) Macrophage Dysfunction Impairs Resolution of Inflammation in the Wounds of Diabetic Mice. *PLoS One* 5 (3), No. e9539.
- (42) Parathath, S., Grauer, L., Huang, L.-S., Sanson, M., Distel, E., Goldberg, I. J., and Fisher, E. A. (2011) Diabetes Adversely Affects Macrophages during Atherosclerotic Plaque Regression in Mice. *Diabetes* 60 (6), 1759–1769.
- (43) Rahman, K., Vengrenyuk, Y., Ramsey, S. A., Vila, N. R., Girgis, N. M., Liu, J., Gusarova, V., Gromada, J., Weinstock, A., Moore, K. J., et al. (2017) Inflammatory Ly6Chi Monocytes and Their Conversion to M2 Macrophages Drive Atherosclerosis Regression. *J. Clin. Invest.* 127 (8), 2904–2915.
- (44) Lavine, K. J., Epelman, S., Uchida, K., Weber, K. J., Nichols, C. G., Schilling, J. D., Ornitz, D. M., Randolph, G. J., and Mann, D. L. (2014) Distinct Macrophage Lineages Contribute to Disparate Patterns of Cardiac Recovery and Remodeling in the Neonatal and Adult Heart. *Proc. Natl. Acad. Sci. U. S. A.* 111 (45), 16029–16034.
- (45) Serhan, C. N., and Levy, B. D. (2018) Resolvins in Inflammation: Emergence of the pro-Resolving Superfamily of Mediators. *J. Clin. Invest.* 128 (7), 2657–2669.
- (46) Serhan, C. N. (2014) Pro-Resolving Lipid Mediators Are Leads for Resolution Physiology. *Nature* 510 (7503), 92–101.
- (47) Dalli, J., and Serhan, C. N. (2012) Specific Lipid Mediator Signatures of Human Phagocytes: Microparticles Stimulate Macro-

phage Efferocytosis and pro-Resolving Mediators. *Blood* 120 (15), No. e60.

(48) Billman, G. E., Kang, J. X., and Leaf, A. (1999) Prevention of Sudden Cardiac Death by Dietary Pure Omega-3 Polyunsaturated Fatty Acids in Dogs. *Circulation* 99 (18), 2452–2457.

(49) GISSI-Prevenzione Investigators (1999) Dietary Supplementation with N-3 Polyunsaturated Fatty Acids and Vitamin E after Myocardial Infarction: Results of the GISSI-Prevenzione Trial. *Lancet* 354 (9177), 447–455.

(50) Wang, J., Yu, J.-C., Kang, W.-M., and Ma, Z.-Q. (2012) Superiority of a Fish Oil-Enriched Emulsion to Medium-Chain Triacylglycerols/Long-Chain Triacylglycerols in Gastrointestinal Surgery Patients: A Randomized Clinical Trial. *Nutrition* 28 (6), 623–629.

(51) Mocellin, M. C., Camargo, C. Q., Nunes, E. A., Fiates, G. M. R., and Trindade, E. B. S. M. (2016) A Systematic Review and Meta-Analysis of the n-3 Polyunsaturated Fatty Acids Effects on Inflammatory Markers in Colorectal Cancer. *Clin. Nutr.* 35 (2), 359–369.

(52) Kiyabu, G. Y., Inoue, M., Saito, E., Abe, S. K., Sawada, N., Ishihara, J., Iwasaki, M., Yamaji, T., Shimazu, T., Sasazuki, S., et al. (2015) Fish, n - 3 Polyunsaturated Fatty Acids and n - 6 Polyunsaturated Fatty Acids Intake and Breast Cancer Risk: The Japan Public Health Center-Based Prospective Study. *Int. J. Cancer* 137 (12), 2915–2926.

(53) Barden, A., Mas, E., Croft, K. D., Phillips, M., and Mori, T. A. (2014) Short-Term n-3 Fatty Acid Supplementation but Not Aspirin Increases Plasma Proresolving Mediators of Inflammation. *J. Lipid Res.* 55 (11), 2401–2407.

(54) Wu, S.-H., Chen, X.-Q., Liu, B., Wu, H.-J., and Dong, L. (2013) Efficacy and Safety of 15(R/S)-Methyl-Lipoxin A(4) in Topical Treatment of Infantile Eczema. *Br. J. Dermatol.* 168 (1), 172–178.

(55) Fiore, S., Maddox, J. F., Perez, H. D., and Serhan, C. N. (1994) Identification of a Human CDNA Encoding a Functional High Affinity Lipoxin A4 Receptor. *J. Exp. Med.* 180 (1), 253–260.

(56) Claria, J., and Serhan, C. N. (1995) Aspirin Triggers Previously Undescribed Bioactive Eicosanoids by Human Endothelial Cell-Leukocyte Interactions. *Proc. Natl. Acad. Sci. U. S. A.* 92 (21), 9475–9479.

(57) Clish, C. B., O'Brien, J. A., Gronert, K., Stahl, G. L., Petasis, N. A., and Serhan, C. N. (1999) Local and Systemic Delivery of a Stable Aspirin-Triggered Lipoxin Prevents Neutrophil Recruitment in Vivo. *Proc. Natl. Acad. Sci. U. S. A.* 96 (14), 8247–8252.

(58) Serhan, C. N. (2005) Lipoxins and Aspirin-Triggered 15-Epi-Lipoxins Are the First Lipid Mediators of Endogenous Anti-Inflammation and Resolution. *Prostaglandins, Leukotrienes Essent. Fatty Acids* 73 (3–4), 141–162.

(59) Markworth, J. F., Maddipati, K. R., and Cameron-Smith, D. (2016) Emerging Roles of Pro-Resolving Lipid Mediators in Immunological and Adaptive Responses to Exercise-Induced Muscle Injury. *Exerc. Immunol. Rev.* 22, 110–134.

(60) McMahon, B., and Godson, C. (2004) Lipoxins: Endogenous Regulators of Inflammation. *Am. J. Physiol. Renal Physiol.* 286 (2), F189–201.

(61) Levy, B. D., Romano, M., Chapman, H. A., Reilly, J. J., Drazen, J., and Serhan, C. N. (1993) Human Alveolar Macrophages Have 15-Lipoxygenase and Generate 15(S)-Hydroxy-5,8,11-Cis-13-Trans-Eicosatetraenoic Acid and Lipoxins. *J. Clin. Invest.* 92 (3), 1572–1579.

(62) Papayianni, A., Serhan, C. N., Phillips, M. L., Rennke, H. G., and Brady, H. R. (1995) Transcellular Biosynthesis of Lipoxin A4 during Adhesion of Platelets and Neutrophils in Experimental Immune Complex Glomerulonephritis. *Kidney Int.* 47 (5), 1295–1302.

(63) Serhan, C. N. (1989) On the Relationship between Leukotriene and Lipoxin Production by Human Neutrophils: Evidence for Differential Metabolism of 15-HETE and 5-HETE. *Biochim. Biophys. Acta, Lipids Lipid Metab.* 1004 (2), 158–168.

(64) Parente, L., and Perretti, M. (2003) Advances in the Pathophysiology of Constitutive and Inducible Cyclooxygenases: Two Enzymes in the Spotlight. *Biochem. Pharmacol.* 65 (2), 153–159.

(65) Fiore, S., and Serhan, C. N. (1990) Formation of Lipoxins and Leukotrienes during Receptor-Mediated Interactions of Human Platelets and Recombinant Human Granulocyte/Macrophage Colony-Stimulating Factor-Primed Neutrophils. *J. Exp. Med.* 172 (5), 1451–1457.

(66) Recchiuti, A., Mattoscio, D., and Isopi, E. (2019) Roles, Actions, and Therapeutic Potential of Specialized Pro-Resolving Lipid Mediators for the Treatment of Inflammation in Cystic Fibrosis. *Front. Pharmacol.* 10 (APR), 252.

(67) Serhan, C. N., Maddox, J. F., Petasis, N. A., Akritopoulou-Zanze, I., Papayianni, A., Brady, H. R., Colgan, S. P., and Madara, J. L. (1995) Design of Lipoxin A4 Stable Analogs That Block Transmigration and Adhesion of Human Neutrophils. *Biochemistry* 34 (44), 14609–14615.

(68) Brennan, E. P., Mohan, M., McClelland, A., Tikellis, C., Ziemann, M., Kaspi, A., Gray, S. P., Pickering, R., Tan, S. M., Tasadaque Ali-Shah, S., et al. (2018) Lipoxins Regulate the Early Growth Response-1 Network and Reverse Diabetic Kidney Disease. *J. Am. Soc. Nephrol.* 29 (5), 1437–1448.

(69) Brennan, E. P., Mohan, M., McClelland, A., de Gaetano, M., Tikellis, C., Marai, M., Crean, D., Dai, A., Beuscart, O., Derouiche, S., et al. (2018) Lipoxins Protect Against Inflammation in Diabetes-Associated Atherosclerosis. *Diabetes* 67 (12), 2657–2667.

(70) Le, Y., Murphy, P. M., and Wang, J. M. (2002) Formyl-Peptide Receptors Revisited. *Trends Immunol.* 23 (11), 541–548.

(71) Bufe, B., Schumann, T., Kappl, R., Bogeski, I., Kummerow, C., Podgorska, M., Smola, S., Hoth, M., and Zufall, F. (2015) Recognition of Bacterial Signal Peptides by Mammalian Formyl Peptide Receptors: A New Mechanism for Sensing Pathogens. *J. Biol. Chem.* 290 (12), 7369–7387.

(72) Börgeson, E., Johnson, A. M. F., Lee, Y. S., Till, A., Syed, G. H., Ali-Shah, S. T., Guiry, P. J., Dalli, J., Colas, R. A., Serhan, C. N., et al. (2015) Lipoxin A4 Attenuates Obesity-Induced Adipose Inflammation and Associated Liver and Kidney Disease. *Cell Metab.* 22 (1), 125–137.

(73) Devos, T., Guillabert, A., D'Haene, N., Berton, A., De Nadai, P., Noel, S., Brait, M., Franssen, J.-D., Sozzani, S., Salmon, I., et al. (2009) Formyl Peptide Receptor-like 2 Is Expressed and Functional in Plasmacytoid Dendritic Cells, Tissue-Specific Macrophage Subpopulations, and Eosinophils. *J. Immunol.* 182 (8), 4974–4984.

(74) Qin, C. X., May, L. T., Li, R., Cao, N., Rosli, S., Deo, M., Alexander, A. E., Horlock, D., Bourke, J. E., Yang, Y. H., et al. (2017) Small-Molecule-Biased Formyl Peptide Receptor Agonist Compound 17b Protects against Myocardial Ischaemia-Reperfusion Injury in Mice. *Nat. Commun.* 8, 14232.

(75) Harada, M., Habata, Y., Hosoya, M., Nishi, K., Fujii, R., Kobayashi, M., and Hinuma, S. (2004) N-Formylated Humanin Activates Both Formyl Peptide Receptor-like 1 and 2. *Biochem. Biophys. Res. Commun.* 324 (1), 255–261.

(76) VanCompernelle, S. E., Clark, K. L., Rummel, K. A., and Todd, S. C. (2003) Expression and Function of Formyl Peptide Receptors on Human Fibroblast Cells. *J. Immunol.* 171 (4), 2050–2056.

(77) Migeotte, I., Communi, D., and Parmentier, M. (2006) Formyl Peptide Receptors: A Promiscuous Subfamily of G Protein-Coupled Receptors Controlling Immune Responses. *Cytokine Growth Factor Rev.* 17 (6), 501–519.

(78) He, H.-Q., and Ye, R. D. (2017) The Formyl Peptide Receptors: Diversity of Ligands and Mechanism for Recognition. *Molecules* 22 (3), 455.

(79) Qin, C., Buxton, K. D., Pepe, S., Cao, A. H., Venardos, K., Love, J. E., Kaye, D. M., Yang, Y. H., Morand, E. F., and Ritchie, R. H. (2013) Reperfusion-Induced Myocardial Dysfunction Is Prevented by Endogenous Annexin-A1 and Its N-Terminal-Derived Peptide Ac-ANX-A1(2–26). *Br. J. Pharmacol.* 168 (1), 238–252.

(80) Li, L., Chen, K., Xiang, Y., Yoshimura, T., Su, S., Zhu, J., Bian, X., and Wang, J. M. (2016) New Development in Studies of Formyl-

Peptide Receptors: Critical Roles in Host Defense. *J. Leukocyte Biol.* 99 (3), 425–435.

(81) Ye, R. D., Boulay, F., Wang, J. M., Dahlgren, C., Gerard, C., Parmentier, M., Serhan, C. N., and Murphy, P. M. (2009) International Union of Basic and Clinical Pharmacology. LXXIII. Nomenclature for the Formyl Peptide Receptor (FPR) Family. *Pharmacol. Rev.* 61 (2), 119–161.

(82) Bozinovski, S., Uddin, M., Vlahos, R., Thompson, M., McQualter, J. L., Merritt, A.-S., Wark, P. A. B., Hutchinson, A., Irving, L. B., Levy, B. D., et al. (2012) Serum Amyloid A Opposes Lipoxin A(4) to Mediate Glucocorticoid Refractory Lung Inflammation in Chronic Obstructive Pulmonary Disease. *Proc. Natl. Acad. Sci. U. S. A.* 109 (3), 935–940.

(83) Gao, J. L., Lee, E. J., and Murphy, P. M. (1999) Impaired Antibacterial Host Defense in Mice Lacking the N-Formylpeptide Receptor. *J. Exp. Med.* 189 (4), 657–662.

(84) Forsman, H., Winther, M., Gabl, M., Skovbakke, S. L., Boulay, F., Rabiet, M.-J., and Dahlgren, C. (2015) Structural Changes of the Ligand and of the Receptor Alters the Receptor Preference for Neutrophil Activating Peptides Starting with a Formylmethionyl Group. *Biochim. Biophys. Acta, Mol. Cell Res.* 1853 (1), 192–200.

(85) Wright, H. L., Moots, R. J., Bucknall, R. C., and Edwards, S. W. (2010) Neutrophil Function in Inflammation and Inflammatory Diseases. *Rheumatology (Oxford, U. K.)* 49 (9), 1618–1631.

(86) Kretschmer, D., Rautenberg, M., Linke, D., and Peschel, A. (2015) Peptide Length and Folding State Govern the Capacity of Staphylococcal β -Type Phenol-Soluble Modulins to Activate Human Formyl-Peptide Receptors 1 or 2. *J. Leukocyte Biol.* 97 (4), 689–697.

(87) Ramon, S., Bancos, S., Serhan, C. N., and Phipps, R. P. (2014) Lipoxin A₄ Modulates Adaptive Immunity by Decreasing Memory B-Cell Responses via an ALX/FPR2-Dependent Mechanism. *Eur. J. Immunol.* 44 (2), 357–369.

(88) Prevete, N., Liotti, F., Marone, G., Melillo, R. M., and de Paulis, A. (2015) Formyl Peptide Receptors at the Interface of Inflammation, Angiogenesis and Tumor Growth. *Pharmacol. Res.* 102, 184–191.

(89) Weiß, E., and Kretschmer, D. (2018) Formyl-Peptide Receptors in Infection, Inflammation, and Cancer. *Trends Immunol.* 39 (10), 815–829.

(90) Hodges, R. R., Li, D., Shatos, M. A., Serhan, C. N., and Dartt, D. A. (2016) Lipoxin A₄ Counter-Regulates Histamine-Stimulated Glycoconjugate Secretion in Conjunctival Goblet Cells. *Sci. Rep.* 6, 36124.

(91) Yao, L., Wang, C., Zhang, X., Peng, L., Liu, W., Zhang, X., Liu, Y., He, J., Jiang, C., Ai, D., et al. (2016) Hyperhomocysteinemia Activates the Aryl Hydrocarbon Receptor/CD36 Pathway to Promote Hepatic Steatosis in Mice. *Hepatology* 64 (1), 92–105.

(92) Zhang, L., Zhang, X., Wu, P., Li, H., Jin, S., Zhou, X., Li, Y., Ye, D., Chen, B., and Wan, J. (2008) BML-111, a Lipoxin Receptor Agonist, Modulates the Immune Response and Reduces the Severity of Collagen-Induced Arthritis. *Inflammation Res.* 57 (4), 157–162.

(93) Paul-Clark, M. J., Van Cao, T., Moradi-Bidhendi, N., Cooper, D., and Gilroy, D. W. (2004) 15-Epi-Lipoxin A₄-Mediated Induction of Nitric Oxide Explains How Aspirin Inhibits Acute Inflammation. *J. Exp. Med.* 200 (1), 69–78.

(94) Scalia, R., Gefen, J., Petasis, N. A., Serhan, C. N., and Lefer, A. M. (1997) Lipoxin A₄ Stable Analogs Inhibit Leukocyte Rolling and Adherence in the Rat Mesenteric Microvasculature: Role of P-Selectin. *Proc. Natl. Acad. Sci. U. S. A.* 94 (18), 9967–9972.

(95) Song, Y., Yang, Y., Cui, Y., Gao, J., Wang, K., and Cui, J. (2019) Lipoxin A₄Methyl Ester Reduces Early Brain Injury by Inhibition of the Nuclear Factor Kappa B (NF-KappaB)-Dependent Matrix Metalloproteinase 9 (MMP-9) Pathway in a Rat Model of Intracerebral Hemorrhage. *Med. Sci. Monit.* 25, 1838–1847.

(96) Brennan, E., Wang, B., McClelland, A., Mohan, M., Marai, M., Beuscart, O., Derouiche, S., Gray, S., Pickering, R., Tikellis, C., et al. (2017) Protective Effect of Let-7 miRNA Family in Regulating Inflammation in Diabetes-Associated Atherosclerosis. *Diabetes* 66 (8), 2266–2277.

(97) Jozsef, L., Zouki, C., Petasis, N. A., Serhan, C. N., and Filep, J. G. (2002) Lipoxin A₄ and Aspirin-Triggered 15-Epi-Lipoxin A₄ Inhibit Peroxynitrite Formation, NF-Kappa B and AP-1 Activation, and IL-8 Gene Expression in Human Leukocytes. *Proc. Natl. Acad. Sci. U. S. A.* 99 (20), 13266–13271.

(98) Levy, B. D., Clish, C. B., Schmidt, B., Gronert, K., and Serhan, C. N. (2001) Lipid Mediator Class Switching during Acute Inflammation: Signals in Resolution. *Nat. Immunol.* 2 (7), 612–619.

(99) Prieto, P., Cuenca, J., Través, P. G., Fernández-Velasco, M., Martín-Sanz, P., and Boscá, L. (2010) Lipoxin A₄ Impairment of Apoptotic Signaling in Macrophages: Implication of the PI3K/Akt and the ERK/Nrf-2 Defense Pathways. *Cell Death Differ.* 17 (7), 1179–1188.

(100) Jadcherla, S. R. (2002) Inflammation Inhibits Muscarinic Signaling in Vivo Canine Colonic Circular Smooth Muscle Cells. *Pediatr. Res.* 52 (5), 756–762.

(101) Maderna, P., and Godson, C. (2009) Lipoxins: Resolutionary Road. *Br. J. Pharmacol.* 158 (4), 947–959.

(102) O'Sullivan, S. (1999) On the Role of PGD₂Metabolites as Markers of Mast Cell Activation in Asthma. *Acta Physiol. Scand. Suppl.* 644, 1–74.

(103) Serhan, C. N., and Chiang, N. (2013) Resolution Phase Lipid Mediators of Inflammation: Agonists of Resolution. *Curr. Opin. Pharmacol.* 13 (4), 632–640.

(104) Kieran, N. E., Doran, P. P., Connolly, S. B., Greenan, M.-C., Higgins, D. F., Leonard, M., Godson, C., Taylor, C. T., Henger, A., Kretzler, M., et al. (2003) Modification of the Transcriptomic Response to Renal Ischemia/Reperfusion Injury by Lipoxin Analog. *Kidney Int.* 64 (2), 480–492.

(105) Leonard, M. O., Hannan, K., Burne, M. J., Lappin, D. W. P., Doran, P., Coleman, P., Stenson, C., Taylor, C. T., Daniels, F., Godson, C., et al. (2002) 15-Epi-16-(Para-Fluorophenoxy)-Lipoxin A(4)-Methyl Ester, a Synthetic Analogue of 15-Epi-Lipoxin A(4), Is Protective in Experimental Ischemic Acute Renal Failure. *J. Am. Soc. Nephrol.* 13 (6), 1657–1662.

(106) Ohkawara, T., Mitsuyama, K., Takeda, H., Asaka, M., Fujiyama, Y., and Nishihira, J. (2008) Lack of Macrophage Migration Inhibitory Factor Suppresses Innate Immune Response in Murine Dextran Sulfate Sodium-Induced Colitis. *Scand. J. Gastroenterol.* 43 (12), 1497–1504.

(107) Janakiram, N. B., Mohammed, A., and Rao, C. V. (2011) Role of Lipoxins, Resolvins, and Other Bioactive Lipids in Colon and Pancreatic Cancer. *Cancer Metastasis Rev.* 30 (3–4), 507–523.

(108) Schottelius, A. J., Giesen, C., Asadullah, K., Fierro, I. M., Colgan, S. P., Bauman, J., Guilford, W., Perez, H. D., and Parkinson, J. F. (2002) An Aspirin-Triggered Lipoxin A₄ Stable Analog Displays a Unique Topical Anti-Inflammatory Profile. *J. Immunol.* 169 (12), 7063–7070.

(109) Chiang, N., Arita, M., and Serhan, C. N. (2005) Anti-Inflammatory Circuitry: Lipoxin, Aspirin-Triggered Lipoxins and Their Receptor ALX. *Prostaglandins, Leukotrienes Essent. Fatty Acids* 73 (3–4), 163–177.

(110) Yu, D., Xu, Z., Yin, X., Zheng, F., Lin, X., Pan, Q., and Li, H. (2015) Inverse Relationship between Serum Lipoxin A₄ Level and the Risk of Metabolic Syndrome in a Middle-Aged Chinese Population. *PLoS One* 10 (11), No. e0142848.

(111) Das, U. N. (1995) Essential Fatty Acid Metabolism in Patients with Essential Hypertension, Diabetes Mellitus and Coronary Heart Disease. *Prostaglandins, Leukotrienes Essent. Fatty Acids* 52 (6), 387–391.

(112) Petri, M. H., Laguna-Fernandez, A., Tseng, C.-N., Hedin, U., Perretti, M., and Back, M. (2015) Aspirin-Triggered 15-Epi-Lipoxin A(4) Signals through FPR2/ALX in Vascular Smooth Muscle Cells and Protects against Intimal Hyperplasia after Carotid Ligation. *Int. J. Cardiol.* 370–372, DOI: 10.1016/j.ijcard.2014.11.010.

(113) Sette, L. H. B. C., and Almeida Lopes, E. P. de. (2014) Liver Enzymes Serum Levels in Patients with Chronic Kidney Disease on Hemodialysis: A Comprehensive Review. *Clinics (Sao Paulo)*. 69 (4), 271–278.

- (114) Das, U. N. (2016) Inflammatory Bowel Disease as a Disorder of an Imbalance between Pro- and Anti-Inflammatory Molecules and Deficiency of Resolution Bioactive Lipids. *Lipids Health Dis.* 15, 11.
- (115) Pouliot, M., Clish, C. B., Petasis, N. A., Van Dyke, T. E., and Serhan, C. N. (2000) Lipoxin A(4) Analogues Inhibit Leukocyte Recruitment to *Porphyromonas Gingivalis*: A Role for Cyclooxygenase-2 and Lipoxins in Periodontal Disease. *Biochemistry* 39 (16), 4761–4768.
- (116) Hashimoto, A., Hayashi, I., Murakami, Y., Sato, Y., Kitasato, H., Matsushita, R., Iizuka, N., Urabe, K., Itoman, M., Hirohata, S., et al. (2007) Antiinflammatory Mediator Lipoxin A4 and Its Receptor in Synovitis of Patients with Rheumatoid Arthritis. *J. Rheumatol.* 34 (11), 2144–2153.
- (117) Ndisang, J. F., and Rastogi, S. (2013) Cardiometabolic Diseases and Related Complications: Current Status and Future Perspective. *BioMed Res. Int.* 2013, 467682.
- (118) Diabetes Canada Clinical Practice Guidelines Expert Committee; Punthakee, Z., Goldenberg, R., and Katz, P. (2018) Definition, Classification and Diagnosis of Diabetes, Prediabetes and Metabolic Syndrome. *Can. J. Diabetes* 42 (Suppl 1), S10–S15.
- (119) L'Heveder, R., and Nolan, T. (2013) International Diabetes Federation. *Diabetes Res. Clin. Pract.* 101 (3), 349–351.
- (120) Huynh, K., Bernardo, B. C., McMullen, J. R., and Ritchie, R. H. (2014) Diabetic Cardiomyopathy: Mechanisms and New Treatment Strategies Targeting Antioxidant Signaling Pathways. *Pharmacol. Ther.* 142 (3), 375–415.
- (121) Papatheodorou, K., Banach, M., Bekiari, E., Rizzo, M., and Edmonds, M. (2018) Complications of Diabetes 2017. *J. Diabetes Res.* 2018 (1), 3086167.
- (122) Niewczasz, M. A., Pavkov, M. E., Skupien, J., Smiles, A., Md Dom, Z. I., Wilson, J. M., Park, J., Nair, V., Schlafly, A., Saulnier, P.-J., et al. (2019) A Signature of Circulating Inflammatory Proteins and Development of End-Stage Renal Disease in Diabetes. *Nat. Med.* 25 (5), 805–813.
- (123) Persson, F., Frimodt-Møller, M., and Rossing, P. (2019) Inflammation Leads the Way on the ROADMAP to Diabetic Kidney Disease. *Kidney Int. Rep.*, 1362–1365, DOI: 10.1016/j.ekir.2019.08.009.
- (124) Anderberg, R. J., Meek, R. L., Hudkins, K. L., Cooney, S. K., Alpers, C. E., Leboeuf, R. C., and Tuttle, K. R. (2015) Serum Amyloid A and Inflammation in Diabetic Kidney Disease and Podocytes. *Laboratory Investigation; A Journal of Technical Methods and Pathology*, 697 DOI: 10.1038/labinvest.2015.38.
- (125) Börgeson, E., and Godson, C. (2012) Resolution of Inflammation: Therapeutic Potential of pro-Resolving Lipids in Type 2 Diabetes Mellitus and Associated Renal Complications. *Front. Immunol.* 3 (17), 318.
- (126) Guilford, W. J., Bauman, J. G., Skuballa, W., Bauer, S., Wei, G. P., Davey, D., Schaefer, C., Mallari, C., Terkelsen, J., Tseng, J.-L., et al. (2004) Novel 3-Oxa Lipoxin A4 Analogues with Enhanced Chemical and Metabolic Stability Have Anti-Inflammatory Activity in Vivo. *J. Med. Chem.* 47 (8), 2157–2165.
- (127) Levy, B. D., Lukacs, N. W., Berlin, A. A., Schmidt, B., Guilford, W. J., Serhan, C. N., and Parkinson, J. F. (2007) Lipoxin A₄ Stable Analogs Reduce Allergic Airway Responses via Mechanisms Distinct from CysLT₁ Receptor Antagonism. *FASEB J.* 21 (14), 3877.
- (128) Fiorucci, S., Wallace, J. L., Mencarelli, A., Distrutti, E., Rizzo, G., Farneti, S., Morelli, A., Tseng, J.-L., Suramanyam, B., Guilford, W. J., et al. (2004) A Beta-Oxidation-Resistant Lipoxin A4 Analog Treats Hapten-Induced Colitis by Attenuating Inflammation and Immune Dysfunction. *Proc. Natl. Acad. Sci. U. S. A.* 101 (44), 15736–15741.
- (129) O'Sullivan, T. P., Vallin, K. S. A., Shah, S. T. A., Fakhry, J., Maderna, P., Scannell, M., Sampaio, A. L. F., Perretti, M., Godson, C., and Guiry, P. J. (2007) Aromatic Lipoxin A4 and Lipoxin B4 Analogues Display Potent Biological Activities. *J. Med. Chem.* 50 (24), 5894–5902.
- (130) Sun, Y.-P., Tjonahen, E., Keledjian, R., Zhu, M., Yang, R., Recchiuti, A., Pillai, P. S., Petasis, N. A., and Serhan, C. N. (2009) Anti-Inflammatory and pro-Resolving Properties of Benzo-Lipoxin A(4) Analogs. *Prostaglandins, Leukotrienes Essent. Fatty Acids* 81 (5–6), 357–366.
- (131) Börgeson, E., Docherty, N. G., Murphy, M., Rodgers, K., Ryan, A., O'Sullivan, T. P., Guiry, P. J., Goldschmeding, R., Higgins, D. F., and Godson, C. (2011) Lipoxin A₄ and Benzo-Lipoxin A₄ Attenuate Experimental Renal Fibrosis. *FASEB J.* 25 (9), 2967–2979.
- (132) Brennan, E. P., Nolan, K. A., Borgeson, E., Gough, O. S., McEvoy, C. M., Docherty, N. G., Higgins, D. F., Murphy, M., Sadlier, D. M., Ali-Shah, S. T., et al. (2013) Lipoxins Attenuate Renal Fibrosis by Inducing Let-7c and Suppressing TGFβ₁. *J. Am. Soc. Nephrol.* 24 (4), 627–637.
- (133) de Gaetano, M., Butler, E., Gahan, K., Zanetti, A., Marai, M., Chen, J., Cacace, A., Hams, E., Maingot, C., McLoughlin, A., et al. (2019) Asymmetric Synthesis and Biological Evaluation of Imidazole- and Oxazole-Containing Synthetic Lipoxin A4 Mimetics (SLXms). *Eur. J. Med. Chem.* 162, 80–108.
- (134) Barnes, P. J. (2006) How Corticosteroids Control Inflammation: Quintiles Prize Lecture 2005. *Br. J. Pharmacol.* 148 (3), 245–254.
- (135) Monaco, C., Nanchahal, J., Taylor, P., and Feldmann, M. (2015) Anti-TNF Therapy: Past, Present and Future. *Int. Immunol.* 27 (1), 55–62.
- (136) Terkeltaub, R. A., Furst, D. E., Bennett, K., Kook, K. A., Crockett, R. S., and Davis, M. W. (2010) High versus Low Dosing of Oral Colchicine for Early Acute Gout Flare: Twenty-Four-Hour Outcome of the First Multicenter, Randomized, Double-Blind, Placebo-Controlled, Parallel-Group, Dose-Comparison Colchicine Study. *Arthritis Rheum.* 62 (4), 1060–1068.
- (137) Perretti, M., Leroy, X., Bland, E. J., and Montero-Melendez, T. (2015) Resolution Pharmacology: Opportunities for Therapeutic Innovation in Inflammation. *Trends Pharmacol. Sci.* 36 (11), 737–755.
- (138) Ba, F., Zhou, X., Zhang, Y., Wu, C., Xu, S., Wu, L., Li, J., Yin, Y., and Gu, X. (2019) Lipoxin A4 Ameliorates Alveolar Fluid Clearance Disturbance in Lipopolysaccharide-Induced Lung Injury via Aquaporin 5 and MAPK Signaling Pathway. *J. Thorac. Dis.* 11 (8), 3599–3608.
- (139) Cai, B., Kasikara, C., Doran, A. C., Ramakrishnan, R., Birge, R. B., and Tabas, I. (2018) MerTK Signaling in Macrophages Promotes the Synthesis of Inflammation Resolution Mediators by Suppressing CaMKII Activity. *Sci. Signaling* 11 (549), eaar3721.
- (140) Chen, Y., Zheng, Y., Xin, L., Zhong, S., Liu, A., Lai, W., Liu, L., Lin, C., Liao, C., Zeng, J., et al. (2019) 15-Epi-Lipoxin A4 Inhibits TNF-Alpha-Induced Tissue Factor Expression via the PI3K/AKT/NF-KappaB Axis in Human Umbilical Vein Endothelial Cells. *Biomed. Pharmacother.* 117, 109099.
- (141) Petasis, N. A., Akritopoulou-Zanze, I., Fokin, V. V., Bernasconi, G., Keledjian, R., Yang, R., Uddin, J., Nagulapalli, K. C., and Serhan, C. N. (2005) Design, Synthesis and Bioactions of Novel Stable Mimetics of Lipoxins and Aspirin-Triggered Lipoxins. *Prostaglandins, Leukotrienes Essent. Fatty Acids* 73 (3–4), 301–321.