Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Evolutionary Medicine of Retroviruses in the Human Genome

Yukako Katsura, PhD and Satoshi Asai, PhD, MD

Division of Pharmacology, Department of Biomedical Sciences,
Nihon University School of Medicine, Tokyo, Japan

ABSTRACT

Humans are infected with many viruses, and the immune system mostly removes viruses and the infected cells. However, certain viruses have entered the human genome. Of the human genome, ~45% is composed of transposable elements (long interspersed nuclear elements [LINEs], short interspersed nuclear elements [SINEs] and transposons) and 5-8% is derived from viral sequences with similarity to infectious retroviruses. If integration of retrovirus occurs in a germline, the integrated viral sequences are heritable. Accumulation of viral sequences has created the current human genome. This article summarizes recent studies of retroviruses in humans and bridges clinical fields and evolutionary genetics. First, we report the repertoires of human-infective retroviruses. Second, we review endogenous retroviruses in the human genome and diseases associated with endogenous retroviruses. Third, we discuss the biological functions of endogenous retroviruses and propose the concept of accelerated human evolution via viruses. Finally, we present perspectives of virology in the field of evolutionary medicine.

Key Indexing Terms: Retrovirus; Human endogenous retrovirus; Disease; Human genome; Evolutionary medicine. [Am J Med Sci 2019;358(6):384–388.]

INTRODUCTION

Currently there are 5,630 confirmed virus species on the earth,¹ and Anthony et al estimated that at least 320,000 undiscovered viruses infect mammalian hosts.² RNA viruses are a major threat to human health, and severe acute respiratory syndrome (SARS) coronavirus, Ebola virus and Middle East respiratory syndrome (MERS) coronavirus are recently known to cause pandemics.³ Of the discovered virus species, 214 viruses are known human-infective RNA viruses and 9 viruses are retroviruses (Table 1).³ The human-infective retroviruses are human T-lymphotropic viruses (HTLV), human immunodeficiency viruses (HIV) and simian foamy viruses (SFV), and consist of 3 genera (Deltaretrovirus, Lentivirus and Spumavirus, respectively; Table 1). The range of hosts of retroviruses is narrow, and HIV can be acquired only from humans although HTLV and SFV are from suspected zoonotic origin and intraspecies transmission to humans after first infection in nonhuman primates.^{3,4} Of the human genome, 5-8% are endogenous retroviruses derived from viral sequences with similarity to the infectious retroviruses. Understanding the endogenous retroviruses is clinically important because the accumulation of viral sequences has created the current human genome, and it can cause diseases associated with endogenous retroviruses.

HUMAN-INFECTIVE RETROVIRUSES

Retroviruses consist of a dimer of single-stranded positive sense RNA enclosed in a capsid of a lipid bilayer

envelope and contain a reverse transcriptase (RT) enzyme.⁵ The retrovirus genome is composed of 3 genes (*gag*, *pol* and *env*) enclosed between 2 long terminal repeats: *pol* (polymerase) has RT and integrase function, *gag* (group antigens) is a polyprotein with processed matrix and core proteins, and *env* encodes envelope proteins (Figure 1A).⁵

Primate T-lymphotropic viruses (PTLVs) are composed of 3 distinct groups (PTLV-1, -2 and -3) and are also called HTLVs in the case of infections in humans. PTLVs are known to cause adult T-cell lymphoma/leukemia, and inflammation due to T lymphocytes in several tissues presents as bronchitis, uveitis and demyelinating diseases called HTLV-1 associated myelopathy/tropical spastic paraparesis. PTLV-1 is the first retrovirus discovered from T-cell lymphoma, and PTLV-2 was found originally in a patient with hairy cell leukemia.^{6,7} Lately, PTLV-3 was discovered in Central Africa,⁸ and Wolfe et al suggested that hunting and eating bushmeat is one of the transmission routes of PTLV beyond species. PTLVs are transmissible via sexual and maternal contact.⁹

HIV destroys lymphoid CD4 T cells by pyroptosis with caspase-1 activation and causes acquired immunodeficiency syndrome.^{10,11} HIV-1 and -2 are classified into 4 groups (M-P) and 8 groups (A-H), respectively, and are further finely grouped into subtypes whose configuration changes frequently.¹² The origin of HIV-1 and -2 is believed to be independent transmission from different primates, since the HIV-1 and -2 sequences are similar to those of chimpanzee simian immunodeficiency virus (SIV) and Old World monkey SIV, respectively.¹³ HIVs are transmissible

TABLE 1. Human-infective retroviruses (Retroviridae species).

Species	Genus	References
PTLV-1	Deltaretrovirus	45
PTLV-2	Deltaretrovirus	46
PTLV-3	Deltaretrovirus	47
HIV-1	Lentivirus	48
HIV-2	Lentivirus	49
SIV	Lentivirus	50
SFVagm	Spumavirus	51
SFVmac	Spumavirus	52
SFV	Spumavirus	53

The table shows human-infective retroviruses: Primate T-lymphotropic virus (PTLV), human immunodeficiency virus (HIV), simian immunodeficiency virus (SIV), African green monkey simian foamy virus (SFVagm), macaque simian foamy virus (SFVmac) and simian foamy virus (SFV).

via direct contact by sexual, iatrogenic, broken skin and maternal routes, and SIV can be transmitted to humans from simian blood cells through broken skin.¹³⁻¹⁵

The association of SFV with specific diseases has not been clarified, although the prevalence of SFV in humans has been well studied.¹⁶ After infection by SFV, it has been observed that foamy degeneration and vacuolization occurs in the infected cells, with formation of numerous cytoplasmic vacuoles, and glycoproteins produced are expressed at the cell surface and result in fusion of cells.¹⁶ SFVs are known to be transmittable among simians only via direct contact with broken skin.¹⁷⁻¹⁹

ENDOGENOUS RETROVIRUSES IN HUMAN GENOME

The reverse transcribed retroviral genome can integrate into a host genome, and the integrated genome is

heritable if the retrovirus infects a germ line of the host. The integrated retroviral sequence, which is 7-11 kb, is called a provirus.⁵ Of the human genome, 5-8% is believed to comprise proviruses with sequence similarity to genes and fragments of retroviruses, and the provirus sequences are also called human endogenous retroviruses (HERVs).²⁰ Some HERVs still have open reading frames with the possibility of protein expression. There have been 3,173 HERV sequences identified from the human genome, and 39 canonical types of HERVs are categorized as classes I, II and III on the basis of sequences similar to different genera of infectious retroviruses (Gammaretrovirus/Epsilonretrovirus, Betaretrovirus and Spumaretrovirus, respectively).^{5,21-22}

It is believed that HERVs are associated with physiological functions and certain diseases based on model animal studies, but the role of HERVs is still under debate.⁵ Multiple sclerosis (MS) is caused by the destruction of myelin and oligodendrocytes, leading to axonal disruption in the brain and spinal cord. In MS patients, it has been reported that expression or abnormal representation of HERV-H, -K and -W and a polymorphism in HERV-Fc1 occurs.²³ Secretion of env proteins coded by HERV-W in oligodendroglial precursor cells reduces oligodendroglial differentiation via activation of Toll-like receptor, and blocking their differentiation results in demyelinated and degenerating axons.²⁴ HERV env proteins are related to complex pathological disorders and can be one of the targets for therapeutic approach.²⁵ A monoclonal antibody against a HERV-W env protein is under clinical trial as a therapeutic approach for MS.^{26,27}

Upregulation of HERV-W was also reported in blood cells and peripheral nerve lesions of chronic inflammatory demyelinating polyradiculoneuropathy patients, and

FIGURE 1. Structure of a retrovirus genome and HERV-K. A, The retrovirus genome has at least 3 genes (*gag*, *pol* and *env*) enclosed between 2 long terminal repeats. B, HERV-K env genes originate 2 proteins (np9 and rec) with a different length, and np9 proteins present 292bp deletion.

chronic inflammatory demyelinating polyradiculoneuropathy might be caused by Toll-like receptor-mediated effects of env proteins on primary Schwann cells.²⁸ In sporadic amyotrophic lateral sclerosis patients, env proteins of HERV-K are specifically expressed in cortical and spinal neurons and can cause cellular degeneration.²⁹ RT proteins encoded by the *pol* gene of HERV-K have been detected in brain and blood tissues, but transmission of amyotrophic lateral sclerosis has not yet been demonstrated.^{23,30}

HERVs are believed to be pathogenic in autoimmune rheumatic diseases and cancers, with evidence of increased expression of HERVs at the protein and RNA levels. Increased expression of HERVs was reported in patients with rheumatoid arthritis, systemic lupus erythematosus, juvenile idiopathic arthritis and Sjogren's syndrome.³¹ Nakkuntod et al reported hypomethylation of HERV-E and -K in systemic lupus erythematosus patients, and a lower methylation level can allow higher expression of HERV genes.³² HERV activation has also been reported in breast cancer, lymphoma, melanoma, ovarian cancer and prostate cancer, and the expression of env proteins can be involved in tumorigenesis via inducing cell-cell fusion.³³ In addition, it is known that sequences derived from HERVs can be a trigger of cancers if partial sequences of HERVs or HERV itself translocates to regulatory regions of oncogenes.³³ Tomlins et al reported that a translocation of HERV-K upstream of ETS translocation variant 1 caused cancerogenesis via enhanced expression of ETS translocation variant 1.³⁴ HERV-K env genes originate 2 proteins (np9 and rec) with a different length by alternative splicing, and np9 proteins present 292bp deletion but rec is not (Figure 1B).²⁵ It has been reported that rec and np9 contribute to tumorigenesis.²⁵ Viral infections induced HERVs transactivation, and the HERVs transactivation causes enhancement of several signal transductions and transcription factors.³⁵ HERV-K transactivated by Kaposi's sarcoma-associated herpes virus is suggested the involvement of tumorigenesis in Kaposi's sarcoma.³⁶

EVOLUTIONARY MEDICINE OF ENDOGENOUS RETROVIRUSES

The endogenous retroviruses remaining in the genome should be neutral or advantageous. If all endogenous retroviruses were harmful, all of the retroviruses would have been excluded from the genome during evolution leading to humans. HERV-K is a case of beneficial endogenous retroviruses and has been integrated stepwise during primate evolution.³⁷ It is known that HIV-1 infection stimulates a T cell response to HERV-K antigens because of protein similarity.³⁸ Monde et al showed that *gag* proteins encoded by HERV-K changed the size and morphology of HIV particles, and these changes caused significant diminishing of the release efficiency and infectivity of HIV viruses.³⁹

Vargiu et al estimated that HERVs diverged in the host genome from 6 to 100 million years ago, and this means the integration of HERVs occurred from after the divergence of Eutheria to the divergence between chimpanzees and humans.²² Syncytin-1 and -2 belong to the HERV-W and -FRD families, respectively, and both of them express env proteins with fusogenic activity and are involved in fusion of trophoblast cells.⁴⁰ The syncytin-1 and -2 sequences are believed to have integrated 12-80 million years ago, and at the same time these sequences obtained mammalian-specific placental function.⁴¹ As in this case, HERVs acquired essential functions that can be evolutionarily preserved.

Retroviral sequences can accelerate the evolution of host genomes. In the host genome, the integrated retroviral elements can be a promotor or enhancer and provide alternative and aberrant sites for splicing of transcripts.⁵ Endogenous retroviruses can enhance recombination and rearrangement of the host genome via long terminal repeats, while crossover between HERV-I loci on the Y chromosome is a cause of male infertility due to deletion of an azoospermia factor-a region.⁴²

CONCLUSIONS

In this review, we focused on known human-infective retroviruses and endogenous retroviruses in the human genome. It was previously believed that only retroviruses can integrate into the host genome, but Horie et al showed that sequences derived from nonretroviral RNA exist in the human genome. Bornaviruses, negative sense single stranded RNA viruses, encode 6 genes (M, X, P, N, G and L), and 2 proteins with high similarity to N genes, called endogenous borna-like N -1 and -2, have been found.⁴³ It is believed that N genes of bornaviruses integrated by RT activation of long interspersed nuclear element, and this mechanism can enhance the integration of any viruses without dependence on retroviruses.⁴³ The mechanism is the same as a system of making a processed gene, which was the evolutionary force to increase a member of a gene family in the human evolution.⁴⁴ If the activation of long interspersed nuclear element enhances the integration of any viruses, there should be many more known and unknown viruses in the human genome. Whole genome sequencing of viruses and infected hosts will help to discover new viral sequences in the human genome, and the number of viral sequences in the genome could be higher than the current estimation based on the sequence similarity of only retroviruses (5-8%). Most viruses can infect specific tissues or cells in the human body, and it is not clear how a virus can recognize specific cells and invade cells. If there is affinity between the virus and cells, the virus might tend to get into the genome of specific cells with high affinity. If there is a niche in the localization of the virus, the niche might allow an entry route of the virus and thereby determine integration.

AUTHOR CONTRIBUTIONS

Y.K. and S.A. conceived and designed the analysis, and Y.K. wrote the paper.

REFERENCES

1. International Committee on Taxonomy of Viruses. <https://talk.ictvonline.org/taxonomy/> [Accessed December 20, 2018].
2. Anthony SJ, Epstein JH, Murray KA, et al. A strategy to estimate unknown viral diversity in mammals. *mBio*. 2013;4: e00598-13.
3. Woolhouse MEJ, Brierley L. Epidemiological characteristics of human-infective RNA viruses. *Sci Data*. 2018;5: 180017.
4. Wolfe ND, Dunavan CP, Diamond J. Origins of major human infectious diseases. *Nature*. 2007;447:279–283.
5. Jern P, Coffin JM. Effects of retroviruses on host genome function. *Annu Rev Genet*. 2008;42:709–732.
6. Poesz BJ, Ruscetti FW, Gazdar AF, et al. Detection and isolation of type C retrovirus particles from fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma. *Proc Natl Acad Sci USA*. 1980;77:7415–7419.
7. Kalyanaraman VS, Sarngadharan MG, Robert-Guroff M, et al. A new subtype of human T-cell leukemia virus (HTLV-II) associated with a T-cell variant of hairy cell leukemia. *Science*. 1982;218:571–573.
8. Calattini S, Chevalier SA, Duprez R, et al. Discovery of a new human T-cell lymphotropic virus (HTLV-3) in Central Africa. *Retrovirology*. 2005;2:30.
9. Wolfe ND, Daszak P, Kilpatrick AM, et al. Bushmeat hunting, deforestation, and prediction of zoonotic disease. *Emerg Infect Dis*. 2005;11:1822–1827.
10. Doitsh G, Galloway NL, Geng X, et al. Cell death by pyroptosis drives CD4 T-cell depletion in HIV-1 infection. *Nature*. 2014;505:509–514.
11. Monroe KM, Yang Z, Johnson JR, et al. IFI16 DNA sensor is required for death of lymphoid CD4 T cells abortively infected with HIV. *Science*. 2014;343:428–432.
12. Sharp PM, Bailes E, Chaudhuri RR, et al. The origins of acquired immune deficiency syndrome viruses: where and when? *Philos Trans R Soc Lond B Biol Sci*. 2001;356:867–876.
13. van't Wout AB, Kootstra NA, Mulder-Kampinga GA, et al. Macrophage-tropic variants initiate human immunodeficiency virus type 1 infection after sexual, parenteral, and vertical transmission. *J Clin Invest*. 1994;94:2060–2067.
14. Visseaux B, Damond F, Matheron S, et al. HIV-2 molecular epidemiology. *Infect Genet Evol*. 2016;46:233–240.
15. Fouchet D1, Verrier D, Ngoubangoye B, et al. Natural simian immunodeficiency virus transmission in mandrills: a family affair? *Proc Biol Sci*. 2012;279:3426–3435.
16. Meiering CD, Linial ML. Historical perspective of foamy virus epidemiology and infection. *Clin Microbiol*. 2001;14:165–176.
17. Schweizer M, Schleier H, Pietrek M, et al. Genetic stability of foamy viruses: long-term study in an African green monkey population. *J Virol*. 1999;73:9256–9265.
18. Calattini S, Wanert F, Thierry B, et al. Modes of transmission and genetic diversity of foamy viruses in a Macaca tonkeana colony. *Retrovirology*. 2006;3:23.
19. Soliven K, Wang X, Small CT, et al. Simian foamy virus infection of rhesus macaques in Bangladesh: relationship of latent proviruses and transcriptionally active viruses. *J Virol*. 2013;87:13628–13639.
20. Belshaw R, Pereira V, Katourakis A, et al. Long-term reinfection of the human genome by endogenous retroviruses. *Proc Natl Acad Sci USA*. 2004;101:4894–4899.
21. Griffiths DJ. Endogenous retroviruses in the human genome sequence. *Genome Biol*. 2001;2:1017.1–1017.5. reviews 1017.
22. Vargiu L, Rodriguez-Tomé P, Sperber GO, et al. Classification and characterization of human endogenous retroviruses; mosaic forms are common. *Retrovirology*. 2016;13:7.
23. Kürty P, Nath A, Créange A, et al. Human endogenous retroviruses in neurological diseases. *Trends Mol Med*. 2018;24:379–394.
24. Kremer D, Schichel T, Förster M, et al. Human endogenous retrovirus type W envelope protein inhibits oligodendroglial precursor cell differentiation. *Ann Neurol*. 2013;74:721–732.
25. Grandi N, Tramontano E. HERV envelope proteins: physiological role and pathogenic potential in cancer and autoimmunity. *Front Microbiol*. 2018;9:462.
26. Diebold M, Derfuss T. The monoclonal antibody GNBAC1: targeting human endogenous retroviruses in multiple sclerosis. *Ther Adv Neurol Disord*. 2019;12: 1756286419833574.
27. Niegowska M, Wajda-Cuszlag M, Stepień-Ptak G, et al. Anti-W_{ENV} antibodies are correlated with seroreactivity against Mycobacterium Avium subsp. paratuberculosis in children and youths at T1D risk. *Sci Rep*. 2019;9:6282.
28. Faucard R, Alexandra Madeira A, Gehin N, et al. Human endogenous retrovirus and neuroinflammation in chronic inflammatory demyelinating polyradiculoneuropathy. *EBioMedicine*. 2016;6:190–198.
29. Li W, Lee MH, Henderson L, et al. Human endogenous retrovirus-K contributes to motor neuron disease. *Sci Transl Med*. 2015;7: 307ra153.
30. Douville RN, Liu J, Rothstein J, et al. Identification of active loci of a human endogenous retrovirus in neurons of patients with amyotrophic lateral sclerosis. *Ann Neurol*. 2011;69:141–151.
31. Tugnet N, Rylance P, Roden D, et al. Human endogenous retroviruses (HERVs) and autoimmune rheumatic disease: is there a link? *Open Rheumatol J*. 2013;7:13–21.
32. Nakkuntod J, Sukkapan P, Avihingsanon Y, et al. DNA methylation of human endogenous retrovirus in systemic lupus erythematosus. *J Hum Genet*. 2013;58:241–249.
33. Bannert N, Hofmann H, Block A, et al. HERVs New role in cancer: from accused perpetrators to cheerful protectors. *Front Microbiol*. 2018;9:178.
34. Tomlins SA, Laxman B, Dhanasekaran SM, et al. Distinct classes of chromosomal rearrangements create oncogenic ETS gene fusions in prostate cancer. *Nature*. 2007;448:595–599.
35. Chen J, Foroozesh M, Qin Z. Transactivation of human endogenous retroviruses by tumor viruses and their functions in virus-associated malignancies. *Oncogenesis*. 2019;8:6.
36. Dai L, Del Valle L, Miley W, et al. Transactivation of human endogenous retrovirus K (HERV-K) by KSHV promotes Kaposi's Sarcoma development. *Oncogene*. 2018;37:4534–4545.
37. Medstrand P, Mager DL. Human-specific integrations of the HERV-K endogenous retrovirus family. *J Virol*. 1998;72:9782–9787.
38. Garrison KE, Jones RB, Meiklejohn DA, et al. T cell responses to human endogenous retroviruses in HIV-1 infection. *PLoS Pathog*. 2007;3: e165.
39. Monde K, Terasawa H, Nakano Y, et al. Molecular mechanisms by which HERV-K Gag interferes with HIV-1 Gag assembly and particle infectivity. *Retrovirology*. 2017;14:27.
40. Blaise S, de Parseval N, Bénit L, et al. Genomewide screening for fusogenic human endogenous retrovirus envelopes identifies syncytin 2, a gene conserved on primate evolution. *Proc Natl Acad Sci USA*. 2003;100:13013–13018.
41. Imakawa K, Nakagawa S, Miyazawa T. Baton pass hypothesis: Successive incorporation of unconserved endogenous retroviral genes for placentalization during mammalian evolution. *Genes Cells*. 2015;20: 771–788.
42. Kamp C, Hirschmann P, Voss H, et al. Two long homologous retroviral sequence blocks in proximal Yq11 cause AZFa microdeletions as a result of intrachromosomal recombination events. *Hum Mol Genet*. 2000;9: 2563–2572.
43. Horie M, Honda T, Suzuki Y, et al. Endogenous non-retroviral RNA virus elements in mammalian genomes. *Nature*. 2010;463:84–87.
44. Katsura Y, Satta Y. Evolutionary history of the cancer immunity antigen MAGE gene family. *PLoS One*. 2011;6:e20365.
45. Poesz BJ, Ruscetti FW, Gazdar AF, et al. Detection and isolation of type C retrovirus particles from fresh and cultured lymphocytes of a patient with cutaneous T-cell lymphoma. *Proc Natl Acad Sci USA*. 1980;77:7415–7419.
46. Kalyanaraman VS, Sarngadharan MG, Robert-Guroff M, et al. A new subtype of human T-cell leukemia virus (HTLV-II) associated with a T-cell variant of hairy cell leukemia. *Science*. 1982;218:571–573.
47. Calattini S, Chevalier SA, Duprez R, et al. Human T-cell lymphotropic virus type 3: complete nucleotide sequence and characterization of the human Tax3 protein. *J Virol*. 2006;80:9876–9888.

48. **Barré-Sinoussi F, Chermann JC, Rey F, et al.** Isolation of a T-lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS). *Science*. 1983;220:868–871.
49. **Kanki PJ, Barin F, M'Boup S, et al.** New human T-lymphotropic retrovirus related to simian T-lymphotropic virus type III (STLV-III/AGM). *Science*. 1986;232:238–243.
50. **Khabbaz RF, Rowe T, Heneine WM, et al.** Simian immunodeficiency virus needlestick accident in a laboratory worker. *Lancet*. 1992;340:271–273.
51. **Schweizer M, Falcone V, Gänge J, et al.** Simian foamy virus isolated from an accidentally infected human individual. *J Virol*. 1997;71:4821–4824.
52. **Brooks JI, Rud EW, Pilon RG, et al.** Cross-species retroviral transmission from macaques to human beings. *Lancet*. 2002;360:387–388.
53. **Achong BG, Mansell PW, Epstein MA.** A new human virus in cultures from a nasopharyngeal carcinoma. *J Pathol*. 1971;103:P18.

Submitted May 10, 2019; accepted September 26, 2019.

This work was supported by JSPS KAKENHI [18K14766](#).

The authors have no conflicts of interest to disclose.

Correspondence: Yukako Katsura, PhD, Prime Research Institute, Kyoto University, Inuyama, Aichi 484-8506, Japan. (E-mail: katsura.yukako.5e@kyoto-u.ac.jp).