Published in final edited form as:

Int J Parasitol. 2020 February ; 50(2): 153–159. doi:10.1016/j.ijpara.2019.12.003.

Effects of temperature and viscosity on miracidial and cercarial movement of *Schistosoma mansoni*: ramifications for disease transmission

K.H. Nguyen^{a,*}, B.J. Gemmell^a, J.R. Rohr^{a,b}

^aDepartment of Integrative Biology, University of South Florida, 4202 East Fowler Avenue, SCA 110, Tampa, FL 33620, USA

^bDepartment of Biological Sciences, Eck Institute for Global Health, and Environmental Change Initiative, University of Notre Dame, Notre Dame, IN, USA

Abstract

Parasites with complex life cycles can be susceptible to temperature shifts associated with seasonal changes, especially as free-living larvae that depend on a fixed energy reserve to survive outside the host. The life cycle of *Schistosoma*, a trematode genus containing some species that cause human schistosomiasis, has free-living, aquatic miracidial and cercarial larval stages that swim using cilia or a forked tail, respectively. The small size of these swimmers (150–350 μm) dictates that their propulsion is dominated by viscous forces. Given that viscosity inhibits the swimming ability of small organisms and is inversely correlated with temperature, changes in temperature should affect the ability of free-living larval stages to swim and locate a host. By recording miracidial and cercarial movement of *Schistosoma mansoni* using a high-speed camera and manipulating temperature and viscosity independently, we assessed the role each factor plays in the swimming mechanics of the parasite. We found a positive effect of temperature and a negative effect of viscosity on miracidial and cercarial speed. Reynolds numbers, which describe the ratio of inertial to viscous forces exerted on an aquatic organism, were <1 across treatments. Q_{10} values were <2 when comparing viscosity treatments at 20°C and 30°C, further supporting the influence of viscosity on miracidial and cercarial speed. Given that both larval stages have limited energy reserves and infection takes considerable energy, successful transmission depends on both speed and lifespan. We coupled our speed data with mortality measurements across temperatures and discovered that the theoretical maximum distance travelled increased with temperature and decreased with viscosity for both larval stages. Thus, our results suggest that *S. mansoni* transmission is high during warm times of the year, partly due to improved swimming performance of the free-living larval stages, and that increases in temperature variation associated with climate change might further increase transmission.

Graphical Abstract

*Corresponding author. Karena.nguyen@gmail.com.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Schistosomiasis; Climate change; Swimming mechanics; Methylcellulose; Reynolds number; Q_{10}

1. Introduction

Seasonal variation in temperature can strongly affect parasite and host abundance, behavior, development, and survival, and thus the spread of emerging and re-emerging infectious diseases (Altizer et al., 2006; Lafferty, 2009). However, the direction of these effects can be additive, antagonistic, or synergistic (Daszak et al., 2001; McMichael et al., 2006; Rohr et al., 2011; Raffel et al., 2012; Cohen et al., 2018), making it difficult to predict net outcomes on disease transmission. For example, high temperatures can accelerate parasite reproduction and development but can also increase parasite mortality and reduce parasite fitness (Lafferty, 2009; Amarasekare and Savage, 2012;). Examining the life history traits of parasites at all stages in response to temperature is therefore crucial to accurately predict host-parasite dynamics under seasonal conditions, which have been shown to influence transmission via rates of host exposure to parasites and parasite infectivity (Shocket et al., 2018).

Temperature has non-linear effects on reproduction, survival, and infectivity of many parasites and hosts (Studer et al., 2010; Dell et al., 2011; Johnson et al., 2015; Mordecai et al., 2013; Cohen et al., 2017; Rohr et al., 2018). Free-living stages of parasites in aquatic systems, which can be larval or intermediate stages, are especially sensitive to changes in temperature because they have finite energy reserves that deplete faster at higher temperatures (Pietroock and Marcogliese, 2003). Q_{10} , a measure of the rate of change of a reaction for every increase in 10°C , is often used to describe how strongly temperature affects a physiological process. Most biological processes have a Q_{10} between two and three, with values less than two indicating that a physiological process is not very temperature-dependent and values greater than two indicating temperature dependency. Q_{10} has been

used to measure the sensitivity of life history traits of parasites and hosts to seasonally relevant temperature ranges (Morley, 2011, 2012; Paull and Johnson, 2011); however, the effects of temperature and viscosity are confounded for small aquatic organisms (Podolsky and Emler, 1993; Fuiman and Batty, 1997; Gemmell et al., 2013), so obtaining accurate Q_{10} measurements for free-living larvae requires parsing out the effects of viscosity.

Viscosity is inversely correlated with temperature and Reynolds numbers are used to describe the ratio of inertial to viscous forces exerted on an organism in a fluid. Small aquatic organisms typically have a Reynolds number less than one, which indicates that viscous forces dominate their locomotive ability and they must overcome resistance (i.e. viscous drag) to move through the water (Purcell, 1977). Conversely, large aquatic organisms with a Reynolds number greater than one are dominated by inertial forces and changes in the viscosity of a fluid will have less of an impact on locomotion (DeMont and Hokkanen, 1992). That is, larger aquatic organisms can generate enough force to resist viscous drag. Reynolds numbers are commonly reported in fluid mechanics literature and less commonly in parasite literature, but may offer additional insights into parasite locomotion and swimming performance across temperatures.

Schistosoma mansoni is a trematode species that causes schistosomiasis, one of the most prevalent neglected tropical diseases of humans (Hotez, 2013). Approximately 200 million people are infected and more than 800 million individuals are at risk of contracting the disease in endemic areas, particularly in sub-Saharan Africa (WHO, 2019). *Schistosoma mansoni* eggs enter freshwater through human feces and hatch into the first larval stage, called miracidia. These ciliated, 150 μm long organisms swim and mostly infect *Biomphalaria* spp. but can also infect other aquatic snails as an intermediate host. Once inside, miracidia develop into cercariae, which are slightly larger than miracidia (250–350 μm) and swim with a forked tail. Cercariae are released diurnally by snails back into freshwater and infect humans via skin penetration.

Few studies have examined the role of temperature and viscosity in parasite motility, even though free-living larval stages of parasites are small, likely operate under a Reynolds number less than one, do not feed, and thus have limited energy for propulsion. The swimming speed of larval stages has been found to increase the likelihood of infection due to increased contact rates with hosts (Fingerhut et al., 2003; Pasternak, 2004), but this will not occur if mortality occurs before infection. Thus, there is a need to quantify the swimming mechanics of miracidia and cercariae in response to changes in temperature and viscosity, and the estimated distance that these larval stages can travel based on swimming speed and survival. The greater the distance these larval stages can travel, the presumed higher probability of transmission given that finding and penetrating a host takes considerable energy.

The main objectives of this study were to i) decouple and quantify the constitutive effects of temperature and viscosity on miracidial and cercarial speed, miracidial power during swimming, and cercarial tail-beat frequency and ii) calculate the theoretical maximum distance miracidia and cercariae can travel across their natural temperature range. To provide additional measures of how strongly temperature and viscosity affect movement, we

calculated Q_{10} values and Reynolds numbers for miracidial and cercarial speed. We expected that higher temperatures would result in higher metabolic rates and thus faster swimming speeds, more power expended for miracidia, and quicker tail-beat frequencies for cercariae. However, we also expected that miracidial and cercarial speed, miracidial power, and cercarial tail-beat frequency would decrease under viscous forces because miracidia and cercariae are small and likely operate at Reynolds numbers less than one.

2. Materials and methods

2.1. Obtaining *Schistosoma mansoni* miracidia and cercariae

We harvested livers from Swiss Webster mice exposed to the NMRI strain of *S. mansoni*. A liver homogenate was passed through a 106 μm sieve (Fisherbrand, ASTM E-11 Standard, USA) and the solution that passed through the sieve was caught in another container. This solution, which contained *S. mansoni* eggs, was subsequently passed through a 45 μm sieve (Fisher Scientific Company, ASTM E-11 Standard, USA). We rinsed the eggs from this sieve with 1.2% saline into a beaker with an artificial freshwater solution modified from Baer et al. (1999) to induce hatching.

To obtain cercariae, *Biomphalaria glabrata* (NMRI strain) were exposed to 10–20 *S. mansoni* miracidia. Individuals were held at room temperature (22°C), fed ad libitum (6 g of fish flakes and 6 g of spirulina in 2.5 g of agar), and tested for infection 8 weeks post-exposure. On the day of the experiment, infected snails were placed under artificial light for 1 h at room temperature (22°C) to induce cercarial emergence. All cercariae were homogenized into one beaker with the same artificial freshwater solution described above.

2.2. Manipulation of temperature and viscosity

Water was either heated using a 200 W submersible aquarium heater or cooled using a 1/10 hp aquarium chiller unit. The water temperatures tested were 10°C, 20°C, and 30°C. All data recordings (filming) were performed in a recirculating water bath to maintain the temperature within $\pm 1^\circ\text{C}$ and reduce the presence of small-scale convective currents within the filming vessel. We manipulated viscosity independently of temperature at 20°C and 30°C by dissolving methylcellulose polymer (25cP) (MC). A Cannon-Fenske routine viscometer was used to determine the concentration of methylcellulose solution that matched the kinematic viscosity of water at 10°C ($\nu = 1.31 \times 10^{-6} \text{ m}^2\text{s}^{-1}$). We had a total of five treatments: 10°C, 20°C, 30°C, 20°C with MC, and 30°C with MC. Previous studies have shown no effect of methylcellulose on the metabolic rates of organisms (Luckinbill, 1973; Fuiman and Batty, 1997).

2.3. High-speed videography of miracidia and cercariae

Once *S. mansoni* miracidia and cercariae were collected, individuals were transferred to a glass observation cuvette (1 cm $\times$ 5 cm $\times$ 5 cm, Model 3-I-40, Starna Cells, Inc., USA) with water at 10, 20, 30°C, 20°C with MC, or 30°C with MC. We used a high-speed camera at 1280 $\times$ 1024 pixel resolution (Edgetronic SC1, USA) to film miracidia and cercariae at 100 frames per second (fps), 250 fps, or 500 fps, depending on the organism's swimming speed (Fig. 1). All videos were imported into ImageJ (Schneider et al., 2012) for kinematic

analyses. We only analyzed individuals that swam in the camera's focal plane over the entire period of quantification to minimize error associated with out of plane movements, which resulted in unequal sample sizes across treatments. We calculated average miracidial speed at 10°C ($n = 17$), 20°C ($n = 24$), 30°C ($n = 36$), 20°C with MC ($n = 11$), and 30°C with MC ($n = 21$) by tracking individuals frame by frame and measuring the distance individuals traveled within that time period. We employed the same method for cercarial speed at 10°C ($n = 8$), 20°C ($n = 10$), 30°C ($n = 16$), 20°C with MC ($n = 10$), and 30°C with MC ($n = 8$). Water was seeded with neutrally buoyant particles (5–20 μm diameter) so for each swimming sequence, the speed of an in-focus nearby particle was also quantified to correct for any effects of convection currents present in the observation cuvette. We calculated net miracidial and cercarial swimming speeds and accounted for any convective currents present in the filming vessel by subtracting the speeds of nearby neutrally buoyant tracer particles. For cercarial tail-beat frequency, we manually counted five tail-beats per individual at five random intervals and calculated the average number of tail-beats per second per individual at 10°C ($n = 10$), 20°C ($n = 10$), 30°C ($n = 21$), 20°C with MC ($n = 10$), and 30°C with MC ($n = 10$). An example of miracidia at 30°C (15fps, Supplementary Movie S1) and cercariae at 30°C (10fps, Supplementary Movie S2) without methylcellulose can be viewed online.

2.4. Statistical analyses

All statistical analyses were conducted in R version 3.2.4 (R Foundation for Statistical Computing, Vienna, Austria, 2017). We applied a $\log_{10}(x+1)$ transformation to miracidial and cercarial speed to adhere to assumptions of normality. To determine the effects of temperature independent of viscosity, we analyzed the miracidial and cercarial data separately and used two-sided t -tests to compare mean miracidial speed, cercarial speed, miracidial power, and cercarial tail-beat frequency between 10°C and 20°C with MC, 10°C and 30°C with MC, and 20°C with MC and 30°C with MC. Due to unequal sample sizes, the variance between some groups were unequal and a Welch's t -test was conducted instead. A sequential Bonferroni correction was then employed to account for multiple comparisons (Benjamini and Hochberg, 1995), with sequential P values of less than 0.05, 0.025, and 0.017 considered significant, respectively. Next, to analyze the overall effects of temperature on these response variables at 10°C, 20°C with MC, and 30°C with MC, we applied a generalized linear model (*glm* function, *stats* package) with temperature as a continuous independent variable. To test for interactions between temperature and viscosity, we excluded the 10°C treatment and used a 2×2 ANOVA (*Anova* function, *car* package) with temperature and viscosity as categorical independent variables. P values less than 0.05 were considered significant for these analyses.

To calculate the amount of power miracidia require to move through the water, we used the following equation for the metabolic cost of motility for ciliated organisms (Crawford, 1992):

$$P = 3\pi Dv^2\eta$$

where D is the organism's diameter (m), v is the organism's velocity (m/s), and η is the water viscosity (N*s/m²). We did not apply the same equation to cercariae given their non-ciliated body morphology and unusual movement patterns.

To measure how strongly temperature affected miracidial and cercarial speed, we calculated Q_{10} values using the following equation:

$$Q_{10} = \left(\frac{R_2}{R_1} \right)^{\frac{10^\circ C}{T_2 - T_1}}$$

where R is the rate of the response variable and T is the temperature (°C).

To measure how strongly viscosity affected miracidial and cercarial speed, we calculated Reynolds number using the following equation:

$$Re = \frac{\rho u L}{\mu}$$

where ρ is the density of the fluid (kg/m³), u is the velocity of the organism (m/s), L is the length of the organism (m), and μ is the dynamic viscosity of the fluid (N*s/m²).

Lastly, to calculate the theoretical maximum distance miracidia and cercariae could travel, we extracted survival data at 10, 20, and 30°C for miracidia from Anderson et al. (1982) and at 20 and 30°C for cercariae from Lawson and Wilson (1980). Cercarial survival data at 10°C was not available for extraction from the literature. We then multiplied longevity values by average swimming speed, assuming constant movement throughout miracidial and cercarial lifespans.

3. Results

3.1. Miracidial and cercarial speed, miracidial power, and cercarial tail-beat frequency

Temperature had a positive significant effect on miracidial speed ($t = 4.71$, $df = 48$, $P < 0.0001$, Table 1) and the 2×2 ANOVA showed a significant effect of temperature ($F_{1,88} = 25.92$, $df = 1$, $P < 0.0001$) and viscosity ($F_{1,88} = 6.99$, $df = 1$, $P = 0.0097$), but no interaction between those two factors ($F_{1,88} = 1.32$, $df = 1$, $P = 0.2534$) (Fig. 2A). A Bonferroni-adjusted t -test revealed that mean miracidial speed was significantly greater at 30°C with MC than 10°C ($P < 0.0001$, Table 2). In fact, miracidia swam 2.5 times faster at 30°C with MC than 10°C, with a difference of 0.17 mm/s (95% Confidence Interval (CI): 0.10, 0.24). All other Bonferroni-adjusted t -tests were not significant (Table 2).

Temperature also had a positive significant effect on cercarial speed ($t = 5.82$, $df = 25$, $P < 0.0001$, Table 1). The 2×2 ANOVA indicated that there was a significant effect of temperature ($F_{1,40} = 69.31$, $df = 1$, $P < 0.0001$) and viscosity ($F_{1,40} = 51.00$, $df = 1$, $P < 0.0001$), but the interaction between those two factors was not significant ($F_{1,40} = 4.00$, $df = 1$, $P = 0.0525$) (Fig. 2B). All Bonferroni-adjusted t -tests for cercarial speed were significant (Table 2).

Temperature had a positive significant effect on miracidial power ($t = 3.62$, $df = 48$, $P = 0.0007$, Table 1) (Fig. 3A). The 2×2 ANOVA showed a significant effect of temperature ($F_{1,88} = 16.31$, $df = 1$, $P = 0.0001$), but not viscosity ($F_{1,88} = 0.42$, $df = 1$, $P = 0.5206$) or the interaction ($F_{1,88} = 0.36$, $df = 1$, $P = 0.5526$). The amount of power generated at 30°C with MC was three times greater than that at 20°C with MC ($P = 0.0150$, Table 2) and more than 18 times greater than that at 10°C ($P = 0.0019$, Table 2). All other Bonferroni-adjusted t -tests were not significant (Table 2).

Temperature also had a positive effect on cercarial tail-beat frequency ($t = 62.90$, $df = 149$, $P < 0.0001$, Table 1). The 2×2 ANOVA indicated there was a highly significant effect of temperature ($F_{1,251} = 1725.95$, $df = 1$, $P < 0.0001$), but not viscosity ($F_{1,251} = 0.28$, $df = 1$, $P = 0.5944$) or the interaction ($F_{1,251} = 2.67$, $df = 1$, $P = 0.1034$) (Fig. 3B). All Bonferroni-adjusted t -tests for cercarial tail-beat frequency were significant (Table 2).

3.2. Q_{10} , Reynolds number, and maximum distance travelled

For miracidial and cercarial speed, Q_{10} values were between 1.05 and 2.65, with cercariae exhibiting the widest range of values. Q_{10} values for miracidial speed were greater than two in treatments where viscosity was not manipulated but decreased to less than two when MC was added. Q_{10} for cercarial speed exhibited similar trends (Table 3). Reynolds numbers for miracidial and cercarial speed were less than one across all treatments, but increased with temperature and decreased with viscosity (Table 3). That is, treatments with MC had smaller Reynolds numbers compared to treatments at the same temperature where viscosity was not manipulated for both larval stages.

Despite dying sooner at higher temperatures, the estimated maximum distance miracidia and cercariae could travel increased significantly with temperature (Fig. 4). When MC was added, the distance miracidia could travel decreased from 39 and 52 m to approximately 33 and 34 m at 20 and 30°C, respectively (Fig. 4A). The distance cercariae could travel decreased from approximately 122 and 127 m to 100 and 81 m at 20°C and 30°C, respectively (Fig. 4B). Overall, the degree to which viscosity decreased the distance cercariae could travel at 30°C versus 20°C with MC was much greater than the decrease between 20°C versus 20°C with MC.

4. Discussion

In this study, we decoupled the effects of temperature and viscosity over a natural seasonal temperature range on the aquatic larval stages of the human parasite *Schistosoma mansoni* to understand the physiological mechanisms underlying their movement. We found that temperature and viscosity have significant effects on miracidial and cercarial swimming speeds and the overall distance that these larval stages can travel. Cercarial tail-beat frequency and miracidial power were strongly affected by temperature but unaffected by changes in viscosity. Reynolds numbers for miracidial and cercarial speed were less than one across all treatments, supporting the notion that both larval stages operate at low Reynolds numbers. Q_{10} values for miracidial and cercarial speed were greater than two when viscosity was not manipulated. These values decreased to values less than two when methylcellulose was added, indicating that the effect of temperature independent of viscosity was weaker

than the effect of these factors combined. Said another way, the relative contribution of temperature to increasing miracidial and cercarial speed was overestimated when temperature and viscosity were coupled. Theoretical maximum distance calculations provided additional evidence that low viscosity at high temperatures strongly impact parasite locomotion.

Swimming speed is an important parameter for estimating exposure rates between the larval stages of *S. mansoni* and their respective hosts. If aquatic larval stages of parasites can swim faster and travel farther before mortality, then the likelihood of encountering and infecting a host increases. Here, we found that miracidial speed nearly doubled with every increase of 10°C, which aligns with previous results that showed that the swimming rate of *S. mansoni* miracidia almost doubled from 4 to 12°C, 12°C to 22°C, and 22 to 34°C (Samuelson et al., 1984). Similarly, we found that cercarial speed more than doubled with every increase of 10°C. These results could be used in mathematical models to more accurately reflect mechanisms underlying disease transmission. For example, a study conducted by Civitello and Rohr (2014) found that models separating exposure and infection rates of *S. mansoni* miracidia produced better predictions of prevalence in intermediate snail host populations compared with models that combined these processes. Incorporating exposure rate as a function of swimming speed and longevity may therefore produce more ecologically relevant predictions of disease risk.

Previous studies have used methylcellulose or similar agents to decouple the effects of temperature and viscosity. All noted that the locomotive ability of free-living organisms with low Reynolds numbers decreased when viscosity was increased at high temperatures; this trend was not seen for larger, high Reynolds number organisms that can more easily overcome viscous forces (Luckinbill, 1973; Linley, 1986; Podolsky and Emlet, 1993; Fuiman and Batty, 1997; Gemmell et al., 2013). Our results agree with these findings, as we found that the swimming performance of miracidia and cercariae was poor at 10°C when viscosity was high. Surprisingly, miracidial speed did not significantly decrease in the 20°C with MC treatment compared with the 20°C treatment while cercarial speed did, suggesting that the larger cercarial stage may be more vulnerable to changes in viscosity at intermediate temperatures. This result was unexpected because miracidia generate thrust through multiple cilia and should be more strongly impacted by viscosity due to a higher surface area to volume ratio. Conversely, cercariae can only generate thrust from a single forked tail and should be less impacted by viscosity due to a lower surface area to volume ratio (Samuelson et al., 1984; Biewener, 2003). However, both miracidial and cercarial speed significantly decreased in the 30°C with MC treatment compared with the 30°C treatment, which demonstrates that free-living larval stages are strongly affected by changes in viscosity at high temperatures.

Miracidia and cercariae have very different body morphologies and means of locomotion, making direct comparisons difficult. Cercarial tail-beat frequency and miracidial power are not directly comparable, but they do provide an indirect measure of the effect of temperature and viscosity on locomotive ability. Viscosity did not significantly hinder cercarial tail-beat frequency and miracidial power production across temperatures in this study. A recent study conducted by Krishnamurthy et al. (2016) found that cercariae generate thrust by

constricting smooth muscles at the anterior and posterior ends of their tail. Importantly, the joints at these ends are flexible and allow cercariae to bend their tails in a way that generates enough thrust to overcome viscous drag. Similar adaptations are present among other low Reynolds number swimmers and could explain why tail-beat frequency and miracidial power are driven more by temperature, which influences metabolic rate and muscle contractions, than changes in viscosity (Crawford, 1992).

For miracidial and cercarial speed, Reynolds numbers increased with temperature and Q_{10} values were greater than two for treatments without MC, supporting the notion that the swimming mechanics of these organisms is temperature-dependent. However, Reynolds numbers for treatments with constant viscosity (i.e. 20°C with MC and 30°C with MC) were much smaller than treatments where viscosity was not manipulated (i.e. 20°C and 30°C), indicating that a decrease in viscosity associated with high temperatures allows miracidia and cercariae to move more efficiently than the effect of temperature alone. Similarly, Q_{10} values were less than two when MC was added, indicating that the influence of temperature on speed was likely overestimated when the effect of viscosity was ignored. This trend has been observed in other aquatic organisms (Podolsky and Emlet, 1993; Fuiman and Batty, 1997).

By independently manipulating temperature and viscosity, we were able to measure the relative effects of temperature and viscosity on the swimming performance of low Reynolds number swimmers. Coupling these findings with temperature-dependent longevity data allowed us to calculate the theoretical maximum distance miracidia and cercariae could travel as a function of energy availability and swimming speed, which decreased significantly with high viscosity (Fig. 4). These results make sense, as the change in viscosity from 30°C ($\nu = 8.00 \times 10^{-7} \text{ m}^2\text{s}^{-1}$) to 10°C ($\nu = 1.31 \times 10^{-6} \text{ m}^2\text{s}^{-1}$) is substantial (64% increase) and will have the greatest impact on small, low Reynolds number swimmers that move in a fluid regime dominated by viscous forces. In contrast, larger organisms would not be impacted greatly by such changes in kinematic viscosity due to the fact that they experience a primarily inertial fluid regime. Overall, our estimates of maximum distance are more easily applied to miracidia, as they are constant swimmers, but represent a conservative estimate for cercariae, as they spend time passively sinking in addition to actively swimming (Graefe et al., 1967; Nuttman, 1974; Brachs and Haas, 2008; Krishnamurthy et al., 2016). Cercariae are likely able to travel farther than our estimate of 127 m at 30°C due to an alternating pattern of swimming and passive sinking in which the passive phase would require little of the stored metabolic energy. For both miracidia and cercariae, increasing temperature may indirectly increase the probability of encountering an intermediate snail host or a human host, respectively, because the cost of increased swim speeds and shorter lifespan is smaller than the cost of decreased swim speeds under viscous conditions. Taken together, these results suggest that the energetic cost of enhanced swimming performance for both larval stages is partially offset by low viscosity at high temperatures and the survival of these organisms may not be as negatively impacted by high temperatures as previously assumed.

We provide many lines of evidence showing that miracidia and cercariae display poor swimming performance due to the additive effects of low temperature, which decreases

metabolic rate, and high viscosity, which decreases swimming efficiency. Conversely, low viscosity at high temperatures reduces the energetic cost of movement for both larval stages and allows them to travel farther than they can at low temperatures despite shorter lifespans. That is, the tradeoff between faster speed and shorter lifespan at high temperatures seems to result in a net benefit for miracidia and cercariae. As such, warming seasonal and diel temperature variation associated with global climate change might increase the exposure rate to potential hosts and thus increase disease transmission (Poulin, 2006; Lafferty, 2009; Mas-Coma et al., 2009). Biotic factors, economic development, urbanization, human migration, mass drug administration, and other factors also influence disease transmission and should not be discounted (Thieltges et al., 2008; Johnson et al., 2009; Blum and Hotez, 2018), but the results presented here offer a novel perspective on why parasitic diseases may increase in the context of global climate change.

This study is one of only a few to apply principles of fluid mechanics to the aquatic larval stages of a human parasite to improve our understanding of how temperature and viscosity affect parasite transmission. Overall, this work provides a fundamental mechanistic understanding of parasite movement across temperatures and explains how increases in seasonal temperature variation might increase disease transmission in host-parasite systems that have free-living larval stages.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Swiss Webster mice infected with *Schistosoma mansoni* were provided by the National Institute of Allergy and Infectious Diseases (NIAID) Schistosomiasis Resource Center of the Biomedical Research Institute (Rockville, MD, USA) through the National Institutes of Health (NIH)-NIAID Contract HHSN272201000005I for distribution through BEI Resources, USA. Funds for the project were provided by grants to J.R.R. from the National Science Foundation, USA (EF-1241889), the NIH, USA (R01TW010286), and the U.S. Department of Agriculture [2009-35102-0543].

References

- Altizer S, Dobson A, Hosseini P, Hudson P, Pascual M, Rohani P, 2006 Seasonality and the dynamics of infectious diseases. *Ecol Lett* 9, 467–484. [PubMed: 16623732]
- Amarasekare P, Savage V, 2012 A framework for elucidating the temperature dependence of fitness. *Am Nat* 179, 178–191. [PubMed: 22218308]
- Anderson RM, Mercer JG, Wilson RA, Carter NP, 1982 Transmission of *Schistosoma mansoni* from man to snail: experimental studies of miracidial survival and infectivity in relation to larval age, water temperature, host size and host age. *Parasitology* 85 (Pt 2), 339–360. [PubMed: 7145476]
- Baer KN, Ziegenfuss MC, Banks SD, Ling Z, 1999 Suitability of high-hardness COMBO medium for ecotoxicity testing using algae, daphnids, and fish. *Bull Environ Contam Toxicol* 63, 289–296. [PubMed: 10475905]
- Benjamini Y, Hochberg Y, 1995 Controlling the False Discovery Rate - a Practical and Powerful Approach to Multiple Testing. *J R Stat Soc B* 57, 289–300.
- Biewener AA, 2003 *Animal Locomotion*. Oxford University Press New York.
- Blum AJ, Hotez PJ, 2018 Global “worming”: Climate change and its projected general impact on human helminth infections. *PLoS Negl Trop Dis* 12, e0006370. [PubMed: 30024871]

- Brachs S, Haas W, 2008 Swimming behaviour of *Schistosoma mansoni* cercariae: responses to irradiance changes and skin attractants. *Parasitol Res* 102, 685–690. [PubMed: 18157546]
- Civitello DJ, Rohr JR, 2014 Disentangling the effects of exposure and susceptibility on transmission of the zoonotic parasite *Schistosoma mansoni*. *J Anim Ecol* 83, 1379–1386. [PubMed: 24702134]
- Cohen JM, Lajeunesse MJ, Rohr JR, 2018 A global synthesis of animal phenological responses to climate change. *Nat. Clim. Change* 8, 224–228.
- Cohen JM, Venesky MD, Sauer EL, Civitello DJ, McMahon TA, Roznik EA, Rohr JR, 2017 The thermal mismatch hypothesis explains host susceptibility to an emerging infectious disease. *Ecol Lett* 20, 184–193. [PubMed: 28111904]
- Crawford DW, 1992 Metabolic cost of motility in planktonic protists: Theoretical considerations on size scaling and swimming speed. *Microb Ecol* 24, 1–10. [PubMed: 24193035]
- Daszak P, Cunningham AA, Hyatt AD, 2001 Anthropogenic environmental change and the emergence of infectious diseases in wildlife. *Acta Trop* 78, 103–116. [PubMed: 11230820]
- Dell AI, Pawar S, Savage VM, 2011 Systematic variation in the temperature dependence of physiological and ecological traits. *Proc Natl Acad Sci USA* 108, 10591–10596. [PubMed: 21606358]
- DeMont ME, Hokkanen JEL, 1992 Hydrodynamics of animal movement, in: Biewener AA (Ed.), *Biomechanics Structures and Systems*. Oxford University Press, New York, pp. 263–284.
- Fingerut JT, Ann Zimmer C, Zimmer RK, 2003 Larval Swimming Overpowers Turbulent Mixing and Facilitates Transmission of a Marine Parasite. *Ecology* 84, 2502–2515.
- Fuiman LA, Batty RS, 1997 What a drag it is getting cold: Partitioning the physical and physiological effects of temperature on fish swimming. *J. Exp. Biol* 200, 1745–1755. [PubMed: 9319652]
- Gemmell BJ, Sheng J, Buskey EJ, 2013 Compensatory escape mechanism at low Reynolds number. *Proc Natl Acad Sci USA* 110, 4661–4666. [PubMed: 23487740]
- Graefe G, Hohorst W, Dräger H, 1967 Forked tail of the cercaria of *Schistosoma mansoni*--a rowing device. *Nature* 215, 207–208. [PubMed: 6049130]
- Hotez PJ, 2013 *Forgotten people, forgotten diseases: the neglected tropical diseases and their impact on global health a development*, Second ed ASM Press Washington, D.C.
- Johnson LR, Ben-Horin T, Lafferty KD, McNally A, Mordecai E, Paaijmans KP, Pawar S, Ryan SJ, 2015 Understanding uncertainty in temperature effects on vector-borne disease: a Bayesian approach. *Ecology* 96, 203–213. [PubMed: 26236905]
- Johnson PT, Lund PJ, Hartson RB, Yoshino TP, 2009 Community diversity reduces *Schistosoma mansoni* transmission, host pathology and human infection risk. *Proc Biol Sci* 276, 1657–1663. [PubMed: 19203926]
- Krishnamurthy D, Katsikis G, Bhargava A, Prakash M, 2016 *Schistosoma mansoni* cercariae swim efficiently by exploiting an elastohydrodynamic coupling. *Nature Physics* 13, 266–271.
- Lafferty KD, 2009 The ecology of climate change and infectious diseases. *Ecology* 90, 888–900. [PubMed: 19449681]
- Lawson JR, Wilson RA, 1980 The survival of the cercariae of *Schistosoma mansoni* in relation to water temperature and glycogen utilization. *Parasitology* 81, 337–348. [PubMed: 7443297]
- Linley JR, 1986 Swimming behavior of the larva of *Culicoides variipennis* (Diptera: Ceratopogonidae) and its relationship to temperature and viscosity. *J Med Entomol* 23, 473–483. [PubMed: 3772953]
- Luckinbill LS, 1973 Coexistence in Laboratory Populations of *Paramecium Aurelia* and Its Predator *Didinium Nasutum*. *Ecology* 54, 1320–1327.
- Mas-Coma S, Valero MA, Bargues MD, 2009 Climate change effects on trematodiasis, with emphasis on zoonotic fascioliasis and schistosomiasis. *Vet Parasitol* 163, 264–280. [PubMed: 19375233]
- McMichael AJ, Woodruff RE, Hales S, 2006 Climate change and human health: present and future risks. *Lancet* 367, 859–869. [PubMed: 16530580]
- Mordecai EA, Paaijmans KP, Johnson LR, Balzer C, Ben-Horin T, de Moor E, McNally A, Pawar S, Ryan SJ, Smith TC, Lafferty KD, 2013 Optimal temperature for malaria transmission is dramatically lower than previously predicted. *Ecol Lett* 16, 22–30. [PubMed: 23050931]
- Morley NJ, 2011 Thermodynamics of cercarial survival and metabolism in a changing climate. *Parasitology* 138, 1442–1452. [PubMed: 21819641]

- Morley NJ, 2012 Thermodynamics of miracidial survival and metabolism. *Parasitology* 139, 1640–1651. [PubMed: 22814411]
- Nuttman CJ, 1974 The fine structure and organization of the tail musculature of the cercaria of *Schistosoma mansoni*. *Parasitology* 68, 147–154. [PubMed: 4826714]
- Pasternak Z, 2004 Host location by larvae of a parasitic barnacle: larval chemotaxis and plume tracking in flow. *J. Plankton Res* 26, 487–493.
- Paull SH, Johnson PTJ, 2011 High temperature enhances host pathology in a snail-trematode system: possible consequences of climate change for the emergence of disease. *Freshwater Biol.* 56, 767–778.
- Pietroock M, Marcogliese DJ, 2003 Free-living endohelminth stages: at the mercy of environmental conditions. *Trends Parasitol.* 19, 293–299. [PubMed: 12855379]
- Podolsky RD, Emlet RB, 1993 Separating the Effects of Temperature and Viscosity on Swimming and Water-Movement by Sand Dollar Larvae (*Dendraster-Excentricus*). *J. Exp. Biol* 176, 207–221.
- Poulin R, 2006 Global warming and temperature-mediated increases in cercarial emergence in trematode parasites. *Parasitology* 132, 143–151. [PubMed: 16393363]
- Purcell EM, 1977 Life at low Reynolds number. *Am. J. Phys* 45, 3–11.
- Raffel TR, Romansic JM, Halstead NT, McMahon TA, Venesky MD, Rohr JR, 2012 Disease and thermal acclimation in a more variable and unpredictable climate. *Nat. Clim. Change* 3, 146–151.
- Rohr JR, Civitello DJ, Cohen JM, Roznik EA, Sinervo B, Dell AI, 2018 The complex drivers of thermal acclimation and breadth in ectotherms. *Ecol Lett* 21, 1425–1439. [PubMed: 30009486]
- Rohr JR, Dobson AP, Johnson PT, Kilpatrick AM, Paull SH, Raffel TR, Ruiz-Moreno D, Thomas MB, 2011 Frontiers in climate change-disease research. *Trends Ecol Evol* 26, 270–277. [PubMed: 21481487]
- Samuelson JC, Quinn JJ, Caulfield JP, 1984 Hatching, chemokinesis, and transformation of miracidia of *Schistosoma mansoni*. *Parasitology* 70, 321–331.
- Schneider CA, Rasband WS, Eliceiri KW, 2012 NIH Image to ImageJ: 25 years of image analysis. *Nat Methods* 9, 671–675. [PubMed: 22930834]
- Shocket MS, Vergara D, Sickbert AJ, Walsman JM, Strauss AT, Hite JL, Duffy MA, Caceres CE, Hall SR, 2018 Parasite rearing and infection temperatures jointly influence disease transmission and shape seasonality of epidemics. *Ecology* 99, 1975–1987. [PubMed: 29920661]
- Studer A, Thieltges DW, Poulin R, 2010 Parasites and global warming: net effects of temperature on an intertidal host–parasite system. *Mar. Ecol. Prog. Ser* 415, 11–22.
- Thieltges DW, Jensen KT, Poulin R, 2008 The role of biotic factors in the transmission of free-living endohelminth stages. *Parasitology* 135, 407–426. [PubMed: 18208633]
- WHO, 2019 Schistosomiasis. World Health Organization Geneva, Switzerland.

Highlights

- High-speed videography quantifies the swimming speed of *Schistosoma mansoni* larvae.
- Swimming speeds of larvae increased with temperature and decreased with viscosity.
- Temperature played a greater role than viscosity in some physiological responses.
- *Schistosoma mansoni* larvae travel farther at higher temperatures despite dying sooner.
- Improved swimming performance in warming climates could increase transmission.

Fig. 1.
Experimental set-up is shown here. Note, this diagram is not drawn to scale.

Fig. 2. Mean (± 1 SE) *Schistosoma mansoni* miracidial speed (A) and cercarial speed (B) under various temperature and viscosity treatments. ● and ▲ represent no methylcellulose (MC) and methylcellulose (MC +) treatments, respectively.

Fig. 3. Mean (± 1 SE) *Schistosoma mansoni* miracidial power (A) and cercarial tail-beat frequency (B) speed across temperature and viscosity treatments. ● and ▲ represent no methylcellulose (MC) and methylcellulose (MC +) treatments, respectively.

Fig. 4. Mean (± 1 SE) estimated maximum distance traveled by *Schistosoma mansoni* miracidia (A) and cercariae (B) across temperature and viscosity treatments, based on active swimming speeds and longevity. Standard errors are derived from miracidial and cercarial speed exclusively because error estimates associated with longevity were not always provided in the original papers from which these data were extracted. ● and ▲ represent no methylcellulose (MC) and methylcellulose (MC +) treatments, respectively.

Table 1.

Speed, power, and tail-beat frequency for *Schistosoma mansoni* miracidia and cercariae as a function of temperature compared across 10°C, 20°C with methylcellulose, and 30°C with methylcellulose. Models are shown below the response variable.

Generalized linear models	Intercept	Slope	Standard error
<i>Miracidial speed</i>			
Log ₁₀ (Speed+1) ~ Temperature	0.014	0.008	0.002
<i>Cercarial speed</i>			
Log ₁₀ (Speed+1) ~ Temperature	0.048	0.009	0.002
<i>Miracidial power</i>			
Power ~ Temperature	-1.257 ⁻¹²	1.229 ⁻¹³	3.391 ⁻¹⁴
<i>Cercarial tail-beat frequency</i>			
Tail-beat frequency ~ Temperature	-5.215	1.135	0.018

Table 2.

Speed, power, and tail-beat frequency for *Schistosoma mansoni* miracidia and cercariae compared at 10°C, 20°C with methylcellulose (MC), and 30°C with methylcellulose (MC). A Welch's *t*-test was conducted for groups with unequal variance. All treatments are listed in descending order of *P* values and are bolded for significance according to sequential Bonferroni corrections.

Comparison	<i>t</i>	<i>df</i>	<i>P</i>
<i>Miracidial speed</i>			
20°C with MC & 30°C with MC	1.90	30.00	0.0674
10°C & 20°C with MC	2.50	11.51	0.0287
10°C & 30°C with MC	4.93	22.10	< 0.0001
<i>Cercarial speed</i>			
20°C with MC & 30°C with MC	3.19	16	0.0057
10°C & 20°C with MC	3.25	16	0.0050
10°C & 30°C with MC	5.36	14	0.0001
<i>Miracidial power</i>			
10°C & 20°C with MC	2.15	10.15	0.0570
20°C with MC & 30°C with MC	2.61	24.88	0.0150
10°C & 30°C with MC	3.58	20.04	0.0019
<i>Cercarial tail-beat frequency</i>			
10°C & 20°C with MC	69.75	82.64	< 0.0001
10°C & 30°C with MC	53.84	53.14	< 0.0001
20°C with MC & 30°C with MC	26.69	59.32	< 0.0001

Table 3.

Reynolds numbers and Q_{10} values for *Schistosoma mansoni* miracidial and cercarial speed across treatments with and without methylcellulose (MC).

Treatment	Miracidia		Cercariae	
	Reynolds number	Q_{10}	Reynolds number	Q_{10}
10°C	0.030	-	0.071	-
20°C	0.091	2.32	0.244	2.65
30°C	0.271	2.35	0.504	2.09
20°C MC	0.059	1.97	0.138	1.95
30°C MC	0.109	1.91 ^a , 1.85 ^b	0.196	1.05 ^a , 1.43 ^b

^a Compared between 30°C MC and 20°C.

^b Compared between 30°C MC and 20°C MC.