Abstracts

3rd International Congress

“Sepsis and Multiorgan Dysfunction”

September 5–8, 2007
Weimar

Organizer
K. Reinhart (responsible)

Guest Editors
K. Reinhart
F. M. Brunkhorst

German Sepsis Society (DSG)
www.sepsis-gesellschaft.de

Contents

Abstracts

Clinical Sepsis Research	3
Abstracts 001, 005–006, 008–009, 013–017, 020–021, 024–025, 027–028, 030–031, 034–038, 041, 043–045, 047, 050, 053–055, 059, 061–062, 067–068, 072, 074–077, 080, 082, 084–092, 095–096	
Pathophysiology	25
Abstracts 002–004, 007, 010–012, 018, 022–023, 029, 032–033, 039, 042, 046, 049, 052, 056–058, 060, 064–066, 070, 078–079	
Satellite Symposia	37
Abstracts 019, 026, 040, 048, 051, 063, 069, 071, 073, 081, 083, 093–094	

Author Index	43
---------------------------	-----------

Infection 2007; 35 (Suppl. II): 1–44

DOI 10.1007/s15010-007-1001-1

Clinical Sepsis Research

001

Infection 2007; 35 (Suppl. II): 3

Investigation of the suitability of Procalcitonin to guide antibiotic therapy in surgical intensive care patients

*Hochreiter M (1), Köhler T (1), Bein B (2), Schweiger A-M (1), von Spiegel T (1), Schröder S (1),

(1) Department of Anesthesiology and Intensive Care, Westwüstenklinikum Heide, Germany (2) Department of Anesthesiology and Operative Intensive Care, Universital Hospital Kiel, Germany

Introduction: The development of resistance by infective bacterial species is an encouragement for us to reconsider the indication and administration of the available antibiotics. Proper recognition of the indication and the correct duration of therapy are particularly important for the use of highly potent substances in intensive care. There has as yet been no clinical chemical parameter which is capable of specifically distinguishing a bacterial infection from a viral or non-infectious inflammatory reaction. It now appears that procalcitonin (PCT) offers this possibility (1).

Objective: The present study is intended to clarify whether PCT can be used to guide antibiotic therapy in surgical intensive care patients.

Methods and results: 87 patients in a surgical intensive care ward were enrolled in this study, with the agreement of the ethics committee. 57 patients were given antibiotic therapy after confirmed infection or a high grade suspicion of an infection. After a daily PCT determination, a new decision – based on a clinical algorithm – was reached each day as to whether the antibiotic therapy should be continued. The control group consisted of 30 patients with a standardised duration of antibiotic therapy over 8 days. There were no significant differences between the control group and the group with PCT-steered antibiotic therapy with respect to the demographic data on age, gender distribution, SAPS II and outcome. The distribution of the diagnoses, the antibiotic class used and the daily costs of antibiotic therapy were comparable in the two groups. The period spent in the intensive care ward was significantly shorter in the group with PCT-steered antibiotic therapy than in the control group. The median period of antibiotic therapy was 6 days in the PCT group, which was significantly shorter than the value of 8 days for the control group. The daily determination of PCT for intensive care patients shortened the duration of antibiotic therapy, with favourable effects on antibiotic costs. There were no unfavourable effects on the outcome.

References: 1. Christ-Crain M and Müller B (2005) Swiss Medical Weekly 135: 451-460

005

Infection 2007; 35 (Suppl. II): 3

Procalcitonin ratio indicates successful surgical treatment of abdominal sepsis

*Novotny A, Emmanuel K, Hueser N, Matevossian E, Kriner M, Ulm K, Bartels H

Department of Surgery, Technical University Munich, Germany

Introduction: Planned relaparotomy and relaparotomy on demand are alternative treatment strategies in the surgical treatment of secondary peritonitis. On-demand relaparotomy was reported to be associated with a slight reduction of mortality when compared with planned relaparotomy. Problems of on-demand relaparotomy are, however, that it must be performed without delay and the first signs of progress of the septic course must be detected before the onset of multiorgan failure. Therefore, a parameter for early detection of progressive sepsis after initial surgical elimination of the source of infection would be

of great benefit. One of the serum biochemical markers that qualifies for this purpose is Procalcitonin (PCT).

Objective: To evaluate serum Procalcitonin as indicator for successful surgical treatment of secondary peritonitis.

Methods: In 104 consecutive patients suffering from secondary peritonitis, Procalcitonin (PCT) serum levels were monitored on day one and two after the initial surgical elimination of the septic focus. The ratio of PCT on day one and two was calculated and correlated to the success of the initial surgical intervention. The initial procedure was considered inadequate if relaparotomies were necessary to eliminate the intra-abdominal infection.

Results: Using classification and regression tree (CART) analysis a cutoff could be calculated at 1.03 for the PCT ratio (d1/d2). Values below that cutoff indicated insufficient elimination of the septic source by the surgical procedure, while values above 1.03 indicated successful surgical treatment of the septic focus. In the study, patients' insufficient treatment could be detected with a specificity of 63.2% and a sensitivity of 95.3%.

Conclusions: The PCT ratio of day one and two after surgical treatment of an intra abdominal source of sepsis is a valuable parameter for deciding if relaparotomies are necessary for the elimination of the septic source.

006

Infection 2007; 35 (Suppl. II): 3

Rifampicin followed by ceftriaxone after one hour reduces the release of toxic and proinflammatory bacterial compounds by *Streptococcus pneumoniae* in comparison to treatment with the bacteriolytic ceftriaxone alone

*Spreer A (1), Stoltefaut V (1), Eiffert H (2), Nau R (1)

(1) Department of Neurology, University Göttingen, Germany,

(2) Medical Microbiology, University Göttingen, Germany

Introduction: Cell wall active antibiotics are important drugs for the treatment of severe bacterial infections such as sepsis or meningitis. However, bacterial cell lysis caused by bacteriolytic antibiotics results in an increased release of proinflammatory bacterial products. Cytoplasmic bacterial compounds including bacterial DNA and hemolysins (e.g. pneumolysin produced by *S. pneumoniae*) or constituents of the cell wall like endotoxin in Gram-negative and lipoteichoic acids in Gram-positive bacteria are important stimulators of the innate immune system. The β -lactam-induced release of bacterial products and the concomitant burst of inflammation can be attenuated by pretreatment with non-bacteriolytic antibiotics as shown for the sequential combination of rifampicin followed by addition of ceftriaxone 6 hours later (Spreer et al. AAC 2003; Nau and Eiffert FEMS Immunol Med Microbiol 2005).

Objective: Because of the high clinical relevance of early adequate and effective antibiotic treatment in sepsis and meningitis, concerns exist about rifampicin monotherapy for 6 hours. Therefore, we evaluated, whether a shorter period of rifampicin pretreatment followed by ceftriaxone also attenuates the release of proinflammatory bacterial compounds in comparison to ceftriaxone treatment alone.

Methods: We analysed in vitro the release of the cytoplasmic pathogenic factor pneumolysin in pneumococcal broth culture under sequential antibiotic treatment with rifampicin followed by ceftriaxone after 1 hour in comparison to ceftriaxone alone. In vivo, we evaluated in a rabbit model of pneumococcal meningitis the influence of the sequential antibiotic treatment on indicators of inflammation in the cerebrospinal fluid.

Results: In vitro, a 1 hour pretreatment of pneumococcal broth culture with rifampicin followed by addition of ceftriaxone reduced the release of pneumolysin into the culture supernatant. Likewise, in vivo,

an antibiotic treatment regimen including rifampicin for 1 hour followed by the addition of ceftriaxone significantly reduced early markers of neuroinflammation in the cerebrospinal fluid, i.e. prostaglandin E2 and total protein 2 hours after initiation of antibiotic therapy. The combined antibiotic treatment resulted in a slight decrease in the speed of bacterial killing, but was nevertheless rapidly bactericidal.

Conclusion: A pretreatment of only 1 hour with a protein-synthesis inhibiting antibiotic before initiation of β -lactam-therapy reduces the release of proinflammatory bacterial products in vitro and attenuates inflammation in vivo. This concept holds promise to reduce inflammation-associated damage in sepsis and meningitis. Therefore, antibiotic regimens relying on a sequential combination of bactericidal protein synthesis inhibitors and β -lactams should be evaluated for therapy of severe bacterial infections.

008

Infection 2007; 35 (Suppl. II): 4

Pro- and anti-inflammatory cytokine response due to blast or explosive trauma with or without secondary sepsis in combat casualties

*Surbatovic M, Jevtic M, Radakovic S, Filipovic N
Military Medical Academy, Belgrade, Serbia

Introduction: Despite the advances made in prehospital and critical care medicine, mortality rate after blast and explosive trauma continues to rise. Severe war injuries and secondary sepsis complicated with multiple organ dysfunction syndrome (MODS) are leading causes of death in combat casualties with mortality rate exceeding 50% (1). Outcome is not determined only by the severity of trauma and/or infection, but also by intensity of immuno-inflammatory response, which is essential for host defense, but uncontrolled leads to the MODS (2). Objective: Aim of this study was to assess the prognostic value of interleukin (IL)-12 interferon (IFN)-gamma, IL-1 receptor antagonist (IL-1ra) and transforming growth factor (TGF)-beta1 in combat casualties.

Methods: Fifty six casualties with severe trauma (blast and explosive) who developed sepsis (mortality 56%) and 20 casualties with same severity of trauma without sepsis (mortality 20%) were enrolled. Fifty five casualties developed multiple organ dysfunction syndrome (MODS), 36 died. Blood was drawn on the first day of trauma. Concentrations of IL-12, IFN-gamma, IL-1ra and TGF-beta1 were determined in plasma using ELISA assays. Results: Mean values of IL-12 were 1.8-fold (n.s.), IFN-gamma 360-fold ($p < 0.01$), and IL-1ra 3.7-fold ($p < 0.01$) higher and TGF-beta1 6.1-fold lower (n.s.) in trauma + sepsis group. Mean values of IL-12 were 1.2-fold (n.s.), IFN-gamma 175.5-fold (n.s.) and IL-1ra 6-fold ($p < 0.01$) higher and TGF-beta1 6-fold lower (n.s.) in MODS group. Mean values of IL-12 were 1.2-fold ($p < 0.05$), IFN-gamma 123-fold (n.s.), TGF-beta1 3-fold (n.s.) higher in survivors and IL-1ra 2.75-fold higher in non-survivors ($p < 0.01$).

Conclusion: In casualties enrolled in this study IFN-gamma and IL-1ra were most specific for diagnosis of combined trauma and sepsis. Concentrations of these cytokines were statistically significantly higher in that group compared to trauma group. Mortality was 2.5 times higher in trauma+sepsis group compared to trauma group although extent of trauma was similar in both groups. Most significant predictor of severity of clinical status and outcome was IL-1ra: higher concentrations predicted MODS and lethal outcome with statistically high significance. Unexpectedly, in our study concentrations of IL-12 and IFN-gamma were higher in MODS group and in survivors. Also, TGF-beta1 had no significance as predictor of severity and outcome what so ever. Further studies are needed to explain that. To our knowledge, next to nothing was done in research of cytokine response to combat trauma with or without sepsis in war time conditions.

References: 1. Surbatovic M, Filipovic N, Radakovic S et al. Mil Med 2007; 172(2):190-95 2. Reinhart K, Meisner M, Hartog C. Diagnosis of sepsis:

009

Infection 2007; 35 (Suppl. II): 4

Nephrotoxicity-associated costs in the treatment of invasive candidiasis with conventional Amphotericin B versus Caspofungin in Intensive Care Medicine in Germany

*Wissing H (1), Müller E (2), Winkler M (3), Weigand MA (4), Wagner A (5), Kaskel P (5), Hoppe-Tichy T (6)
(1) Klinik für Anästhesiologie, Intensivmedizin und Schmerztherapie, Klinikum der Johann-Wolfgang-Goethe-Universität, Frankfurt/Main, Germany. (2) Klinik für Anästhesiologie, Intensivmedizin und Schmerztherapie, Knappschafts-Krankenhaus Bochum-Langendreer Universitätsklinik, Bochum, Germany. (3) Klinik für Allgemein-Viszeral- und Transplantationschirurgie, Medizinische Hochschule Hannover, Hanover, Germany. (4) Klinik für Anästhesie, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany. (5) Medical Department, Outcomes Research, MSD SHARP & DOHME GMBH, Haar (Munich), Germany, (6) Apotheke, Universitätsklinikum Heidelberg, Heidelberg, Germany

Background: Drug-induced renal impairment can significantly prolong treatment and length of hospital stay. In a European study amphotericin B-associated nephrotoxicity (defined as $\geq 50\%$ increase in baseline creatinine level) increased mean length of hospital stay by 5.3 days (95% confidence interval [CI] 1.6; 9.1, $p = 0.004$), though $< 1\%$ of affected patients required renal replacement therapy (Ullmann et al., 2006). In a double-blind randomized clinical trial (RCT) in 224 patients with invasive candidiasis (80% APACHE II score ≤ 20), 24.8% of patients treated with conventional amphotericin B (AmBd) vs. 8.4% with caspofungin developed nephrotoxicity ($p = 0.02$; Mora-Duarte et al., 2002), though caspofungin was non-inferior to AmBd in terms of a favorable overall response (resolution of all symptoms and signs of candida infection and culture-confirmed eradication).

Objective: Based on the RCT data, to calculate the probability of nephrotoxicity and its complications with use of AmBd vs. caspofungin in patients with invasive candidiasis in intensive medical care in Germany.

Methods: Our mathematical model was based on (i) the RCT nephrotoxicity rates; (ii) the nephrotoxicity-associated prolongation of hospital stay (5.3 days); and (iii) the direct intensive care unit cost per day (€1156, inflation-adjusted for 2007; Billing et al., 2004). Bootstrapping and Monte Carlo simulations were performed (SAS 9.1.3, WinBUGS 1.4.1) based on individualized patient doses (in mg, price for bulk purchaser) and RCT treatment duration (caspofungin 12.1 days vs. AmBd 11.7 days).

Results: The number needed to treat for one patient to be harmed for the occurrence of nephrotoxicity with AmBd vs. caspofungin was 7 (95% CI 3; 17). Prolongation of hospital stay because of renal impairment was 0.87 days (95% CI 0.34; 1.39) for each patient treated with AmBd. The cost of treatment with AmBd was €640 inclusive of VAT. For each patient treated with AmBd the additional economic cost (hidden cost) due to AmBd-associated renal impairment was €1001 (95% CI €393; €1609), equivalent to 1.6 times the AmBd drug cost. The cost of treatment with caspofungin was €5315. For caspofungin there is a supplementary reimbursement (Zusatzentgelt), which amounts to €5151 in 2007.

Conclusion: Based on the RCT data, the observed treatment cost inclusive of hidden cost was €1642 (95% CI €1034; €2250) for each AmBd-treated patient with invasive candidiasis. Unlike with caspofungin (for which a supplementary reimbursement applies), use of AmBd and the hidden cost due to AmBd-associated side effects cannot be documented via the supplementary reimbursement (Zusatzentgelt) scheme and thus affect the base rate of the individual hospital. In the RCT caspofungin led to a substantial reduction in the risk for nephrotoxicity in patients with invasive candidiasis compared to AmBd while showing similar efficacy.

Reference: Novel and Conventional Parameters. Advances in sepsis 2001; 1(2):42-51

013

Infection 2007; 35 (Suppl. II): 5

Septic disseminated intravascular coagulation treated with the direct thrombin inhibitor argatroban

*Beyer J, Halbritter K, Weise M, Schellong S

Department of Angiology, University Dresden, Germany

Introduction: Septic DIC is characterized by systemic intravascular activation of coagulation, deposition and inadequate removal of fibrin, microvascular thrombosis and severe bleeding due to the excessive consumption of coagulation factors. Since excessive formation of thrombin is present in DIC direct thrombin inhibitors could be suitable to modulate the coagulation system. Argatroban selectively and reversibly inhibits free and clot-bound thrombin.

Objective: We tested the influence of the direct thrombin inhibitor argatroban on coagulation parameters in a patient with septic DIC.

Methods: A 75 year old woman (known CHF NYHA III, diabetes type II, chronic renal failure) presented with dyspnea and fever since 4 days due to urinary infection with *E. coli* (>10⁵ /ml; procalcitonine 61 ng ml⁻¹; CRP 317 mg L⁻¹) and pneumonia. Within hours septic MOD developed (APACHE II: 26; acute renal, hepatic and cardiopulmonary failure; deterioration of coagulation parameters). Increasing lactate and arterial hypotension required noradrenaline (0.4-1 mg h⁻¹) and dobutamine (15-30 mg h⁻¹). The initial platelet count (119 Gpt L⁻¹) decreased rapidly to 71 Gpt L⁻¹ and the ISTH score of DIC increased from 2 to 4. Substitution of 1500 IE AT III (45%) had no effect. In the palliative situation (age, poor quality of life; patients rejection of organ support) and poor prognosis, we decided on a healing attempt with argatroban to influence DIC and for thromboprophylaxis, starting with a bolus (0.5 mg IV), continuous infusion (0.5-1 mg h⁻¹) and dose adjustments according to aPTT and thrombin time. Results: Triple substitution with 500 IE AT III (0; 3 and 5 hours of argatroban) led to a rise (maximum 78%) and remained there. The platelet count stabilized at 64-74 Gpt L⁻¹. Argatroban was stopped after 20 hours and AT III and platelet count decreased (52 % and 47 Gpt L⁻¹ respectively). After resetting of argatroban (0.2-0.5 mg h⁻¹), platelets raised to 61 Gpt L⁻¹ and substitution of AT III resulted in an increase to 84%. Inflammation parameters decreased (CRP to 150 mg L⁻¹, PCT to 6.3 ng ml⁻¹), amelioration of the renal function was observed and noradrenalin could be discontinued. In spite of this success in modulation of coagulation system the cardiac and hepatic functions and the respiratory insufficiency deteriorated. The woman died on day 7 of cardio-respiratory failure.

Conclusion: Despite the fatal outcome for the multimorbid patient our findings point to a positive influence of direct thrombin inhibitor argatroban on septic DIC, since platelet count and AT III levels stabilized indicating an interruption of the activation-consumption-cascade. The unique pharmacological properties (close dose-response-relation, short half-life and hepatic elimination) allow application in acute or chronic renal failure which is often present in septic DIC. Intensive monitoring of aPTT, thrombin time, fibrinolytic activity and dose adjustments are necessary to reduce bleeding risk.

014

Infection 2007; 35 (Suppl. II): 5

Disseminated intravascular coagulation in neonates with congenital heart disease after cardiopulmonary bypass – treatment with antithrombin III

*Bartkowski R, Wojtalik M, Sniatkowska-Bartkowska A,

Henschke J, Wodzinski A, Westerski P, Mrówczyński W

Department of Pediatric Cardiac Surgery, University School of Medicine, Poznan, Poland

Background: Disseminated Intravascular Coagulation (DIC) is a multi-etiological syndrome with a broad spectrum of clinical manifestation.

Methods: The study was performed in the Department of Pediatric Cardiac Surgery Chair of the Cardiothoracic Surgery K. Marcinkowski University of Medical Sciences Poznań Poland between 1998 and 2004. The study included all 19 neonates with disseminated intravascular coagulation who were operated on congenital heart diseases in this period. We performed specific haematologic tests included: platelet count, PT and APTT, FDP, plasma fibrinogen level, antithrombin III level. The tests were performed and repeated daily until discharge from Intensive Care Unit or until the death of patients. All patients were properly managed for their original condition of admission. All patients with DIC received plasma substitution and heparine therapy. Eight patients received AT III substitution and eleven received only plasma substitution and heparine therapy. Statistical analysis was carried out using descriptive measures: mean, t-student, and chi square test.

Results: Results showed that among eight patients with DIC, who get AT III, only one died, and in the group without AT III (n=11) 6 patients died. There is statistically significant difference between these two groups (p<0.005). In all patients we noticed elevated fibrinogen degradation products (FDP), prolonged prothrombin time and partial thromboplastin time. In all patients we noticed also reduced plasma fibrinogen concentration and thrombocytopenia. We noticed the difference between platelet count in the group of patient who died and who survived. In the group of patients who died mean platelet count was lower than in the other group. In the first day of DIC the difference in mean platelets count between two groups is the smallest and it is getting higher each day and is the highest in the fifth day of DIC.

Conclusion: We can conclude that AT III substitution plays beneficial role in the treatment of DIC in neonates after cardiopulmonary bypass. The platelets count may predict mortality.

015

Infection 2007; 35 (Suppl. II): 5

Role of thioredoxin in sepsis

*Rosenhagen C, Hofer S, Nakamura H, Yodoi J, Martin E,

Breitkreutz R, Weigand MA

Department of Anaesthesiology, University Heidelberg, Germany

Introduction: Thioredoxin is known to be beneficial in acute inflammatory diseases because it blocks neutrophil chemotaxis into site of infection and its potent antioxidant properties.

Objective: To evaluate the role of TRX in sepsis.

Methods: we measured TRX levels in septic patients compared to healthy individuals. In an acute peritonitis mouse model (CLP) we examined the effect of neutralisation of TRX as well as treatment with TRX in two different dosages.

Results: We found that serum plasma levels of TRX are significantly higher in patients with sepsis compared to controls. Interestingly non-survivors show even higher TRX levels than survivors of sepsis. The CLP septic mouse model revealed 3 interesting results: 1) neutralisation of TRX impairs survival of septic mice, 2) high dosage treatment of TRX (1g/mouse) proved to be deleterious in the septic mouse model, 3) a low dose treatment of TRX (50µg/mouse) after CLP significantly enhanced the survival of mice.

Conclusion: These results demonstrate a critical role of TRX in the septic inflammatory response and suggest TRX as a potential therapeutic option for septic shock.

016

Infection 2007; 35 (Suppl. II): 6

Soluble RAGE – a new diagnostic and prognostic early sepsis marker?

*Bopp C (1), Hofer S (1), Bierhaus A (2), Martin E (1), Weigand MA (1)

(1) Department of Anaesthesiology, University Heidelberg, Germany, (2) Department of Medicine I and Clinical Chemistry, University Heidelberg, Germany

Background: The Receptor of advanced glycation endproducts (RAGE), as a member of the immunoglobulin superfamily, is a new inflammation perpetuating multiligand receptor. Recent data suggest (i) that RAGE perpetuate and amplify inflammation and (ii) that targeting this receptor might attenuate hyperinflammation. RAGE has soluble isoforms referred to as total sRAGE, that comprise the extracellular ligand-binding domain but are lacking the cytosolic and transmembrane domains. Actually, there is a lack of knowledge regarding the role of sRAGE in human sepsis. In view of these data, the aim of this work was (i) to investigate whether total pool of sRAGE is increased in plasma of septic patients and (ii) to assess the ability of sRAGE to predict mortality in patients with severe sepsis or septic shock.

Design: Observational clinical study.

Setting: Surgical intensive care unit of the University Hospital of Heidelberg, Germany.

Patients: Patients admitted to the intensive care unit over a 6-month period with clinical evidence of severe sepsis or septic shock and eight healthy controls.

Interventions: None.

Measurements and Main Results: Twenty-nine intensive care patients were enrolled in the study within the first 24 h after onset of severe sepsis or septic shock. Eight healthy controls served as controls. Plasma sRAGE concentrations were elevated in septic patients compared with healthy volunteers (1.764 ± 138 vs. 1.026 ± 177 , $p < 0.05$). Additionally, nonsurvivors after 28 days have had higher plasma sRAGE concentrations than survivors (2.302 ± 189 vs. 1.326 ± 112 , $p < 0.001$). ROC curve analysis of plasma sRAGE concentrations of septic patients showed a specificity of 75 % and a sensitivity of 84.6 % with 1596 pg/ml as cutoff.

Conclusion: This is the first study showing elevated plasma sRAGE concentrations in septic patients. Noteworthy, nonsurvivors had higher plasma sRAGE concentrations than survivors suggesting that sRAGE is related to severity and outcome of septic patients and may serve as a new sepsis marker.

Reference: 1. Liliensiek B, Weigand MA, Bierhaus A et al. Receptor for advanced glycation end products (RAGE) regulates sepsis but not the adaptive immune response. *J Clin Invest* 2004; 113: 1641-1650.

017

Infection 2007; 35 (Suppl. II): 6

Venous catheter bacteriological monitoring. Necessity or habit?

*Smuszkievicz P (1), Trojanowska I (1), Tomczak H (2)

(1) Department of Anesthesiology and Intensive Therapy, University Poznan, Poland (2) Department of Clinical Microbiology, University Poznan, Poland

Introduction: Catheter in the central venous system may be a source of generalized infection, due to its invasive nature, maintaining time and the patient life-threatening condition, causing prolongation of ICU and hospital stay, increasing mortality and the entire costs of treatment. In septic patients, especially if the bacteremia is suspected, intravenous catheter (IVC) is removed and tip is microbiologically examined.

Objective: We focused on the bacteriological analysis of IVC tips and the assessment of the frequency of the catheter-related blood infections (CR-BSI).

Our goal was also to estimate the economical and clinical purposefulness of microbiological IVC tips analysis.

Methods: The study was conducted at an adult 7-bed surgical/medical ICU between 1.X.2004 – 31.VII.2006. We retrospectively studied IVC tips and blood culture obtained from 120 septic patients (32,9% of all patients treated during that period). Maximal barrier precautions (MBP) – hand washing, sterile site preparation, use of hat, mask, gloves, gown and sterile dressing application were always implemented when introducing IVC. CR-BSI was defined according to the CDC guidelines – positive IVC tip cultures with an associated positive blood culture (collected from separate approach) and the exclusion of different source of bacteremia. Catheter removing was performed in case of prolonged time of catheter insertion or when the clinical state of patient indicated that catheter was the possible source of infection or when the inflammation of the puncture site was observed. Cost information was obtained from microbiological laboratory.

Results: We examined 238 IVC tips. In 123 IVC tips (51,6%) we didn't culture any pathogens. The rest of studied 115 (48,4%) IVC tips was positive. Of these 115 tips cultured, we isolated in 89 cases 1 type of pathogen, in 21 cases 2 and in 5 cases we found 3. Of 723 blood samples cultured, 181 (24,1%) indicated bacteremia and 168 was collected in the same time as IVC was removed. It was found that in 25 cases (3% of blood samples) taken from 20 patients, the same pathogens were isolated from IVC tips. We cultured: *Staph. epidermidis* (8 cases), *P. mirabilis* (5), *E. cloacae* (3), *Staph. aureus* (2), *Ps. aeruginosa* (2), *C. albicans* (2), *C. parapsilosis* (2), *Staph. haemolyticus* (1), *Ac. baumannii* (1) and *Ent. faecalis* (1) (in 2 cases 2 pathogenes were isolated). In 12 cases the same pathogens in blood and IVC tips were also cultured in other places – BAL (6), abdomen (3), bed-sore (2), urinary tract (1). In 13 cases (5,5% of tips) IVC's were the source of CR-BSI. Positive predictive value (PPV) and negative predictive value (NPV) amount to 11 % and 91 %, respectively. Sensitivity and specificity of the method is 54 % and 52 %, respectively. Time of maintenance of IVC amounted to 5-21 days. The laboratory cost of processing a negative tip culture was 11 €, while a positive tip culture 20 – 45 €, depending on the number of pathogens. The total cost of IVC tips microbiological examination amounted to about 4000 €.

Conclusions: 1. Bacterial colonisation of IVC is rarely responsible for CR-BSI. 2. Relatively low PPV (11 %) makes a IVC tips culture as a method of diagnosing CR-BSI useless. 3. Economically, routine bacterial monitoring of IVC tips seems to be not justifiable. 4. Taking all these into account we decided to discontinue the routine culture of IVC tip for the diagnosis of CR-BSI.

020

Infection 2007; 35 (Suppl. II): 6

Implementing a sepsis therapy concept conforming to regulatory requirements in a regular hospital setting and establishment of a local sepsis therapy network

Bier M

Klinik für Anästhesie, Intensivmedizin und Schmerztherapie, Kreiskrankenhaus Ottweiler

Introduction: In recent years, numerous innovations in diagnosis and therapy of sepsis have been integrated into normal hospital routine operations (1,2). Some remarkable research (3) has clearly demonstrated the possibility to significantly reduce lethality rates related to septic shocks through proper implementation of those evidence based innovations in sepsis therapy and diagnosis. However, this research was conducted in university hospital settings with above average resources compared to regular hospitals.

Methods: The remaining challenge has been to implement appropriate methods to enable proper treatment of patients in regular hospital settings conforming to regulatory requirements. That challenge could be successfully met by implementing the following methods: 1) Implementation of SOP's, Distribution and Publication on the Intranet, Training of Medical Personnel, Establishment of a qualified 'On-Call' Function 2) Cooperation with neighbouring hospitals spe-

cialized in sepsis therapy and diagnosis and specialized treatment/diagnosis centres to achieve the following goals: -Exchange of Information -Access to specialized and expensive pharmaceuticals (Xigris®, Voriconazole, etc.) and diagnostics (e.g. Septifast®) -Possibility of patient transfers if own hospital resources are not sufficient (e.g. Liver replacement therapy, specialized surgery) 3) Organization of training courses and implementation of quality review circles bringing together people from all hospital settings with different background and resources (e.g. local hospitals and specialized care centres)

Results: Through implementation of the methods described above, modern sepsis diagnosis and therapy concepts were successfully implemented in regular hospital settings conforming to regulatory requirements. Those methods mentioned above were further supplemented by the following measures: Improvement of logistics of specialized pharmaceuticals and diagnostics to facilitate access Implementation of Continuous renal replacement therapy, Weaning protocols and , education of the medical staff (nursery and physician), qualified On Call Function. The close cooperation with neighbouring intensive care centres resulted in a better utilization of available resources across all levels of patient care. Avoidance of unnecessary inter-hospital patient transfers, early transfer to appropriate hospital based on indication.

Conclusion: A modern and innovative therapy and diagnosis of sepsis conforming to guidelines is also possible in a regular hospital setting

References: 1) Surviving Sepsis Campaign Guidelines for Management of Severe Sepsis and Septic Shock. Crit Care Med 2004;32(3):858-873 2) Diagnosis and therapy of sepsis – Guidelines of the German Sepsis Society Inc. and the German Interdisciplinary Association for Intensive and Emergency Medicine. A&I 2006, 11: S157-S180 3) Implementation of an evidence-based “standard operating procedure” and outcome in septic shock.

021

Infection 2007; 35 (Suppl. II): 7

Correlation between procalcitonin, EuroSCORE and SOFA score during the intensive care unit course in patients subjected to cardiac surgery

*Maravic-Stojkovic V (1), Spasic T (1), Jovic M (1), Borzanovic M (1), Djukanovic B (1), Brunner D (2)
(1) Dedinje Cardiovascular Institute, Belgrade, Serbia,
(2) Interlight, Lozana, Swaziland

Introduction: We recently reported the clinical course of patients with good left ventricle ejection fraction (LVEF) by assessing Sequential Organ Failure (SOFA) score and procalcitonin (PCT) serum concentrations as prognostic markers early after cardiac surgery.

Methods: In this study we observed patients with systolic heart failure (HF), prospectively collecting clinical and laboratory data: B-type natriuretic peptide (BNP), PCT, and SOFA score in patients admitted to intensive care unit. Aim To examine whether PCT levels differ in patients with and without HF subjected to the open heart surgery. Methods In RCT study, 307 patients (age 59.84±9.55 years, 65.5% male) subjected to CABG, valve procedures or combined CABG and valve operations were divided in three groups: A (n=103) with echocardiographically estimated LVEF>30% on standard care as control; B (n=102) for procalcitonin guided antibiotic treatment; C (n=102) with LVEF<30% for BNP analysis. The predicted surgical risk was calculated according to EuroSCORE. The significance of the immune activation was assessed by PCT and C-reactive protein (CRP) serum concentrations. PCT and CRP were measured preoperatively, 6h, 24h and 48h postoperatively. BNP was measured preoperatively in patients with HF. SOFA score was assessed daily as SOFA1, SOFA2, SOFA3, SOFAAUC. Operation data, laboratory data, and clinical outcomes were observed.

Results: The average ICU stay in group A was 5.74±11.49 days, in group B 6.97±11.61 days, and 4.38±6.59 in group C. SOFA1 in A vs B vs C group was 1.31±1.67 vs 1.62±1.65 vs 5.07±2.67. SOFA2 was 0.97±1.56 vs 1.34±1.67 vs 4.89±2.60 days. SOFA3 was 0.63±1.11 vs 1.13±1.68 vs 4.33±2.52, respectively. In the group C plasma BNP levels

(mean: 667.65±1058.45 ng/L) correlated to EuroSCORE (r=0.409, p<0.001), PCT levels measured postoperatively (PO) on day 1 and 2 (r=0.3, p<0.005), and SOFA1,2, and 3 (r=0.404, p<0.01). EuroSCORE was in correlation with age, operation duration, CPB time, cross-clamping time, PCT measured on 2nd POD (r=0.199, p<0.05), ICU stay, SOFA1,2, and 3 score (r=0.46, p<0.001), and entire hospital stay. As BNP, it is in inverse correlation to LVEF (r=-0.23, p<0.02), the time waiting for operation, and CRP (r=-0.26, p<0.05). Postoperative hospital stay was 12.08±11.28 vs 12.93±10.73 vs 12.19± 9.29 days (p>0.05) in A vs B vs C group.

Conclusions: We have found that PCT concentrations measured 1-2 days after surgery was predictive for poor prognosis in patients with HF, as it was shown previously in patients without it. The SOFA score has predicting potential as well as PCT determined 24 hours after surgery.

024

Infection 2007; 35 (Suppl. II): 7

Multimodal management of sepsis due to the bronchial stump insufficiency – video presentation of the case

Rolle A (1), *Pereszlenyi A (1), Krassler J (2), Münchow St (1), Bis B (1)

(1) Department of Thoracic and Vascular Surgery, Centre of Pneumology and Thoracic Surgery, Coswig (Dresden), Germany, (2) Department of Anaesthesiology and Intensive Care, Zentralklinikum Suhl, SRH Kliniken AG, Suhl, Germany

Objective is to present a management of the sepsis due to bronchial stump insufficiency (BSI) empyema in a video demonstration. The film shows all the technical details of the management in a step by step manner.

Methods: A 64-years old female patient with a history of 3 years chronic postlobectomy empyema due to BSI and failed repeated chest tube drainages was referred to our Institution. Initially, due to her poor condition and sepsis, the empyema cavity (after the right lower lobectomy) was drained under the CT control; a wide-spectrum combination of antibiotics was applied too. After the significant detoxication but still poor performance status, the BSI was tried to close endoscopically by endovascular coils and tissucol. The stump closure was obtained for several days, but was reopened under the spontaneous breathing and coughing acts of the patient. The same procedure was than repeated with a fibrin-glue application, but BSI appeared after 7 days again. 3 months after these unsuccessful endobronchial interventions, the limited thoracoplasty together with myoplasty (m.latissimus dorsi, m. serratus ant.) and debridement, re-closure of the bronchial stump was than performed.

Results: Only after this active surgical access it was possible to discharge the patient with healed bronchial stump home. She left the hospital at the 35th postoperative day and today she is 20 Mo after the procedure without any signs of septic complication or recurrence of BSI.

Conclusion: The post-interventional empyema with the sepsis after the BSI is still the feared complication with high morbidity and mortality rate. Endobronchial procedures for BSI are reported with controversial results in the literature. Due to the poor performance status, sepsis of the patient, the definitive endoscopic closure of the bronchial stump can only be hardly obtained. The definitive solution – surgical BSI closure combined by the myoplasty -and limited thoracoplasty (after the detoxication and stabilization of the patient) – is the recommended treatment for sepsis due to BSI empyema.

025

Infection 2007; 35 (Suppl. II): 8

Einsatz von niedrig dosiertem Hydrocortison bei Patienten mit schwerer Sepsis und septischem Schock – eine Analyse aus Daten der Regensburger Sepsiskohorte 2003–2004

Robl F (1), Glück T (1), Ehrenstein B (1), Salzberger B (1), Schölmerich J (1), *Audebert F (2)

(1) Department of Internal Medicine I, University Regensburg, Germany (2) Department of Internal Medicine II, Hospital 'Barmherzige Brüder' Regensburg, Germany

Einleitung: Seit Jahrzehnten wird der Einsatz von Hydrocortison als supportive Therapiemaßnahme bei Patienten mit septischem Schock diskutiert. In bisher publizierten Studien (Briegel, Bolleart, Annane et al.) wurden in Subgruppen teilweise Vorteile für das Überleben und die Reversibilität der Katecholaminpflichtigkeit ohne gravierende Nebenwirkungen beschrieben. Die Ergebnisse der europäischen CORTICUS-Studie zum Einsatz von niedrig-dosiertem Hydrocortison beim septischen Schock stehen noch aus.

Fragestellung: Welche Auswirkungen hat der Einsatz von niedrig dosiertem Hydrocortison auf die 14- bzw. 30-Tages-Letalität, den Bedarf an Katecholaminen (Noradrenalin) und Insulin, sowie auf ausgewählte Laborparameter (Leukozyten, CRP, Natrium, Kalium, Serumglukose).

Methoden: 2003 bis 2004 wurden 88 Patienten mit schwerer Sepsis und septischem Schock in die Kohorte aufgenommen. Die Patienten wurden in 3 Gruppen stratifiziert: frühzeitige, adäquat dosierte (120mg-240mg/d) und rechtzeitige Hydrocortison-Gabe (HC+), keine Hydrocortison-Gabe (HC-) und inadäquate Hydrocortison-Gabe bzgl. Zeitpunkt, Dauer und Dosierung (HC+/-). Über 14 Tage wurden Laborparameter, Sepsis-Scores, BZ-Mittelwerte, Katecholamin- und Insulinbedarf dokumentiert sowie die 14- bzw. 30-Tages-Letalität erfasst und dann verglichen.

Ergebnisse: 33 Patienten erhielten eine adäquate (HC+), 25 erhielten keine (HC-) und 30 Patienten eine inadäquate (HC+/-) Gabe von Hydrocortison. Es zeigten sich bei Aufnahme in der Intensivstation zwischen diesen Gruppen keine stat. signifikanten Unterschiede beim durchschnittlichen Apache-II-Wert (gesamt 27,2; HC+ 28,7; HC- 25,4; HC+/- 27,1) und SAPS-II-Wert (gesamt 50,4; HC+ 53,4; HC- 47,8; HC+/- 49,2), dem BZ-Mittelwert (gesamt 150,2 mg/dl; HC+ 149,9 mg/dl; HC- 139,9 mg/dl; HC+/- 159,3 mg/dl), dem mittleren Insulinbedarf (HC+ 71,8 IE/24h; HC- 62,8 IE/24h; HC+/- 59,9 IE/24h) sowie im mittleren Noradrenalinbedarf (HC+ 32,5 mg/24h; HC- 15,3mg/24h; HC+/- 18,9 mg/24h). Mit 36,4% (HC+) vs. 24% (HC-) lag eine signifikant erhöhte 14-Tages-Letalität der HC+-Gruppe vor. Nach 30 Tagen war der Unterschied nicht mehr statistisch signifikant (48,5% vs. 32%). Natrium, Kalium, CRP und Leukozyten zeigten keine signifikanten Unterschiede. Eine nach SAP-II und APACHE-II Score adjustierte Analyse zeigt weiterhin eine erhöhte Letalität der HC+ Gruppe (nicht signifikant).

Zusammenfassung: Die 3 Behandlungsgruppen zeigten bzgl. der Erkrankungsschwere bei Einschluss sowie des initialen Katecholaminbedarfs keine signifikanten Unterschiede. Auffällig war eine statistisch signifikant erhöhte Frühletalität innerhalb der ersten 14 Tage in der Gruppe, die ab Tag1 mit niedrig dosiertem Hydrocortison behandelt wurde, gegenüber der anderen, die kein Hydrocortison erhielt. Somit konnte bzgl. Hydrocortisongabe kein Überlebensvorteil für Patienten mit schwerer Sepsis gezeigt werden. Der Einsatz von niedrig dosiertem Hydrocortison kann aus unserer Sicht aktuell nicht empfohlen werden.

027

Infection 2007; 35 (Suppl. II): 8

Computer aided adequate dosing of antibiotics for patients with instable renal function

*Baumgaertel M, Fessele K, Siemonsen S, Zahn R, Department of Internal Medicine 8, Hospital 'Nürnberg Süd', Nürnberg, Germany

Introduction: Antibiotics are too often used without consideration of the creatinine clearance. Especially nephrotoxic antibiotics can lead to a further decline of renal function and sometimes even to the need of renal replacement therapy.

Methods: For an easy calculation of the creatinine clearance and the adequate dosage of antibiotics and other drugs, we created the computer application www.intensivdocs.de, which is available everywhere through the internet. For most of the calculations and recommendations it is just necessary to enter the patients sex, age, height, weight and creatinine, using e.g. the Cockcroft-Gault equation for creatinine clearance.

Results: Within seconds an patient individualized dosing schedule recommendation is given. The application also warns the user of potential hazards, for instance restrictions of use with declining renal function or important interactions with comedication, e.g. QT-time-prolongation. The included module for antibiotic therapy gives the senior intensivist the possibility to tailor the antibiotic therapy to his ICU, taking prevalence of multi resistant bacteria and local resistance statistics into account. Thereby he can share his knowledge with the junior ICU staff and keep a high standard 24 hours/7 days a week.

Discussion: Especially for junior staff this application provides the necessary information fast and makes this information, through direct adjustment to the patient's needs, better applicable in an ICU environment than conventional means. The benefit for the patient is a safer, faster and more effective treatment.

Conclusion: This application is helpful in maximizing the benefit and reducing the risk of an antibiotic treatment for critically ill patients.

028

Infection 2007; 35 (Suppl. II): 8

Evaluation of clinical condition of the patients with severe sepsis treated with activated protein C: an observational study

*Lipinska-Gediga M (1), Adamik B (2)

(1) Department of Surgery, Medical University of Wroclaw, Poland, (2) Department of Anesthesiology and Intensive Therapy, Medical University of Wroclaw, Poland

Introduction: Early treatment with Drotrecogin alpha, recombinant human activated protein C (APC) with anti-inflammatory and anticoagulant properties may improve outcome in severe sepsis.

Objective: Our goal was to evaluate the data in patients with severe sepsis and septic shock who received APC.

Methods: The study was conducted at the medical and surgical intensive care unit of the 596-bed university hospital of Wroclaw, Poland. The study was observational, we retrospectively evaluated the clinical changes in patients' condition during 96 hour of APC administration. Patients' demographics, type of infection, and the number of organ failure were obtained at the baseline. APACHE II (Acute Physiology and Chronic Health Evaluation), SOFA (Sequential Organ Failure Assessment) score were calculated at baseline and after termination of APC infusion (96 hours). Other parameters like CRP (c-reactive protein), FDP (fibrinogen degradation products), fibrinogen level, WBC (white blood cells count) were recorded daily. Hospital mortality was primary outcome and patients were divided into two groups: nonsurvivors (NS, 5 patients) and survivors (S, 10 patients).

Results: APC was administered to 15 patients with mean organ failure 3,8(3-5 organs) in S group and 5,2(5-6 organs) in NS group. Survivors

had significantly lower mean SOFA score at baseline, comparing to non-survivors ($8 \pm 2,8$ vs. $12 \pm 1,6$). There was no significant difference in APACHE II score between groups at the entry to the study ($20,7 \pm 6,0$ in S group vs. $23,2 \pm 5,2$ in NS group); APACHE II score decreased to $9,8 \pm 4,5$ in S group and remained high $24,6 \pm 8,5$ in NS group at the end of APC administration. CRP level was elevated at baseline in both groups ($212,7 \pm 88,1$ in S and $226,1 \pm 124,7$ in NS). Marked decrease in CRP level was observed only in S group after 96h ($74,8 \pm 56,2$). Mean CRP value was also lower in NS group but the result was not statistically significant. At the entry to the study, elevated FDP level in all patients decreased in S group and increased in NS group (from 5852 ± 2346 to 2865 ± 1299 in S group vs. 6353 ± 1099 to 8623 ± 6706 in NS group at baseline and after 96h, respectively).

Conclusions: Both groups were characterized by similar clinical condition at baseline (APACHE II score), but in S group multiorgan failure was less severe (consistent with SOFA score). Delayed admission to ICU because of the late recognition of severe sepsis/septic shock was associated with a low efficiency of APC treatment and high mortality. Early recognition of severe sepsis and treatment with APC provides more benefit for the patients and is associated with a lower mortality.

030

Infection 2007; 35 (Suppl. II): 9

Increased circulating endothelial progenitor cells in septic patients: correlation with survival

Rafat N.[°] (1), *Hanusch C.[°] (1), Brinkkoetter P.T. (2), Schulte J (1), Zijlstra J. G. (4), van der Woude F. J. (2), van Ackern K. (1), Yard B. A. (2), Beck G. C. (1)

(1) Department of Anaesthesiology and Critical Care Medicine, Universital Hospital Mannheim, University Heidelberg, Germany (2) Department of Internal Medicine V, Universital Hospital Mannheim, University Heidelberg, Germany (3) Department of Internal Medicine, University Medical Centre, Groningen, Netherlands

[°] Neysan Rafat and Christine Hanusch have equally contributed to this work.

Introduction: Endothelial damage and detachment of endothelial cells is known to occur in septic patients. Thus, recruitment of circulating endothelial progenitor cells (cEPC's) to these lesions might have a beneficial effect on the clinical course in septic patients. Therefore, we were interested if EPC's, detected by flowcytometry, are increasingly mobilized during sepsis and if this mobilization is associated with clinical outcome.

Methods: Patients with (n=32) and without (n=15) sepsis and healthy volunteers (n=15) were included in this study. Peripheral blood mononuclear cells (PBMC) were isolated by Ficoll density gradient centrifugation and cEPC's were characterized by three-colour-fluorescence flowcytometry using antibodies against CD133, CD34 and vascular endothelial growth factor-Receptor (VEGF-R)-2. Serum-concentrations of vascular endothelial growth factor (VEGF), granulocyte macrophage-colony stimulating factor (GM-CSF) and erythropoietin (EPO) were determined by ELISA. Severity of sepsis was assessed according to APACHE II scoring.

Results: In septic patients the number of cEPC's was significantly higher than in non-septic ICU-patients ($p < 0,05$) and healthy controls ($p < 0,02$). Non-survivors (n=8), defined as death within 28 days after onset of sepsis, had significant lower numbers of cEPC's than survivors (n=24) ($p < 0,0001$). The number of cEPC's was correlated with survival in septic patients. Serum-VEGF-concentrations were significantly higher in septic patients compared to non-septic ICU-patients and healthy controls ($p < 0,01$), and correlated with the cEPC's numbers ($p < 0,0001$). Similar findings were observed for GM-CSF and EPO.

Conclusions: Our data suggest that cEPC enumeration in peripheral blood of septic patients might be a valuable marker to assess the clinical outcome in these patients.

031

Infection 2007; 35 (Suppl. II): 9

Fulminant Streptococcal Toxic Shock-Like Syndrome (STSLS): a case report

*Schott M, Freudenhammer, M., Jantzen JP

Department of Anaesthesiology and Intensive Care, Hospital Hannover Nordstadt, Klinikum Region Hannover, Germany

Introduction: Streptococcal Toxic Shock-Like Syndrome (STSLS) is a rare but life-threatening complication of Group A Streptococcal infection. Based on case report, we present the clinical course of the disease as well as current recommendations for treatment and differential diagnoses.

Case report: A 41 year-old white female was admitted to our clinic with extreme lower back pain. The primary challenge was presented by rapidly developing septic shock and associated multi-organ failure. The presence of a long-dwelling vaginal tampon and an intra-uterine device (IUD) together with a recently-developed erythema suggested a diagnosis of Toxic-Shock-Syndrome. The initial antimicrobial treatment with meropenem and metronidazole was subsequently adjusted to include clindamycin and ceftriaxone. Despite aggressive therapy with volume replacement, catecholamines, hydrocortisone and activated protein C, the patient died 17 hours after admission from therapy-resistant septic shock. An autopsy revealed severe bacterial sepsis. Postmortem microbiological blood cultures and vaginal swabs were positive for group A Streptococci and confirmed the working diagnosis of Streptococcal Toxic Shock-Like Syndrome.

Conclusion: Toxic Shock Syndrome is an important differential diagnosis for fulminant sepsis. The rapid progression and high mortality rate, particularly for the streptococcal shock syndrome, necessitate a rapid search for and radical removal of the infective focus as well as targeted antimicrobial, haemodynamic and supportive therapy of the sepsis.

034

Infection 2007; 35 (Suppl. II): 9

Sepsis-Diagnostik mit Hilfe des SeptiFast®-Systems: Erfahrungen auf einer internistischen Intensivstation

*Sahre H (1), Vogel S (2), Park J-W (1), Weber S (2)

(1) Department of Internal Medicine I, Klinikum Hoyerswerda gGmbH, Germany, (2) Institut der mikrobiologie and Hospital Hygiene, Klinikum Hoyerswerda gGmbH, Germany

Introduction: SeptiFast® ist ein neues diagnostisches System zur Schnell diagnose von Bakterien-DNA im Patientenblut. Dabei wird eine Multiplex-PCR gegen 25 verschiedene Erreger (grampositive und gramnegative Bakterien, Pilze) in Realtime (Light-Cycler) durchgeführt. Nach 4 – 6 h sind Ergebnisse verfügbar.

Objective: In der vorliegenden Studie untersuchen wir den Nutzen des Nachweises für eine internistische Intensivstation eines Schwerpunkt-krankenhauses.

Methods: Multiplex-Realtime PCR mit dem Light-Cycler-System

Results: Insgesamt untersuchten wir 119 Proben von 73 Patienten im Zeitraum von Dezember 2006 bis Juni 2007. Bei 32 (26,9) Proben war kein Vergleichsmaterial (Blutkulturen) vorhanden, sodass diese von der Auswertung ausgeschlossen wurden. Bei den restlichen 87 Proben war das Ergebnis im SeptiFast® in 58 Fällen (66,7%) gleich mit der Kultur, in 14 Fällen (16,1 %) nur im SeptiFast® und in 3 Fällen (3,4 %) nur in der Kultur erzielt worden. Bei 12 Untersuchungen (13,8 %) erhielten wir ein divergierendes Ergebnis. Bei den Gruppen, bei denen nur der SeptiFast® oder beide ein Ergebnis erbrachten, kam es

in der Therapie zu einer früheren Anpassung der Antibiotikatherapie oder zum Beginn einer Therapie bei bisher nicht bekanntem Erreger (z.B. Pilze). In der Gruppe mit positiver Blutkultur wurden in unserer Studie nur Bakterien isoliert, die für das klinische Bild des Patienten keine Relevanz hatten (1 x *Bacillus* sp., 1 x *Propionibacterium* sp., 1 x *S. epidermidis*). Die Wertung eines divergierenden Ergebnisses (SeptiFast® entspricht nicht der Kultur) ist zur Zeit noch unklar und bedarf weiterer Studien.

Conclusion: Die Multiplex-PCR führt zu einer beschleunigten Diagnostik sepsis-relevanter Erreger. In unserer Studie konnte deswegen eine Therapieanpassung früher und effizienter erfolgen. Damit ergab sich klinisch ein Vorteil gegenüber der konventionellen Blutkulturdiagnostik. Zum gegenwärtigen Zeitpunkt ist der SeptiFast® nach unserer Studie eine sinnvolle Ergänzung der konventionellen Sepsisdiagnostik, er kann diese aber keinesfalls ersetzen. Ein Teil der SeptiFast®-Befunde deckt sich nicht mit Blutkulturbefunden, wobei die Relevanz dieser Ergebnisse noch unklar ist. Um die finanziellen Vorteile des SeptiFast®-Systems zu untersuchen, bedarf es größerer Studien mit entsprechenden Vergleichsgruppen.

035

Infection 2007; 35 (Suppl. II): 10

Implementation of severe sepsis bundles by the Intensive Care Units in Poland – preliminary results

*Kübler A, Duszyńska W, Barteczko B, Grotowska M
Department of Anaesthesiology and Intensive Care, The Medical University of Wrocław, Poland

Introduction: Surviving Sepsis Campaign (SSC) proposed 6-hour resuscitation and 24-hour management bundles as a measure to reduce mortality in severe sepsis. This approach was recently adopted for use in Intensive Care Units (ICUs) in Poland and the first results of bundles' application are presented.

Methods: At the beginning of January 2007 the sepsis bundle protocol was introduced as an additional option to the website-based severe sepsis registry program functioning in Poland since 2003.

Results: Till June 2007 the first 120 cases of bundles' implementation were registered. The rate of compliance with the 6-hour resuscitation bundle was 11% and with the 24-hour management bundle was 36%. Despite the small number of patients and low compliance rate there was observed a distinct difference in mortality between compliant and non-compliant groups. The mortality rate in patients compliant with the resuscitation bundle was 38% and non-compliant 50%. Respectively, mortality in patients compliant and non-compliant with the management bundle was 37% and 55%.

Conclusion: The preliminary results of sepsis bundles' implementation showed marked difference in mortality between compliant and non-compliant groups, that was beyond expectation for such a small population of patients. The bundle approach is now recommended for routine use in patients treated for severe sepsis in ICUs. In the nearest future it should provide more comprehensive and detailed analysis of data

036

Infection 2007; 35 (Suppl. II): 10

Four years of severe sepsis surveillance in Poland

*Kübler A, Durek G, Duszyńska W, Barteczko B, Grotowska M
Department of Anaesthesiology and Intensive Care, Medical University of Wrocław, Poland

Introduction: Severe sepsis is a common clinical syndrome treated in Intensive Care Units (ICUs) and the main cause of ICU mortality. Long-term epidemiological studies of severe sepsis are recommended to assess the clinical course of this syndrome and the efficacy of new diagnostic and therapeutic measures.

Methods: In 2003 there was introduced in Poland a voluntary registration of severe sepsis cases treated in ICUs. Registered data include: type and site of infection, severity of disease, methods and results of the therapy. After 4 years the results of surveillance were analysed.

Results: There were 3984 cases of sepsis registered till May 2007. The mean age was 54 years. Mean APACHE II score on admission was 24 and mean one-day TISS-24 value was 38. Severe sepsis developed more frequently in men (58) and due to surgical disorders (54%). Mean period of stay in the ICU was 17 days and average mortality 54%. Comparison of the results obtained in successive years showed high consistency of data including age, gender, severity scores and type of infection. The progressive decline in mortality was observed: 2004-56%, 2005-51% and 2006-47%. The decrease in the relative rate of mortality during last 3 years amounted to 16%.

Conclusion: In the population of patients with severe sepsis in the last 3 years there was observed a marked decrease in mortality (16%). It can hardly be denied that the improvement can result from earlier diagnosis of the syndrome and better adherence to the severe sepsis therapy guidelines.

037

Infection 2007; 35 (Suppl. II): 10

Severe respiratory failure due to an abdominal compartment syndrome and immediate decompressive laparotomy as a lifesaving intervention in a patient with acute pancreatitis

Völz J-E, Möller M, Hansen C, Neuzner J
Department of Internal Medicine II (Cardiology and Internal Intensive Care, Hospital Kassel, Germany

Introduction: Abdominal hypertension and its severe form of abdominal compartment syndrome (ACS), are characterized by elevated intraabdominal pressure (IAP) leading to renal, cardio-pulmonary and intestinal dysfunction, thereby promoting multiorgan failure (MOF) and death. We report on a 45 year-old male who were presented with an acute necrotizing pancreatitis, complicated by a life-threatening respiratory insufficiency and MOF due to the development of massive abdominal congestion with markedly elevated IAP after vigorous volume resuscitation. Surgical abdominal decompression (AD) immediately reversed abdominal hypertension, leading to improved respiratory function as well as hemodynamic parameters and renal function.

Objectives: A Report of the lifesaving effect due to decompressive laparotomy (DL) in a near-suffocation caused by ACS.

Methods: Case report

Results: A 45-years-old overweight man (BMI 32) was referred from another hospital because of a necrotizing pancreatitis with MOF. He had been admitted two days before with diffuse abdominal pain, nausea and vomiting. CT-scan showed edematous and necrotizing pancreatitis. He rapidly deteriorated. Massive volume resuscitation (20l/20h) were needed. Increasing doses of catecholamines (NA 3 mg/h; Dob 30 mg/h) were given. He was intubated. The Ventilation was insufficient, even with high inspiratory pressure by severely impaired oxygenation (see list). On admission he was analgosedated and paralyzed. HR was 150/min, MAP 73 mmHg. Lung sounds were only audible in the upper third of the chest. The abdomen was tense and distended with no bowel sounds. Apache II score was 28. Pulmonary function was severely impaired FiO2 1.0; PaO2/FiO2 ratio 62 (see list). The patient was anuric. Ultrasound (US) of the thorax and abdomen revealed an elevated diaphragm with bilateral atelectasis and minimal pleural effusion but only minimal ascites in the pelvic region. The IAP was markedly elevated (29 mmHg, urinary bladder). The diagnosis of ACS with predominant respiratory failure due to pulmonary compression by the displaced diaphragm was made. Decompressive laparotomy (DL) with a temporary abdominal closure was performed. The IAP (14 mmHg) as well as pulmonary function (PaO2/FiO2 ratio: 133) improved immediately (see list). Catecholamines could be reduced

(NA 0,6 mg/h, Dob 30 mg/h) and urine output started during the next few hours again. Lung-protective ventilator settings could be instituted.

Conclusion: Multiple organ dysfunction is a wellknown sequella of ACS. In our case a markedly elevated IAP resulted in compression of the lung with progressive reduction of total lung capacity and severe respiratory failure. The immediate DL was lifesaving in this case und could be considered in pat. with a severe ACS. List: Admission: IAP 29 mmHg, pCo₂: 60,4 mmHg, FIO₂: 1,0, PaO₂/FIO₂: 62, Insp. pressure: 50-38-20, I:E:1:2, TV: 500 ml. 4 h post OP: IAP 14 mmHg, pCo₂: 38,8 mmHg, FIO₂: 0,6, PaO₂/FIO₂: 133, Insp. pressure: 36-27-20, I:E:1:1, TV: 460 ml.

038

Infection 2007; 35 (Suppl. II): 11

RIFLE – a newly developed classification for patients with renal injury

Albert T. Schumann M

Department of Anaesthesiology and Intensive Care, University Leipzig, Germany

Introduction: Acute Renal Failure is a frequent complication in septic patients and contributes significantly to the mortality of sepsis. To date, there are more than 30 definitions of Acute Renal Failure, resulting in a great variety of published data concerning incidence and outcome. In order to establish a uniform definition, the Acute Dialysis Quality Initiative developed the RIFLE classification that defines several grades of increasing severity of kidney dysfunction.

Objective: After describing in detail how to classify according to RIFLE we assigned it to our patients to compare between the several classes (Risk, Injury, Failure) with regard to admission to ICU, start of the Renal Replacement Therapy, ICU mortality, development of the patients (changing the classes) during their ICU stay, non-renal SOFA-Score as a measure for severity of illness and to length of ICU stay.

Methods: We analysed retrospectively the electronic hospital database of 55 adult patients that were admitted to the surgical and anaesthesiological ICU of the University of Leipzig Medical Center and who had got Renal Replacement Therapy.

Results: We found out that there is a connection between the RIFLE classes (as a measure of severity of renal dysfunction) and mortality on ICU although the non-renal SOFA-Score was similar for all classes. Patients without Acute Renal Failure showed a lower non-renal SOFA-Score than patients who had developed a renal dysfunction. Furthermore there was no association between age or length on ICU stay and the separated RIFLE classes.

Conclusion: RIFLE is a newly developed classification for kidney dysfunction which is – although the non-renal dysfunctions are similar – associated with an increased mortality rate on ICU.

041

Infection 2007; 35 (Suppl. II): 11

Comparison of two training tools for severe sepsis: simulation training versus papercase scenario

*Bröhl K, Kersten R, Marz S, Haase U, Schröder T
Institutionen

Introduction: Improved training of professionals in intensive care medicine [1, 2] is an important element in the reduction of the high sepsis mortality rate (30-50%), along with basic science and the search for new therapeutic strategies. To reach this aim, the Berlin Simulation Centre has introduced a training- tool to improve diagnose and therapy of severe sepsis, over the last four years. During the 50th Kasseler Symposium (June 2007) we performed a sepsis simulation training, with some modifications to the normal setting [3]. Goals of the

workshop were to increase the knowledge in the detection of a beginning sepsis, evidence based therapy of severe sepsis and septic shock as well as the efficient implementation of scientific findings in the clinical routine. The intention of this survey was to compare the gain of knowledge between sepsis simulation training and a papercase scenario of septic patients.

Methods: The 24 participants were divided into two groups (A and B), and assigned to an instructor. A patient history was presented to each group: While three participants out of group A had the opportunity to actively perform on the simulator, group B was given with a papercase scenario. After 45 minutes group A and B switched. Questionnaires to assess the gain of knowledge were performed in advance of the course, before switching groups and after the entire course. Each questionnaire consisted of nine identical questions to capture and compare learning outcomes after simulation training and papercase scenario.

Results: Out of 24 participants 14 (seven of each group) completed their forms correctly and were analysed. There was no significant difference between the groups concerning epidemiological data such as age, sex or working experience. All candidates showed a significant gain in knowledge after the workshop. For both groups the improvement of tested knowledge after the simulation training was significant ($p < 0,05$); no significance could be established for the papercase scenarios in neither groups ($p > 0,05$).

Conclusion: The survey showed clear limitations in the gain of knowledge due to the short duration of the workshop and the small sample-size. The simulator supported training of intensive care professionals is an effective way to improve knowledge and superior to the results of a papercase scenario in the setting of a three hour workshop.

043

Infection 2007; 35 (Suppl. II): 11

Successful treatment of severe streptococcal toxic shock syndrome

*Schrauzer T, Forster C, Willam C, Eckardt KU, John S

Department of Internal Medicine IV (Nephrology and Hypertensiology), University Erlangen-Nürnberg, Germany

The streptococcal toxic shock syndrome (STSS) is a rare disease with an increasing incidence and high mortality caused by beta-hemolytic streptococci. A 35-year-old intubated patient with septic shock and multi-organ failure was transferred to our intensive care unit after 20 hours of intensive care management in an external hospital. For the previous week he had complained about influenza-like symptoms. He had a history of hypertension and right-sided nephrectomy after reflux nephropathy. On admission we saw an obese patient (BMI 38kg/m²) with a blood pressure of 80/40 mm Hg (MAP 50 mm Hg), a sinus tachycardia of 200 bpm, a respiratory rate of 30/min and a rectal temperature of 41.3°C. Beside acrocyanosis he showed marked reticular livedo of the lower limbs with progressing into extensive desquamation of the skin of both hands and lower limbs. He had septic shock and consecutive renal and liver dysfunction, disseminated intravascular coagulation, rhabdomyolysis and acute respiratory distress syndrome. Initial hemodynamic monitoring (PiCCO) revealed markedly reduced systemic vascular resistance index (SVRI 699 DSm²/cm⁵) and elevated cardiac index (CI 5.26 L/min/m²) while treatment with noradrenaline (up to 15mg/h) and dobutamine (up to 20mg/h). After aggressive fluid resuscitation of 50L crystalloids during the first 72 hours (increase in body weight from 110 to 160kg !), high-dose catecholamines, systemic cooling by continued venovenous hemofiltration (CVVHF), administration of activated protein C and antibiotic treatment with meropenem and moxifloxacin (according to the guidelines of the German sepsis society) the patient's condition stabilized. Administration of catecholamines could be finished on day 6. Additionally a negative water balance could be reached beginning on day 7. Blood cultures were positive for group A streptococci (M type 3.1, T type 3). Penicillin was added following antibiotic testing. PCR confirmed expression of super antigens SpeA (ETA), Spe F and d Ssa. According to the criteria of the "Working Group on Severe Strepto-

coccal Infections" STSS was diagnosed. Despite extensive diagnostics no portal of entry could be identified. Following extubation on day 17, renal and liver function recovered from day 29. Amputations could be avoided. The patient was transferred to rehabilitation center on day 41. Because of the severe endothelial dysfunction in patients with STSS, an early and aggressive fluid resuscitation is essentially.

044

Infection 2007; 35 (Suppl. II): 12

Sepsis and Procalcitonin: Different initial PCT-levels depending on sepsis focus

Fricke R, *Kubitza S

Department of Intensive Care Medicine, HELIOS Klinikum Berlin-Buch, Berlin, Germany

Introduction: Clinical experience with septic medical-ICU patients suggests that PCT-levels not only depend from severity of illness but also from infectious focus and microbial agent.

Method: Retrospective analysis from patients with septic ICD-code (A40.0-A41.9) documented in the hospital information system. From all patients fulfilling the ACCP/SCCM criteria for sepsis, initial SAPS II and daily SOFA-scores were calculated, initial and maximum CrP, daily PCT-levels and proven or suspected infectious focus and microbial agent are recorded. Linear regression analysis was calculated for different combinations.

Results: From a total of 80 patients retrieved from the hospital information system, 40 were fulfilling the ACCP/SCCM-sepsis-criteria whereof 20 until now were susceptible for further analysis. In 6 of them a urogenital or a bronchopulmonary focus of sepsis could be detected, in 8 patients an abdominal or no specified focus could be detected. Gram-stain positive or negative bacterial specimens were found in 11 and 8 patients respectively. While no correlation could be detected between initial PCT-levels and maximum SOFA-score depending on gram-stain-character ($R^2 < 0.2$), a good correlation could be found in this relationship depending on sepsis-focus. Patients with a bronchopulmonary focus had much higher severity of illness with lower initial PCT-levels ($y = 0.1319x + 9.5718$, $R^2 = 0.9557$) than patients with urogenital focus ($y = 0.026x + 4.2097$, $R^2 = 0.7755$). For patients with suspected abdominal focus of sepsis, no correlation could be detected. Initial SAPS II-score and maximum SOFA-score also had a sufficient correlation but could not differentiate between pulmonary and urogenital focus. No correlation could be found between initial PCT and SAPS II and between initial and maximum CrP and maximum SOFA.

Conclusion: Initial PCT-levels may probably allow to early predict the severity of illness depending on the suspected focus of infection. Further prospective evaluations are necessary to confirm this hypothesis.

045

Infection 2007; 35 (Suppl. II): 12

The activity of C-Protein in infants undergoing correction of congenital heart disease with the use of cardiopulmonary bypass – pilot study

*Szpecht D, Wojtalik M, Mrówczyński W, Poprawski G, Kempka-Dobra E

Department of Pediatric Cardiac Surgery, University School of Medicine, Poznan, Poland

Subject: The protein C can influence the postoperative outcome in neonates undergoing correction of congenital heart disease. The aim of this study was to: 1. analyze the activity of protein C among neonates undergoing CPB, 2. analyze the influence of pre- and perioperative infection risk factors on C protein activity changes, and 3. in-

vestigate C protein activities in children with and without postoperative infection.

Material and methods: Twelve infants were operated with the use of CPB in deep and moderate hypothermia. The blood samples were drawn in 6 time points: before general anaesthesia, 5 minutes after beginning of CPB, 5 minutes before disconnecting from CPB, 15 minutes after administering protamine, in second and third postoperative day. The activity of C protein was measured with CoagChrom 3003 Coagulometer.

Results: The mean baseline C protein activity was below the norm, decreased immediately after the institution of CPB and gradually raised to reach starting values at the end of observation. There changes were significant ($p < 0.0001$). There were not significant statistical differences among children with and without infection risk factors. Similarly present infection did not influence C protein activity.

Conclusions: CPB exacerbates the preoperative deficit of C protein. The subsequent recovery is visible, however C protein activities do not reach normal ranges for the age group.

047

Infection 2007; 35 (Suppl. II): 12

Clinical application of SeptiFast, a PCR method for the detection of bacteraemia, in intensive care patients

Klemm M (1), Prinz M (1), Nowak A (2), Morgenstern T (2), Meisner M (3), Rothe KF (2), *Demant T (1)

(1) Institut of Clinical Chemistry and Laboratory Medicine, Hospital Dresden-Friedrichstadt, Germany, (2) Department of Anaesthesiology and Intensive Care, Hospital Dresden-Friedrichstadt, Germany, (3) Department of Anaesthesiology and Intensive Care, Hospital Dresden-Neustadt, Germany

Introduction: Sepsis is a major complication in critically ill patients with a mortality rate as high as 50%. Crucial for successful treatment is the rapid identification of the causative agent in order to facilitate specific antimicrobial therapy.

Objective: We evaluated a new PCR system designed to detect clinically relevant pathogens causing septicaemia in intensive care patients. Results were compared with conventional blood cultures (BC) and correlated with procalcitonin (PCT) as a clinical marker of sepsis.

Methods: Patients enrolled in this study had evidence for a new focus of infection and a clinical suspicion for sepsis. Inclusion criteria were fever ($> 38^\circ\text{C}$) or hypothermia ($< 36^\circ\text{C}$) and additionally leucocytosis ($> 12.000/\mu\text{l}$), leucopenia ($< 4.000/\mu\text{l}$), leucocyte left shift ($> 10\%$), tachycardia ($> 90/\text{min}$), tachypnoea ($> 20/\text{min}$) or hyperventilation ($\text{pCO}_2 < 33 \text{ mmHg}$). Blood ($2 \times 10 \text{ ml}$) for BC was incubated under aerobic and anaerobic conditions using an automated CO2 detection system (BacTec, BD). For PCR an EDTA-blood sample ($2 \times 3.5 \text{ ml}$) was drawn simultaneously and analysed using the SeptiFast-test on the LightCycler® system for real-time PCR (Roche Diagnostics.).

Results: 56 pairs of samples from 44 individuals were analysed. PCR results were usually available on the same day (min. 6.5 h) while definitive BC results were reported at least two days later. BC and PCR were both negative in 30 (group A) and both positive in 13 cases (group B). Pathogens detected by both methods were K. pneumoniae, E. coli, S. aureus, E. faecium and C. albicans. In addition, K. pneumoniae, E. coli and E. cloacae were determined by PCR but not by BC. In 8 patients PCR was negative but BC was positive (group C) with 4 samples contaminated with dermal Streptococci deliberately not reported by the SeptiFast interpretive software. In another 3 cases two BC were drawn simultaneously with only one turning positive, in the remaining one patient E. faecium was detected by BC but not by PCR. In 5 patients PCR revealed E. faecium, E. faecalis, E. coli or K. pneumoniae while the corresponding BC remained sterile (group D). PCT concentrations (median) were markedly higher in positive compared to negative PCRs ($p = 0.0001$) and less so in positive compared to negative BCs ($p = 0.038$). They were also significantly higher in group B vs. A ($16.4 \text{ vs. } 0.5 \text{ ng/ml}$, $p = 0.001$) and in group D vs. C ($4.8 \text{ vs. } 0.8 \text{ ng/ml}$, $p = 0.040$). Conclusions: SeptiFast results were provided in less

than 15% of the time required for BC. They were in excellent agreement with BC in 43 cases (77%). Discrepant results occurred in 5 patients with false-negative BC and unequivocally in one possibly in 3 subjects with false-negative PCR.

Conclusions: In patients with pathogens detected by PCR PCT was significantly increased independent of BC results indicating clinically relevant bacteraemia. In contrast, patients with positive BC but negative PCR had only marginally increased PCT compatible with a lesser bacterial burden.

050

Infection 2007; 35 (Suppl. II): 13

Heat shock protein 70 (HSP 70) levels as prognostic marker of septic patients

*Hoffmann U, Klank D, Holzinger T, Liebetrau C, Lang S, Brückmann M, Borggreffe M
1st Department of Medicine, Faculty of Clinical Medicine
Mannheim, University of Heidelberg, Germany

Introduction: Inducible heat shock protein 70 (HSP 70) is a stress protein whose expression is upregulated when the cell or organism is placed under conditions of stress. HSP 70 is essential for cellular recovery, survival and maintenance of normal cellular function. It is also a molecular chaperone that prevents protein aggregation and refolds damaged proteins in response to cellular stress caused by environmental insults, pathogens and disease. Heat-shock proteins are synthesized by cells in response to heat, as well as to various other stressful stimuli associated with severe sepsis. We measured levels of HSP 70 in septic patients treated with and without Drotrecogin alfa (activated).

Methods: We determined levels of HSP 70 in the serum of septic patients treated with Drotrecogin alfa (activated) (n=7) and septic patients without treatment (n=6) on day 1 of sepsis with ELISA-method. Statistical analysis was performed with ANOVA.

Results: 5 Patients with treatment of Drotrecogin alfa (activated) died (71,4%) and 3 patients without Drotrecogin alfa (activated) treatment (50 %). We measured significantly higher levels of HSP 70 on day 1 in non-survivors (mean=24,378 ng/ml $\pm$ SEM=8,618) compared to survivors (mean=4,172 ng/ml $\pm$ SEM=1,703; $p<0.001$). Levels of HSP70 were also higher in non-survivors treated with Drotrecogin (alfa) activated compared to non-survivors not treated with Drotrecogin (alfa) activated.

Conclusions: HSP 70 levels are elevated in the early course of septic patients. Non-Survivors of sepsis had higher levels of HSP 70 than survivors. Furthermore, HSP 70 may be related to the prognosis and clinical outcome of septic patients. Further in vitro and in vivo investigations are needed to confirm these results.

053

Infection 2007; 35 (Suppl. II): 13

Decline of ADAMTS13 activity depends on the severity of the inflammatory response

Bockmeyer CL (1), Lösche W (1), Budde U (2), Schneppenheim R (3), Kentouche K (4), Bauer M (1), Reinhart K (1), Brunkhorst FM (1), *Claus RA (1)

(1) Department of Anaesthesiology and Intensive Care Medicine, Friedrich-Schiller-University Jena, Germany, (2) Lab-Association Prof. Arndt and Partners, Coagulation Laboratory, Hamburg, Germany, (3) Department of Pediatric Hematology and Oncology, Children's University Hospital Hamburg-Eppendorf, Germany, (4) Department of Pediatrics, University Jena, Germany.

Background and Aims: ADAMTS13 is a protease that specifically acts on the multimeric von Willebrand factor (VWF). Congenital or acquired deficiency of ADAMTS13 can result in the appearance of ultra-large VWF multimers with subsequent activation of circulating

blood platelets, thrombus formation in the microcirculation and (multi) organ failure. There is some evidence that systemic inflammation also results in restriction of microcirculation and that the development of multiple organ dysfunction syndrome (MODS) is associated with decreased ADAMTS13 activity and increased VWF levels.

Methods: In a prospective study on patients with various degrees of systemic inflammation we aim to provide evidence for a crucial role of an imbalance between ADAMTS13 activity and VWF level in the development of inflammation induced MODS. Four groups of patients were studied: healthy volunteers with moderate SIRS after strenuous physical exercise, heart surgery patients with low risk for systemic inflammation and MOF, heart surgery patients with SIRS and moderate MODS, and patients with severe sepsis or septic shock. Consecutive plasma samples were analysed for ADAMTS13 activity and VWF and compared with various markers of systemic inflammation and activated coagulation as well as with parameters of describing the extent of MODS.

Results: After physical stress, we found a marked decrease of ADAMTS13. With the exception of pre-operative values, in all patient groups ADAMTS13 was found to be significantly decreased compared to values measured in age- and sex-matched controls. The decrease in ADAMTS13 activity was related to the extend of systemic inflammation and was most pronounced in patients with MODS, specifically in patients who died in septic shock. Exemplarily we demonstrate an association of low ADAMTS13 activity/ MOF with the appearance of ULVWF multimers. Furthermore, some of the patients with low ADAMTS13 activity and high VWF levels revealed evidence of DIC and drastic changes in platelet count.

Conclusion: Our data provide further evidence that systemic inflammation may cause an imbalance between ADAMTS13 activity and VWF level, and that this imbalance may contribute to inflammation-mediated MODS.

054

Infection 2007; 35 (Suppl. II): 13

Vasopressin-induced impairment of microcirculation in patients with septic shock is associated with a drop-off in ADAMTS13 activity

*Claus RA, Klinzing S, Bockmeyer CL, Reinhard C, Sakr Y, Simon TP, Schürholz T, Reinhart K, Marx G
Department of Anaesthesiology and Intensive Care, Friedrich Schiller University Jena, Germany

Background: Arginine vasopressin (AVP) as the strongest endogenous vasopressor in terms of maximal contractile effect induced and its duration has found growing interest in the therapy of vasoplegic states, e.g. patients with septic shock. In parallel, in these patients a decrease of metalloprotease (ADAMTS13) with inactivating activity against the ultra-large, highly thrombogenic form of Von Willebrand Factor (ULVWF) has been described. It is also known, that (i) subsequent formation of VWF-rich microthrombi impairs the microcirculatory state and (ii) administration of a stabilized analogue of AVP results in a decrease of ADAMTS13 by unknown mechanisms and the appearance of ULVWF in plasma of healthy individuals.

Objective: to test the effect of AVP given in septic shock on mucosal microcirculation, hemodynamic parameters, urine output, haemostaseological parameters including ADAMTS13 and platelet count.

Methods: 20 patients with septic shock were studied in a consecutive design (12 male, 8 female; aging 63; 35-78 years). AVP was administered in a dosage of 2 IU/kg/h over 2 hours. During the study period, norepinephrine infusion rate was reduced to keep mean arterial blood pressure constant. Hemodynamic and haemostaseological parameters, platelet count and urine output were analyzed. Oral mucosal tissue oxygen saturation (SO₂), microcirculatory blood flow and flow velocity were measured with a laser Doppler flowmetry and remission spectroscopy system (O2C, LEA Medtechnik, Giessen). ADAMTS13 activity was determined using the Kokame method.

Results: AVP administration did not change cardiac output, but resulted in significant decrease of oral mucosal oxygen saturation, blood flow and a significant decrease of flow velocity. Also we analyzed a significant decline of ADAMTS13 activity (0.92; 0.69-1.15 U/mL vs. 0.51; 0.39-0.59 U/mL) and of platelet count after 4 hours (75; 64-100 vs. 71; 55-87 Gpt/l). During AVP administration, norepinephrine dosage and heart rate dropped down significantly. Furthermore, urine output increased (125; 20-480 vs. 210; 30-690 ml/h).

Conclusions: AVP administration causes a deterioration of oral mucosal blood flow independent of macrocirculatory conditions. This was accompanied by a further decrease of ADAMTS13 activity from declined baseline levels in patients with septic shock, which may result in secondary AVP induced platelet aggregation, collateral formation of VWF-rich microthrombi and subsequent occlusion of microvessels, which may be reflected by platelet consumption and impaired microcirculation after AVP infusion. Therefore, besides its beneficial vasoconstrictive effects, AVP administration may be reconsidered due to potential aggravation of organ dysfunction subsequent to triggering thrombotic-microangiopathic complications.

055

Infection 2007; 35 (Suppl. II): 14

Anti-platelet drugs reduce mortality in surgical and non-surgical ICU patients

*Lösche W (1), Neumann J (2), Winning J (1), Claus RA (1), Reinhart K (1), Bauer M (1)

(1) Department of Anaesthesiology and Intensive Care Therapy, University Jena, Germany, (2) Department for Anaesthesiology and Intensive Care, Clinical Centre Meiningen, Germany

Introduction: Platelet activation in sepsis may contribute to fatal outcome via thrombotic microangiopathy and modifying inflammatory reactions. Anti-platelet drugs such as acetylsalicylic acid or clopidogrel are used for secondary prevention of cardiovascular events and their effect lasts for up to 7 days after withdrawal.

Objectives: We studied whether pre-hospital use of anti-platelet may reduce mortality in critically ill patients of either non-surgery or surgery/traumatic cause.

Methods: 361 non-surgery and 254 surgery patients who were admitted to the intensive care unit (ICU) within 24 h of hospital treatment were included in a retrospective study. Elective surgery or use of statins were exclusion criteria.

Results: 155 of the 615 patients were on anti-platelet drugs prior to hospitalisation. 84% were on acetylsalicylic acid, 6 % on clopidogrel and 10 % on a combination of ASS and clopidogrel. 45 % of patients developed sepsis, and the overall ICU mortality was 38 %. Pre-hospital use of anti-platelet drug was associated with lower minimum and maximum platelet counts that were reached at days 4 and 15, respectively.

In both non-surgery and surgery patients the use of anti-platelet drugs was associated with higher age (70±11 vs 58±20 and 70±13 vs 49±20, respectively) and APACHE II score (27±9 vs 21±9 and 24±9 vs 17±8, respectively). Use of anti-platelet drugs was also associated with an insignificant increase in transfusion-requiring bleeding in non-surgery and surgery patients (27% vs 14% and 69% vs 55%, respectively).

Compared to patients without anti-platelet drugs, the use of anti-platelet drugs was associated with a slightly reduced mortality (40% vs 50%) in non-surgery patients and a slightly increased mortality in surgery patients (31% vs 24%), but the differences were not significant. However, in patients on anti-platelet drugs the predicted mortality rates calculated on the basis of the individual APACHE II scores was remarkably higher than the actual mortality rates (non-surgery patients 58±24%, surgery patients 48±26%).

When matching patients (non-surgery and surgery) according to their individual APACHE II scores, the use of anti-platelet drugs was associated with a relative risk of mortality of 0.51 (95% CI 0.28 – 0.85) in the 3rd quartile of APACHE II scores (21 – 26) and 0.73 (0.58 – 0.92) in the 4th quartile (> 26) of the scores.

Multivariate regression indicated that the use of anti-platelet drugs was a significant negative predictor of mortality in non-surgery as well as surgery patients (p<10-7 and p<10-3, respectively), whereas APACHE II and age were positive predictors of mortality.

Conclusion: The use of anti-platelet drugs for secondary prevention of cardiovascular disease clearly decreased the mortality rate in both non-surgically and surgically ICU-patients. Therefore, anti-platelet drugs could be a novel therapeutic option to reduce organ failure in these patients.

059

Infection 2007; 35 (Suppl. II): 14

Breakdown of sphingomyelin by secreted sphingomyelinase and formation of ceramide enriched macrodomains in patients with systemic inflammation and organ failure

Bunck AC, Hupe D, Dorer M, Acht B, Brunkhorst FM, Deigner HP, *Claus RA

Department of Anaesthesiology and Intensive Care Therapy, Friedrich-Schiller University Jena, Germany

Introduction: Bioactive lipids such as ceramide (Cer) have significant effects on cells relevant for inflammation, which may therefore be regarded as candidate mediators in systemic inflammatory response syndrome (SIRS). In pts with sepsis, Cer concentration in mononuclear cells has been identified as a marker to predict multiple organ failure. In order to elucidate the role of Cer formation in the development and progression of systemic sequelae of SIRS, we addressed the question whether there is a difference in the sphingolytic activity of the secreted isoform of sphingomyelinases (pSMPD1) in plasma of pts with various degrees of SIRS or sepsis of different origin.

Methods: Plasma samples were obtained from pts with SIRS after off pump coronary bypass surgery through a mini-left anterior thoracotomy (MIDCAB), after CABG using an extracorporeal circuit (cardiopulmonary bypass – CPB) as well as from patients with sepsis. Activity was determined by the hydrolysis of fluorescently labelled sphingomyelin, plasma presence of the enzyme was identified by immunoblotting. In endothelial cells exposed to plasma, we determined the sphingomyelin turnover and the resulting Cer pattern by chromatographical separation as well as the formation of Cer enriched macrodomains by fluorescence microscopy.

Results: Plasma activity of sphingomyelinase (pSMPD1) in samples of pts with sepsis (median 262.3 pmol/(ml*h)) were higher than those of age matched controls (median 123.6 pmol/(ml*h); p < 0.005). In pts with fatal outcome (n=7) sphingolytic activity increased during the study period (+ 77.4 pmol/(ml*h)), while a decrease in the subgroup of sepsis survivors was observed (- 252.1 p < 0.02). Low and high pSMPD1 activity levels were paralleled by equally low or high values of established clinical severity markers such as SOFA score or proCT value. Comparing the absolute increase of pSMPD1 activity 24 hours after either MIDCAB surgery or CABG with CPB we found the increase of enzyme activity to be significantly lower in both the MIDCAB group as well as in patients without SIRS at the first day post op. Beyond immunological detection of increased pSMPD1 in septic patients, we found an increase in breakdown of sphingomyelin in endothelial cells after stimulation with patients' plasma as well as endotoxin or prototypic pro-inflammatory cytokines such as TNF. Also we found formation of ceramide enriched macrodomains by immunostaining using specific antibodies directed against Cer, CD14, Fas-receptor and TNF-Receptor1.

Conclusion: Here we demonstrate an increased pSMPD1 activity in pts with SIRS and sepsis. Together with data from in vitro experiments and animal models, the results provide the first demonstration of a bio-functional relevant activity of pSMPD1 resulting in an altered signal transduction in systemic inflammation. The association with sepsis related mortality suggests that increased pSMPD1 activity may be involved in the complex network of processes finally resulting in an unfavourable outcome.

061

Infection 2007; 35 (Suppl. II): 15

Clinical utility of a new PCR-based assay for rapid pathogen detection (SeptiFast®) in patients with clinical sepsis compared to standard blood culture

*Bingold TM (1), Hunfeld K (2), Scheller B (1), Rönneberg T (1), Sartorius S (1), Wahrmann M (1), Klösel S (1), Zwißler B (1), Wissing H (1)

(1) Clinic for Anaesthesiology, Intensive Care and Pain Therapy, University Frankfurt, Germany, (2) Institute of Medical Microbiology & Infection Control, University Frankfurt, Germany

Introduction: Early specification of bacterial and fungal pathogens plays a crucial role in sepsis therapy. A new real-time PCR system (SeptiFast®, Roche Diagnostics, Mannheim) allows a more rapid identification for 25 of the most common pathogens compared to standard blood culture (BC). The test specifically targets the ITS -region of bacteria and fungi in 3 mL K-EDTA blood and can be completed in less than 6 h.

Objective, Methods, Results: We were interested whether the new real-time PCR system would indicate an advantage in early identification of common pathogens compared to standard BC. After approval of the study by the local ethics committee and having obtained informed consent, we included 21 surgical patients with severe sepsis and septic shock according to the S2 guidelines of the German Society of Sepsis. All patients had significant elevated inflammatory parameters (PCT, IL-6, LBP) at study entry. Blood samples for PCR and BC were taken at admission to the anaesthesiological/surgical ICU and at up to six following time points, every 24 h. Samples were taken by sterile venous puncture or from the arterial line. Whenever possible, swabs were taken from suspicious sites for completion of microbiological diagnostics. Samples were analysed by the Institute of Medical Microbiology. Results: In total 134 paired blood samples were obtained from 21 patients. In 62 samples PCR was positive, and in 20 samples blood cultures (table 1). In 4 positive BCs the pathogen must be considered as contamination (CONS). In 10 samples PCR and BC detected the same pathogens. Additionally in 11 samples pathogens detected by PCR were also identified by swabs. Table 1: 134 paired samples: PCR pos PCR neg BC pos (+swap) 10 (21) 10 BC neg (+swap) 52 (42) 62 In these 21 septic patients BC was positive in 10 cases, whereas PCR was positive in 16 cases. In three patients pathogens could not be detected in the blood stream by any method at all. Table 2: Pathogen detection in 21 patients PCR Pos PCR Neg BC pos 8 2 BC neg 8 3 As shown in Table 3, PCR detected pathogens at earlier time points than BC. Table 3: First Detection of pathogen Time Point PCR + BC+ Accordance I. 6 3 2 II. 7 2 1 III. 1 1 IV. 2 0

Conclusion: PCR provides detection of pathogens more frequently and much earlier than by standard BC and swabs in patients with severe sepsis and septic shock. Even in these patients merely 15% of BC became positive, whereas 46% of PCR detected a pathogen. Both methods failed in three patients. The gain of time is not only achieved by a shorter time to report (6h PCR vs. 24-48 h BC) but also by detection at earlier timepoints (table 3). So we conclude, that real-time PCR SeptiFast® could be a useful tool to improve early, adequate antibiotic treatment.

062

Infection 2007; 35 (Suppl. II): 15

Detection of bacterial DNA in critically ill patients using the SeptiFast PCR system

*Lehmann LE (1), Book M (1), Weber SU (1), Schewe JC (1), Hunfeld KP (2), Wissing H (3), Seifert H (ORT), Hoeft A (1), Stüber F (1)

(1) Department of Anaesthesiology and operative Intensive Care Medicine, University Bonn, Germany, (2) Institute of Medical Microbiology & Infection Control, University Frankfurt, Germany, (3) Department of Anaesthesiology, Intensive Care and Pain Therapy, University Frankfurt, Germany

Introduction: Persistent bacteraemia is indicative of an invasive infection that may lead to sepsis. Rapid administration of adequate antimicrobial therapy improves outcome. However, pathogen identification by conventional microbiological methods usually takes up to 36 h. Two-step nucleic acid based techniques comprising of an initial broad-range PCR followed by a separate pathogen identification step have been introduced.

Objective: Our study evaluates a newly developed single step real-time PCR based method (SeptiFast) in a cohort of sepsis patients and patients prior to elective surgery without clinical or laboratory signs of infection and compares it to conventional blood cultures. Methods: After approval of the local ethics committee and acquisition of written and informed consent of the patient or a legal guardian, patients fulfilling the ACCP/SCCM sepsis criteria were included in the study. For each comparative analysis 20 mL whole blood was drawn for blood culture analysis using a BACTEC system according to the manufacturers recommendations and 3 mL whole blood was collected for PCR analysis using the SeptiFast system according to the manufacturers recommendation.

Results: 453 blood samples of 108 patients study patients and 206 blood samples of 103 control patients were compared to parallel blood cultures. SeptiFast resulted in 2 fold more positive findings (n=114) than blood cultures (n=58). In 40 of 58 positive blood cultures findings were identical to SeptiFast. The negative predictive value of SeptiFast against blood culture was 0.95; sensitivity: 0.69; specificity: 0.81. 18 samples of positive blood cultures showed different SeptiFast results. 13 of these 18 cases could be resolved by being not in the panel detectable by SeptiFast or being flagged as contaminations. In 206 samples of the control group no positive SeptiFast results were found. 22 blood culture findings were positive for CoNS, after prolonged culturing: these results were regarded as contaminations. SeptiFast results were also analyzed against a constructed microbiologic gold standard comprised of the presence of pathogens in blood cultures or other additional patient specimen (e.g. BAL, urine, swabs) taken on the same day as the SeptiFast sample. SeptiFast detected 74 pathogens, which were negative by blood culture findings. 51 of these 74 SeptiFast results matched with microbiologic results from additional specimen. This increased the sensitivity of SeptiFast compared to the constructed gold standard to 0.83; the specificity increased to 0.93; the negative predictive value was 0.95.

Conclusion: Our study demonstrates concordance of PCR results with blood cultures. We confirmed increased sensitivity of PCR based diagnosis of bloodstream infection. Our results are indicative of potential clinical utility of PCR based pathogen identification. Studies addressing changes in clinical decision-making, outcome and cost-benefit by the utilization of PCR results are needed.

067

Infection 2007; 35 (Suppl. II): 16

The ventilation strategies influence outcome in patients with severe sepsis – results from a national prospective multicenter study

Schädler D (1)*, Schmitz G (1), Engel C (2), Bogatsch H (2), Frerichs I (1), Kühlen R (3), Quintel M (4), Rossaint R (5), Scholz J (1), Brunkhorst FM (6), Loeffler M (2), Reinhart K (6), Weiler N (1) for the German Competence Network Sepsis (SepNet)
(1) Department of Anaesthesiology and Intensive Care Medicine, University Medical Centre Schleswig-Holstein, Campus Kiel, Germany, (2) Institute for Medical Informatics, Statistics and Epidemiology, University Leipzig, Germany, (3) Department of Intensive Care Medicine, RWTH University Aachen, Germany, (4) Centre of Anaesthesiology, Emergency and Intensive Care Medicine, University Hospital Göttingen, Germany, (5) Department of Anaesthesiology, RWTH University Aachen, Germany, (6) Department of Anaesthesiology and Intensive Care Medicine, Friedrich-Schiller-University Jena, Germany

Introduction: Mechanical ventilation plays an important role in the management of septic patients. It still remains unclear whether the pressure controlled ventilation (PCV), volume controlled ventilation (VCV) or assisted ventilation (AV) is the optimum ventilatory strategy.

Objectives: The objectives of this study were to 1) describe the actual ventilatory strategies applied in the clinical practice, 2) analyse the association of the different ventilatory strategies with mortality and 3) identify the potential risk factors for death associated with the ventilator therapy and the respiratory system in patients with severe sepsis and septic shock.

Methods: In the “Epidemiology of Sepsis in Germany” study, a prospective observational cross-sectional study, data from 454 intensive care units (ICU) in 310 randomly selected hospitals in Germany were collected by local one-day visits of trained physicians from SepNet’s 17 regional study centres. Variables associated with the respiratory system and mortality were analysed in patients ventilated with PCV, VCV and AV as well as in patients classified as acute lung injury (ALI), acute respiratory distress syndrome (ARDS) and non-ALI. Multivariate analysis was applied to identify possible risk factors for death.

Results: In a cohort of 415 patients with severe sepsis, 20% did not need mechanical ventilation. Non-invasive ventilatory support was used in 5% of the ventilated patients, 32% were ventilated via tracheostoma, 63% were either orally or nasally intubated. Hospital mortality did not differ between PCV (57%), VCV (71%) and AV (51%) ($p=0.23$). Peak pressures above 35 cm H₂O were associated with higher mortality rates ($p=0.05$) whereas there was no association between the tidal volume and hospital mortality ($p=0.57$). ALI occurred in 42% and ARDS in 27% of the ventilated patients. Significantly higher hospital mortality was found in patients with ALI (57%) and ARDS (69%) when compared with non-ALI patients (41%; $p=0.015$). The estimated respiratory system compliance, renal dysfunction and APACHE II score were identified as independent risk factors for hospital mortality.

Conclusions: In this study we did not find an association between the ventilatory strategy and mortality of patients with severe sepsis or septic shock. The combination of ALI and sepsis or septic shock significantly worsened the patients’ outcome.

The study was supported by the Federal Ministry of Education and Research, Berlin, Germany (BMBF, Grant No. 01 KI 0106) and an unrestricted grant from Lilly Germany Inc., Bad Homburg, Germany.

068

Infection 2007; 35 (Suppl. II): 16

The attributable risk of parenteral nutrition to mortality in septic patients – results from a national prospective multicenter study

Schmitz G. (1), Schädler D. (1)*, Engel C. (2), Bogatsch H. (2), Frerichs I. (1), Ragaller M. (3), Scholz J. (1), Brunkhorst F.M. (4), Löffler M. (2), Reinhart K. (4), Weiler N. (1), for the German Competence Network Sepsis (SepNet)
(1) Department of Anaesthesiology and Intensive Care Medicine, University Medical Centre Schleswig-Holstein, Campus Kiel, Germany, (2) Institute of Medical Informatics, Statistics and Epidemiology, University of Leipzig, Germany, (3) Department of Anaesthesiology and Intensive Care Medicine, University Hospital of the Technical University of Dresden, Germany, (4) Department of Anaesthesiology and Intensive Care Medicine, University Hospital of the Friedrich-Schiller-University of Jena, Germany

Introduction: The provision of an appropriate nutritional support is an accepted part of the standard intensive care in septic patients. Nonetheless, it is unclear whether the route of nutrition has an influence on the clinical outcome in this subgroup of critically ill patients.

Objectives: To describe current clinical practice of nutrition and evaluate its association with mortality in patients with severe sepsis or septic shock in Germany.

Methods: In the “Epidemiology of sepsis in Germany” study, a prospective observational cross-sectional study, data from 454 intensive care units from a representative sample of 310 randomly selected hospitals in Germany were collected by trained physicians from the SepNet’s 17 regional study centers during randomly scheduled one-day visits over a period of one year. 415 patients with severe sepsis or septic shock were identified (according to ACCP/SCCM consensus conference criteria). Data were evaluated concerning nutrition modalities. Logistic regression analysis was used to identify predictors for death.

Results: Valid data on nutrition were available for 399 of 415 patients. 20.1% of these patients received exclusively enteral nutrition while 35.1% were nourished exclusively parenterally. Mixed nutrition (parenteral and enteral) was administered to 34.6 % of patients, 10.3% were not fed at all. Patients with septic shock, mechanical ventilation or a possible contraindication to enteral nutrition were less likely to receive exclusively enteral nutrition. Median APACHE II score but not SOFA score was significantly different among the nutrition groups. Overall hospital mortality in septic patients was 55.2%. Hospital mortality was significantly higher in patients receiving exclusively parenteral (62.3%) or mixed nutrition (57.1%) than in patients with exclusively enteral nutrition (38.9%) ($p=0.005$). By multivariate analysis, presence of parenteral nutrition (odds ratio, 2.09; 95% confidence interval, 1.29 – 3.37), APACHE II score (odds ratio, 1.05; 95% confidence interval, 1.02-1.09) and renal dysfunction (odds ratio, 2.07; 95% confidence interval, 1.30 – 3.31) were significantly predictive of mortality.

Conclusions: Patients with severe sepsis or septic shock in German intensive care units receive preferentially parenteral or mixed nutrition. In this study, the use of parenteral nutrition was directly associated with an increased risk of death.

The study was supported by the Federal Ministry of Education and Research, Berlin, Germany (BMBF, grant no. 01 KI 0106) and an unrestricted grant from Lilly Germany Inc., Bad Homburg, Germany.

072

Infection 2007; 35 (Suppl. II): 17

Supplementation with fish oil of parenteral nutrition in sepsis patients

*Köhler J, Abel P, Friesecke S

Intensive Care Unit, Department of Internal Medicine, University Greifswald, Germany

Background: It has been shown, that n-3 polyunsaturated fatty acids (PUFA) contained in fish oil may modulate immune function. Depending on availability, n-6 PUFAs are metabolized to become rather potent inflammatory mediators, whereas n-3 PUFAs are a source of less active inflammatory mediators.

Objective: This study was designed to examine whether an increase in the proportion of n-3 : n-6 PUFAs from 1:7 in conventional lipid emulsions to 1:2 in a lipid mixture containing fish oil would have an effect on inflammatory processes and clinical course of sepsis.

Design: Placebo-controlled, double-blind, parallel-group, randomized controlled trial.

Patients: 62 consecutive patients treated for sepsis, severe sepsis, or septic shock on the medical ICU of a university hospital if parenteral nutrition was required for at least 3 days. Intervention and Outcome Measures Patients were randomly assigned to receive either a mixture of fish oil and a soybean-oil based standard MCT/LCT lipid emulsion (intervention group), or the standard MCT/LCT lipid emulsion (placebo group). For 7 days, laboratory parameters of systemic inflammatory IL-6, PCT, and CrP were determined. Incidence of nosocomial infections, duration of mechanical ventilation, length of stay on ICU, mortality on ICU, survival at 28 days, and daily SOFA score were recorded.

Results: Of the 62 patients, 24 were in sepsis, 6 in severe sepsis, and 32 in septic shock. Mean age of both groups was 63 years, mean SAPS II was 53 in the intervention group vs. 55 in the placebo group (n. s.). A decline of mean IL-6, PCT, CrP, and SOFA score was noted, but did not differ between intervention and placebo groups. Incidence of nosocomial infections, mean duration of mechanical ventilation (28 vs. 24 days, n. s.), mean length of ICU stay (35 vs. 27 days, n. s.), ICU mortality (55 vs. 53%, n. s.), and survival at 28 days (47 vs. 39%, n. s.) did not differ significantly between intervention and placebo groups, either.

Conclusions: In the study population – unselected sepsis patients of a medical ICU –, supplementation of parenteral nutrition with fish oil could not be demonstrated to have a significant effect on parameters of inflammation, or clinical outcome. This may be due to the heterogeneity of medical patients concerning stage and severity of sepsis.

074

Infection 2007; 35 (Suppl. II): 17

1,3-beta-D-glucan in critically ill patients with severe sepsis and SIRS after CABAG

Otto G, Bloos F, Brunkhorst FM

Department of Anesthesiology and Intensive Care Medicine, University Jena, Germany

Background: The saccharide 1,3-beta-D-glucan is a main component of the fungal cell wall. Markedly elevated serum levels of 1,3-beta-D-glucan were found in patients with systemic fungal infections [1,2]. Our aim was to evaluate 1,3-beta-D-glucan concentrations in plasma of critically ill patients with severe sepsis and SIRS.

Methods: 26 patients with severe sepsis and 22 patients after coronary artery bypass grafting (CABG) were included in this pilot study. Plasma samples were collected daily in the sepsis group or on day 1 after surgery in the CABG-group. 1,3-beta-glucan was measured with a turbidometric assay (Wako Pure Chemical Inc.). A cut-off level of 11 pg/ml was used.

Results: 1,3-beta-glucan serum levels were elevated after uncomplicated CABG and differed significantly to the sepsis group (figure). Median concentrations were in the normal range in severe sepsis patients but fraction of elevated 1,3-beta-glucan serum levels tend to increase with length of stay.

Conclusions: This first observational study demonstrated consistent results of serum 1,3-beta-glucan measurements in different populations of critically ill patients. Bacterial translocation has been suspected as reason for SIRS after CABG, however, this has never been associated with fungemia. This finding warrants further research.

References: [1] Obayashi T et al. Lancet 1995;345:17-20. [2] Miyazaki T et al. J Clin Microbiol 1995; 33: 3115-3118.

Grant acknowledgement: The study was supported by Wako Pure Chemical Industries, Osaka, Japan

075

Infection 2007; 35 (Suppl. II): 17

Prospective assessment of parameters of hepatic functions in severe human sepsis

*Kortgen A. (1), Werth M. (2), Paxian M. (1), Bauer M. (1)

(1) Klinik für Anästhesiologie und Intensivtherapie, Friedrich-Schiller-Universität Jena (2) Klinik für Anästhesiologie und Intensivmedizin, Universitätsklinikum des Saarlandes, Homburg/Saar

Introduction: Sepsis results from the host response to infection and sepsis-associated tissue injury is modulated by the reticuloendothelial system making the liver a primary target and due to its central role in metabolism to a primary effect organ.

Objective: To assess incidence and course of impaired hepatic functions in severe human sepsis.

Material and methods: Markers of hepatocellular integrity, protein synthesis, and cholestasis were measured daily in 48 consecutive patients fulfilling criteria for severe sepsis. In addition, plasma disappearance rate of the anionic dye indocyanine green (PDRICG) was determined. Demographic data, APACHE II-, MOD- and SOFA-scores, values obtained from laboratory tests and COLD measurements at inclusion were evaluated with unpaired t-test or Mann-Whitney rank sum test for continuous variables or Fisher's exact test for categorical variables. Analysis of variance (ANOVA) was used for serial measurements. For receiver operating characteristics areas under the curve (AUC) were calculated. All p-values are two-tailed. Statistical significance is defined as p<0.05.

Results: All-cause mortality was 37.5% and incidence of liver dysfunction 42 % and 74% as assessed by hyperbilirubinemia or PDRICG, respectively. While conventional markers for liver injury (e.g. ALT: AUC 0.48, $p=0.084$; bilirubin: AUC 0.43, $p=0.412$) failed to predict poor outcome, a PDRICG<8% (AUC 0.81, $p=0.006$) predicted death with a sensitivity of 81% and specificity of 70%. Weak but significant correlations were obtained between PDRICG, surrogates of altered blood flow, and inflammation.

Conclusion: In contrast to the biphasic model of liver impairment in sepsis with an early ischemic hepatitis followed by delayed cholestasis, our data suggest early occurrence of, and a key role for excretory dysfunction in severe sepsis, which is underestimated by conventional laboratory markers. PDRICG better reflects liver dysfunction in the light of diagnostic uncertainty regarding clinical signs of liver failure, such as coagulopathy and encephalopathy in the septic host. Excretory function is important for biotransformation of xenobiotics. This might be an hitherto underestimated problem in pharmacotherapy in patients presenting with severe sepsis potentially leading to accumulation of cytotoxic xenobiotics in hepatocytes and secondary drug-induced liver injury.

076

Infection 2007; 35 (Suppl. II): 18

IL-6 as a severity marker of serious sepsis and septic shock

*Kajtor I, Kisvarga Z, Marjanek Z

Department of Anaesthesiology and Intensive Care, Jávorszky Ödön Hospital Vác, Hungary

Introduction: The degree of systemic inflammatory response against bacterial invasion considerably correlates with the number of organ dysfunctions and amount of released mediators. We examined IL-6 levels in serious sepsis and septic shock at survivor and non-survivor patients. We analysed the kinetics of IL-6 and its relationship with PCT level.

Patients: We randomised 13 patients with serious sepsis (3 organ failure) and 15 patients with septic shock. We identified pulmonary infection at 19 cases and intraabdominal infection at 9 cases as septic focus. We measured IL-6 and PCT levels, counted APACHE-II and SOFA scores at the recognition of serious septic process (0h) and in the 6th, 12th, 24th, 48th and 72th hours.

Results: We couldn't find any significant difference among 0h IL-6 levels. 6h IL-6 levels were significantly higher at patients with septic shock. 12h IL-6 levels were significantly decreased in both groups, although IL-6 levels remained higher at non-survival patients. We marked a peak of APACHE-II, SOFA score and IL-6 in the 6. hour. We detected over 10ng/ml PCT levels from 6th hour and decrease from 36th hour in both groups.

Conclusion: IL-6 is suitable for estimation of sepsis severity and usable at diagnostics of septic shock. Its permanently higher level is a predictive marker of outcome.

077

Infection 2007; 35 (Suppl. II): 18

Toxic shock syndrome

*Marjanek Z, Ortutay A, Kisvarga Z

Jávorszky Ödön Hospital Vác, Department of Anaesthesiology and Intensive Care, Hungary

Case report: The toxic shock syndrome (TSS) is a high mortality disease. Pyrogen endotoxins generated by *Staphylococcus aureus* and *Streptococcus pyogenes* (TSST-1, SPEA, SPEA) effect excessive cytokine production as superantigens stimulating the T-lymphocytes on non-specific way. TSS is clinically characterized by rapid onset and progression, fever, shock and MOF. Early and adequate therapy determines the outcome of TSS.

We compare *Staphylococcus aureus* and *Streptococcus pyogenes* TSS presenting two case reports (68 yrs old male and 48 yrs old female). We demonstrate criterias, diagnostic differences and therapeutical management according to CDC definitions.

Although the incidence of the disease is considerably low, the importance of TSS is based on

- 1) the huge mortality, which compromises the healthy population,
- 2) the lack of natural immunity against virulence factors in public communities,
- 3) unchanged mortality in spite of modern diagnostic and therapeutical management.

080

Infection 2007; 35 (Suppl. II): 18

Plasma separation and adsorption treatment of patients with severe sepsis and septic shock using an adsorber with an arginine ligand

*Umgelter A (1), Reindl W (1), Huber W (1), Ronco C (2), Lutz J (3), Frank H (3), Heemann U (3)

(1) II. Medizinische Klinik und Poliklinik, Klinikum rechts der Isar der Technischen Universität München, Germany, (2) Department of Nephrology Dialysis & Transplantation San Bortolo Hospital, Vicenza, Italy, (3) II. Medizinische Klinik und Poliklinik, Abteilung für Nephrologie, Klinikum rechts der Isar der Technischen Universität München, Germany

Introduction: Endotoxemia leading to endothelial activation and loss of vascular resistance by increased production of nitric oxide is an important factor in the development of distributive shock and multi-organ failure in septic patients. The removal of endotoxin from the circulation has thus been advocated as a treatment option in sepsis. We report the results of the first application of a novel Endotoxin adsorption column (ET-Adsorber, Gambro, Sweden) in septic patients.

Methods: Patients were included if they met the following criteria: 2 or more SIRS criteria, a plasma-endotoxin concentration of 0.1 EU/ml in the chromogenic Limulus Amebocyte Lysate (LAL) test, a clinical indication for the placement of a dialysis catheter. Plasma separated from blood in a plasma filter and pumped through the adsorption column made of a hydrophilic vinyl copolymer with an arginine ligand, and then returned to the blood stream. Five treatments per patient were performed on consecutive days. Clinical and laboratory, as well as hemodynamic data were recorded and values before and after treatment sessions were compared.

Results: Ten consecutive patients were included (8m, 2f, age 56.2 ± 10.7 years). They had sepsis resulting from nosocomial pneumonia ($n=3$), spontaneous bacterial peritonitis ($n=2$), soft-tissue-abscesses ($n=2$), peritonitis ($n=2$) and without obvious focus in one case. Initial APACHE II score was 29.7 ± 8.7 , and the mean SOFA score before the first treatment was 14.5 ± 4.5 . Plasma-endotoxin before the first treatment had a median of 0.70 EU/ml (range: 0.11 – 2.4). Hospital

mortality was 40%. 49 treatment sessions were performed. Comparing measurements before and after adsorption treatments, mean arterial pressure increased from 80.76 ± 15.7 mmHg to 88.16 ± 16.7 mmHg and cardiac power index from 0.72 ± 0.22 to 0.81 ± 0.22 while norepinephrine (NE) could be reduced from 4.0 ($0 - 29$) $\mu\text{g}/\text{min}$ to 2.2 ($0-29$) $\mu\text{g}/\text{min}$. There was an increase in Global Enddiastolic Volume Index (GEDVI) from 744.6 mL m⁻² to 784 mL m⁻². In between treatments, changes were partially reversed. Changes in Cardiac Index (CI), and Systemic Vascular Resistance (SVRI) were not significant. Proapoptotic activity of plasma and the amount of advanced oxidation protein products decreased during the adsorption treatments but there was no change in reactive carbonyl compounds. There were no complications attributable to catheters or extracorporeal circulation.

Conclusion: In septic patients with endotoxin adsorption the dose of vasopressors could be significantly reduced while MAP levels increased suggesting an improved vasotonus and improved myocardial contractility. The treatment appears to be safe and efficient and randomized studies should be performed.

082

Infection 2007; 35 (Suppl. II): 19

Immune effects of hydrocortisone in septic shock – results from the CORTICUS Berlin Study Group

*Engelhardt T (1), Ahlers O (1), Sprung CL (2), Briegel J (3), Volk HD (4), Annane D (5), Moreno R (6), Singer M (7), Gerlach (8), Keh D (1,9)

(1) Anesthesiology and Critical Care Medicine (ACCM), Charité Universitätsmedizin, Berlin, Germany, (2) ACCM, Hadassah Hebrew University Medical Center, Jerusalem, Israel, (3) Anesthesiology, Ludwig-Maximilians-Universität, Munich, (4) Medical Immunology, Charité, Berlin, Germany, (5) ICU, Raymond Poincaré Hospital, Garches, France, (6) UCIP, H.St. António dos Capuchos, C. Hosp. de Lisboa Central EPE., Lisbon, Portugal, (7) Medicine and Wolfson Institute of Biomedical Research, University College, London, United Kingdom, (8) ACCM, Vivantes Klinikum Neukoelln, Berlin, (9) ACCM on behalf of the CORTICUS Berlin Study Group, Charité, Berlin, Germany

Introduction: We investigated immune effects of hydrocortisone (HC) administration in septic shock with regard to adrenal function.

Methods: Patients were enrolled in the randomized study of HC in septic shock (CORTICUS). A 250 μg ACTH-test was performed before administration of HC (50 mg q 6h for 5 days, tapered until day 11) or placebo (PL); plasma cortisol levels were measured in a central laboratory. Responders (R) were defined by an incremental cortisol increase >9 $\mu\text{g}/\text{dl}$ and non-responders (NR) ≤ 9 $\mu\text{g}/\text{dl}$. LPS-stimulated release of Interleukin(IL)6 and IL10 were measured at baseline, day 3 (D3) as well as at day 6 (D6), and IL6/IL-10-ratio was calculated. Values on D3 and D6 were adjusted for baseline and analyzed with GLM for repeated measurements.

Results: From March 02 to November 05, 84 patients were enrolled in 13 sites: 42 HC (26 R and 16 NR) and 42 PL (30 R and 12 NR). Mean baseline values were: IL6: HC 773 pg/ml (595-950), PL 658 (95%CI: 515-800), R 703 (568-837), NR 746 (527-965); IL10: HC 32,6 pg/ml (21,3-43,9), PL 37,6 pg/ml (20,2-55,0), R 32,9 (21,9-44,0), NR 39,9 (17,1-62,6); IL6/IL10-ratio: HC 46,6 (23,7-69,4), PL 43,6 (25,3-62,0), R 45,2 (27,8-62,6), NR 44,9 (17,7-72,2). Mean changes from baseline were: IL6: HC -38% (D3) and +4% (D6), PL +44% (D3) and +105%, $p<0.005$; IL10: HC +82% (D3) and +106% (D6), PL +84% (D3) and +239% (D6), n.s.; IL6/IL10-ratio: HC -23% (D3) and -12% (D6), PL +102% (D3) and +33% (D6), $p<0.05$; Within the HC-group, changes from baseline were not significantly different between R and NR for all parameters. Within the PL-group, IL6/IL-10-ratio decrease was significantly higher in NR ($p<0.05$) whereas no differences could be observed for other parameters.

Conclusion: No significant differences between R and NR were found at baseline. HC attenuated stimulated IL6 production in monocytes significantly and thereby caused a significantly different decrease of IL6/IL10-ratio. The observed immune effects of HC were independent of adrenal function at baseline.

Grant Acknowledgement: Deutsche Forschungsgemeinschaft (KE870/1-1,2), European Union, European Society of Intensive Care Medicine, International Sepsis Forum

084

Infection 2007; 35 (Suppl. II): 19

Clinicians' perception on glycemic control – results of a multicenter ICU survey

Brunkhorst FM, Bloos F, Hartog C, Braune A, Reinhart K, for the German Competence Network Sepsis (SepNet)
SepNet Office, Department of Anaesthesiology and Intensive Care Medicine, Friedrich-Schiller-University Jena, Germany

Background: Strict glycemic control may lower mortality and morbidity in critically ill patients. The perceptions concerning glycemic control among health care workers in Germany are not known. We conducted a multicentric survey to determine beliefs and attitudes about glycemic control in ICU patients from health care workers of the ICU team.

Methods: We formulated items along the lines of the questionnaire administered by McMullin et al [1] which addresses four main objectives: 1) Perception of threshold values for hypoglycaemia or hyperglycemia; 2) identification of patient groups benefiting most of euglycemia; 3) concern about the use and accuracy of glucometers and 4) strategies for optimal glucose management. We used the same self-administered format with binary answers of yes/no and ten-point scales of 1 mmol/l increments for glucose values. The questionnaire was sent to 17 ICUs of regional SepNet centers. Participation was voluntary. Answers depicted “nurse”, “resident” or “attending physician”, but were otherwise anonymous. The survey was conducted from October 2005 – January 2006.

Results: Questionnaires were returned from 342 nurses, 98 residents and 34 attending physicians. Nurses and residents reported to feel concern about hypoglycemia from 3.0 mmol/l (median, IQR 3-4 mmol/l) downwards, consultants reported a threshold of 3.5 mmol/l (3.0-4.0 mmol/l). Hyperglycemia in non-diabetic patients was a concern from 8.0 mmol/l upwards (8.0-9.0 mmol/l for nurses, 8.0-10.0 mmol/l for residents, 7.0- 8.0 mmol/l for attending physicians). The threshold for hyperglycemia in diabetic patients was reported by residents and attending physicians as 8.0 mmol/l (8.0-10.0 mmol/l for residents, 8.0-8.75 mmol/l for attending physicians) and by nurses as 9.0 mmol/l (8.0-10.0 mmol/l). When asked for conditions in which patients would benefit most from glycemic control, attending physicians most often named stroke and heart surgery, residents named stroke and sepsis, and nurses sepsis and diabetes. Nearly all diseases in question except diabetes were rated with varying degree of importance by the different members of the ICU health care team. Strategies to improve glycemic control were also rated variably.

Conclusions: This survey in German ICUs shows that attitudes and belief on the practice of glycemic control differ widely among nurses, residents and attending physicians. Generally, attending physicians reported tighter blood sugar thresholds, perceived more benefit in given disease conditions and asked for more frequent measurements, reminders and i.v. insulin therapy than residents or nurses.

References: [1] McMullin J et al. Intensive Care Med 2004, 30(5):798-803.

Acknowledgement: This study was supported by the Federal Ministry of Education and Research (BMBF) grant No: 01 KI 0106.

085

Infection 2007; 35 (Suppl. II): 20

Lipopolysaccharide binding protein in a surgical ICU: A marker of sepsis?

Sakr Y, Brunkhorst FM, Burgett U, Nacul FE, Reinhart K
Department of Anesthesiology and Intensive Care Medicine,
University Jena, Germany

Background: We investigated the time course of LBP plasma concentrations in surgical intensive care unit (ICU) patients, their value in discriminating sepsis from systemic inflammatory response syndrome (SIRS), and their association with severity of sepsis and outcome in these patients compared with interleukin (IL)-6, C-reactive protein (CRP), and procalcitonin (PCT).

Methods: All 327 consecutively admitted patients to our surgical units were included. Blood samples were obtained daily for LBP, IL-6, CRP, and PCT measurements.

Results: Serum LBP concentrations were higher in patients who had severe sepsis/septic shock on ICU admission than in patients who never had sepsis (20.5[8.1-38.8] vs. 14.2[7.7-22.2] µg/mL, $p<0.05$) but were similar in patients with sepsis without organ failure and those who never had sepsis. After three days, LBP levels were similar in all groups. In a receiver operating characteristic curve analysis, LBP concentrations moderately discriminated sepsis from SIRS (AUC=0.66) and severe sepsis from sepsis without organ failure (AUC=0.71). IL-6 had the highest AUC in discriminating sepsis from other conditions (AUC=0.76) and PCT had the highest AUC for discrimination of severe sepsis from sepsis (AUC=0.86). LBP concentrations on admission and during the first week, were similar in patients with Gram-positive and those with Gram-negative infections (15.9[11-26.7] and 37.2[25.1-62.4] vs. 16.3[5.3-31.6] and 31.6[13.4], µg/mL, $p>0.2$). LBP concentrations on admission were similar in non-survivors and survivors and did not discriminate ICU mortality. However, the maximum LBP concentration during the first 3 days in the ICU discriminated moderately between survivors and non-survivors.

Conclusions: In the surgical ICU, LBP moderately discriminated patients without infection from patients with severe sepsis but not from patients with sepsis without organ dysfunction. LBP concentrations did not distinguish between Gram-positive and Gram-negative infections. The correlation of LBP concentrations with disease severity and outcome is weak compared to other markers and it is of little value to the clinician in this patient population.

Keywords: LBP, IL-6, postoperative patients, sepsis.

Acknowledge: This study was supported by an unrestricted grant from DPC Biermann GmbH, Bad Nauheim, Germany

The goal of this study was to test whether CABG is associated with significant bacteremia or DNAemia

Methods: Patients scheduled for elective CABG were included. Three groups were investigated: Patients with off-pump coronary artery bypass surgery (group: OPCAB), with left ventricular ejection fraction (LVEF) >50% undergoing conventional CABG (group: HIEF), patients with LVEF <30% undergoing CABG (group: LOEF). A blood culture was obtained and 10 ml of EDTA blood were taken by sterile venous puncture during the following time points: After induction of anesthesia, after surgery, 1st and 2nd postoperative day. Microbial DNA was measured by PCR with the LightCycler System 2.0® (Roche Diagnostics).

Results: Patients were included into the OPCAB-group (n=11 patients, age 62.7±9.7 years), the HIEF-group (n=20, age 64.1±10.1 years), LOEF-group (n=21, age 67.1±9.8 years). Procalcitonin (PCT) levels were elevated in all groups and did not differ significantly ($p=0.60$; figure). Postoperatively, 1 (0.8 %) of all blood cultures and 5 (4 %) of all PCRs revealed positive results.

Conclusions: Postoperative inflammation was shown by increase of PCT. However, elevation of PCT was not associated with relevant bacteremia or DNAemia. Therefore, bacterial translocation did not seem to play a significant role in these patients after elective CABG with and without cardiopulmonary bypass.

Acknowledgements: This study was sponsored by Roche Diagnostics, Penzheim, Germany.

086

Infection 2007; 35 (Suppl. II): 20

Frequency of bacteremia and DNAemia after elective coronary artery bypass grafting

Senderrek M (1), Bloos F (1), Straube E (2), Sachse S (2),
Marx G (1), Bauer M (1), Reinhart K (1)
(1) Department of Anesthesiology and Intensive Care Medicine,
University Jena, Germany, (2) Department of Microbiology,
University Jena, Germany

Introduction: Coronary artery bypass grafting (CABG) is associated with a significant inflammatory reaction. Bacterial translocation during cardiopulmonary bypass may contribute to such inflammation.

087

Infection 2007; 35 (Suppl. II): 21

Relevance of testing microbial DNA in comparison with blood culture in critically ill patients without evidence of infection

Bloos F (1), Senderrek M (1), Straube E (2), Sachse S (2), Bauer M (1), Marx G (1), Reinhart K (1)

(1) Department of Anesthesiology and Intensive Care Medicine, University Jena, Germany, (2) Department of Microbiology, University Jena, Germany

Introduction: Measuring microbial DNA may be a useful addition to standard microbiological diagnostic techniques since DNA detection by PCR is feasible within one day. However, being a very sensitive method, PCR may even detect amounts of DNA not relevant for diagnosis of infection. The goal of this study was to evaluate the frequency of false positive PCR results in critically ill patients without infection.

Methods: Patients were included into the study if they had undergone major elective thoracic or oral surgery. A blood culture was obtained and 10 ml of EDTA blood were taken by sterile venous puncture. Further samples were taken 2 and 3 days after enrolment. Microbial DNA was measured by PCR with the LightCycler System 2.0® (Roche Diagnostics).

Results: 36 patients (age: 60.3 ± 13.0 years; 98 samples) were included into the study. 5.1 % of blood cultures and 4.1 % of the PCRs tested positive. Following microorganisms had been detected:

Blood culture	frequency	PCR	frequency
Staph. epidermidis	2	Staph. epidermidis	3
Staph. aureus	1	Staph. aureus	1
Staph. cohnii	1	Klebsiella	1
Micrococcus luteus	1		

Overall concordance of the PCR results with the blood culture results was 0.91. None of the microorganisms mentioned in the table were detected in the concomitantly taken sample for the other technique.

Conclusions: Positive results in this study were most likely due to contamination and occur in patients without evidence of infection in similar frequencies in blood cultures and in the PCR.

Acknowledgements: This study was sponsored by Roche Diagnostics, Penzheim, Germany.

088

Infection 2007; 35 (Suppl. II): 21

Relevance of testing for microbial DNA in patients with severe sepsis and septic shock

Bloos F (1), Straube E (2), Sachse S (2), Bauer M (1), Marx G (1), Reinhart K (1)

(1) Department of Anesthesiology and Intensive Care Medicine, University Jena, Germany, (2) Department of Microbiology, University Jena, Germany

Introduction: Blood culture is an important method for identifying the underlying microorganism causative of sepsis. However, blood culture results may take up to 48 hours possibly delaying appropriate antibiotic therapy. The goal of this study was to test whether measuring microbial DNA in the blood is a meaningful addition to the microbiological examination in septic patients.

Methods: Patients were included into the study if the patients either fulfilled criteria of severe sepsis or septic shock. A blood culture was

obtained and 10 ml of EDTA blood were taken by sterile venous puncture. Additional samples were obtained during new septic episodes. Microbial DNA was measured by PCR with the LightCycler System 2.0® (Roche Diagnostics).

Results: 68 patients (age: 60.6 ± 15.6 ; 110 samples) were included. 15.5 % of the blood cultures and 37.3 % of the PCR revealed a positive result. Procalcitonin levels were greater in sepsis patients, that tested PCR positive, compared to patients with negative PCR (Figure). A similar difference was found in patients with positive compared to negative blood cultures; however, difference in PCT-values did not reach statistical significance.

Conclusions: In patients with sepsis, PCR tests positive more often than the blood culture. These positive results are meaningful because they are associated with higher PCT values. Therefore, measurement of microbial DNA may add meaningful information additional to blood culture in patients with clinical diagnosis of sepsis.

Acknowledgements: This study was sponsored by Roche Diagnostics, Penzheim, Germany.

089

Infection 2007; 35 (Suppl. II): 22

Serum levels of programmed cell death in organ failure

Paxian M (1), Brunkhorst FM (1), Paxian G (1), Masuch M (1), Krenn CG (2), Reinhart K (1), Bauer M (1)

(1) Department of Anesthesiology and Critical Care Medicine, University Jena, Germany, (2) Department of Anesthesiology and General Intensive Care, Medical University of Vienna, General Hospital, Vienna, Austria

Introduction: Apoptosis and related forms of cell death have been suggested to contribute to a dysregulated immune response in shock and sepsis while their role in the development of end organ failure is discussed controversially.

Methods: For a systematic evaluation of an activation of distinct cascades of apoptotic signaling, we analyzed serum levels of members of the TNF/TNFR family and the caspase cleavage product cytokeratin-18 neopeptide (CK-18) in patients with SIRS (n=49) or sepsis (n=38), which were subsequently correlated to disease severity.

Results: Serum concentrations of markers of the TNF/TNFR family were higher in infectious as opposed to non-infectious systemic inflammation, albeit only sFas was increased above while sFasL and TRAIL levels were below levels observed in healthy volunteers. Increased serum levels of the caspase-3 cleavage product CK-18 were restricted to overt multi organ failure. A detailed exemplary analysis of apoptotic signaling correlating with liver dysfunction revealed a similar restriction of CK-18 release to most severe forms of liver failure along with the appearance of cytochrome c, but lack of sFasL in serum indicative of hepatocellular apoptotic injury via the intrinsic / mitochondrial pathway.

Conclusion: Our data are consistent with effective protection of solid organs against apoptotic injury in mild and moderate organ dysfunction while programmed cell death seems to contribute to overt organ failure, e.g. of the liver.

Acknowledgement: This study was supported by a competitive, peer-reviewed R&D grant from the Thuringian Ministry of Science (TMWFK)

090

Infection 2007; 35 (Suppl. II): 22

Viral enteritis mimicking septic necrotizing enterocolitis in very preterm neonates

Gortner L, *Wurm D

Department of Pediatrics and Neonatology, University Hospital of the Saarland, Homburg/Saar, Germany

Background: Necrotizing enterocolitis (NEC) is believed to occur secondary to local reduced immunity, impaired splanchnic perfusion and bacterial overgrowth resulting in sepsis, often caused by gram-negative bacteria. NEC is the leading cause of death from gastrointestinal disease in preterm newborns and the most common reason for neonatal surgery in case of intestinal perforation.

Methods: 4 very preterm neonates (birth weight 450 – 1550 g, gestational age 26+3 to 31+4 weeks) developed signs of late onset neonatal infections with poor capillary refill, hypotension, abdominal distension, feeding intolerance and bloody stools. The infants were on partial enteral nutrition (Alphare) at the time of first symptoms of NEC (6th to 52nd day of life). Bacterial blood cultures and stool analyses were negative in all preterms, however, viral diagnostics revealed Astrovirus in one, Astrovirus plus Norovirus in two and Adenovirus plus Norovirus in one patient. Negative virologic findings were obtained after a mean of 5 days after manifestation of viral enterocolitis. All preterms survived without sequelae.

Conclusion: Virus-induced gastroenteritis may mimic bacterial enterocolitis, however, the clinical course was without need for surgical intervention and thus more benign compared to typical necrotizing enterocolitis causing late onset neonatal sepsis.

091

Infection 2007; 35 (Suppl. II): 22

Posttraumatic stress disorder and depression in long-term survivors of sepsis

*Rothaug J, Zimmer A, Mescha S, Marx G, Meißner W

Department of Anaesthesiology and Intensive Care Medicine, University Jena, Germany

Introduction: In Germany about 45.000 patients survive sepsis per year. With the increasing number of survivors the long-term quality of life becomes more and more an issue. ICU treatment is associated with physical and psychological stress. In some patients this stress leads to the development of posttraumatic stress disorder (PTSD) (1). PTSD and depression were assessed in long-term survivors of sepsis.

Methods: 63 patients were approached who survived severe sepsis or septic shock (ACCP/SCCM) on our ICU. 2 screening instruments were used: for assessment of PTSD the IESR (Impact of Event Scale, Revised) was used, consisting of the three sub-scales 'intrusion', 'avoidance' and 'arousal'. The IESR is event independent, therefore patients were asked in the introduction to describe the most aggravating events they experienced on the ICU. For screening of depression the ADS (Allgemeine Depressions Skala) was applied.

Results: 36 patients, 25 males, 11 females returned the questionnaires. Mean age was 63 ± 16 years. The median follow-up time was 2 years after discharge. In the IESR mostly mentioned as traumatising events were nightmares, helplessness and dependence from staff, and pain. Subjects were asked to answer the IESR with regard to these most traumatising events. 24 subjects filled in the IESR. Subjects with a positive score above 0.0 are likely to suffer from PTSD. 5 patients (20.8%) achieved positive scores. Neither age (t-test: p=0.393) nor gender (Chi2 test, p=0.608) had an influence on the results. 29 patients returned the ADS. The cut-off score for depressive symptoms is 16, with increasing scores pointing out increasing intensity of symptoms. 62% (24 patients) scored 16 or higher. 5 patients (17%) had scores from 16-20, 10 patients (35%) scored from 21-30, and 3 patients (10%) scored higher than 30. Neither age (t-test, p=0.623) nor gender (Chi2 test, p=0.197) had an impact on the result.

Discussion: 20% of the observed sepsis survivors are likely to suffer from PTSD. As the IESR is only a screening instrument, clinical interviews would be necessary to confirm this diagnosis. The ADS scores show that many patients who underwent treatment in our ICU suffer from depressive symptoms. Again, to support this diagnosis, clinical interviews would be required. Recent publications support these findings (2). The median follow-up time of 2 years suggests that both PTSD and depression are already in a state of chronification. It remains unclear whether the subjects were burdened with depressive symptoms already before their admission to ICU. However, with regard to PTSD the symptoms could be attributed to the ICU stay, as the subjects filled in the IESR considering the events during their stay.

Conclusion: In a screening survey of sepsis survivors 20% indicated symptoms of PTSD, 62% displayed acute depressive symptoms. To prevent chronification of these psychiatric disorders early psychotherapeutic treatment should be available for ICU.

Reference: Critical Care (2007) 11: R27

092

Infection 2007; 35 (Suppl. II): 23

Changes in action of muscle and nerves activity in studies of clinical neurophysiology in patients with advanced sepsis. Observations based on three different cases

Huber J (1), Bryl A (1), Trojanowska I (2), Śniatkowska-Bartkowska A (3), Smuszkiewicz P (2), Witkowska A (1)

(1) Department of Pathophysiology of Locomotor Organs, Karol Marcinkowski University of Medical Science in Poznań, Poland, (2) Department and Clinic of Anaesthesiology, Intensive Therapy and Pain Treatment, Karol Marcinkowski University of Medical Science in Poznań, Poland, (3) Department of Pediatric Anaesthesiology and Intensive Therapy, Karol Marcinkowski University of Medical Science in Poznań, Poland

In patients with sepsis a precise neurophysiological diagnosis allows to specify the early symptoms of disease within muscle and nervous system, monitoring its progress and directing towards treating as well as exposing additional injuries not linked to the sepsis itself. Effects of sepsis were differently described as evoking symmetrical miogenic rather than neurogenic changes in more proximal than distal muscles. Sepsis originated polyneuropathy was defined as symmetric abnormalities expressed in distal parts of motor and sensory nerve fibers. Observed in the sepsis changes are mostly axonal, with the greater expression in the sensory fibers. Little attention was paid to the advancement of this disease and pathological disturbances accompanied by the patients age related alterations in the functioning of the nervous system.

This study concerns three cases of advanced sepsis in patients aged 14, 22 and 48 years treated more than three months. Besides neurological examinations, neurophysiological tests including electromyography (EMG), electroneurography (ENG; M, F and H-wave studies as well as sensory conduction studies-SCV) within upper and lower extremities has been performed.

Our findings shows the picture of changes in the muscle motor units activity and nerve transmission as depending primarily to the patient age with minor influence of the sepsis advancement. Miogenic changes can be found in muscles only within 1 month after beginning of the incident, later (usually after 2 months) they are preferably neurogenic and they can be clearly seen only in distal muscles. Sensory disturbances are early, serious and constant both in proximal and distal parts of extremities. Demyelinating than axonal symmetrical changes were mainly found in distal parts of motor fibers. Result of neurological examinations exactly overlap with neurophysiological findings.

095

Infection 2007; 35 (Suppl. II): 23

Results from the HAN-D-OUT (Hannover Dialysis Outcome) study: Comparison of intensified extended dialysis versus conventional extended dialysis in patients with acute kidney injury and sepsis on intensive care unit

*Hafer C, Jahr N, Vahlbruch J, Fliser D, Haller H, *Faulhaber-Walter R
Abteilung Nephrologie, Medizinische Hochschule Hannover

Introduction: Increasing the dose of renal replacement therapy (RRT) has been shown to improve survival in patients with AKI for both, intermittent hemodialysis (IHD) and continuous renal replacement therapy (CRRT). Data from AKI patients with CRRT suggest that this improvement of survival is not seen beyond a certain dose of RRT. Evidence based data for extended dialysis (ED), which combines the advantages of IHD and CRRT, are missing.

Objective: To compare mortality of patients with acute renal failure (AKI) and sepsis on the intensive care-unit (ICU) randomized to receive standard extended dialysis (SED) or intensified extended dialysis (IED).

Methods: We performed a prospective randomized study between 07/2002 and 05/2006 in the ICU at the Hannover Medical School. 157 patients with acute kidney injury were randomized to receive either SED dosed to maintain plasma urea levels between 20-25 mmol/l, or IED dosed to maintain plasma urea levels near-normal (<15 mmol/l). In the latter group patients received at least two daily treatments within the first 2 days after initiation of RRT. We used the same dialysis system (Genius™) and high-flux polysulphone dialyzers for all RRTs. The outcome measures were survival at day 14 (primary), and survival and renal recovery (RR), as defined by no need of RRT at day 28 (secondary) after initiation of RRT. Here we present the analysis of a subgroup of patients with sepsis (n= 71, 45.2%), defined according to guidelines of the Deutsche Sepsis Gesellschaft. Statistical analysis was intention-to treat and done by f-test, t-test, Kaplan-Meier curve and log rank test. Significance was accepted as p<0.05.

Results: 66 patients were eligible for final analysis (SED: n=27 vs. IED n=39). Both groups were matched for baseline variables like gender, age, BMI, APACHE 2 or kidney function. The IED group received a significantly higher dose of dialysis (p<0.05) resulting in a significantly lower plasma urea already at 24h after initiation of RRT (i.e. at 72h: SED: 24.5±9.2 mmol/l vs. IED: 11.3±4.8 mmol/l, p<0.0001) and remaining significant for two weeks. We detected neither differences in survival at day 14 and day 28 after initiation of RRT nor for RR (survival at day 14: SED: 59.3% vs. IED 56.4%, n.s.; survival at day 28: SED: 48.1% vs. IED: 41%, n.s., RR: SED: 61.5% vs. IED 50%, n.s.).

Conclusion: IED vs. SED does not improve survival or renal recovery within 28 days after initiation of RRT in septic patients with AKI. Increasing the dose of RRT aiming to normalize plasma urea is not associated with a further reduction of mortality in patients with sepsis and AKI.

096

Infection 2007; 35 (Suppl. II): 23

Effects of implementation of the Surviving Sepsis Campaign's Sepsis Bundles on quality of care

Bloos F, Marx G, Reinhart K.

Department of Anaesthesiology and Intensive Care Medicine, University Hospital Jena, Germany

The Surviving Sepsis Campaign (SSC) has released its guidelines in 2004 [1]. However, implementation of such guidelines may be incomplete due to their number and complexity. The sepsis bundles are a concept to improve application of guidelines [2]. Goal of this study was to assess the effects of the sepsis bundles on quality of care on a single intensive care unit (ICU).

Methods: All patients, who developed severe sepsis or septic shock between Nov 2005 and March 2007, were included into the study. In Aug. 2006, a standard operating procedure (SOP) based on the sepsis bundles of the SSC has been implemented. The SSC's Chart Review Database was used to monitor the application of the bundles [3].

Results: Before the sepsis bundles were introduced, performance was inadequate regarding fast application of a broad spectrum antibiotic, obtaining blood cultures, and implementation of our ICU policy checking the indication for drotrecogin alfa (activated). All three items were improved after implementation of the SOP while maintenance of blood glucose levels decreased after SOP introduction. There was no difference in outcome between the two groups.

Data are expressed as frequencies or median (25%, 75% percentile). **Conclusion:** Assessment of quality of care with tools such as the Chart Review Database can identify shortcomings in the care of sepsis patients. Implementation of the sepsis bundles improved performance in items identified as inadequate.

References: [1] Dellinger RP et al. Crit Care Med 2004; 32: 858 [2]
 Levy MM et al. Crit Care Med 2004; 32: S595 [3]
http://www.survivingsepsis.org/manual_database

	Before SOP N = 115	After SOP N = 71	p
<i>Resuscitation Bundle</i>			
Lactate measured within 6 h	99.1 %	100 %	1.000
Blood culture before broad spectrum antibiotics	49.6 %	63.4 %	0.071
Broad spectrum antibiotics within 1 h	52.2 %	71.8 %	0.009
Fluid bolus applied	96.5 %	94.2 %	0.478
CVP >8 mmHg within 6 h	75.4 %	84.1 %	0.195
ScvO ₂ >65 % within 6 h	54.4 %	59.4 %	0.541
<i>Management Bundle</i>			
Indication for low dose hydrocortisone checked within 24 h	80.4 %	83.3 %	0.693
Indication for drotrecogin alfa (activated) checked within 24 h	60.0 %	88.7 %	<0.0001
Blood glucose maintenance within 24 h	80.9 %	67.6 %	0.052
Median inspiratory plateau pressure <30 cm H ₂ O	85.8 %	85.1 %	1.000
<i>Outcome</i>			
Hospital mortality	51.3 %	50.7 %	1.000
ICU length of stay [days]	15 (6; 24)	16 (5; 25)	0.944
Hospital length of stay [days]	34 (21; 50)	34 (21; 55)	0.608

Pathophysiology

002

Infection 2007; 35 (Suppl. II): 25

Phagocytosis induced cell death in neonatal macrophages is diminished after group-B-streptococci (GBS) challenge

*Gille C (1), Leiber A (1), Spring B (1), Kempf V (2), Spellerberg B (3), Poets CF (1), Orlowski TW (1)

(1) Children's Hospital, University Tuebingen, Germany (2) Department of Medical Microbiology and Hygiene, University Tuebingen, Germany (3) Institut für Mikrobiologie und Immunologie, University Ulm, Germany

Introduction: Apoptotic-inflammatory imbalance plays an essential role in the pathophysiology of sepsis, the most common cause of neonatal mortality. Phagocytosis and intracellular degradation are early events in pathogen-host interaction, resulting in a proinflammatory activation. Phagocytosis induced cell death (PICD) may counteract an exaggerated inflammation. Neonatal monocyte derived macrophages (CBMF) are equivalent to MF of adults in phagocytosing and killing GBS.

Objective: Compared to PBMF, CBFM are less sensitive towards apoptotic signals mediated by phagocytosis of GBS.

Methods: PBMF and CBFM were stimulated with green fluorescent protein (gfp)-labelled GBS for 60 minutes. After removal of free bacteria cells were either fixed, stained with anti-CD14 monoclonal antibodies and phagocytic activity was analysed by flow cytometry or cultured again for up to three hours to monitor intracellular degradation of GBS by loss of gfp-fluorescence. MF apoptosis was detected after four hours by quantification of phosphatidylserine externalisation via Annexin-V-staining, hypodiploid DNA content was analysed after 24 hours. Results: 60 minutes post infection $31 \pm 18\%$ of PBMF vs. $33 \pm 32\%$ of CBFM were gfp-positive, mean gfp-fluorescent intensity (MFI) was 43 ± 15 vs. 33 ± 12 (all $p > 0.05$).

Results: Three hours after removal of bacteria less than 10% of PBMF and CBFM remained gfp-positive. After four hours, PBMF up-regulated Annexin-V binding by 5-fold after bacterial challenge while there was no enhanced Annexin-V binding on CBFM ($p < 0.05$). After 24 hours, 23% of PBMF vs. 5% of CBFM showed a hypodiploid DNA content ($p < 0.05$) although MF of both groups underwent apoptosis in the presence of staurosporine (32% vs. 39%; $p > 0.05$).

Conclusion: CBFM were found less sensitive towards apoptotic signals after phagocytosis of GBS than PBMF while staurosporine induced apoptosis was equivalent. This may play a role in hyperinflammation during neonatal sepsis.

003

Infection 2007; 35 (Suppl. II): 25

Proinflammatory cytokines cause downregulation of renal urea transporters during sepsis

*Schmidt C, Kurt B, Bucher M

Department of Anaesthesiology, Universital Hospital Regensburg, Germany

Introduction: The pathogenesis of endotoxemic tubular dysfunction with failure in urine concentration is poorly understood. Urea plays an important role in the urinary concentrating mechanism and expression of the urea transporters UT-A1, UT-A2, UT-A3, UT-A4 and UT-B is essential for tubular urea reabsorption.

Objective: The present study attempts to assess the regulation of renal urea transporters during severe inflammation in vivo.

Methods: By agreement of the animal protection committee C57BL/6J mice were injected with lipopolysaccharides (LPS, 10 mg/kg) or proinflammatory cytokines. Hemodynamic, renal parameters and the expression of renal urea transporters were investigated. To clarify the

role of cytokines and renal ischemia in the regulation of renal urea transporters, experiments with cytokine knock-out mice, mice treated with low dose LPS (1,5 mg/kg) as sepsis model without induction of hypotension, glucocorticoid-treated mice, and mice with renal artery clipping serving as model for renal ischemia were performed.

Results: LPS-injected mice (10 mg/kg) presented with reduced GFR, fractional urea excretion and inner medulla osmolality associated with a marked decrease in expression of UT-A1, UT-A2, UT-A3, UT-A4 and UT-B. Similar alterations were observed after application of TNF-alpha, IL-1beta, IFN-gamma or IL-6. LPS-induced downregulation of urea transporters was not affected in knockout mice with deficient TNF-alpha, IL-receptor-1, IFN-gamma or IL-6. Glucocorticoid treatment inhibited LPS-induced increases of tissue TNF-alpha, IL-1beta, IFN-gamma or IL-6 concentration, diminished LPS-induced renal dysfunction and attenuated the downregulation of renal urea transporters. Injection of low dose LPS (1,5 mg/kg) also led to renal dysfunction paralleled by a downregulation of renal urea transporters without alterations in blood pressure. Renal ischemia induced by renal artery clipping did not influence the expression of urea transporters.

Conclusion: Our findings demonstrate downregulation of renal urea transporters which likely accounts for tubular dysfunction during sepsis and suggest that downregulation of renal urea transporters during LPS-induced ARF is mediated by proinflammatory cytokines and is independent from renal ischemia due to sepsis-induced hypotension. Furthermore, our results contribute to understand the failure of single anti-cytokine strategies on one hand and the beneficial effects of glucocorticoids on the other hand in the therapy of septic patients.

Acknowledgements: The study was supported by grants from the German Research Foundation (SFB 699; B3/B4).

004

Infection 2007; 35 (Suppl. II): 25

Inhibition of Caspase-3 activation by Drotrecogin alfa (activated) in a murine model of sepsis

*Schürholz T (1), Wagner T (1), Saenger J (2), Claus RA (1), Simon TP (1), Reinhart K (1), Marx G (1)

(1) Department of Anesthesiology and Intensive Care, Friedrich-Schiller-University, Jena, Germany (2) Institute of Pathology Bad Berka, Germany

Introduction: Since Drotrecogin alfa (activated) (DAA; recombinant activated Protein C) was introduced for therapy in severe sepsis several mechanisms on the mode of action were discussed. Yet, some effects were investigated only in cell culture [1]. Recently, anti-apoptotic effects of DAA were shown on circulating mononuclear cells of septic shock patients [2]. Caspase activation may directly alter cell function of cardiomyocytes. A possible relationship between endotoxin-induced caspase activation and reduced contractile reserve was described [3]. However, a direct effect of DAA on solid organ tissue in-vivo is not described.

Objective: To investigate whether there is an effect of DAA on Caspase-3 in myocardial tissue in a murine model of sepsis.

Methods: Male NMRI-mice ($n=21$, body weight $30 \pm 3g$) were randomized to sham-treatment, sepsis by cecal ligation and puncture (CLP) and vehicle infusion, or CLP sepsis with DAA infusion ($24 \mu g/kg/hr$). 48 hrs prior to CLP all mice were given a permanent i.v.-line (self-made) and an arterial transmitter (PA-C10, St. Paul, MN, USA) to measure heart rate (HR) and mean arterial pressure (MAP). Activity was recorded by scoring from 1 (healthy) to 5 (agony) [4]. CLP was adjusted to survive 24 hrs. All mice were killed 12 hrs after CLP after scoring, and recording of vital data. Hearts were fixed in formalin and embedded in paraffin. Immunohistochemical analysis of the paraffin sections was performed using the avidin-biotin-peroxidase-complex (ABC) method. For Caspase-3 analysis a rabbit polyclonal anti-Caspase-3 Cleaved antibody (CPP32, Zytomed, Berlin, Germany) was

used (dilution 1:100) after heat pretreatment. Anti-Caspase-3 positive cells were counted in 10 fields in light microscopy (original magnification $\times 625$) of each heart and the average was recorded.

Results: 12 hrs after CLP scoring revealed significantly higher activity in sham-treated (score 1.0 ± 0 ; $p=0.001$) and DAA-treated mice (score 1.8 ± 0.7 ; $p=0.015$) compared to control (score 3.1 ± 0.9). HR was non-significantly higher in sham and to a lesser degree in DAA mice (Sham 650 ± 38 /min; DAA 547 ± 175 /min; Control 530 ± 143 /min). MAP was significantly higher in sham group ($p=0.031$) and non-significantly higher in DAA group when compared to control (Sham 137 ± 15 mmHg; DAA 111 ± 26 mmHg; Control 97 ± 4 mmHg). Anti caspase-3 positive cells were significantly lower in sham-treated mice (2.1 ± 0.6 cells; $p=0.001$) and DAA mice (4.9 ± 2.7 cells, $p=0.01$) compared to controls (9.6 ± 4.0 cells).

Conclusion: To our knowledge this is the first description that treatment with DAA leads to a lower quantity of active Caspase-3 in myocardial tissue in murine sepsis. This may be a sign of an organ-protective, anti-apoptotic effect of DAA in sepsis.

References: [1] Joyce DE et al., J Biol Chem 2001; 276(14): 11199. [2] Bilbault P et al., Crit Care Med 2007; 35(1): 69. [3] Lancel S et al., Circulation 2005; 111(20): 2596. [4] van Griensven M et al., Shock 2002; 18(5): 445.

007

Infection 2007; 35 (Suppl. II): 26

T-cell-activation by enteric glia cells: Primary origin for intestinal bacterial translocation in sepsis?

*Hollenbach E (1), Maher T (2), Zöller M (3), Stremmel W (4), Neumann M (5), Malfertheiner P (6), Vallance B (7), Rühl A (8), Engelmann L (1)

(1) Medical ICU, University Leipzig, Germany, (2) Harvard Medical School and Massachusetts Institute of Technology (MIT), Department of Pharmacology and Allied Health Sciences, Cambridge, Massachusetts, USA, (3) German Cancer Research Institute, Department of Tumor Immunology, University Heidelberg, Germany, (4) Department of Gastroenterology and Hepatology, University Heidelberg, Germany, (5) Institute of Experimental Internal Medicine, University Magdeburg, Germany, (6) Department of Gastroenterology and Hepatology, University Magdeburg, Germany, (7) British Columbia Research Institute, University of British Columbia, Vancouver, Canada, (8) Institute of Human Biology, Technical University Munich, Germany

Introduction: Enteric glia (EGC) are the most numerous cell type in the enteric nervous system. It has been previously shown by our group that EGC participate in intestinal immune responses via their phagocytosis, induction of MHCII/ICAM1 by pro-inflammatory signals, conspicuous antigen-specific stimulation of syngeneic T-cells, and their production of proinflammatory cytokines.

Methods: We have assessed putative important pro-inflammatory signaling pathways in purified EGC cultures from rat intestine after exposure to bacterial peptidoglycans (PGN). To selectively expose EGC to PGN, we have used an *H. pylori* based model of intracellular translocation of PGNs.

Results: A strong peak of IkB α -degradation and nuclear translocation of p65 occurred 60-90 min after infection with the *H. pylori* wild-type (HPWT) strain proving NF- κ B activation by immunoblot, whereas no significant IkB α -degradation could be detected after EGC infection with an isogenic mutant *H. pylori* (HPM) strain that is unable to convey the intracellular translocation of PGN. These results were confirmed by EMSA. To further analyze the signaling pathways induced by immune-activation of EGC, we studied activation of NOD in EGC upon PGN challenge. RICK (a downstream kinase of NOD leading to NF- κ B activation) activity after intracellular PGN translocation was assessed by *in vitro* kinase assays. These assays demonstrated a dramatically upregulated RICK activity in EGC after PGN challenge with a maximum at 60min after infection. In contrast, HPM-strain could not activate RICK in enteric glia. *In vitro* data were verified and

reproduced in an *in vivo* infection mouse model using salmonella typhimurium. Surprisingly, immunohistochemical analysis revealed nuclear translocation of phospho-p65 in EGC in the colon 24h after infection, while the maximum of NF- κ B activation in "commonly" immunoreactive cells like macrophages were detected after 48h.

Conclusion: In summary, evidence was provided that EGC strongly and early respond to intracellular PGN with activation of RICK/NF- κ B. These data support consideration of EGC as another cell-type in gastrointestinal immunology. EGC may play a major role in the initiation process of intestinal bacterial translocation in sepsis by increasing the permeability of the intestinal wall through an induction of bacterial triggered inflammation.

010

Infection 2007; 35 (Suppl. II): 26

Functional coronary autoregulation as a likely cause of a reduced ischemia-reperfusion injury in septic hearts

*Zausig YA (1), Binder L (2), Löffler EK (1), Missler G (1), Sinner B (1), Zink W (1), Graf BM (1)

(1) Department of Anaesthesiology, Emergency and Intensive Care Medicine, Georg August University of Goettingen, Germany (2) Department of Clinical Chemistry, Georg August University of Goettingen, Germany

Introduction: After ischemia septic hearts show mechanical and morphological reperfusion injury to a lesser extent than control hearts (McDonough 1994). Existing data show a different coronary reaction compared to control hearts. In this regard, several causes of cardio protection are discussed, e.g. up regulation of iNOS or modulation via adenosine (Westerhof 2006).

Objective: We postulated a functioning coronary autoregulation as a potential cause for a reduced ischemia-reperfusion injury in septic hearts.

Methods: Polymicrobial sepsis was induced in male Wistar rats (230-260 gr.) via cecal ligation and single puncture (CLP, $n=15$). Control hearts received only a laparotomy (Sham, $n=13$). After 24 hours of incubation hearts were isolated, retrograde perfused in Langendorff apparatus and paced at controlled rate of 350 beats per minute. After stabilization, a 40 minute lasting no-flow ischemia was induced followed by 40 min of reperfusion. Mechanical performance, perfusion pressure (PP), and troponin T release were investigated at constant coronary flow (CF, 10 ml/min) in one group (Sham $n=7$, CLP $n=9$); and coronary autoregulation (intrinsic capacity of an organ to regulate its own blood flow when perfusion pressure is challenged) was determined by changing perfusion pressure (10 mmHg step by step) in another group (Sham $n=6$, CLP $n=6$). Autoregulation index was calculated by $ARI = (1 - (CF/CF)) / (PP/PP)$ to quantify the autoregulatory capacity of the coronary flow. Coronary reserve was examined by an intracoronary bolus of adenosine before and after ischemia. All results are presented as mean $\pm$ standard deviation. $P < 0.05$ was taken as significant.

Results: Before ischemia CLP hearts showed a significant reduction in left ventricular pressure (LVP, CLP: 63 ± 5 vs. Sham: 115 ± 5 mmHg) and contractility ($+dp/dt$, CLP: 1712 ± 154 vs. Sham: 3003 ± 90 mmHg/sec). After reperfusion CLP hearts showed no difference in LVP (58 ± 12 mmHg) and $+dp/dt$ (1539 ± 340 mmHg/sec) compared to pre-ischemic values; while Sham hearts showed a significant reduction. Only Sham hearts showed a significant increase in troponin T levels. Postischemic coronary vascular resistance expressed as an increase in perfusion pressure significantly increased in Sham hearts (74 ± 5 mmHg), but not in CLP hearts (54 ± 3 mmHg). Coronary reserve was preserved in CLP hearts, but not in Sham hearts. Autoregulation Index showed an intact coronary autoregulation at perfusion pressure of 60 – 120 mmHg after ischemia only for septic hearts. Conclusion: After 40 minutes of ischemia septic hearts show a lesser reduction in mechanical performance compared with Sham hearts. Additionally, coronary vascular reaction in septic hearts after ischemia differs explicit from the reaction of Sham hearts. The preserved coronary re-

serve and autoregulation might be the cause of a lesser ischemic reperfusion injury in septic hearts. Additionally, coronary reaction on ischemia and adenosine was delayed in time in septic hearts.

011

Infection 2007; 35 (Suppl. II): 27

Dobutamine, Dopamine, and Epinephrine, but not Levosimendan improve the cardiac performance in isolated septic depressed rat heart

*Zausig YA (1), Geilfuss D (1), Löffler EK (1), Hofer S (2), Sinner B (1), Weigand MA (2), Zink W (1), Graf BM (1)
(1) Department of Anaesthesiology, Emergency and Intensive Care Medicine, Georg August University of Goettingen, Germany,
(2) Department of Anaesthesiology, Ruprecht Karls University of Heidelberg, Germany

Introduction: The recent guidelines of the German Sepsis Society advocate dobutamine to improve cardiac performance in severe sepsis or septic shock (S2 Guideline DSG 2006). But this recommendation is based on a low evidence based level (Grade E – expert opinion). Additionally, there is only a limited amount of experimental studies comparing the effects of inotropic drugs in isolated septic hearts. In addition, the calcium-sensitizer, levosimendan offers a new therapeutic approach in septic myocardial depression (Morelli 2005).

Objective: The aim of our study was to compare the direct cardiac effects of dobutamine, dopamine, epinephrine, and levosimendan in the isolated septic rat heart.

Methods: Polymicrobial sepsis was induced in male Wistar rats (220-250 gr.) via cecal ligation and single puncture (CLP, n=40). After 20 hours of incubation hearts were isolated and retrograde perfused at constant pressure (55 mmHg) in Langendorff apparatus. After stabilization the hearts were randomly assigned by lottery to four groups (10 hearts each) and received dopamine, dobutamine, epinephrine or levosimendan at concentrations of 10^{-9} to 10^{-3} M with one of these drugs for a period of 15 min followed by a drug-free washout period. Mechanical performance (LVP: left ventricular pressure, +dp/dt: contractility, -dp/dt: lusitropie) and heart rate (HR) were recorded. O_2 tension, electrolytes, and lactate in the coronary outflow were measured to determine oxygen delivery (DO_2), percentage oxygen extraction (O_2Ext), myocardial oxygen consumption (MVO_2), and cardiac efficiency. Raw data from each functional and metabolic variable were compared by analysis of variance (ANOVA) with repeated measures. $P < 0.01$ was considered to be statistically significant.

Results: Except for levosimendan, all tested drugs showed over a range from 10^{-9} to 10^{-3} M concentration dependent significant positive chronotropic, inotropic and lusitropic effects. Dobutamine, dopamine, and epinephrine increased maximal HR (21, 26, 30%, respectively), +dp/dt (187, 194, 260%, respectively), and -dp/dt (202, 211, 244%, respectively). This was accompanied by an increase in MVO_2 (118, 113, 142%, respectively), and an adequate gain in coronary flow (56, 62, 50%, respectively). Levosimendan, only showed a trend towards a positive inotropic effect, but this was not significant. Although, this was accompanied by a significant increase in MVO_2 (47%). High concentrations of dobutamine ($>10^{-5}$ M), dopamine ($>10^{-3}$ M), epinephrine ($>10^{-6}$ M), and levosimendan ($>10^{-5}$ M) showed negative chronotropic and inotropic effects.

Discussion: Dobutamine, dopamine und epinephrine showed a dose dependent increase in heart rate, left ventricular pressure, contractility and relaxation in the isolated septic heart. In contrast, an improvement of cardiac performance in septic depressed heart via a short-term intervention with levosimendan showed no beneficial effect in this model.

012

Infection 2007; 35 (Suppl. II): 27

Direct cardiac effects of ketamine and propofol in isolated septic rat heart

*Zausig YA (1), Busse H (1), Löffler EK (1), Busch C (2), Sinner B (1), Weigand MA (2), Zink W (1), Graf BM (1)
(1) Department of Anaesthesiology, Emergency and Intensive Care Medicine, Georg August University of Goettingen, Germany,
(2) Department of Anaesthesiology, Ruprecht Karls University of Heidelberg, Germany

Introduction: Throughout their stay on the intensive care unit septic patients continuously receive sedatives to reduce anxiety and agitation. Sedatives show different haemodynamic characteristics, e.g. ketamine showed negative cardiac effects in healthy guinea pig hearts (Graf 1995). These effects might influence cardiac performance negatively, especially in patients with septic depressed heart function. At this time there is no recommended sedative for patients in severe sepsis or septic shock (Vender 2004), and there is only a limited amount of experimental studies focusing on direct cardiac effects of sedatives in septic hearts.

Objective: The aim of our study was to compare the direct cardiac effects of ketamine and propofol in the isolated septic rat heart.

Methods: Polymicrobial sepsis was induced in male Wistar rats (220-250 gr.) via cecal ligation and single puncture (CLP, n=20). After 20 hours of incubation hearts were isolated and retrograde perfused at constant pressure (55 mmHg) in Langendorff apparatus. After stabilization the hearts were randomly assigned by lottery to two groups (10 hearts each) and received ketamine or propofol at concentrations of 10^{-7} to 10^{-4} M with one of these drugs. Mechanical performance (LVP: left ventricular pressure, +dp/dt: contractility, -dp/dt: lusitropie), and heart rate (HR) were recorded. O_2 tension, electrolytes, and lactate in the coronary outflow were measured to determine oxygen delivery (DO_2), percentage oxygen extraction (O_2Ext), and myocardial oxygen consumption (MVO_2), and cardiac efficiency. Raw data from each functional and metabolic variable were compared by analysis of variance (ANOVA) with repeated measures. $P < 0.05$ was considered to be statistically significant.

Results: Propofol and ketamine showed over a range from 10^{-7} to 10^{-5} M concentration dependent no significant effect on LVP, +dp/dt or HR. But propofol showed at 10^{-4} M significant negative inotropic (LVP: -30%; +dp/dt: -38%) and chronotropic (-29%) effects. This was accompanied by a reduction in MVO_2 of 16% at 10^{-4} M. Ketamine only showed a slight but significant reduction in HR (-14%) at 10^{-4} M compared to control values. Even though this reduction was to a lesser extent compared to propofol at equimolar concentration (10^{-4} M).

Discussion: Propofol, but not ketamine showed significant dose-dependant negative inotropic as well as chronotropic effects at concentrations $>10^{-5}$ M in the isolated septic heart. Septic depressed heart function might deteriorate with these high plasma concentrations of propofol achieved by bolus administration or long-term infusion (Kim 2007) in sedated patients in severe sepsis or septic shock, especially in case of multiple organ failure accompanied by a decreased hepatic and renal metabolism. Ketamine showed at concentration from 10^{-7} to 10^{-4} M a safe administration regarding influence on inotropic performance in septic rat heart.

018

Infection 2007; 35 (Suppl. II): 28

Early microcirculatory failure in the septic rat brain assessed by the neurovascular coupling and cerebral autoregulation

*Rosengarten B (1), Klatt S (1), Hecht M (1), Schermuly R (2), Grimminger F (2), Kaps M (1)

(1) Department of Neurology, University Giessen, Germany,

(2) Department of Internal Medicine IV, University Marburg, Germany

Introduction: Early microcirculatory failure is a motor of sepsis related organ dysfunction. The brain is very sensitive against an inappropriate blood supply. Two vasoregulative mechanisms assure adequate cerebral blood flow. The neurovascular coupling describes the relation of cortical activity and the local blood supply whereas the cerebral autoregulation maintains constant cerebral blood flow despite perfusion pressure changes.

Objective: A sepsis syndrome might affect both vasoregulative mechanisms. The neurovascular coupling allows simultaneous investigation of neuronal as well as vascular function and is ideally suited to investigate early changes in both systems. Independent from cortical function the cerebral autoregulation gives information on vasoregulative function.

Methods: Administering 1 or 5mg/kg i.v. LPS in 10 rats we compared somatosensory evoked potentials (SEP) and evoked LDF responses utilizing electrical forepaw stimulation and assessed cerebral autoregulation by repetitive carotid compression. Cerebral blood flow, blood pressure, evoked electrical and hemodynamic responses as well as pH and lactate levels were recorded for 3.5h under sepsis progression.

Results: All rats developed a severe sepsis syndrome. Microcirculatory failure as assessed by neurovascular coupling and autoregulation preceded changes in evoked potential amplitudes by 2h.

Conclusion: Microcirculatory failure precedes changes in evoked potentials and thus might play a considerable role in development of septic encephalopathy. Cerebrovascular risk cannot be assessed referring only to macrocirculatory parameters or knowledge of overall brain perfusion. Although SEPs indicate an early sepsis syndrome diagnosis might be improved by addressing also microcirculatory function.

022

Infection 2007; 35 (Suppl. II): 28

Pharmacodynamics and organ storage of hydroxyethyl starch in acute hemodilution in pigs

*Rosenhagen C (1), Schönfeld AH, Vogt N (2), Pfenninger E (3), Eisenbach C (4), Weigand MA (1)

(1) Department of Anaesthesiology, University Heidelberg, Germany, (2) Cardioanaesthesiology, Department of Surgery, University Ulm, Germany, (3) Department of Anaesthesiology, University Ulm, Germany, (4) Department of Internal Medicine IV, University Heidelberg, Germany

Introduction: HES is one of the most frequently used plasma substitutes. A variety of different HES solutions exist worldwide, which differ greatly in their pharmacologic properties.

Objective: To determine the differential influence of molecular weight and the degree of substitution of HES solutions on pharmacodynamics and pharmacokinetics including organ storage in a model of acute hemodilution in pigs.

Methods: After bleeding, 20ml/kg, animals were substituted with 6% HES preparations (200/0.62, 200/0.5, and 100/0.5). Quantitative analysis of HES concentrations in body fluids (plasma, urine) was performed and samples of liver, spleen, kidney, lung and lymph nodes were extracted.

Results: We did not observe any significant differences in the ability to sufficiently achieve plasma volume expansion and restoration of

macrocirculation, nor maintenance of indicators of microcirculation between the groups. Urine production was significantly higher in HES-treated animals and highest in animals substituted with HES 100/0.5. Plasma clearance was measured under steady-state conditions with significantly reduced clearance for the HES 200/0.62 group compared with HES 100/0.5 and HES 200/0.5 (6.6 vs. 13.2 and 13.9ml/min; $P \leq 0.001$), thus being dependent on the degree of substitution. Even after only 6h, the amount of infused HES not detectable in either blood or urine was significantly higher in HES 200/0.62-treated animals (50.7% compared with HES 200/0.5 (28.8%), $P=0.020$ and HES 100/0.5 (28.4%), $P=0.018$), with its proportion rising over time. Finally, we could demonstrate considerable amounts of all HES solutions being stored in liver, kidney, lung, spleen and lymph nodes.

Conclusion: All preparations analyzed sufficiently restored macro- and microcirculation; however, for all solutions relevant tissue storage of HES was observed after only 6h.

023

Infection 2007; 35 (Suppl. II): 28

PPARgamma inhibits formation of reactive oxygen in primary mouse spleen macrophages

*von Knethen A, Soller M, Johann AM, Brüne B

Institute of Biochemistry, I-Pathobiochemistry, Faculty of Medicine, University Frankfurt, Germany

Introduction: Monocyte/macrophage desensitization is a marker of the hypo-inflammatory phase of sepsis. We propose that alternative macrophage activation in response to phagocytosis of apoptotic cells (POAC) is one mechanism provoking an anti-inflammatory macrophage phenotype, thus causing immune paralysis. Using RAW 264.7 mouse macrophages which have been retrovirally transduced with a dominant negative PPARgamma1 mutant we demonstrated recently that inhibition of the oxidative burst in response to POAC is PPARgamma dependent. Because it is well known that PPARgamma dominant negative mutants also affect signaling by other PPARs such as PPARalpha and PPARbeta/delta, it is our intention to generate a PPARgamma conditional knockout mouse system. Thus, PPARgamma expression should be knockedout in primary murine macrophages, to finally verify a PPARgamma-dependent mechanism in primary cells.

Objective: Verification of PPARgamma-dependent macrophage desensitization using primary PPARgamma knock-out macrophages derived from spleen of conditional PPARgamma knockout mice.

Methods: mxCre PPARgamma^{fl/fl} mice were treated i.p. for 1 week with pIpC [250 µg/mouse] at day 1, 4, and 7 to induce PPARgamma deletion in leukocytes. MACS separation for CD11b-positive cells from spleen suspensions was performed using an autoMACS. Enrichment was verified by FACS analysis. Deletion of PPARgamma was verified using genomic DNA- and RT-PCR. For ROS detection cells were divided into 4 FACS tubes/mouse. For 1 h the PPARgamma agonist rosiglitazone [10 µM] was added to tubes 1 and 2, whereas tubes 3 and 4 remained as controls. After 1 h PMA [1 µM] was added to tubes 1 and 3 and incubation continued for 30 min. Afterwards, 20 nM of the redox-sensitive dye hydroethidine was added to all samples for 20 min. ROS production was determined by FACS analysis. In addition, cytopins were performed to follow PKCalpha localization. Results FACS analysis in PPARgamma negative cells revealed no alteration of PMA-elicited ROS production after adding the PPARgamma agonist rosiglitazone. In line in spleen derived macrophages of control mice, expressing PPARgamma, PPARgamma activation inhibited PMA-mediated ROS production. Moreover, PKCalpha translocation was analyzed as well in this experimental setup. We noticed that PKCalpha translocation was restored in PPARgamma knockout macrophages whereas translocation was inhibited in control cells.

Conclusion: In primary spleen derived mouse macrophages activation of PPARgamma provokes desensitization by inhibiting PMA mediated ROS formation. Superoxide production was restored in PPAR-

gamma knockout macrophages derived from spleens of conditional PPARGamma knockout mice. Taking these data into consideration we propose, that PPARGamma mediated alternative macrophage activation in response to phagocytosis of apoptotic cells, which is a marker of sepsis. This might be important for sepsis progression.

029

Infection 2007; 35 (Suppl. II): 29

Cecal ligation and incision: an acute onset model to study severe sepsis in rats

*Scheiermann P (1), Hoegl S (1), Revermann M (1), Ahluwalia D (1), Zander J (2), Boost KA (1), Nguyen T (1), Muhl H (3), Zwißler B (1)

(1) Department of Anaesthesiology, Intensive Care and Pain Therapy, Hospital of the Johann Wolfgang Goethe University Frankfurt/Main, (2) Institute of Medical Microbiology and Infection Control, Hospital of the Johann Wolfgang Goethe University Frankfurt/Main (3) Pharmazentrum/ZAFES, Hospital of the Johann Wolfgang Goethe University Frankfurt/Main

Introduction: Sepsis is a leading cause of death among critically ill patients. It is characterized by infectious origin, systemic inflammation and organ dysfunction. Cecal ligation and puncture (CLP) with subsequent bacterial peritonitis is a common way of inducing sepsis in rodents. However, sepsis progression in the CLP model is slow. Up to this date, severe sepsis with acute onset in animals has been induced mainly through intraperitoneal (i.p.) or intravenous (i.v.) injection of single bacteria species or endotoxin. It has been argued, however, that these models are somewhat artificial and do not adequately mirror the situation in septic patients on ICU.

Objective: We therefore aimed to establish an animal model of severe sepsis characterized by acute onset and high mortality based on the widely accepted CLP model.

Methods: After governmental approval of this prospective randomized study, 28 male Sprague-Dawley rats (weight 527[17] g, Median[Semi-Inter Quartile Range], Rank Sum Test) were anaesthetized by i.p. injection of pentobarbital/fentanyl (p/f). Following tracheotomy, rats were mechanically ventilated. Anaesthesia was maintained by i.v. infusion of p/f. After baseline arterial blood gas analysis (BGA), the cecum was exteriorized through midline laparotomy. In the SHAM-group (14 rats), the cecum was replaced into the abdomen after gentle manipulation. In the other group (14 rats), the cecum and the mesenteric blood vessels were ligated below the ileocecal valve. The cecum was opened through a blade incision on the anti-mesenteric side (cecal ligation and incision, CLI). Subsequently, the cecum was replaced into the abdomen. The abdominal wall was closed in two layers in both groups after i.p. fluid resuscitation. Mean arterial blood pressure (MAP) and BGA were measured hourly for 6.5 h. At the end of observation time or in case of prior death, plasma samples (for ELISA), peritoneal lavage (for cell count), and abdominal swabs (for bacterial culture) were obtained.

Results: Within 6.5 h, mortality was 0% (SHAM) and 50%* (CLI), respectively [$p < 0.05$]. MAP was 125[18]/71[16] mmHg*. Arterial base excess (BE) was -2[1]/-17[4] mmol/l*. Arterial pH was 7.45[0.03]/7.17[0.06]*. Plasma Interleukin-(IL-)1 β level was 10[35]/435[164] pg/ml*. Plasma IL-6 level was 832[824]/19718[9384] pg/ml*. In the peritoneum, mononuclear population was 4[1]/42[36] cells/ml*. Abdominal swabs from SHAM-animals were sterile on agar and showed *E. faecalis*, *P. mirabilis* and *E. coli* growth in the CLI-group.

Conclusion: Bacterial growth proved the infectious origin in this animal model. Systemic inflammation was present in CLI-animals as indicated by an increase in IL-1 β and IL-6 plasma levels and by leukocyte migration into the peritoneum. Organ dysfunction in CLI-animals was reflected by a decrease in MAP, BE, and pH. Thus, we have successfully established an acute onset model of severe sepsis in rats by CLI, which will provide the base for future experimental studies.

032

Infection 2007; 35 (Suppl. II): 29

Evaluation of respiratory parameters in the course of murine sepsis

*Thomas J, Wagner F, Hoffarth B, Georgieff M, Senfleben U
Dept. of Anesthesiology, University of Ulm, Germany

Introduction: The mouse model of sepsis offers extensive possibilities to study pathophysiological principles. Moreover, the use of genetically engineered animals enables us to accomplish the bridging from molecules to physiology. Transgenic mice harbouring well described particular defects of signaling pathways provide a tool to investigate immunological and physiological functions of these pathways during critical diseases like sepsis.

Objective: The murine sepsis models, such as CLP, need to be carefully and well-characterized regarding pathophysiological changes over time. For this purpose, techniques are needed which are as little invasive as possible. We therefore examined respiratory parameters of unrestrained septic animals in order to correlate respiratory changes with the severity of sepsis.

Methods: Respiration of unrestrained adult female C57BL/6 mice was measured using a whole body plethysmograph. Upon isoflurane anesthesia induction CLP surgery (double puncture with 18G-, 20G-, 22G-, 26G-needles) and laparotomy only (Sham) was performed in 4-8 mice for each group. Subsequently, mice received 1ml 0.9% NaCl s.c. once and buprenorphine 25ng/g s.c. twice a day. Minute volume (MV), tidal volume (TV) and breathing rate (f) were detected for 30 min after the operation followed by recurrent measurements every 12 hours for 5 days.

Results: Basic parameters: MV 1.9 ± 0.2 ml/min/g, TV 6.1 ± 0.3 μ l/g, f 237 ± 75 /min. The 5-day-mortality rate of the 5 groups was similar to studies performed earlier in our lab. (Sham: 0%, 26G: 16%, 22G: 25%, 20G: 37%, 18G: 100%). Post-op all groups showed similar respiratory parameters. At 12hrs after CLP MV in the 18G- and 20G group was decreased by 70 % and 35%, respectively. Interestingly, only those animals in the 20G-group that survived showed an increase of MV at 24h and 48h after CLP. The other groups demonstrated inconstant respiratory parameters. Thereafter, 20G-, 22G-, and 26G-treated mice showed similar reduced MV compared to sham mice, which exhibited normal values from 72hrs post-surgery on. From 84hrs on all surviving CLP mice started to increase their MV and fully recovered at 120hrs. Interestingly, the pattern of respiratory frequency closely resembled the course of MV in all groups throughout the observation period. TV did not show clear differences between the groups. It has to be noted that telemetric observation of 22G- and 26G-mice revealed that their over-all activity is suppressed for 72hrs upon sepsis-induction.

Conclusion: This technique provides a suitable tool to study the spontaneous course of respiration during murine sepsis. MV is mainly determined by changes of f and not TV. Especially during the early period severe grades of sepsis are indicated by reduced MV which possibly indicates fatal outcome. Presumably, increased activity after 72hrs leads to enhanced oxygen consumption and, hence, to increased MV. At this time, surviving septic mice appear to recover from this disease.

033

Infection 2007; 35 (Suppl. II): 29

Statins regulate the constitutive expression of HLA-DR by a post-transcriptional mechanism

*Weber SU, Schewe JC, Lehmann LE, Book M, Hoeft A, Stüber F
Department of Anesthesiology and Intensive Care Medicine,
Medical Center, University of Bonn, Germany

Introduction: Epidemiological studies indicate a possible protective effect of statins in the prevention of sepsis. Apart from lowering cholesterol by inhibition of 3-hydroxy-3-methylglutaryl-coenzyme-A

(HMG-CoA) reductase they display pleiotropic immune effects. A lack of HLA-DR (human leukocyte antigen), a component of the major histocompatibility complex (MHC) class II inhibits monocyte function which is a crucial event in immunosuppression during sepsis. Possibly, statins act as anti-inflammatory agents.

Objective: The objective of our study was to test if statins affect the constitutive expression of HLA-DR on monocytes.

Methods: Monocytes of healthy subjects (with approval of the ethics committee) and the monocyte cell line Mono Mac 6 (MM6) were incubated with statins (simvastatin, mevastatin, pravastatin, lovastatin and fluvastatin) as well as mevalonate and their influence on MHC class I (HLA ABC) and class II (HLA-DR1) was observed by flow cytometry after 24h. To characterize possible mechanisms, the expression of CD74 (invariant chain) cathepsin S, interleukin 10 and its receptor, annexin V binding and caspase-3 activation were analyzed. mRNA expression was quantified by real-time rtPCR. ANOVA was used for statistical analysis. results.

Results: Monocyte HLA-DR was downregulated by simvastatin at 500 nM ($-13.51 \pm 4.02\%$, $p < 0.05$) with increasing potency up to 20 μ M ($-26.86 \pm 8.35\%$, $p < 0.01$). mevastatin, pravastatin, lovastatin and fluvastatin downregulated HLA-DR as well ($p < 0.05$). The expression of MHC class I instead was not affected by statins. Mevalonate, the product of HMG-CoA reductase can overcome the inhibition of the pathway. In the presence of mevalonate, simvastatin did not affect HLA-DR expression. Investigating possible mechanisms we analyzed induction of apoptosis by statins. While statins indeed induced apoptosis in MM6 cells, as confirmed by annexin V and caspase-3 staining, HLA-DR downregulation could be observed in the non-apoptotic annexin V and caspase-3 negative populations. Statins did not induce IL-10 production and did not change IL-10 receptor expression. mRNA expression of HLA-DR was not inhibited by statins. Therefore, a post transcriptional mechanism was suspected. No change was observed in cathepsin S expression, a processing enzyme. The expression of the HLA-DR chaperon CD74 was markedly decreased by simvastatin (MFI 122.71 ± 4.16 vs. control 163.70 ± 1.52 , $p < 0.01$).

Conclusion: Statins regulate the constitutive expression of HLA-DR in monocytes. This effect is dependent of HMG-CoA reductase inhibition. A posttranscriptional mechanism is suspected. Possibly, the downregulation of the chaperon CD74 inhibits the intracellular processing of HLA-DR. The reduction of monocyte HLA-DR expression emphasises the anti-inflammatory action of statins.

References: Hackam DG et al. The Lancet 2006;367:413-418. 2. Doecke WD et al. Clinical Chemistry 2006;51:2341-23

039

Infection 2007; 35 (Suppl. II): 30

Beta-defensin gene copy number variations in patients with severe sepsis

*Zhang XH (1,2), Book M (1), Chen QX (2), Lehmann LE (1), Weber (1), S, Schewe J-C (1), Hoeft A (1), Stuber F (1)

(1) Department of Anaesthesiology and Intensive Care Medicine, University of Bonn, Germany (2) School of Medicine, Zhejiang University, Hangzhou, P. R. China

Introduction: Severe sepsis is a serious complication following trauma, major surgery or infectious diseases. Little is known about genetic background which influences the susceptibility to severe sepsis. Human beta-defensins expressed by epithelial cells and peripheral white blood cells are cationic antimicrobial peptides, which also display modulating effects on other immune cells. Protein levels in plasma are detectable in patients with severe sepsis. Recent studies identified that the genes coding for human beta-defensin 2 (DEB4), human beta-defensin 3 (DEFB103) and human beta-defensin 4 (DEFB104) showed variation in copy number. The effects of beta-defensins in immune system predestine beta-defensins genes to be candidate genes for genetically determined interindividual differences in host defence against invading pathogens.

Objective: This study investigated the association of human beta-defensin 4 gene copy number variation with the susceptibility to severe sepsis.

Methods: 265 patients with severe sepsis and 231 healthy individuals were enrolled in the study. DNA was extracted from peripheral blood. Gene copy number was quantified by using real-time quantitative PCR.

Results: There are one- to four-copy number variants of human beta-defensin 4 gene among patients and healthy controls. The frequencies of copy number variants are not significantly different between sepsis group and control group ($p = 0.294$, chi-square test).

Conclusion: There is no association of copy number variation of human beta-defensin 4 gene with the susceptibility to severe sepsis in this study. A larger case-control study may be required to achieve adequate statistical power.

042

Infection 2007; 35 (Suppl. II): 30

Interleukin 13 but not Interleukin 4 exerts Akt-mediated survival of cardiomyocytes

*Rees W (1), Pöling J (1), Kubin T (2), Hein S. (3), Warnecke H (1), Braun T (2)

(1) Department of Cardiac Surgery, Schüchtermann-Klinik Bad Rothenfelde, Germany, (2) Max-Planck-Institut, Bad Nauheim, Germany, (3) Department of Cardiac Surgery, Kerckhoff-Klinik Bad Nauheim, Germany

Introduction: Cardiac remodeling is relevant in cardiovascular diseases and in heart failure (HF). It is widely accepted that pro-inflammatory cytokines play a major role in the development of HF, but few is known about the role of anti-inflammatory cytokines. Interleukin IL-13 is an anti-inflammatory cytokine secreted predominantly by activated Th2 cells. A recent report shows that IL-13 and IL-4 are markedly increased during various cardiac diseases. Moreover, an IL-13 receptor, IL-13R 1 as well as IL-4R are markedly expressed on human cardiac muscle cells. However, the meaning of these cytokines was only considered in respect of activation of immune cells in the development of heart diseases.

Methods: The purpose of the present study was to determine survival effects of anti-inflammatory cytokines on cultured cardiac myocytes, if modulated by IL-4 or IL-13. Parameters of description were cell size, survival, protein synthesis, ATP content and myofibril development. Leukemia inhibitory factor (LIF) was used as a control as this IL-6 class is considered to exert hypertrophic as well as survival effects.

Results: For the first time marked increases in protein synthesis, ATP-content and number of surviving cells induced only by IL-13 but not by IL 4 could be demonstrated in cultured myocardial cells. In contrast to leukemia inhibitory factor (LIF), which served as a control, the elevation in protein synthesis was rather due to an improved viability in IL-13 treated cells. The signal transduction pathways regulating this induction were investigated. Activation of Akt, Janus kinases (JAK), signal transducer / activator of transcription (STAT 6) and cAMP-independent pathways were shown to modulate the cell function.

Conclusions: These findings indicate that IL-13 can directly support myocyte function in addition to the inactivation of immune cells. Our results and the existence of IL-13 receptor in the heart suggest that IL-13 might have a protective role in cardiac myocytes.

046

Infection 2007; 35 (Suppl. II): 31

A new isolated perfused rat small bowel model for analysis of intestinal oedema formation

*Lautenschläger I (1), Kuchenbecker S-C (1), Dombrowsky H (1), Frerichs I (1), Scholz J (1), Steinfath M (1), Zabel P (2), Uhlig S (3), Weiler N (1)

(1) Department of Anaesthesiology and Intensive Care Medicine, University Medical Centre Schleswig-Holstein, Campus Kiel, Germany, (2) Department of Clinical Medicine, Research Centre Borstel, Leibniz – Centre for Medicine and Biosciences, Borstel, Germany, (3) Department of Pharmacology and Toxicology, RWTH Aachen University, Aachen, Germany

Introduction: Septic multi-organ failure is a life-threatening condition in critically ill patients. Intestinal malperfusion leading to bowel oedema with disturbed permeability and bacterial translocation may initiate the development of sepsis (1).

Objective: Up to now, there exists no experimental model allowing a systematic analysis of molecular mechanisms of intestinal oedema formation and possible targets for therapeutic options. Therefore, we developed a new ex situ model of an isolated perfused rat small bowel for studying intestinal oedema formation.

Methods: The small bowel was isolated in female Wistar rats (body weight: 220-250 g). Cannulas were inserted into the proximal and distal small intestine as well as into the aorta close to the superior mesenteric artery and portal vein. The totally isolated organ was then transferred into a specially designed temperature-controlled, moist chamber with a built-in weighing system. The bowel weight was continuously measured as no organ bath was used. Ileal compatible medium enriched with glutamine, glucose and lactose was used for luminal bowel perfusion at a rate of 0.15 ml/min. Vascular perfusion was performed with modified oxygenated Krebs-Henseleit buffer in a flow-controlled mode at a rate of 7.5 ml/min. Fluorescein isothiocyanate (FITC) albumin and norepinephrine were added to the vascular perfusion medium. Vascular and intraluminal pressures and flows were continuously recorded. Lymphatic vessels were opened and the lymphatic fluid collected allowing the quantification of lymphatic fluid production. Variations in FITC albumin content added to perfusion media allowed the characterization of fluid fluxes among different compartments. After the establishment of the model based on experiments on 180 rats, an initial experimental series testing the effect of platelet activating factor (PAF) on intestinal function was performed on 8 rats.

Results: In this new model, vascular, luminal and lymphatic flows, arterial, venous and intraluminal pressures and bowel weight can be monitored simultaneously. The initial experiments revealed that the bowel was vital and stable for at least 140 min. This was confirmed by histological techniques and by measuring the oxygen and glucose consumptions as well as the lactate-to-pyruvate ratio. The first experiments with administration of PAF showed vasoconstriction and a shift of fluid from vessels to the luminal compartment while generation of lymph was increased for a short time period of about 10 min. This indicates a rise in permeability evoked by PAF comparable to former findings in the isolated lung (2).

Conclusion: The presented model of an isolated perfused small bowel allows the separation of the vascular, luminal, extracellular and lymphatic compartments. We expect the model to be suitable for establishing the molecular mechanisms of bowel oedema formation.

References: 1. Fink Curr Opin Crit Care 2003; 9, 143-51 2. Göggel et al Nat Med 2004; 10, 155-60

049

Infection 2007; 35 (Suppl. II): 31

Patterns of cytokine inhibition by Drotrecogin alfa (activated) and ethacrynic acid in a murine model of sepsis

*Wagner T, Schürholz T, Claus RA, Simon TP, Reinhart K, Marx G
Clinic of Anesthesiology and Intensive Care, Friedrich-Schiller-University Jena, Germany

Introduction: Interaction between coagulation and inflammation offers potential of therapeutic intervention. One of those possible targets is the inhibition of NF- κ B. Both Drotrecogin alfa (activated) (DAA; recombinant activated Protein C) [1] and ethacrynic acid (EA) were shown to inhibit NF- κ B in vitro [2].

Objective: To investigate whether there is an anti-inflammatory effect of DAA or EA in a murine model of sepsis.

Methods: Sepsis was induced by cecal ligation and puncture (CLP) in male NMRI-mice (n=20, body weight 30 $\pm$ 3g). Animals were randomly assigned to vehicle infusion, or CLP sepsis with DAA infusion (24 μ g/kg/hr) or CLP sepsis with EA infusion (50 μ M). 48 hrs prior to CLP all mice were given a permanent i.v.-line (self-made) and an arterial transmitter (PA-C10, St. Paul, MN, USA) to measure heart rate (HR) and mean arterial pressure (MAP). Activity was recorded by scoring from 1 (healthy) to 5 (agony) [3]. CLP was adjusted to survive 24hrs. After 12hrs of sepsis activity score and vital data of all mice were recorded prior to killing. Plasma samples were collected and frozen following manufacturers guidelines until measurement. Determination of Interleukin (IL)-6, IL-10, monocyte chemoattractant protein-1 (MCP-1), Interferon-gamma (IFN- γ), tumor necrosis factor alpha (TNF- α) and IL12-p70 was performed using a cytometric bead array (CBA, BD Biosciences, Heidelberg, Germany). CBA is combined by six bead populations with distinct fluorescence intensities with specific capture antibodies. These bead populations are quantitatively measured using a flow cytometer. Differences between groups were evaluated by 1-way ANOVA.

Results: 12 hrs after CLP activity score was significantly higher in DAA-treated mice (score 1.8 $\pm$ 0.7 (mean $\pm$ SD); p=0.015) compared to control (score 3.1 $\pm$ 0.9). HR tended to be higher in DAA and EA mice (DAA 547 $\pm$ 175/min; EA 551 $\pm$ 110/min; Control 530 $\pm$ 143/min). MAP tended to be higher in DAA group and was similar after EA treatment when compared to control (DAA 111 $\pm$ 26mmHg; EA 98 $\pm$ 18mmHg; Control 97 $\pm$ 4mmHg). IL-6 (DAA 2305 $\pm$ 2211 pg/ml; EA 10612 $\pm$ 4670 pg/ml; Control 7953 $\pm$ 2408; p<0.05), IL-10 (DAA 74 $\pm$ 86 pg/ml; EA 662 $\pm$ 596 pg/ml; Control 1485 $\pm$ 1429; p<0.01) and MCP-1 (DAA 1621 $\pm$ 1084 pg/ml; EA 4960 $\pm$ 2552 pg/ml; Control 5207 $\pm$ 4193; p<0.05) were significantly lower after treatment with DAA but not with EA compared to control. All other investigated cytokines were not reduced by DAA.

Conclusion: In this model of murine sepsis DAA effectively inhibited different cytokines. EA treatment did not inhibit the investigated cytokines. DAA in experimental murine sepsis is capable to attenuate pro-inflammatory effects in a dose similar to those in humans.

References: [1] Joyce DE et al. Crit Care Med 2002; 30(5 Suppl): S288-293. [2] Han Y et al. Shock 2005; 23(1): 45-53. [3] van Griensven M et al. Shock 2002; 18(5): 445-449.

052

Infection 2007; 35 (Suppl. II): 32

Expression and secretion of proteolytically active ADAMTS13 is inhibited during proinflammatory stimulation

Bockmeyer CL, Sieber MW, Conradi F, Bauer M, Lösche W, *Claus RA

Department for Anaesthesiology and Intensive Care Medicine, Friedrich-Schiller University Jena, Germany

Background and Aims: ADAMTS13 is a plasma protease, primarily synthesised in hepatic stellate cells and the only enzyme held responsible for proteolytic degradation of Von Willebrand Factor regulating the molecular weight of the multimeric protein. ADAMTS13 activity is dramatically decreased in patients with inflammation and infection. High molecular weight VWF activates circulating platelets and may contribute to the development of thrombotic microangiopathy. Beyond a hypothesized inhibition of the protease by enormously elevated substrate levels under these conditions, an association between inflammation and ADAMTS13 expression/ secretion is not known.

Methods: We determined transcript level of ADAMTS13 by real time PCR experimentation. Results were normalized to unvaried housekeeping genes. Incubation media secreted activity was determined by a modified method of Kokame. As in vitro models, we used an immortalized hepatic stellate cell line (HSC) and human microvascular endothelial cells (HMEC-1), cultivated and stimulated with proinflammatory cytokines under standard conditions for 24 hrs. Additionally, as an ex vivo model, we stimulated human precision cut liver slices (HPLS) obtained from surgical waste material.

Results: On transcriptional level, we found ADAMTS13 in unstimulated HSC, HMEC-1 and HPLS. In all settings, incubation media secreted proteolytic activity was determined. After 24h, proinflammatory stimulation with interleukin 1beta, tumor necrosis factor and endotoxin in concentrations relevant to clinical conditions caused significant decrease of ADAMTS13-mRNA in dependence of the used normalisation transcript, which was mirrored in a decrease in proteolytic activity in incubation media (-64 %). Also in HPLS, we found a significant decrease in both, mRNA and mature protein.

Conclusions: To our knowledge for the first time we demonstrate in a human ex vivo setting a significant decrease in both, mRNA-synthesis and secretion of ADAMTS13 under proinflammatory conditions, which might contribute to the presence of high molecular weight VWF in plasma followed by activation of circulating platelets and the development of thrombotic microangiopathy.

056

Infection 2007; 35 (Suppl. II): 32

Normalisation strategy of gene expression studies after inflammatory insults

*Sieber MW (1,2), Guenther M (1), Witte OW (2), Frahm C (2), Claus RA (1)

Departments of (1) Anaesthesiology and Intensive Care Medicine, and (2) Neurology, University Hospital Friedrich-Schiller University, Jena, Germany.

Background: It is possible to detect and quantify mRNA levels of a specific transcript in tissue or cell cultures by use of the powerful technique combining reverse transcription of extracted RNA with real-time PCR. Accuracy of the results strongly depends on methodical factors such as sampling, storage conditions, RNA isolation or conduct of reverse transcription and real-time PCR. In order to neutralise these unwanted effects resulting from the above mentioned factors it is necessary to validate the results of internal standardisation, most commonly employing constantly expressed genes, reference genes or housekeeping genes. Normalisation using one, general housekeeping

gene, invariably expressed in a broad panel of tissue and under variable biological conditions, is controversially discussed.

Objective: To determine two housekeepers from the frequently used reference genes consisting of glyceraldehyd-3-phosphate dehydrogenase (GAPDH) and beta-actin (ACTB) as well as hydroxymethylbilane synthase (HMBS), hypoxanthine guanine phosphoribosyl transferase 1 (HPRT1) and glucuronidase beta (GUSB) in mouse brain subjected to middle cerebral artery occlusion (MCAO) and reperfusion as a model of systemic inflammation.

Results: This study demonstrated that the traditionally used housekeeping gene GAPDH is up-regulated up to 10-fold after inflammatory insult. Analysis of transcription profiles using GeNorm and Normfinder software showed that the expression level of GAPDH is strongly variable in comparison to other potential reference genes in the two conditions ischemia and ageing. The most suitable pair of genes for gene expression studies focussing on ageing effects, are GUSB and ACTB. The most eligible reference transcripts for the analysis of gene expression under ischemic conditions as well as in ageing, are ACTB and HMBS.

Conclusion: The comparison of the most commonly used reference genes demonstrated that there is no general housekeeper under variable biological conditions. ACTB and HMBS could be identified as reference genes representing appropriate internal standards for normalisation of mRNA quantification under inflammatory and ischemic conditions as well as in the study of age-dependent effects.

057

Infection 2007; 35 (Suppl. II): 32

Proinflammatory stimulation decreases expression and binding of ADAMTS13 to human microvascular endothelial cells

*Sossdorf M, Wanderer J, Bockmeyer CL, Lösche W, Claus RA

Department of Anaesthesiology and Intensive Care Medicine, University Jena, Germany.

Background and aims: The metalloproteinase ADAMTS13 plays a crucial role in haemostasis by cleaving the highly thrombogenic, ultra-large form of von-Willebrand-Factor (ULVWF), primarily immediately after endothelial secretion of ULVWF. Deficient activity of ADAMTS13 results in formation of vWF-rich thrombi characterised by a variety of clinical symptoms such as haemolysis, thrombocytopenia and micro-circulatory disturbances. Hepatic stellate cells are the principal site of ADAMTS13-biosynthesis, but mRNA was also detected in other cell types and tissues. In this study we examine the intracellular localisation of ADAMTS13 in the endothelial cell line HMEC-1 and its expression following stimulation with inflammatory cytokines.

Methods: Immortalised human microvascular endothelial cells (HMEC-1) and hepatic stellate cells (LX-2) as positive control were analysed by flow cytometry and confocal immunofluorescence microscopy. Confluent cultures plated on dishes or chamber slides were stimulated with pro-inflammatory mediators (i.e. TNF-alpha, LPS, IL1-beta) and labelled with endothelial cell-specific primary and secondary antibodies (ICAM-1, CD62P, vWF). The mean fluorescent intensity (MFI) and the percentage of positive cells of intracellular and surface-bound fluorescent molecules were analysed by FACScan flow cytometer. Fluorescent imaging of HMEC-1 by using ADAMTS13-monooclonal antibodies was employed to localise ADAMTS13.

Results: The adhesion molecule ICAM-1 was up-regulated after stimulation with LPS, IL1-beta, TNF-alpha in a time and concentration-dependent manner. Interestingly, no CD62P (P-Selectin) signal on the surface of the endothelial cells was detectable. ADAMTS13 is expressed in HMEC-1, however fluorescence intensity is very weak in comparison to the endothelial marker protein vWF. Stimulation reduces the intracellular level of ADAMTS13 measured by the amount of antibody bound.

Conclusions: The large surface area makes the endothelium an important source of ADAMTS13. Our results illustrate the synthesis and storage of ADAMTS13 in HMEC-1 and its down-regulation by inflammatory stimuli. These findings suggest that the plasma pool of ADAMTS13 is partially determined by endothelial cells and, furthermore, that reduced endothelial expression of the VWF-cleaving protease may contribute to progression of thrombotic microangiopathy, as observed in sepsis.

058

Infection 2007; 35 (Suppl. II): 33

The HemeOxygenase1 System in human sepsis and organ failure

*Sponholz C (1,2), Huse K (2), Platzer M (2), Claus RA (1), Bauer M (1)

(1) Department for Anaesthesiology and Intensive Care Medicine, Friedrich-Schiller University Jena, Germany, (2) Leibniz-Institut für Altersforschung – Fritz-Lipmann-Institut e.V. (FLI), Jena, Germany

Introduction: Hemoxygenases are enzymes converting heme into equal amounts of carbon monoxide (CO), biliverdin and free iron. Expression of the inducible isoform HMOX1 is up regulated in different tissues, including heart, lung, liver and monocytes by heme, cytokines, hypoxia, and other stimuli. The HMOX1 gene is encoded on chr22 and according the current annotation the gene structure is organized in 5 exons. However, expressed sequence tags (EST) support evidence for an additional exon upstream of exon1. Of major interest is a microsatellite in the form of a GT-dinucleotide repeat within the promoter region varying in lengths. The findings that (i) HMOX1 cleaves the toxic agent heme into anti-inflammatory agents (CO and biliverdin), (ii) the expression is up regulated by cytokines and (iii) the HMOX1 polymorphism may influence inflammatory response represents relevant factors to investigate the role of HMOX1 in sepsis and multiple organ failure (MOF).

Methods: DNA was separated from leukocytes of septic patients (n=122) from either Jena or Homburg university hospital and clinical and laboratory data, including APACHE and SOFA scores, mortality rate were recorded on an observation period of five days. PCR, with primers flanking the GT-repeat, was performed either on patient DNA or a human DNA pool serving as an unaffected control. Fragment analysis was performed for evaluation of lengths of the (GT)n-Polymorphism. RNA from different tissue panels were used to demonstrate a „new“ exon1 upstream the „former“ exon1 of the HMOX1 gene by PCR, cloning and sequencing experiments.

Results: The distribution of the (GT)n-length polymorphism displayed two peaks, at 23 and 30 repeat units in patients with sepsis and unaffected controls, therefore number of (GT)n-repeats were categorized into short (<25) and long (>25) repeat units. Patients carrying at least one short allele [(s/s, n=13) or (s/l, n=52)] exhibited lower SOFA scores on day 5 (6,93 vs. 9,89; p=0,005) and enhanced difference between day 5 and 1 (-2,33 vs. -0,45; p=0,011) compared to (l/l, n=57). The SOFA difference between day 5 and 1 was greater in (s/s) and (s/l) than in (l/l) (-2,55 vs. -0,45; p=0,062 and -2,28 vs. -0,45; p=0,026, respectively). In non-survivors, carrier of at least one short allele had significantly lower SOFA scores on day 5 (8,67 vs. 12,42, p=0,027) and an enhanced difference between day 5 and 1 (-1,83 vs. +0,62, p=0,05) compared to (l/l). (l/l) had higher SOFA scores than (s/l) at day 5 (12,42 vs. 8,93, p=0,05), but there was no difference neither between (s/s) and (l/l) nor between (s/s) and (s/l). The SOFA difference between day 5 and 1 within the non-survivors was significantly greater in (s/s) and (s/l) than in the (l/l) group (-3,00 vs. +0,62, p=0,039 and -1,50 vs. +0,62, p=0,014 respectively). cDNA sequencing analysis in different tissues supported evidence for a „new“ exon1 upstream the „former“ exon1 and alternative splicing variants of HMOX1 gene, with or without intron retention of the (GT)n-polymorphism, could be found.

Conclusion: The (GT)n-length polymorphism of HMOX1 may influence sepsis course and outcome depending on the number of the

(GT)n-repeat length, however due to the limited number of patients no information is available on a varied predisposition for sepsis. The structure of the HMOX1 gene needs to be reevaluated by inserting a novel exon1 upstream of the former exon1. Splicing variants include the (GT)n-length polymorphism by intron retention, probably depending in the number of repeat units.

060

Infection 2007; 35 (Suppl. II): 33

Expression profiles of hepatocellular transporter proteins after pro-inflammatory stimulation – a molecular mechanism for the development of sepsis-induced liver dysfunction

*Recknagel P (1), Claus RA (1), Müller D (2), Bauer M (1)

(1) Department of Anaesthesiology and Intensive Care Medicine, Friedrich-Schiller University Jena, Germany, (2) Institute of Pharmacology and Toxicology, Friedrich-Schiller University Jena, Germany

Introduction: Hepatic failure still represents the prominent prognosis limiting factor in patients with severe sepsis. Hypoxia caused by septic shock and perfusion failure, as well as inflammatory mediators, released by several kinds of immuno-competent cells, are both initial factors for sepsis-induced liver dysfunction. In addition to the ensuing progressive loss of a critical number of metabolically active cells through necrosis and apoptosis, the differential regulation of hepatic transporter proteins (e.g. OATP1, BSEP, MRP2, etc.) is also thought to cause a disturbance in hepato-cellular clearance function and thus to result in the pathophysiological pattern of liver dysfunction. This is of particular interest regarding the hepato-biliar clearance of a variety of pharmacological substances administered under these conditions.

Methods: To assess the expression-modulating effects of either hypoxia or inflammatory mediators in cultured human precision liver slices (HPLS) as an ex-vivo model mimicking the hepato-cellular environment under septic conditions. To achieve prototypical cellular events during sepsis, human precision liver slices prepared from surgical waste material were incubated for 0, 6, and 24 hours under standard conditions with i) mixed inflammatory mediators or ii) 2,6-dimethyl-2,5-heptadien-4-one (depletion of the cellular glutathione content) to simulate septic, pro-inflammatory conditions and sepsis-induced oxidative stress, respectively. Transcription levels of selected transporters were determined by Real-time PCR run on an iCycler™ (Biorad). Results were normalised to the unvaried housekeeping gene hypoxanthine ribosyl transferase (HPRT).

Results: In both settings, i.e. inflammatory stimulation and oxidative stress we demonstrated differential changes in gene expression of individual basolateral and canalicular transporter proteins as well as group-associated differences in the expression of basolateral as opposed to canalicular transporters. Although some transporters displayed robust or even increased expression (e.g. OATP1, NTCP), the majority of the investigated transporter proteins showed steady state transcript concentrations that were substantially decreased: Particularly sensitive proteins were the basolateral transporter proteins MRP3 and OATP2 as well as the canalicular transporters MRP2, MDR3 and the bile salt export pump (BSEP), with the canalicular transporters overall more sensitive to either intervention than the basolateral ones.

Conclusion: With this study we demonstrated directed steady state transcript variations to noxious stimuli in cultured HPLS for various transporters. These expression patterns might reflect a mechanism for a disturbance in hepato-cellular excretory function and could, therefore, lead to non-uptake from sinusoidal blood flow or accumulation of potentially toxic products within the hepatocytes, hence promoting the development of sepsis-induced cholestasis or liver dysfunction.

064

Infection 2007; 35 (Suppl. II): 34

Argatroban Improves Intestinal Microcirculation in Experimental Sepsis

*Lehmann C (2), Fuchs C (1), Ladwig E (1), Behrend K (1), Gründling M (1), Pavlovic D (1)

(1) Department of Anaesthesiology and Intensive Care Medicine, Ernst Moritz Arndt University Greifswald, Germany (2) Department of Anesthesia, Dalhousie University, Halifax, Nova Scotia, Canada

Introduction: Successful treatment of severe sepsis and septic shock remains to be a major challenge in critical care medicine. The anticoagulant activated protein C remarkably improved the outcome of septic patients (1). The aim of our study was to investigate the effects of another anticoagulant substance – the direct thrombin inhibitor argatroban (ARG) – on the intestinal microcirculation during experimental sepsis in rats (2) and its direct vascular effects in vitro.

Methods: A total of 40 male Lewis rats were randomly assigned to 4 groups (n=10): sham surgery (SHAM), experimental sepsis (colon ascendens stent peritonitis – CASP), CASP+ARG, and SHAM+ARG. Following 15 hours observation time the animals of the CASP+ARG and the SHAM+ARG groups received 2 mg/kg argatroban intravenously. 2 hours later all animals underwent intravital microscopy of the intestinal wall. In separate experiments, in preparations taken from healthy animals, direct effect of argatroban on isometric tension of rat aortal rings in vitro was also examined.

Results: Argatroban administration in CASP rats as compared to untreated animals increased mucosal (+48%) and muscular (longitudinal: +39%; circular: +42%) functional capillary density and attenuated the rolling behaviour (V1 venules: -55%; V3 venules: -45%) and the number of firmly adhering leukocytes (V1 venules: -47%; V3 venules: -24%) in the intestinal submucosa. In vitro argatroban had no effects on precontracted rat aortal rings (with 20 or 40 mM KCl, or 5x10⁻⁸ M phenylephrine). In contrast, the incubation (30 min) in argatroban (10⁻⁶M) diminished sensitivity of the aortal rings to phenylephrine (EC50: 7.48 (0.06) vs. 7.19 (0.08); means (SEM), controls v.s. incubation, n=8, p=0.008).

Conclusions: Argatroban administration during experimental sepsis preserved intestinal microvascular perfusion and exerted anti-inflammatory effects by reducing leukocyte adherence to the endothelium in submucosal venules. Some of these effects could be mediated by its weak vasodilating property. It should be examined whether argatroban could have anti-inflammatory or vasoactive effects in human microcirculation.

References: (1) Bernard GR et al.: Efficacy and safety of recombinant human activated protein C for severe sepsis. *N Engl J Med.* 2001;344:699-709. (2) Miyahara S et al.: Argatroban attenuates leukocyte- and platelet-endothelial cell interactions after transient retinal ischemia. *Stroke.* 2003;34:2043-9.

065

Infection 2007; 35 (Suppl. II): 34

Effects of antiplatelet drugs in sepsis: data from a clinical and animal study

*Winning J (1), Lösche W (1), Baranyai J (1), Hamacher J (2), Eisenhut Y (2), Kohl M (3), Deigner HP (3), Claus RA (1), Bauer M (1)

(1) Anaesthesiology and Intensive Care Medicine, University Hospital Jena, Germany, (2) Division of Pulmonology, University Hospital Homburg, Germany, (3) SIRS-Lab GmbH, Jena, Germany

Introduction: Sepsis is known to be associated with platelet activation that may contribute to organ failure and modify the systemic inflammatory response.

Methods: 224 patients with community acquired pneumonia (CAP) were studied for an association between pre-hospital treatment acetylsalicylic acid (n= 36) or clopidogrel (n=8) and clinical outcome. Patients on statins were excluded. BALB/c mice were pre-treated with clopidogrel for 4 days prior to intraperitoneal endotoxin injection (LPS from *E. coli* 0111:B4).

Results: Although CAP patients with anti-platelet drugs (n=44, 69±7 years) were not different in main laboratory and clinical parameters when admitted to hospital, they were less likely to be admitted to intensive care unit (ICU) than control patients (n=180; 58±13 years; ICU admission: 9.1% vs 26.1%; p<0.02). Within the first 48 h of endotoxin shock we did not see any significant differences between clopidogrel and control animals (n = 26 each) in haemodynamic alterations as well as lethality (14 and 13 survivors, respectively). Clopidogrel abolished the LPS-induced drop in platelet count and reduced the number of thrombi in the lung (6.1±2.3 vs 11.5±4.4 thrombi/screen). There were no differences in pO₂ or pCO₂, but higher pH values and bicarbonate levels were measured in the clopidogrel animals (7.01±0.01 vs. 6.93±0.04, and 10.2±0.14 vs 7.3±0.14 mmoles/l). Gene expression analysis in blood leukocytes by DNA microarray indicated suppressed LPS-induced up-regulation of inflammatory genes, including arachidonate-5-lipoxygenase activating protein and leukotriene B₄ receptor 1, in clopidogrel-treated animals. Between 2 and 7 days after LPS treatment, 59 % of control and 29 % of clopidogrel-treated animals died.

Conclusions: Anti-platelet drugs may have a beneficial effect in systemic inflammation and sepsis and could be a novel therapy option, at least in patients on low bleeding risk. Their effect seems to be due to an antithrombotic and anti-inflammatory action and a prevention of organ failure.

066

Infection 2007; 35 (Suppl. II): 34

Anti-inflammatory effects of glucocorticoids require the dimerization of glucocorticoid receptor in myeloid cells

*Kleyman A (1), Neumann A (1), Hübner S (1), Illing A (1), Winning J (2), Schmid W (3), Richter W (4), Reichardt H (5), Schütz G (3), Tuckermann J (1)

(1) Division of Tissue specific hormone action, Leibniz Institute of Age Research – Fritz Lipmann Institute, Jena, Germany; (2) Anaesthesiology and Intensive Care Medicine, University Hospital Jena, Germany (3) Division for Molecular Biology of the Cell Deutsches Krebsforschungszentrum Heidelberg, Germany; (4) Klinikum der FSU, Elektronenmikroskopisches Zentrum, Jena, (5) Division of Cellular and Molecular Immunology University of Göttingen, Germany.

Introduction: Besides severe side effects Glucocorticoids (GCs) are widely used for treatment of inflammatory diseases. However, the precise mechanism, how GCs repress inflammation, is not completely understood.

Objective: GCs bind to the Glucocorticoid Receptor (GR) – an ubiquitously expressed ligand-induced transcription factor. GR regulates expression of target genes through different mechanisms: 1) GR binds directly to a specific DNA sequence – Glucocorticoid Response Element (GRE) – and regulates the expression of nearby located genes. This mechanism depends on dimerization of GR. 2) GR interacts as a monomer with other transcription factors such as NF-κB, AP1, STAT5, and modulates their activity. It is believed that the second mechanism of GR action mediates the anti inflammatory activity of GCs.

Methods and results: By the use of conditional GR knockout mice (GR^{lysMcre} mice) we could show that the presence of the GR in myeloid cells is required for anti-inflammatory effects of GCs in septic shock model. Furthermore in knock-in mice that express a dimerization defective GR (GR^{dim}), we could demonstrate that dimerization of glucocorticoid receptor is necessary for the repression of inflammation by GCs. We found that GCs could not completely repress LPS

induced activation of GRdim bone marrow derived macrophages (BMDM), in contrast to wild type BMDM. Analysis of expression profiles of wild type and GRdim BMDM reveal new GR target genes responsible for anti inflammatory action of GCs. We found that GCs up regulate the expression of several Krüppel-like transcriptional factors (KLFs) in dimerization dependent manner. The absence of KLFs regulation in GRdim BMDM probably leads to misregulation of important pro inflammatory mediators. Besides that we showed that upon LPS treatment GRdim mice as well as GRlysMcre mice have severe defects in integrity of endothelial barrier and cardio-vascular system that could explain the higher susceptibility of mutant mice to the LPS induced septic shock.

Conclusion: These data challenge the initial concept that inhibition of pro-inflammatory transcription factors activity is sufficient for anti-inflammatory action of GC.

070

Infection 2007; 35 (Suppl. II): 35

Recombinant human activated protein increases the number of endothelial progenitor cells in vitro

*Brückmann M, Glasl C, Schmidt J, Lang S, Hoffmann U, Borggreffe M

Department of Medicine I, Faculty of Medicine Mannheim, University of Heidelberg, Mannheim, Germany

Introduction and objective: Endothelial progenitor cells (EPCs) play an important role within the process of new blood vessel formation, regeneration of ischemic tissues and maintenance of vascular endothelium integrity. Non-anticoagulant biological activities, such as a reduction of endothelial cell permeability, enhanced endothelial cell survival as well as improvements of microcirculatory disorders have been proposed for recombinant human activated protein C (rhAPC; drotrecogin alfa (activated)). Aim of the present study was to examine the effect of rhAPC on EPC differentiation in vitro.

Methods: Total mononuclear cells (MNCs) were isolated from buffy coats of healthy donors by density gradient centrifugation. MNCs were plated in endothelial basal medium on fibronectin coated plates in the presence of supplements (FCS 20%, EGF 10µg/ml, bovine brain extract 3µg/ml, hydrocortisone 1µg/ml) to promote differentiation and growth of EPCs. Cells were stimulated daily with rhAPC (50ng/ml or 5µg/ml) or alternatively with VEGF (50ng/ml) as a positive control. After 4 days of culture adherent cells were characterized as EPCs by identification of CD34, CD31, VEGF-receptor-2 and vWF-expression by FACS analysis. Two independent investigators evaluated the number of EPCs per well by counting three randomly selected fields. Statistical analysis from at ten independent experiments was performed by student's t-test.

Results: RhAPC (50ng/ml and 5µg/ml) increased EPC numbers after 4 days in culture versus unstimulated control cells (+43% for rhAPC 50ng/ml, $p < 0.05$; +22% for rhAPC 5µg/ml, $p = 0.05$). This effect was comparable to VEGF in its capacity to increase EPC numbers (+43%, $p < 0.05$).

Conclusion: The ability of rhAPC to upregulate the number of EPCs in vitro may contribute to vessel regeneration and vascular integrity and may provide a new therapeutic concept to improve EPC numbers in patients with severe sepsis.

078

Infection 2007; 35 (Suppl. II): 35

Glutamine improves microcirculation in experimental endotoxemia

*Lehmann C, Wuttke U, Saeger D, Pavlovic D, Wendt M, Gründling M

Department of Anaesthesiology and Intensive Care Medicine, Ernst Moritz Arndt University, Greifswald, Germany

Introduction: Successful treatment of severe sepsis and septic shock remains to be a major challenge in intensive care medicine. Recently, experimental glutamine (GLN) administration has shown to improve remarkably the outcome of septic rats, by reducing cytokine release, organ damage and mortality. The intestine serves as a pathologically significant circulatory region in cases of clinical sepsis.

Objective: The aim of the present study was to evaluate the effects of GLN and its galenic formulation l-alanine-l-glutamine dipeptide (DIP) on the intestinal microcirculation during experimental endotoxemia using intravital fluorescence microscopy (IVM). It was also attempted to determine whether GLN administration affected the cytokine release by evaluating the blood levels of the cytokines TNF- α and IL-1 β , IL-6 and IL-10.

Methods: Rats were randomly divided in six groups (each n=10): CON (control), LPS (5 mg/kg lipopolysaccharide i.v.), GLN+LPS (LPS animals pre-treated with 0.75g/kg GLN i.v.), DIP+LPS (LPS animals pre-treated with DIP i.v., 0.75g/kg GLN content), LPS+GLN (LPS animals post-treated with 0.75g/kg GLN i.v.) and LPS+DIP (LPS animals post-treated with DIP i.v.; 0.75g/kg GLN content). Two hours following endotoxin challenge microcirculation of terminal ileum was studied using IVM. Blood samples were drawn at the beginning and the end of the experiment in order to determine cytokine release (ELISA).

Results: Two hours following endotoxin challenge, significant reductions in the functional capillary density (FCD) were observed in all layers of the intestinal wall in untreated LPS animals ($p < .001$). GLN and DIP pre-treatment, respectively, prevented the LPS induced FCD decrease in the same extent ($p < .001$). Intestinal FCD also improved following post-endotoxin administration of GLN and DIP, respectively. The number of activated, adherent leukocytes in the submucosal collecting venules was attenuated following GLN/DIP pre-treatment, but elevated following post-treatment. In the smaller, postcapillary venules also GLN post-treatment reduced significant leukocyte adherence ($p < .01$). At the beginning of the experiment the IL-1 α , IL-6, IL-10 and TNF- α ; concentration did not differ between the groups while they all significantly increased two hours following the LPS challenge ($p < .001$). GLN/DIP administration did not reduce the cytokine release. IL-6 release was even increased following DIP pre- and post-treatment, respectively ($p < .05$).

Conclusion: Treatment of endotoxemic animals with glutamine and l-alanyl-l-glutamine dipeptide, respectively, improved the intestinal microcirculation by increasing functional capillary density in all layers of the intestinal wall (pre- and post-treatment). Leukocyte activation was more effectively suppressed by GLN/DIP pre-treatment. Because of the crucial role of the microcirculation in the pathogenesis of sepsis and multiple organ dysfunction syndrome, these effects may be of clinical importance.

079

Infection 2007; 35 (Suppl. II): 36

Genetic predisposition for adverse outcomes in trauma patients: First stage of a genome wide approach

*Schewe JC (1), König IR (2), Book M (1), Weber SU (1), Lehmann LE (1), Lüpert M (1), Kabir K (1), Wirtz DC (1), Henrich D (3), Marzi I (3), Ertel W (4), Hostmann A (4), Bühren V (5), Woltmann A (5), Muhr G (6), Friese J (6), Pohlemann T (7), Bouillon B (8), Maegele M (8), Biberthaler P (9), Bogner V (9), Faist E (10), Trentzsch H (10), Van Griensven M (11), Schade U (12), Stüber F (1) (1) University Bonn Medical Centre, (2) University Hospital Schleswig-Holstein, Campus Lübeck, (3) University Hospital Frankfurt, (4) University Medical School, Charité, Campus Benjamin Franklin, (5) BG Trauma Centre Murnau, (6) BG Trauma Centre Bergmannsheil Bochum, (7) University Clinic of the Saarland, (8) University of Witten/Herdecke, Hospital Cologne Merheim, (9) University Hospital of Surgery – City, Ludwig Maximilians University, (10) Klinikum Grosshadern, Ludwig Maximilians University of Munich, (11) Ludwig Boltzmann Institut of Traumatology, Vienna, (12) University Hospital Essen

Introduction: Injury remains a major health problem worldwide. Surviving the initial event the patient is at risk for subsequent life-threatening complications, which include sepsis and multiple organ dysfunction syndrome (MODS). Evolving evidence suggests that various genetic polymorphisms are associated with increased susceptibility to infection and poor outcomes. Single nucleotide polymorphisms (SNPs) refer to common genetic variants that may be associated with disease susceptibility.

Objective: The aim of the project is to identify genetic susceptibility factors contributing to adverse outcomes such as severe sepsis and related organ dysfunction in trauma patients. The combination of a genome wide scan and multiple candidate gene analysis will increase informativity and help define a possible new diagnostic tool for risk

assessment. This study represents the first genome wide approach to test the hypothesis of genetic predisposition for adverse outcomes in a large cohort of trauma patients.

Methods: A prospective, consecutive entry study was developed to assure inclusion of severely injured trauma patient within a network of 13 German major trauma centres. The study has been positively been approved by the Ethics Committee of Bonn University Medical Centre and by local Ethics Committee of collaborating centres. Trauma was defined based on the Injury Severity Score (ISS). Inclusion criteria were ISS>16, age =>18 and ICU treatment > 24 hours. Genome wide scan analysis was performed with the Affymetrix 500K Array Set covering a total of 500.568 SNP's. Data are anonymized and evaluated according to the guidelines of the DAE/AKW. Patient data documentation is conducted involving the German trauma registry. The study is based on a multi-stage genetic epidemiologic design. Genetic association analysis was performed using SNP wise two-sided Cochran-Armitage trend-test.

Results: Up to now 600 patients were included, 250 of the patients with sepsis and related organ dysfunction have been genotyped in the first step. Median age was 39 in male and 40 years in female patients (median ISS=27). Incidence of sepsis showed no correlation with initial ISS ($p>0.05$). Overall incidence of sepsis and related organ dysfunction in the recruited 600 patients was 40%. There was a clear correlation of organ dysfunction respectively MODS and sepsis ($p<0.0001$). Based on a critical p-value of 0.00001, 21 SNP's in 18 genomic regions were identified with a significantly association to severe sepsis related organ dysfunction in trauma patients when compared to a large population control group.

Conclusion: This is the first step of a study to test hypothesis of genetic predisposition for adverse outcomes in a large cohort of trauma patients. Preliminary data show that adverse outcomes of trauma patients can be related to genomic polymorphisms of candidate genes. Completion of recruitment, genome-wide scanning, replication of the results and association study for relevant SNPs is in progress.

Satellite Symposia

019

Infection 2007; 35 (Suppl. II): 37

Septifast PCR (SF) for microbiological documentation of blood culture (BC)-negative infections in febrile neutropenic patients (Pts)

Lamoth F (1), Jatton-Ogay K (2), Prodhom G (2), Senn L (1), Bille J (2), Marchetti O (1), *Calandra T (1)

(1) Infectious Diseases Service, CHUV and University of Lausanne, Switzerland, (2) Institute of Microbiology, CHUV and University of Lausanne, Switzerland

Introduction: BC are the gold standard of microbiological diagnosis in febrile neutropenic Pts. No causative agent is identified in 2/3 of febrile episodes. SF (Roche) is a new PCR test, which may detect bacterial and fungal DNA in blood of Pts with BC-negative infections.

Objective: To assess the utility of SF for the microbiological documentation in febrile neutropenic Pts.

Methods: Blood samples for BC and SF were prospectively drawn in 49 adult neutropenic cancer Pts at D0 (onset of fever) and D3 (if persistent fever). Febrile episodes were classified as microbiologically (MDI) or clinically documented infection (CDI) and fever of unknown origin (FUO).

Results: 133 samples were analyzed in 85 febrile episodes (27 MDI, 25 CDI, 33 FUO). BC and SF were positive in 24 (28%) and 42 (49%) episodes, respectively. The pathogens were: i) BC: G+ 48%, G- 52%; ii) SF: G+ 51%, G- 40%, fungi 9%. Using BC as reference, sensitivity of SF was 59% (54% in G+, 60% in G-), specificity 48%, PPV 30%, NPV 76%. 50% of pathogens not detected by SF were not included in the test panel. Performance of BC and SF in different settings:

	BC-positive	SF-positive	P
No antibiotics at time of sampling (n = 47)	13 (28%)	16 (34%)	NS
Antibiotics at time of sampling (n = 86)	13 (15%)	46 (53%)	< 0.05
Onset of fever, D0 (n = 31)	7 (23%)	13 (42%)	0.2
Persistent fever, D3 (n = 55)	6 (11%)	33 (60%)	< 0.05

SF revealed previously undetected pathogens in 25/33 (76%) positive samples on D3. In 11/13 (85%) SF-pos CDI the pathogen was consistent with the site of infection (enterocolitis 7, mucositis 3, sinusitis 1). Fungi were detected in 4 BC-neg episodes of possible invasive mycoses (Candida 3, Aspergillus 1).

Conclusion: Septifast PCR is a promising additional tool for the microbiological documentation of blood culture-negative febrile neutropenic episodes. Septifast may be particularly useful in patients receiving antibiotics at time of blood sampling or with persistent fever.

026

Infection 2007; 35 (Suppl. II): 37

Fungal infections in critically ill ICU patients

*Maschmeyer G

Hematology and Oncology, Department of Internal Medicine, Hospital 'Ernst von Bergmann', Potsdam, Germany

Critically ill patients undergoing intensive care treatment bear a significant risk of acquiring an invasive fungal infection. Their immune defense may be severely compromised by corticosteroids or other immunosuppressive agents, and their skin and mucosal integrity is often markedly disrupted by open wounds (particularly in burn patients) and invasive devices. Catheter-associated invasive *Candida* infections are among the most frequently observed infectious complications. Early diagnosis, prompt intervention with systemic antifungal therapy and removal of intravascular devices have been shown to be of significant benefit in terms of morbidity and mortality. The local epidemiology, especially the relevance of fluconazole-resistant *Candida* spp., should be carefully taken into consideration when the antifungal agent for first-line intervention is selected. Antifungal treatment should be continued for at least 14 days after the first negative blood culture. In high-risk surgical ICU patients, antifungal prophylaxis may be beneficial. Organ transplant recipients, severely immunocompromised and neutropenic patients, such as those undergoing allogeneic hematopoietic stem cell transplantation or intensive antileukemic chemotherapy, as well as burn patients and patients with severe chronic pulmonary diseases are at risk for invasive pulmonary aspergillosis. Early CT scans may show typical lesions, i.e., nodules surrounded by glass-ground opacities ("halo" sign) or an air-crescent sign indicating a more advanced disease. Bronchoscopy plus bronchoalveolar lavage in order to obtain samples for culture, histopathology, *Aspergillus* antigen and eventually PCR may help to prove the filamentous fungal origin. Paranasal sinuses may be affected as well as the central nervous system, so that these sites should be examined thoroughly and eventually taken into account for obtaining additional samples for diagnostic procedures. Early intervention with a highly effective systemic antifungal is mandatory. When choosing the appropriate antifungal agent, co-medications interacting with cytochrome P450 isoenzymes have to be considered. Pharmacokinetic properties, particularly with respect to tissue penetration, and safety profiles of available antifungal should be carefully taken into consideration when treatment choices are made. It is important to know that in patients with invasive pulmonary aspergillosis, response in terms improvement of CT findings may not be expected until 10 to 14 days of full-dose antifungal therapy have been given.

040

Infection 2007; 35 (Suppl. II): 37

Septic liver failure – pathophysiology and therapy concepts

Bauer M

Department of Anesthesiology and Critical Care Therapy, University Jena, Germany

Due to its central role in metabolism and immune function, the liver is both, target and promoter of systemic inflammation and multiple organ failure. Cytokine release, perfusion disorders and oxygen radicals may lead to direct and indirect hepatocellular injury. Impaired excretion seems to reflect the predominant disorder of parenchymal cells while function of non-parenchymal cells remains unclear in the clinical setting. Traditional laboratory parameters underestimate liver dysfunction and are not helpful in this patient collective – neither to direct therapy nor to estimate prognosis. Dynamic tests, such as assessment of plasma disappearance rate of indocyanine green are bet-

ter suited in this context. Early initiated causal therapy – i.e. antibiotic therapy and source control in case of sepsis – accompanied by sufficient oxygen supply to the hepatocytes are basic therapeutic strategies in this setting. Proved specific therapies for liver dysfunction with influence on mortality do not exist. Liver transplantation is typically not an option for these patients. Extracorporeal liver support is increasingly used in clinical practice although it is not evaluated in case of secondary liver failure with respect to evidence based criteria.

048

Infection 2007; 35 (Suppl. II): 38

Rapid PCR-based pathogen detection in patients with clinical sepsis: a microbiologist's perspective

Hunfeld K-P

Institute of Medical Microbiology & Infection Control, University Frankfurt, Germany

Currently, blood culture (BC) is regarded the gold standard for diagnosing blood stream infection (BSI) but, commonly, 24 to 48 hours are needed before a precise pathogen identification and susceptibility test results become available. More rapid pathogen identification and treatment, however, are crucial for survival in sepsis patients. Recently, two European multi-center studies focused on the laboratory evaluation and technical feasibility of a new real-time PCR-based system (LightCycler SeptiFast test, ROCHE) developed for faster pathogen detection in patients with BSI. The test specifically targets the ITS-region of bacteria and fungi and can be completed in less than 6 hours from K-EDTA blood. Subsequent pathogen detection in PCR positive samples is performed by use of specific hybridization probes. Here, data on the robustness and diagnostic reliability of the test as obtained for 1186 blood samples from 396 ICU patients with suspected sepsis and 137 controls are presented. In addition, supplementary microbiological test results generated throughout these studies are discussed. PCR starting from 1.5 ml and from 3 ml blood were equally sensitive and specific. However, performance of PCR starting from 1.5 ml instead of 3 ml blood turned out more robust. Obviously, the average time-to-report is much longer for conventional BC compared to PCR while PCR detection rate for various microorganisms is significantly higher compared to BC. In a subgroup of 53 healthy controls and 65 patients with suspected sepsis that was analyzed at our institution from 06/2005 to 02/2006 by PCR and conventional cultures under routine laboratory conditions, the concordance of positive BCs with positive PCR results was 71.4%. Concordance of negative PCR results with negative BC results was 75.3% mainly due to the higher rate of PCR positives. Analytical specificity of PCR in patients without evidence of sepsis was 98.9%. If constructed gold standards (clinical judgment, additional microbiological data) were used to adjust for the potentially lower sensitivity of BC, the correlation with PCR was 87.5% and 90.4% for positive and negative results, respectively, in the 118 patients investigated at our centre. Compared with conventional BC, PCR, obviously, detects more microorganisms in patients with BSI while the results in the control group suggest high specificity. Reflecting on a meta-analysis of currently available study data the potentials and limitations of this new technology will be discussed and commented on from the perspective of a clinical microbiologist in the context of additional clinical and microbiological information.

051

Infection 2007; 35 (Suppl. II): 38

Sepsis therapy – where does the money go?

*Martin J (1), Franck M (2), Schleppers A (3)

(1) Department of Anaesthesiology, operative Intensive Care and Pain Management, Klinik am Eichert, Göppingen, Germany, (2) Department of Anaesthesiology and operative Intensive Care, Charité – universitätsmedizin Berlin, Germany, (3) Department of Anaesthesiology and operative Intensive Care, University Mannheim, Germany

Intensive care medicine is next to the operating room the most expensive division of a hospital. Ca. 15-20% of the costs for inpatient therapy are spent in intensive care medicine, while only 5 % of the patients need intensive care treatment [1]. The prevalence study of the German Sepsis Society showed, that ca. 140.000 patients per year fall ill with sepsis; of it 75.000 patients with severe sepsis [2]. The direct costs for treatment amount to ca. 1.8 billion Euro per year. With 60.000 cases of death per year sepsis is the third leading cause of death after coronary artery disease and acute myocardial infarction. In the last years enormous progress in the therapy of sepsis was achieved. Although the 28-day-mortality in the prevalence study [2] was still at 54%, e.g. the work of Kortgen et al. [3] showed, that with stringent compliance to the sepsis guidelines and implementation of the 6- and 24-hours therapy bundles the mortality may be lowered significantly below 30%. The therapy costs per day for intensive care medicine amount to ca. 1.000 € per patient day [4]. More than 40% of the costs are for personnel. Therefore the goal should be to lower the time spent in the intensive care unit, and also in the hospital, through optimized therapy according to the guidelines, while at the same time improving mortality and morbidity. Garland et al. [5] showed, that second to length of stay as the greatest independent cost factor in an intensive care unit, the individual treatment of the physician is the next greatest independent cost factor for intensive care therapy. In conclusion these studies indicate that for therapy of sepsis it is absolutely imperative, to implement the sepsis bundles according to the guidelines. With it not only mortality and morbidity are improved, but with shortened length of stay costs decrease. In the work of Nguyen [6] was explained, that with an appropriate training program for all employees of the intensive care unit an implementation of the sepsis guidelines succeeds. As Shorr et al. [7] showed, this also leads to a reduction of direct costs. In summary it may be said that with costs of 1.000 € per day for intensive care therapy, the treatment of severe sepsis belongs to the most expensive syndromes. Still with consistent implementation of the current existing guidelines in the daily treatment routine the quality may increase and therapy cost may decrease.

References: 1.Barkow D. Ärztl. Fortbild. Qualitätssich. 2000; 94: 828-833 2.Engel C, Brunkhorst FM et al. Intensive Care Med 2007; 33: 606–618 3.Kortgen A, Niederprüm P, Bauer M: Crit Care Med 2006; 34: 943-949 4.Prien Th, Groll O, Geldner G, Martin J, Weiler Th, Dahmen KG, Sorgatz H, Bach A. Anästhesiologie & Intensivmedizin 2002; 43: 244-254 5.Garland A et al. Am J Respir Crit Care Med 2006; 174: 1206–1210 6.Nguyen HB et al. Crit Care Med 2007; 35: 1105-1112 7.Shorr AF et al. Crit Care Med 2007; 35: 1257-1262

063

Infection 2007; 35 (Suppl. II): 38

Can we learn from children?

Sasse M

Kinderklinik, Abt. Kinderkardiologie und pädiatrische Intensivmedizin, Medizinische Hochschule Hannover, Germany

Children hospitals in Germany are often in a financial deficiency. They are mostly not able to compete with the great departments of adult surgery or internal medicine. On one hand the amount of patients is low, otherwise the reference of the costs in the DRG-System

is not adequate. Children hospitals are commonly interdisciplinary medical facilities. The units are small with an extraordinary expense of health personnel and technical equipment. The poor financial situation is legitimated as an investment in the future of the population. About the small dimension of the single pediatric departments, intensive networking with tertiary children hospitals and other clinics in the surrounding is essential and has a long tradition. The mentioned conditions and deficits had maintained to some changes in the working procedures. Because of the inner children-hospital availability of widespread competence, the communication channel is short and fast. Usually there is no significant time delay between recognition of the seriously ill child and the integration of different pediatric departments in diagnosis and therapy. The strict centralisation in the pediatric networks benefits a very early integration of the tertiary centers in the treatment of severe illness. Methods for adequate rescue therapies are mostly available only in these centers. This is a reason for fast transfer of the seriously ill child and therefore a better efficiency of the central hospitals. The short delays to the optimal treatment reduce costs and shorten the hospital stay. The retransfer after the successful intervention to the hospital of admission release new capacity of the tertiary center. Basic requirement for this procedure is a barrier free communication, collective continuing training and engaging therapy bundles. The competence of the tertiary center and the possibility of a transport-system have to be available 24 hour per day. In our pediatric intensive care network this system has carried to a significant improvement for example in the treatment of the Waterhouse-Friderichsen Syndrome. All patients were treated with the common therapy algorithm. The gap till the first consultation of the tertiary center was between 15 to 60 min. Therapy bundles could work very effective. The primary and continuous investment of training and communication is high, but much lower than the advantages of optimal therapy. The collaboration in our pediatric network was uncomplicated and fast. The implementation of the therapy strategies was effected in a short time. As a result of the optimized algorithm the medical costs could be reduced significantly. All participants of the network, including the patients, assessed the therapy bundle as a win-win situation. The typical interdisciplinary medical facilities in children hospitals and the traditional barrier free communication systems between pediatricians could be a good example for optimized medical structure.

069

Infection 2007; 35 (Suppl. II): 39

Rapid PCR-based pathogen identification in critically ill patients – an intensivist's perspective

Reinhart K

Department of Anaesthesiology and Intensive Care Medicine, University Jena, Germany

Adequacy of antibiotic therapy is crucial to reduce mortality in critically ill patients with infection and sepsis. Since identification of the underlying microorganisms by standard microbiological diagnostic procedures takes several days, initial antibiotic therapy must be empirically chosen. Uncertainty of antibiotic appropriateness may be improved if the underlying pathogen can be detected earlier. Microbial DNA from human blood can be detected by polymerase chain reaction (PCR) within one day. Recently, such a PCR based test has become commercially available (SeptiFast® test, Roche Diagnostics). However, little experience is available whether such a diagnostic tool can deliver clinical meaningful results in the intensive care setting.

In 36 critically ill patients without evidence of infection, 4.1 % out of 98 PCR samples tested positive compared to 5.1 % of the blood cultures. Thus, frequency of false positive results was similar in blood culture and in the PCR. In another study, where blood cultures and PCR were sampled in 68 patients with severe sepsis or septic shock, PCR revealed more positive results (37.3 % out of 110 samples) than the blood cultures (15.5 %). Higher rates of positive results in the PCR compared to blood culture confirm data from other four not yet published studies. Patients with a positive PCR for bacterial DNA had similar procalcitonin levels as patients with positive blood culture. Current studies so far did not prospectively address therapeutic impact of the pathogen detection by PCR. However, we and others retrospectively compared the positive PCRs of the sepsis patients with other microbiological findings and appropriateness of antibiotic (ABx) therapy (figure). 36.7 % of the positive PCR results suggested a different antimicrobial therapy. While two of the positive results were unlikely to be caused by infection but rather by contamination, antibiotic treatment would have needed adjustment in the remaining 13 patients.

In conclusion, PCR based pathogen detection is not only technically feasible but also inheres biological plausibility as bacterial DNA in the circulation is associated with signs of severe sepsis. Furthermore, current data are strongly suggestive that pathogen detection by measurement of bacterial DNA has impact on the clinical decision making. However, prospective studies are necessary to further elucidate the clinical utility and cost effectiveness of this intriguing new diagnostic approach.

071

Infection 2007; 35 (Suppl. II): 39

Holistic haemodynamic monitoring: could Minimally-invasive CCO and continuous ScvO₂ monitoring be practical combination?

Marx G

Department of Anaesthesiology and Intensive Care, University Jena, Germany

In the initial treatment of a critically ill patient, blood pressure, heart rate, urine output, and central venous pressure guide resuscitation. Despite normalization of these variables, global tissue hypoxia may still persist when the cardiorespiratory system is unable to cover metabolic demand adequately. In order to achieve adequate oxygen delivery, haemodynamic monitoring is necessary in order to provide appropriate cardiovascular support. Although measurements of regional perfusion and oxygenation are gaining increasing interest, cardiac output (CO) measurement is still regarded as the key haemodynamic variable in the assessment of cardiac function and guidance of therapy of critically ill patients. Flow, as gauged by CO, is the key measure of how well the circulation is delivering oxygen and nutrients to the vital organs and is the focus of all resuscitation efforts.

Compared to thermodilution technique requiring the insertion of a pulmonary artery catheter the minimally-invasive continuous CO (CCO) represents an alternative, less-invasive technique for CO-measurement. This method of determining CO uses the arterial pres-

sure waveform analysis. Edwards Lifesciences, LLC (Irvine, CA) has introduced a system for monitoring cardiac output continuously (Vigileo/FloTrac) that does not require thermodilution or dye dilution. Instead, it calculates the cardiac output by using arterial waveform characteristics in conjunction with patient demographic data. The accuracy and precision of the algorithm used by the Vigileo/FloTrac system has been evaluated satisfactorily (1, 2, 3).

On the other hand sustained tissue hypoxia is one of the most important cofactors in the development of multiorgan failure. Central venous oxygen saturation reflects the balance between oxygen requirements and oxygen delivery, and thus may be used to assess the adequacy of tissue oxygenation. The beauty of central venous oximetry is that it only needs the insertion of a central venous catheter. Early goal directed therapy for patients with severe sepsis or septic shock, which includes treatment goals for mean arterial pressure, central venous pressure, and central venous oxygen saturation, was able to increase survival in these patients.

The combined minimal invasive continuous monitoring of CO using FloTrac and of oxygen requirements using central venous oxygen saturation (using e.g. PreSep) can easily be applied in septic patients to evaluate the adequacy of tissue oxygenation and consumption. Thus, it facilitates immediate assessment of unpredictable rapid changes of the haemodynamic status in sepsis, thereby offering specific therapeutic interventions in cardiovascular insufficiency.

Literatur

Manecke Jr GR, Auger WR. Cardiac Output Determination From the Arterial Pressure Wave: Clinical Testing of a Novel Algorithm That Does Not Require Calibration. *J Cardiothorac Vasc Anesth* 21; 2007: 3-7.

de Waal EE, Kalkman CJ, Rex S, Buhre WF. Validation of a new arterial pulse contour-based cardiac output device. *Crit Care Med*. 2007 [Epub ahead of print].

Breukers RM, Sepehrkhoy S, Spiegelenberg SR, Groeneveld J. Cardiac Output measured by a new arterial pressure waveform analysis method without calibration compared with thermodilution after cardiac surgery. *J Cardiothorac Vasc Anesth* in press.

073

Infection 2007; 35 (Suppl. II): 40

Anticoagulation with citrate for continuous renal replacement therapy in the ICU

Brauer M

Department of Anaesthesiology and Intensive Care, University Jena, Germany

For many years trisodiumcitrate has been used as anticoagulant for packed red cells or blood samples. It can also be used for anticoagulation in continuous renal replacement therapy (CRRT). Citrate is infused into the arterial line of the hemodialysis kit and blocks coagulation through its ability to chelate ionized calcium. The sieving coefficient for citrate is 0.87. About 35 to 50 % of the citrate-calcium complex is filtered by dialysis. When the blood is given back into the patient through the venous line, citrate gets suddenly diluted and then metabolized mainly by the liver, thereby coagulation of the patient is not affected by citrate. The metabolism of 1 mmol of trisodiumcitrate yields 3 mmol NaHCO_3 . Although the basic principle is simple the occurrence of side effects has prevented its widespread acceptance till now. When using citrate for anticoagulation, some issues must be considered: A special replacement solution or dialysate is needed containing less sodium, because the citrate infusion already contains some sodium. The replacement solution or dialysate should be free of calcium in order to reduce the necessary dose of citrate, but calcium must be substituted into the venous line. The replacement solution or dialysate usually contains only little magnesium, therefore it must be supplemented. The acid base balance should be regularly monitored, as patients tend to become alcalotic under citrate anticoagulation because of NaHCO_3 generation. Hepatic function should be near normal, as citrate must be metabolized. Inadequate metabolism of citrate

is signalled by an increasing gap between total and ionized calcium. If these requirements are considered, citrate anticoagulation can be safely applied. Meanwhile different commercial available systems are on the market which make the practice of citrate anticoagulation much easier in practice. The most important advantage of citrate anticoagulation is the lacking risk of bleeding because citrate is only active in the extracorporeal circuit, making CCRT possible even in patients with a high risk of bleeding like neurosurgical patients. An additional advantage is the longer filter life longer using citrate: $80,2 \pm 60$ h, compared with heparin $30,2 \pm 30$ h. This compensates for the higher costs of citrate (daily costs approximately 72 € for citrate compared with 3,18 € for heparin) and hemodialysis-kit (up to 40 € more expensive compared with the standard kit).

081

Infection 2007; 35 (Suppl. II): 40

Procalcitonin (PCT) for diagnosing and postoperative monitoring of abdominal infections and sepsis

**Rau BM, Klar E*

Department of General, Thoracic, Vascular, and Transplantation Surgery, University of Rostock, Rostock, Germany

Introduction: Infections and sepsis are most serious complications in abdominal inflammatory disorders and after surgical interventions. Early and accurate diagnosis of septic complications remains a compelling issue for clinicians and novel approaches to overcome this problem are in demand. Among a large array of inflammatory parameters procalcitonin (PCT), the 116 amino acid pro-peptide of calcitonin, has emerged as valuable tool to identify patients at risk to develop infection and sepsis.

Methods: A review of the most important and recent literature about the usefulness of PCT determinations in patients with severe inflammatory abdominal disorders is presented.

Results: Although the term "sepsis" parameter does not fully embrace the integral properties of PCT a large number of clinical studies have outlined the pivotal role of this parameter in the host response to microbial and fungal infections. In patients with secondary peritonitis PCT is a reliable means to assess septic multisystem organ failure (MODS) and overall prognosis with sensitivity and specificity rates of more than 80%. In acute pancreatitis PCT allows early assessment of clinically relevant pancreatic infections and prognostic estimation alike yielding a sensitivity and specificity of 79% and 93%, respectively. Postoperative monitoring of PCT proved as a helpful tool to identify patients with evolving or persisting septic complications after elective and emergency abdominal surgery and may guide therapeutic decision-making such as the use of antibiotics. Irrespective of the underlying disease PCT outperformed the reference variable C-reactive protein (CRP).

Conclusions: Compared with established laboratory routine variables PCT significantly contributes to an earlier and better stratification of patients at risk to develop septic complications. Moreover, PCT provides excellent prognostic assessment in severe abdominal inflammation. The available test systems meet all demands for the use of PCT under clinical routine and emergency conditions.

083

Infection 2007; 35 (Suppl. II): 40

Microangiopathy and intravascular coagulopathy in severe sepsis

Brunkhorst FM

Department of Anaesthesiology and Intensive Care Medicine, University Jena, Germany

Background: Activation of the coagulation system in severe sepsis has been characterised by widespread intravascular fibrin deposition and

platelet aggregation (disseminated intravascular coagulation, DIC) with subsequent microvascular and tissue injury. However, the contributing role of platelets in the pathophysiology of sepsis is not entirely clear, although the degree and duration of thrombocytopenia, as well as the net change in the platelet count, are important determinants for survival. A reduction in platelet count may occur in the absence of a significant alteration in coagulation factors (as measured by prothrombin time or activated partial thromboplastin time), indicating that mechanisms other than DIC may be involved.

Objective(s): von-Willebrand Factor (VWF) is synthesized in endothelial cells and megakaryocytes and circulates in plasma as multimers up to 20,000 kDa in size, and is essential for platelet adhesion and thrombus formation. Its deficiency or dysfunction causes an inherited bleeding disorder, von Willebrand disease, whereas high plasma levels are associated with an increased risk of death from severe sepsis. ADAMTS13 is the main physiological modulator of the size and aggregability of VWF in plasma. In patients with thrombotic thrombocytopenic purpura (TTP), a congenital or immuno-mediated deficiency of ADAMTS-13 reduces or abolishes the degradation of ultralarge multimers of VWF (ULVWF) that cause the formation of intravascular platelet thrombi (thrombotic microangiopathy, TMA), resulting in multiorgan failure very similar to severe sepsis. There is an increasing evidence from the literature, that a decrease in ADAMTS13 activity is a common in patients with severe sepsis. This decrease may have prognostic implications, since it is associated with the progression of organ dysfunction.

Conclusions: The predictive value of these variables for disease severity and mortality in patients with severe sepsis warrants further evaluation in larger patient cohorts. Assessment of the functional proteolytic activity of ADAMTS13 and the detection of ULVWF may be of major clinical relevance, since plasma exchange with enzyme containing plasma preparations such as fresh frozen plasma (FFP), cryoprecipitate-poor plasma or the application of recombinant ADAMTS13 may restore the capacity to cleave ULVWF in the circulation.

093

Infection 2007; 35 (Suppl. II): 41

New technology: Preliminary results of continuous SCVO₂ Monitoring in pediatric patients

Ranucci M

Director of Clinical Research in the Department of Cardiothoracic and Vascular Anesthesia/Intensive Care, IRRCS Policlinico S. Donato, Milan, Italy

The evaluation of cardiac output in neonates and pediatric patients is impossible with traditional thermodilution methods, being pulmonary catheters unavailable for this small-sized patients. Pulse contour methods are not validated for pediatric patients, and echocardiography remains strongly operator-dependent.

Therefore, in clinical practice the cardiac output (or, better, the adequacy of cardiac output to the metabolic demands) in neonates and pediatric patients still remains assessed on the basis of indirect signs: urine output, peripheral pulses, temperature gradient, and SvO₂.

This last index is validated for the adequacy of cardiac output after cardiac operations, and is an independent predictor of mortality in this clinical condition. However, it requires serial blood samples to allow a strict monitoring of cardiac output adequacy. In small-sized patients this may lead to excessive blood removal and need for transfusions; moreover, it is not an on-line monitoring, and rapid changes of the hemodynamic status may be missed.

A recent new central venous catheter (PediSat®, Edwards Lifesciences, Irvine, CA) with continuous SvO₂ monitoring has entered in clinical use. We have studied this new device in 15 pediatric patients undergoing cardiac surgery at our Institution, with the aim of investigating its reliability in clinical practice.

The characteristics of these patients are shown in the table.

The SvO₂ values measured with the PediSat have been compared with the correspondent values obtained with standard oxymetry. Bland – Altman analysis was used to assess bias and error of the method.

When using an adequate protocol with calibration at the beginning of cardiopulmonary bypass and for every change in hemoglobin value (> 2 g/dL), we could obtain a bias of 1.2 mmHg, a precision of 3.6 mmHg, and a percentage error of 19%.

These values, according to the clinical practice, are highly acceptable as surrogate for standard laboratory assessment. However, we must consider, from this preliminary experience, that the application of a

Case	Age	Weight	Pathol	Site	Size	Length	CPB
1	4 ys	22 kgs	ASD	RIJV	5.5	8	Yes
2	2 ms	8 kgs	TOF	RIJV	4.5	5	Yes
3	9 ys	27 kgs	COA	FV	4.5	8	No
4	7 ms	8.7 kgs	TOF	RIJV	4.5	5	Yes
5	3.5 ys	18 kgs	ASD	RIJV	5.5	8	Yes
6	4 ys	14 kgs	TOF	FV	5.5	8	Yes
7	3 ys	15 kgs	TGA+PS	RIJV	5.5	8	Yes
8	1.5 ys	9.5 kgs	TOF	RIJV	4.5	5	Yes
9	8 ys	23 kgs	PAPVR	LIJV	5.5	8	Yes
10	4 ys	13 kgs	TOF	RIJV	5.5	8	Yes
11	7 ms	7.4 kgs	VSD	RIJV	4.5	5	Yes
12	3 ms	10 kgs	TOF	RIJV	4.5	5	Yes
13	6 days	3 kgs	TAPVR	RIJV	4.5	5	Yes
14	7 ys	23 kgs	AnCorArt	RIJV	5.5	8	Yes
15	1.5 ys	8.9 kgs	VSD	RIJV	4.5	8	Yes

strict protocol is needed for achieving a good reliability. Intraoperative measurements had the higher level of error; conversely, in the more steady conditions of the Intensive Care Unit, a better reliability was obtained

094

Infection 2007; 35 (Suppl. II): 42

The benefits of the pulmonary artery catheter in septic patients

Vincent J-L

Dept of Intensive Care, Erasme Hospital, Free University of Brussels, Belgium

The pulmonary artery (PA) catheter is an invasive tool to monitor acutely ill patients who need complex management. It provides three types of information, about intravascular pressures, measurement of cardiac output, and measurement of mixed venous oxygen saturation (SvO₂).

Several prospective, randomized, controlled studies have indicated that using a PA catheter does not improve outcomes. These results are worrisome, because there can be only two interpretations: One is that the use of the PA catheter does not influence therapy (and thus cannot influence outcomes); the other is that it does influence therapy but that the changes in therapy do not improve outcome (i.e., our interventions are useless).

Taking the second scenario, it seems unlikely that all ICU interventions are useless, first because there would simply be no need for our specialty if that were the case, but second, because there are a number of studies indicating that the intervention of an intensivist can indeed

positively influence outcomes. Therefore, if our interventions can be beneficial, perhaps it is the way in which the PA catheter findings are interpreted and used which is inadequate and I suggest that we need to revisit the basics of haemodynamic management and reassess the way in which the PA catheter is used, insisting on a strict three step process:

1. Correct measurement: pay attention to zeroing, calibration, elimination of artefacts, proper reading of the values – unfortunately, many errors are made in data collection.
2. Correct interpretation: this is based on a good knowledge of physiology and a complete integration of the three types of elements (pressures, cardiac output, and SvO₂) – unfortunately, there are still many clinicians who neglect simple elements like the pressure difference between the PA diastolic pressure and the PA occlusion pressure (PAOP), or between the PAOP and the right atrial pressure (RAP), or who measure cardiac output without concurrent SvO₂...
3. Correct application: data collection and interpretation is still useless if the results are not correctly applied to guide management. Unfortunately, even when data are correctly collected and interpreted, it is not uncommon to find that they have not been correctly applied to the individual patient and therapy with, e.g., fluids and inotropic agents remains suboptimal.

Some people may argue that to improve application we need therapeutic protocols. However, the PA catheter is indicated only in patients with complex issues influenced by multiple components; designing global protocols to guide therapy for such patients would be a difficult task and would represent an attempt to replace the good physician at the bedside. Just as interpretation of chest X-rays cannot be automated, proper haemodynamic management requires a good doctor at the bedside to integrate the individual factors present in each patient into a patient-specific management package. Let us concentrate our efforts on the training of such doctors.

AUTHOR INDEX

- Abel P 17
 Acht B 14
 Adamik B 8
 Ahlers O 19
 Ahluwalia D 29
 Albert T 11
 Annane D 19
 Audebert F 8

Baranyai J 34
 Barteczko B 10, 10
 Bartels H 3
 Bartkowski R 5
 Bauer M 13, 14, 17, 20, 21, 21, 22, 32, 33, 33, 34, 37
 Baumgaertel M 8
 Beck G C 9
 Behrend K 34
 Bein B 3
 Beyer J 5
 Biberthaler P 36
 Bier M 6
 Bierhaus A 6
 Bille J 37
 Binder L 26
 Bingold TM 15
 Bis B 7
 Bloos F 17, 19, 20, 21, 21, 23
 Bockmeyer CL 13, 13, 32, 32
 Bogatsch H 16, 16
 Bogner V 36
 Book M 15, 29, 30, 36
 Boost KA 29
 Bopp C 6
 Borggreffe M 13, 35
 Borzanovic M 7
 Bouillon B 36
 Brauer M 40
 Braun T 30
 Braune A 19
 Breitkreutz R 5
 Briegel J 19
 Brinkkoetter P T 9
 Brühl K 11
 Brückmann M 13, 35
 Brüne B 28
 Brunkhorst FM 13, 14, 16, 16, 17, 19, 20, 22, 41
 Brunner D 7
 Bryl A 23
 Bucher M 25
 Budde U 13
 Bühnen V 36
 Bunck AC 14
 Burgett U 20
 Busch C 27
 Busse H 27

Calandra T 37
 Chen QX 30
 Claus RA 13, 13, 14, 14, 25, 31, 32, 32, 32, 33, 33, 34
 Conradi F 32

Deigner HP 14, 34
 Demant T 12
 Djukanovic B 7
 Dombrowsky H 31
 Dorer M 14
 Durek G 10
 Duszynska W 10, 10

Eckardt KU 11
 Ehrenstein B 8
 Eiffert H 3
 Eisenbach C 28
 Eisenhut Y 34
 Emmanuel K 3

 Engel C 16, 16
 Engelhardt T 19
 Engelmann L 26
 Ertel W 36

Faist E 36
 Faulhaber-Walter R 23
 Fessele K 8
 Filipovic N 4
 Fliser D 23
 Forster C 11
 Frahm C 32
 Franck M 38
 Frank H 18
 Frerichs I 16, 16, 31
 Freudenhammer M 9
 Fricke R 12
 Frieze J 36
 Friesecke S 17
 Fuchs C 34

Geilfuss D 278
 Georgieff M 29
 Gerlach 19
 Gille C 25
 Glasl C 35
 Glück T 8
 Gortner L 22
 Graf BM 26, 27, 27
 Grimminger F 28
 Grotowska M 10, 10
 Gründling M 34, 35
 Guenther M 32

Haase U 11
 Hafer C 23
 Halbritter K 5
 Haller H 23
 Hamacher J 34
 Hansen C 10
 Hanusch C 9
 Hartog C 19
 Hecht M 28
 Heemann U 18
 Hein S 30
 Henrich D 36
 Henschke J 5
 Hochreiter M 3
 Hoeft A 15, 29, 30
 Hoegl S 29
 Hofer S 5, 6, 27
 Hoffarth B 29
 Hoffmann U 13, 35
 Hollenbach E 26
 Holzinger T 13
 Hoppe-Tichy T 4
 Hostmann A 36
 Huber J 23
 Huber W 18
 Hübner S 34
 Hueser N 3
 Hunfeld K 15
 Hunfeld KP 15, 38
 Hupe D 14
 Huse K 33

Illing A 34

Jahr N 23
 Jantzen JP 9
 Jatón-Ogay K 37
 Jevtic M 4
 Johann AM 28
 John S 11
 Jovic M 7

Kabir K 36

 Kajtor I 18
 Kaps M 28
 Kaskel P 4
 Keh D 19
 Kempf V 25
 Kempka-Dobra E 12
 Kentouche K 13
 Kersten R 11
 Kisvarga Z 18, 18
 Klank D 13
 Klar E 40
 Klatt S 28
 Klemm M 12
 Kleyman A 34
 Klinzing S 13
 Klösel S 15
 Kohl M 34
 Köhler 3
 Köhler J 17
 König IR 36
 Kortgen A 17
 Krassler J 7
 Krenn CG 22
 Kriner M 3
 Kubin T 30
 Kubitz S 12
 Kübler A 10, 10
 Kuchenbecker S-C 31
 Kuhlen R 16
 Kurt B 25

Ladwig E 34
 Lamoth F 37
 Lang S 13, 35
 Lautenschläger I 31
 Lehmann C 34, 35
 Lehmann LE 15, 29, 30, 36
 Leiber A 25
 Liebetrau C 13
 Lipinska-Gediga M 8
 Loeffler M 16
 Löffler EK 26, 27, 27
 Löffler M 16
 Lösche W 13, 14, 32, 32, 34
 Lüpertz M 36
 Lutz J 18

Maegele M 36
 Maher T 26
 Malfertheiner P 26
 Maravic-Stojkovic V 7
 Marchetti O 37
 Marjanek Z 18, 18
 Martin E 5, 6
 Martin J 38
 Marx G 13, 20, 21, 21, 22, 23, 25, 31, 39
 Marz S 11
 Marzi I 36
 Maschmeyer G 37
 Masuch M 22
 Matevosian E 3
 Meisner M 12
 Meißner W 22
 Mescha S 22
 Missler G 26
 Möller M 10
 Moreno R 19
 Morgenstern T 12
 Mrówczyński W 5, 12
 Muhl H 29
 Muhr G 36
 Müller D 33
 Müller E 4
 Münchow St 7

Nacul FE 20
 Nakamura H 5

- Nau R 3
 Neumann A 34
 Neumann J 14
 Neumann M 26
 Neuzner J 10
 Nguyen T 29
 Novotny A 3
 Nowak A 12
- Orlikowsky TW 25
 Ortutay A 18
 Otto G 17
- Park J-W 9
 Pavlovic D 34, 35
 Paxian G 22
 Paxian M 17, 22
 Pereszlenyi A 7
 Pfenninger E 28
 Platzner M 33
 Poets CF 25
 Pohlemann T 36
 Pöling J 30
 Poprawski G 12
 Prinz M 12
 Prodhom G 37
- Quintel M 16
- Radakovic S 4
 Rafat N. 9
 Ragaller M 16
 Ranucci M 41
 Rau BM 40
 Recknagel P 33
 Rees W 30
 Reichardt H 34
 Reindl W 18
 Reinhard C 13
 Reinhard K 13, 13, 14, 16, 16, 19, 20, 20, 21, 21, 22, 23, 25, 31, 39
 Revermann M 29
 Richter W 34
 Robl F 8
 Rolle A 7
 Ronco C 18
 Rönneberg T 15
 Rosengarten B 28
 Rosenhagen C 5, 28
 Rossaint R 16
 Rothaug J 22
 Rothe KF 12
 Rühl A 26
- Sachse S 20, 21, 21
 Saeger D 35
 Saenger J 25
 Sahre H 9
 Sakr Y 13, 20
- Salzberger B 8
 Sartorius S 15
 Sasse M 38
 Schade U 36
 Schädler D 16, 16
 Scheiermann P 29
 Scheller B 15
 Schellong S
 Schermuly R 28
 Schewe JC 15, 29, 30, 36
 Schleppers A 38
 Schmid W 34
 Schmidt C 25
 Schmidt J 35
 Schmitz G 16, 16
 Schneppenheim R 13
 Schölmerich J 8
 Scholz J 16, 16, 31
 Schönfeld AH 28
 Schott M 9
 Schrauzer T 11
 Schröder S 3
 Schröder T 11
 Schulte J 9
 Schumann M 11
 Schürholz T 13, 25, 31
 Schütz G 34
 Schweiger A-M 3
 Seifert H 15
 Senderrek M 20, 21
 Senftleben U 29
 Senn L 37
 Sieber MW 32, 32
 Siemonsen S 8
 Simon TP 13, 25, 31
 Singer M 19
 Sinner B 26, 27, 27
 Smuszkiewicz P 6, 23
 Sniatkowska-Bartkowska A 5, 23
 Soller M 28
 Sossdorf M 32
 Spasic T 7
 Spellerberg B 25
 Sponholz C 33
 Spreer A 3
 Spring B 25
 Sprung CL 19
 Steinfath M 31
 Stoltefaut V 3
 Straube E 20, 21, 21
 Stremmel W 26
 Stüber F 15, 29, 30, 36
 Surbatovic M 4
 Szecht D 12
- Thomas J 29
 Tomczak H 6
 Trentzsch H 36
 Trojanowska I 6, 23
- Tuckermann J 34
- Uhlig S 31
 Ulm K 3
 Umgelter A 18
- Vahlbruch J 23
 Vallance B 26
 van Ackern K 9
 van der Woude F J 9
 Van Griensven M 36
 Vincent J-L 42
 Vogel S 9
 Vogt N 28
 Volk HD 19
 Völz J-E 10
 von Knethen A 28
 von Spiegel T 3
- Wagner A 4
 Wagner F 29
 Wagner T 25, 31
 Wahrmann M 15
 Wanderer J 32
 Warnecke H 30
 Weber S 9, 30
 Weber SU 15, 29, 36
 Weigand MA 4, 5, 6, 27, 27, 28
 Weiler N 16, 16, 31
 Weise M 5
 Wendt M 35
 Werth M 17
 Westerski P 5
 Willam C 11
 Winkler M 4
 Winning J 14, 34, 34
 Wirtz DC 36
 Wissing H 4, 15, 15
 Witkowska A 23
 Witte OW 32
 Wodzinski A 5
 Wojtalik M 5, 12
 Woltmann A 36
 Wurm D 22
 Wuttke U 35
- Yard B A 9
 Yodoi J 5
- Zabel P 31
 Zahn R 8
 Zander J 29
 Zausig YA 26, 27, 27
 Zhang XH 30
 Zijlstra J G 9
 Zimmer A 22
 Zink W 26, 27, 27
 Zöller M 26
 Zwißler B 15, 29