Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Original Article

Viral Coinfection in Childhood Respiratory Tract Infections[☆]

A. Martínez-Roig,^a M. Salvadó,^b M.A. Caballero-Rabasco,^{a,*} A. Sánchez-Buenavida,^a
N. López-Segura,^a M. Bonet-Alcaina^a

^a Servicio de Pediatría, Hospital del Mar, Barcelona, Spain

^b Departamento de Genética y Microbiología, Laboratorio de Referencia de Cataluña, Barcelona, Spain

ARTICLE INFO

Article history:

Received 7 April 2013

Accepted 27 January 2014

Available online 1 December 2014

Keywords:

Coinfection

Respiratory tract infections

Respiratory syncytial virus

ABSTRACT

Introduction: The introduction of molecular techniques has enabled better understanding of the etiology of respiratory tract infections in children. The objective of the study was to analyze viral coinfection and its relationship to clinical severity.

Methods: Hospitalized pediatric patients with a clinical diagnosis of respiratory infection were studied during the period between 2009 and 2010. Clinical and epidemiological data, duration of hospitalization, need for oxygen therapy, bacterial coinfection and need for mechanical ventilation were collected. Etiology was studied by multiplex PCR and low-density microarrays for 19 viruses.

Results: A total of 385 patients were positive, 44.94% under 12 months. The most frequently detected viruses were RSV-B: 139, rhinovirus: 114, RSV-A: 111, influenza A H1N1-2009: 93 and bocavirus: 77. Coinfection was detected in 61.81%, 36.36% with two viruses, 16.10% and 9.35% with three to four or more. Coinfection was higher in 2009 with 69.79 vs 53.88% in 2010. Rhinovirus/RSV-B on 10 times and RSV-A/RSV-B on five times were the most detected coinfections. Hospitalization decreased with greater number of viruses ($P < .001$). Oxygen therapy was required by 26.75% (one virus was detected in 55.34% of cases). A larger number of viruses resulted in less need for oxygen ($P < .001$). Ten cases required mechanical ventilation, four patients with bacterial coinfection and five with viral coinfection ($P = .69$).

Conclusions: An inverse relationship was found between the number of viruses detected in nasopharyngeal aspirate, the need for oxygen therapy and hospitalization days. More epidemiological studies and improved quantitative detection techniques are needed to define the role of viral coinfections in respiratory disease and its correlation with the clinical severity.

© 2013 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

Coinfección vírica en las infecciones respiratorias infantiles

RESUMEN

Introducción: Las técnicas moleculares han permitido un mejor conocimiento de la etiología de las infecciones respiratorias infantiles. El objetivo del estudio fue analizar la coinfección viral y su relación con la gravedad clínica.

Métodos: Se estudió a pacientes pediátricos hospitalizados con diagnóstico clínico de infección respiratoria durante el periodo comprendido entre 2009 y 2010. Se recogieron datos clínicos, epidemiológicos, duración de la hospitalización, necesidad de oxigenoterapia, coinfección bacteriana y necesidad de ventilación mecánica. Etiología estudiada con técnica PCR múltiple y microarrays de baja densidad para 19 virus.

Resultados: Un total de 385 pacientes presentaron resultados positivos, 44,94% menores de 12 meses. Los virus más detectados fueron: VRS-B: 139, rhinovirus: 114, VRS-A: 111, influenza A H1N1-2009: 93 y bocavirus: 77. Se detectó coinfección en el 61,81%, un 36,36% con 2 virus, 16,10% con 3 y 9,35% con 4 o más. La coinfección fue superior en 2009 con 69,79 frente 53,88% en 2010. Rhinovirus/VRS-B en 10 ocasiones y VRS-A/VRS-B en 5 fueron las coinfecciones más detectadas. Menor hospitalización a mayor número de virus detectados ($p < 0,001$). Necesitaron oxigenoterapia el 26,75% (en 55,34% se aisló un virus), objetivando a mayor número de virus menor necesidad de oxígeno ($p < 0,001$). Precisarons ventilación mecánica 9 casos, 4 de ellos con coinfección bacteriana y 5 con coinfección vírica ($p = 0,69$).

Palabras clave:

Coinfección

Infecciones de vías respiratorias

Virus respiratorio sincitial

[☆] Please cite this article as: Martínez-Roig A, Salvadó M, Caballero-Rabasco MA, Sánchez-Buenavida A, López-Segura N, Bonet-Alcaina M. Coinfección vírica en las infecciones respiratorias infantiles. Arch Bronconeumol. 2015;51:5–9.

* Corresponding author.

E-mail address: 97512@hospitaldelmar.cat (M.A. Caballero-Rabasco).

Conclusiones: Objetivamos una relación inversamente proporcional entre número de virus detectados en aspirado nasofaríngeo, necesidad de oxigenoterapia y días de hospitalización. Se necesitan más estudios epidemiológicos y mejoría en las técnicas de detección cuantitativa para definir el papel de las coinfecciones víricas en la enfermedad respiratoria y su correlación con la gravedad clínica.

© 2013 SEPAR. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Respiratory viral infection is a major source of morbidity and mortality in childhood. Just over one-third of preschool-aged children develop lower respiratory infections during their first year of life. Between 1% of 2% of these patients require admission to hospital.^{1,2}

The etiological agent is not always identified in these patients. Children with respiratory infections are generally treated as outpatients and the etiology of their disease is not investigated. If they are hospitalized, the commonly employed evaluation techniques are often not sufficiently sensitive.³

Experience and understanding of the role of viral coinfections in respiratory infections has grown in recent years, thanks to the introduction of molecular techniques.^{4–6} At present, the clinical data available on coinfection, in terms of both the number of viruses involved and the severity of the condition, are variable and even, at times, contradictory. These discrepancies may be due to factors that affect the various etiological agents, such as geographical region and methods of detection employed.

The aim of this study was to analyze the viral etiology of respiratory infections and the real clinical implications of coinfection, and its possible relationship to clinical severity in a population of hospitalized patients in a general hospital located in Barcelona with a secondary-level pediatric department. Etiology was determined using multiplex polymerase chain reaction (PCR) and low density microarrays.

Patients, Materials and Methods

Patients

Pediatric patients, aged between 7 days and 15 years, hospitalized for respiratory infections were included in a prospective study conducted from February 1, 2009 until December 31, 2010 in the Pediatric Department of Hospital de Mar, Barcelona. The basic inclusion criterion was an initial diagnosis of (a) respiratory tract infection (rhinopharyngitis, laryngitis, bronchitis or pneumonia), with clinical signs suggesting viral infection; (b) influenza-like syndrome; (c) suspected pertussis; or (d) pneumonia with clinical signs suggesting bacterial infection or empyema, poor progress during admission and suspected viral coinfection.

Personal and epidemiological data were collected together with the reason for admission and clinical progress, duration of hospitalization, oxygen requirements in cases with O₂ saturations $\leq 92\%$, and need for admission to the Pediatric Intensive Care Unit (PICU) for mechanical ventilation were recorded.

Cases were stratified by age groups (younger than 12 months, between 1 and 3 years, between 3 and 5 years and older than 5 years), and probable diagnosis at time of admission, days of hospitalization (pooled as less than 4 days, between 5 and 10 days and longer than 10 days), need for oxygen therapy, need for transfer to PICU for mechanical ventilation, number of viruses detected and presence of bacterial coinfection were recorded.

Samples for Virological Study

Nasopharyngeal aspirate was collected from all patients during the first 12 h of admission. In patients with clinical suspicion of

bacterial pneumonia, a nasopharyngeal aspirate specimen was also obtained at any time when poor progress was suspected to be due to viral or bacterial coinfection and not to a poor response to treatment or virulence of the primary etiological agent. Laboratory tests were performed on the same day that the specimen was obtained, except at weekends. Samples were stored at 2–8 °C until analysis.

Extraction of Nucleic Acids

Nucleic acid (RNA/DNA) extraction was performed using MagnaPure LC, Roche Diagnostics.

Amplification and Detection

The CLART® *PneumoVi* kit from Laboratorios Genómica, Madrid, Spain was used. This method can detect and characterize the 19 most common types and subtypes of human viruses causing respiratory infections, including: adenovirus; bocavirus; coronavirus; enterovirus (echovirus); influenza virus A (human H3N2, human H1N1, B, C and H1N1/2009 subtypes); metapneumovirus (A and B subtypes); parainfluenza virus 1, 2, 3 and 4 (A and B subtypes); rhinovirus; respiratory syncytial virus type A (RSV-A); respiratory syncytial virus type B (RSV-B).

Viruses were detected using a multiplex polymerase chain reaction (PCR) technique after reverse transcription of viral RNA (RT-PCR) for amplification of a specific 120–330 base-pair fragment of the viral genome. The amplified product was visualized using the low-density microarray-based technological platform, CLART® (Clinical Array Technology).

Each amplification tube had an internal control to monitor amplification.

In 2009, detection of the influenza A virus H1N1/2009 was not included in the system, but was added in 2010. Until this time, identification was performed using real-time PCR (RT-PCR) that detected the M2 and HA1 genes, using the kit marketed by Roche.

Statistical Analysis

Data were stored in an Excel-supported database and the statistical analysis was performed using SPSS software. First, a univariate descriptive analysis was performed. Qualitative data were presented by the frequency distribution of percentages in each category. In the analytical statistics phase, the number of viruses detected was analyzed to determine any relationship with days of hospitalization, age and need for oxygen therapy. The association between these factors was investigated using hypothesis testing methods, and proportions were compared using the Pearson's Chi-squared test. Results with *P*-value $< .05$ were considered significant.

Results

Samples from 463 pediatric patients admitted to the Pediatrics Department of the Hospital del Mar, Barcelona, for community-acquired respiratory infections were analyzed. Results were positive in 385 (83.15%) samples. Age distribution was as follows: younger than 12 months, 173 (44.94%), from 12 months to 3 years, 143 (37.14%), from 3 to 5 years, 53 (13.77%) and older than 5 years, 16 (4.15%); 44.94% of the cases were under the age of 12. None of the patients had risk factors for respiratory diseases. Clinical diagnoses

Fig. 1. Viruses isolated in 2009 and 2010 in 385 children hospitalized in the Pediatric Department of the Hospital del Mar, Barcelona, for community-acquired respiratory tract infections.

of the positive patients were as follows: 100 cases of bronchiolitis, 92 pneumonia, 81 bronchitis, 80 upper respiratory tract infection, 25 influenza-like syndromes, six pertussis and one newborn with risk of infection (Table 1).

Viruses detected are shown in Fig. 1. The five most common viruses in both study periods were RSV-B (139 cases), rhinovirus (114), RSV-A (111), influenza virus serotype A H1N1/2009 (93) and bocavirus (77). The percentages of coinfections in each period were: (a) in 2009, 69.79% of infections were caused by more than 1 viral agent (134 of 192 infections detected); (b) in 2010, the proportion was 53.88% (104 of 193 infections detected). The most common coinfections were rhinovirus+RSV-B in 10 cases and RSV-A+RSV-B in 5.

Coinfection was found in 61.81% of samples from positive patients. Of these, two viruses were found in 36.36%, three viruses in 16.10% and four or more viruses in 9.35%.

We found a shorter hospital stay in patients with a greater number of viruses ($P<.001$), as reflected in Table 2; the association was linear.

The 103 patients who required oxygen therapy (basal saturations $\leq 92\%$) are described in Table 3. When these data were correlated, it was revealed that the younger the patient, the more frequent the need for oxygen therapy, a trend that was statistically significant ($P<.001$), and the greater the number of viruses detected the less need for oxygen therapy ($P<.001$).

Admission to the PICU for mechanical ventilation was required in nine cases. Age, clinical presentation, viruses detected and diagnosis are shown in Table 4. The predominant etiological agent in these patients was RSV. No statistically significant differences were found when viral coinfection was correlated with the need for mechanical ventilation in the PICU ($P=.69$).

Table 1
Relationship Between Number of Viruses Detected and Diagnoses.

Diagnosis	Number of viruses detected			Total
	1 virus	2 viruses	≥ 3 viruses	
Bronchiolitis	50	41	9	100
Pneumonia	29	36	27	92
Bronchitis	24	32	25	81
URTI	32	23	25	80
Flu-like syndrome	5	8	12	25
Pertussis	5	1	0	6
Neonatal risk of infection	1	0	0	1
Total	147	140	98	385

URTI: upper respiratory tract infection.

A clinical suspicion of bacterial coinfection based on the patients' clinical picture and acute phase reactant values was determined in 37 cases (8.50%). However, this was confirmed in only two cases, in which *Streptococcus pneumoniae* was confirmed in a positive blood culture or PCR. One virus was detected in 20 cases and two viruses in 14 cases. Hospital stay for 27 patients was between 1 and 4 days and between 5 and 10 days for 10 patients. In the eight cases with a diagnosis of atypical pneumonia on serology, viral coinfection with a single virus was detected in 5.

Discussion

The first finding of significance is the high rate of positive results in the population without risk factors (83.15% of samples studied) and the high rate of coinfection (61.81%), particularly in the first year of the study (69.79%). In the literature, coinfection rates range from 8% to 76%, depending on the series.^{7–16} With regard to the severity of our cases, it is important to note that a greater number of viruses is not synonymous with greater severity. According to our results, the number of viruses detected is in inverse proportion to the need for oxygen therapy and days of hospital admission; this relationship was statistically significantly ($P<.001$). However, of the nine cases admitted to the PICU, more presented coinfection, but this was not statistically significant.

Some studies have reported no clinical differences between patients with respiratory infections caused by a single agent and those caused by multiple viruses detected in nasopharyngeal aspirates from hospitalized children.^{8,11,13,17–23} Of interest is the study performed by Camargo²² in Sao Paulo during the H1N1 influenza epidemic of 2009, in which 21.9% of cases had coinfection not associated with greater morbidity or mortality. In contrast, other studies show that coinfection is a risk factor for poor progress.^{14,23–28} One of the most important of these was performed in Vienna by Aberle et al.,¹⁴ between 2000 and 2004. They reported that airway obstruction was more severe when RSV contributed to the coinfection, and mean hospital stay lengthened if rhinovirus was involved. In another study, Grensill et al.²⁶ showed that when the coinfection consisted of RSV and metapneumovirus, progress was poorer. Furthermore, Richard et al.,²⁸ who studied the relationship of coinfection with need for PICU admission, concluded that the presence of two or more viruses increased the likelihood of PICU admission 2.7-fold. However, in another study conducted in patients admitted to the PICU, Ghani et al.²⁹ found that only bacterial coinfection was associated with a longer hospital stay, with no increase in mortality.

Table 2
Correlation Between Hospital Stay and Detection of Respiratory Viruses.

Hospital stay (days)	Number of viruses detected			Total cases by hospital stay (%)
	1 virus	2 viruses	≥3 viruses	
1–4	67	76	77	220(57.1)
5–10	67	53	10	130(33.8)
>10	13	11	11	35(9.1)
Total cases with number of viruses detected (%)	147(38.2)	140(36.4)	98(25.5)	385(100)

P<.001.

Table 3
Correlation Between Age and Number of Viruses Detected in Patients Requiring Oxygen Therapy.

Age	Number of viruses detected			Total No. (%)
	1 virus	2 viruses	≥3 viruses	
<12 m	42	22	10	74(71.84)
1–3y	13	7	3	23(22.33)
>3–5y	1	3	1	5(4.85)
>5y	1	0	0	1(0.98)
Total n (%)	57(55.34)	32(31.07)	14(13.59)	103

m: months; y: years.
P<.001.

Table 4
Characteristics of Patients Admitted to the Pediatric Intensive Care Unit.

Age	Sex	Diagnosis	Type of infection
21 days	Male	Bronchiolitis/pneumonia	RSV-A+enterovirus+adenovirus+suspected bacterial coinfection
25 days	Male	Bronchiolitis/atelectasis	RSV-B
1 month	Female	Bronchiolitis	RSV-A
2 months	Female	Bronchiolitis	RSV-B
3 months	Female	Bronchiolitis	RSV-A+influenza B+adenovirus
3 months	Male	Bronchiolitis	Adenovirus
9 months	Male	Bronchitis	Rhinovirus+bocavirus
20 months	Female	Empyema	RSV-B+influenza B <i>S. pneumoniae</i>
4 years	Male	Empyema	RSV-A+H1N1-2009 <i>S. pneumoniae</i>

We should also remember that the detection of more than one virus may be due to the presence of viral fragments persisting for up to 5–6 weeks after onset of symptoms,¹⁴ which may not have any real role in the pathogenicity of the clinical process.³⁰ Relationships may be established with the help of epidemiological techniques, but the use of quantitative techniques would be much more useful for interpreting the contradictions produced by attributing greater or lesser severity to coinfections.

One important limitation of this study of the role of coinfection in upper respiratory tract infections is the small number of cases evaluated, compared to the high number seen in normal pediatric practice. The vast majority of cases do not need hospitalization, and this limits our knowledge of the real situation and prevents us from extrapolating our data to the general population. One final issue that remains unclear is the long-term effect that these coinfections may have on the respiratory system and the development of chronic lung disease.

Conclusions

Although an inverse relationship was found between the number of viruses detected in nasopharyngeal aspirate, the need for oxygen therapy and hospital stay, the results of our study differ widely from the data published by other groups. More epidemiological studies and improved quantitative detection techniques are needed to define the role of viral coinfections in respiratory disease and how they correlate with clinical severity.

Conflict of Interests

The authors state that they have no conflict of interests.

References

- Hall CB, Weinberg GA, Iwane MK, Blumkin AK, Edwards KM, Staat MA, et al. The burden of respiratory syncytial virus infection in young children. *N Engl J Med*. 2009;360:588–98.
- Madhi SA, Kuwanda L, Cutland C, Klugman KP. Five-year cohort study of hospitalization for respiratory syncytial virus associated lower respiratory tract infection in African children. *J Clin Virol*. 2006;36:215–21.
- Jin Yoo S, Kuak EY, Shin BM. Detection of 12 respiratory viruses with two-set multiplex reverse transcriptase-PCR assay using a dual priming oligonucleotide system. *Korean J Lab Med*. 2007;27:420–7.
- Mahony J, Chong S, Merante F, Yaghoubian S, Shina T, Lisle C, et al. Development of respiratory virus panel test for the detection of twenty human respiratory viruses by using multiplex PCR and a fluid microbead-based assay. *J Clin Microbiol*. 2007;45:3056–62.
- Brunstein JD, Cline CL, McKinney S, Thomas E. Evidence from multiplex molecular assays for complex multipathogen interactions in acute respiratory infections. *J Clin Microbiol*. 2008;46:97–102.
- Kumar RM. The widely used diagnostics DNA microarrays – a review. *Am J Infect Dis*. 2009;5:207–18.
- Frobert E, Escuret V, Javouhey E, Casalegno JS, Bouscambert-Duchamp M, Moulinier C, et al. Respiratory viruses in children admitted to hospital intensive care units: evaluating the CLARTR pneumovir DNA array. *J Med Virol*. 2011;83:150–5.
- Peng D, Zhao D, Liu J, Wang X, Yang K, Xicheng H, et al. Multipathogen infections in hospitalized children with acute respiratory infections. *Virol J*. 2009;6:155.
- Raymond F, Carbonneau J, Boucher N, Robitaille L, Boisvert S, Wu WK, et al. Comparison of automated microarray detection with real-time PCR assays for detection of respiratory viruses in specimens obtained from children. *J Clin Microbiol*. 2009;47:743–50.

10. Renois F, Talmud D, Huguenin A, Moutte L, Strady C, Cousson J, et al. Rapid detection of respiratory tract viral infections and coinfections in patients with influenza-like illnesses by use of reverse transcription-PCR DNA microarray systems. *J Clin Microbiol*. 2010;48:3836–42.
11. Suryadevara M, Cummings E, Bonville CA, Bartholoma N, Riddell S, Kiska D, et al. Viral etiology of acute febrile respiratory illnesses in hospitalized children younger than 24 months. *Clin Pediatr (Phila)*. 2011;50:513–7.
12. Calvo C, García-García ML, Blanco C, Vázquez MC, Frías ME, Pérez-Breña P, et al. Multiple simultaneous viral infections in infants with acute respiratory tract infections in Spain. *J Clin Virol*. 2008;42:268–72.
13. Marguet C, Lubrano M, Gueudin M, le Roux P, Deschildre A, Forget C, et al. In very young infants severity of acute bronchiolitis depends on carried viruses. *PLoS ONE*. 2009;4:e4596.
14. Aberle JH, Aberle SW, Pracher E, Hutter HP, Kundi M, Popow-Kraupp T. Single versus dual respiratory virus infections in hospitalized infants: impact on clinical course of disease and interferon-gamma response. *Pediatr Infect Dis J*. 2005;24:605–10.
15. Bonzel L, Tenenbaum T, Schrotten H, Schildgen O, Schweitzer-Krantz S, Adams O. Frequent detection of viral coinfection in children hospitalized with acute respiratory tract infection using a real-time polymerase chain reaction. *Pediatr Infect Dis J*. 2008;27:589–94.
16. Meerhoff TJ, Houben ML, Coenjaerts FE, Kimpen JL, Hofland RW, Schellevis F, et al. Detection of multiple respiratory pathogens during primary respiratory infection: nasal swab versus nasopharyngeal aspirate using real-time polymerase chain reaction. *Eur J Clin Microbiol Infect Dis*. 2010;29:365–71.
17. Papadopoulos NG, Moustaki M, Tsolia M, Bossios A, Astra E, Prezerakou A, et al. Association of rhinovirus infection with increased disease severity in acute bronchiolitis. *Am J Respir Crit Care Med*. 2002;165:1285–9.
18. Xepapadaki P, Psarras S, Bossios A, Tsolia M, Gourgiotis D, Liapi-Adamidou G, et al. Human metapneumovirus as a causative agent of acute bronchiolitis in infants. *J Clin Virol*. 2004;30:267–70.
19. Canducci F, Debiaggi M, Sampaolo M, Marinozzi MC, Berrè S, Terulla C, et al. Two-year prospective study of single infections and co-infections by respiratory syncytial virus and viruses identified recently in infants with acute respiratory disease. *J Med Virol*. 2008;80:716–23.
20. Nascimento MS, Souza AV, Ferreira AV, Rodrigues JC, Abramovici S, Silva Filho LV. High rate of viral identification and coinfections in infants with acute bronchiolitis. *Clinics (Sao Paulo)*. 2010;65:1133–7.
21. Martin ET, Kuypers J, Wald A, Englund JA. Multiple versus single virus respiratory infections: viral load and clinical disease severity in hospitalized children. *Influenza Other Respir Viruses*. 2012;6:71–7.
22. Camargo C, Guatura SB, Bellei N. Respiratory viral coinfection among hospitalized patients with H1N1 2009 during the first pandemic wave in Brazil. *Braz J Infect Dis*. 2012;16:180–3.
23. Huguenin A, Moutte L, Renois E, Leveque N, Talmud D, Abely M, et al. Broad respiratory virus detection in infants hospitalized for bronchiolitis by use of a multiplex RT-PCR DNA microarray system. *J Med Virol*. 2012;84:979–85.
24. Flores CJC, Vizcaya AC, Araos BR, Montecinos PL, Godoy MP, Valiente-Echeverría F, et al. Human bocavirus in Chile: Clinical characteristics and epidemiological profile in children with acute respiratory tract infections. *Rev Chilena Infectol*. 2011;28:504–11.
25. Marcone DN, Ellis A, Videla C, Ekstrom J, Ricarte C, Carballal G, et al. Viral etiology of acute respiratory infections in hospitalized and outpatient children in Buenos Aires, Argentina. *Pediatr Infect Dis J*. 2013;32:E105–10.
26. Greensill J, McNamara PS, Dove W, Flanagan B, Smyth RL, Hart CA. Human metapneumovirus in severe respiratory syncytial virus bronchiolitis. *Emerg Infect Dis*. 2003;9:372–5.
27. Paranhos-Baccalà G, Komurian-Pradel F, Richard N, Vernet G, Lina B, Floret D. Mixed respiratory virus infections. *J Clin Virol*. 2008;43:407–10.
28. Richard N, Komurian-Pradel F, Javouhey E, Perret M, Rajoharison A, Bagnaud A, et al. The impact of dual viral infection in infants admitted to a pediatric intensive care unit associated with severe bronchiolitis. *Pediatr Infect Dis J*. 2008;27:213–7.
29. Ghani AS, Morrow BM, Hardie DR, Argent AC. An investigation into the prevalence and outcome of patients admitted to a pediatric intensive care unit with viral respiratory tract infections in Cape Town, South Africa. *Pediatr Crit Care Med*. 2012;13:e275–81.
30. Brieu N, Guyon G, Rodière M, Segondy M, Foulongne V. Human bocavirus infection in children with respiratory tract disease. *Pediatr Infect Dis J*. 2008;27:969–73.