Published in final edited form as:

ACS Catal. 2020 February 7; 10(3): 2308–2313. doi:10.1021/acscatal.9b05383.

Highly Stereoselective Synthesis of Fused Cyclopropane- γ -Lactams via Biocatalytic Iron-Catalyzed Intramolecular Cyclopropanation

Xinkun Ren[†], Ajay L. Chandgude[†], Rudi Fasan^{*}

Department of Chemistry, University of Rochester, 120 Trustee Road, Rochester, NY 14627, United States

Abstract

We report the development of an iron-based biocatalytic strategy for the asymmetric synthesis of fused cyclopropane- γ -lactams, which are key structural motifs found in synthetic drugs and bioactive natural products. Using a combination of mutational landscape and iterative site-saturation mutagenesis, sperm whale myoglobin was evolved into a biocatalyst capable of promoting the cyclization of a diverse range of allyl diazoacetamide substrates into the corresponding bicyclic lactams in high yields and with high enantioselectivity (up to 99% *ee*). These biocatalytic transformations can be performed in whole cells and could be leveraged to enable the efficient (chemo)enzymatic construction of chiral cyclopropane- γ -lactams as well as β -cyclopropyl amines and cyclopropane-fused pyrrolidines, as valuable building blocks and synthons for medicinal chemistry and natural product synthesis.

Graphical Abstract

^{*}Corresponding Author: rfasan@ur.rochester.edu.

[†]Author Contributions

X.R and A.L.C. contributed equally to this work.

ASSOCIATED CONTENT

Supporting Information

Experimental procedures, additional Figures and Tables, compound characterization data, NMR spectra, chiral GC/SFC chromatograms, crystallographic data for **2b** and **2k**.

Notes

The authors declare no competing financial interest.

Keywords

Myoglobin; carbene transfer; cyclopropanation; protein engineering; lactams

Cyclopropanes fused to lactam and pyrrolidine units constitute key pharmacophores in many pharmaceuticals, such as cyproximide,¹ boceprevir,² amitifadine,³ and trovafloxacin⁴ (Figure 1). These structural motifs are also found in a number of biologically active natural compounds,^{5, 6} including the ergot alkaloid cycloclavine⁷ and the antibiotic rachelmycin and indolizomycin.⁸ In nature, these motifs are generated by action of SAM-dependent enzymes via SAM-mediated methyl transfer to an olefinic group of the substrate, followed by cyclization of the resulting carbocationic intermediate (e.g., indolizomycin).⁵ As revealed by the biosynthesis of cycloclavine, an alternative biosynthetic strategy involves the participation of a non-heme iron- α -keto-glutarate-dependent oxidase to catalyze the formation of a bicyclic dihydropyrrole core which is then reduced by a imine reductase to give a cyclopropyl-pyrrolidine.⁹ In this case, formation of the cyclopropane ring is believed to proceed via an hydroxylation (or halogenation) step followed by enamine-mediated intramolecular S_N2 or, more plausibly, via a ring-closing radical mechanism (Scheme 1a).^{9–11} Unfortunately, given the high substrate specificity of these biosynthetic enzymes, the scope of these enzymatic transformations beyond the native substrates remains very limited.

Toward introducing a biocatalytic strategy to access valuable cyclopropane-fused lactam scaffolds, we envisioned the possibility to promote the cyclization of N-allyl-diazoacetamides via an intramolecular cyclopropanation reaction catalyzed by an engineered myoglobin. In recent years, engineered myoglobins^{12–16} along with engineered P450s^{17–20} and artificial metalloenzymes^{21–28} have emerged as promising biocatalysts for catalyzing intermolecular olefin cyclopropanations via carbene transfer processes. More recently, the scope of myoglobin-based carbene transferases could be extended to the cyclization of diazoacetates, thus demonstrating the possibility to execute sterically demanding intramolecular cyclopropanation reactions within the ‘active site’ of this hemoprotein.²⁹ Building upon this progress, we envisioned the opportunity to exploit the myoglobin scaffold for directing the asymmetric synthesis of fused cyclopropane- γ -lactams starting from readily accessible diazoacetamide substrates. Synthetic methods to execute this transformation have been notably scarce, and limited to the use of rare and expensive metals (Rh, Ru).^{30–33} Furthermore, achieving broad substrate tolerance and/or high levels of enantioselectivity using these protocols has been notoriously challenged by competing side-reactions and the occurrence of *trans/cis* isomerism across the amide bond of diazoacetamides.³¹ Alternative methods to access chiral γ -lactams via Pd-catalyzed cyclopropane functionalization have also been reported.³⁴ Complementing these chemocatalytic approaches, we describe herein the development of an iron-based biocatalytic strategy for the highly enantioselective construction of fused cyclopropane- γ -lactams via the intramolecular cyclopropanation of allyl diazoacetamides (Scheme 1b). This approach offers an efficient and sustainable route to afford highly enantioenriched bicyclic lactams of prominent value for medicinal chemistry and/or natural product synthesis.

In initial studies, we tested wild-type sperm whale myoglobin (Mb) along with a panel of other hemoproteins (e.g., P450_{BM3}, catalase, cytochrome *c*) for their ability to catalyze the intramolecular cyclopropanation of (*E*)-2-diazo-*N*-(3-(4-fluorophenyl)allyl)-*N*-methylacetamide (**1a**) to give **2a** (Table 1). However, all of these proteins showed poor to no detectable activity (0–13% conversion) and very low enantioselectivity (1–9% *ee*) in this reaction (Table S1). We also tested Mb(H64V,I107S), a Mb variant previously optimized for the intramolecular cyclopropanation of allyl α -diazoacetates.²⁹ This biocatalyst showed improved activity for formation of **2a** compared to Mb (32% vs. 13% conv.) but significantly reduced enantioselectivity (45% *ee*; Table 1, entry 2) compared to the ester counterpart (99% *ee*)²⁹, highlighting the need to optimize the Mb scaffold for the transformation targeted here. This requirement can be attributed to the inherently different steric demands and conformational properties of the allyl diazoacetamide substrates compared to allyl diazoacetates as a result of the *N*-substituted amide vs. ester bond present in these molecules.

To develop a more efficient and selective biocatalyst for the present reaction, we screened an in-house mutability landscape library of Mb variants based on the distal histidine variant H64V in which the shape of the heme pocket was systematically varied by substituting each of the active site amino acid residues (i.e., Leu 29, Phe43, Val68, Ile107) to any of the other 19 amino acids (Table S2).¹³ Encouragingly, this strategy led to the identification of Mb(H64V,V68A) and Mb(H64V,V68G) as promising catalysts for converting **1a** to the bicyclic product **2a**, exhibiting significantly improved efficiency (57–59% yield) and enantioselectivity (81–82% *ee*; Table 1, entries 3–4) compared to Mb or Mb(H64V,I107S). The (1*R*,5*S*,6*S*)-configuration of **2a** was assigned based on crystallographic analysis of **2b** and **2k** (Fig. S4–S5; Tables S9–S10). Mb(H64V,V68A) was then subjected to site-saturation mutagenesis of the yet unaltered active site residues Phe43, Ile107 and Leu29, followed by screening of the resulting libraries in whole cells using **1a** as the substrate. The most promising ‘hits’ were validated by *in vitro* experiments with purified protein prior to the next round of directed evolution. Gratifyingly, progressive improvement of both the activity and stereoselectivity of the enzyme for the synthesis of **2a** was achieved through optimization of position 43 and 107 after two rounds of mutagenesis and library screening (Figure 2, Table S3). Ultimately, this process led to the development of Mb(F43Y,H64V,V68A,I107V) as an optimal biocatalyst for this reaction, enabling the quantitative conversion of **1a** into **2a** with 99% *ee* (Table 1, entry 5). Interestingly, the large majority of the beneficial mutations accumulated in this Mb variant entail an expansion of the distal heme pocket (i.e., His64→Val; Val68→Ala/Gly, Ile107→Val) compared to our previously reported Mb-based catalysts for intermolecular olefin cyclopropanation with EDA (Mb(H64V,V68A))¹² or intramolecular cyclopropanation of allyl diazoesters (Mb(H64V,I107S))²⁹, which is consistent with the increased steric demands associated with the cyclization of the *N*-substituted diazoacetamide substrate.

Further investigations with Mb(F43Y,H64V,V68A,I107V) demonstrated its compatibility with whole-cell reactions (Table S4) as well as its tolerance to higher substrate loadings (i.e., 5 mM), without noticeable loss in product conversion (94%) and enantioselectivity (98% *ee*; Table 1, entry 6). Under these conditions, the reaction reaches >80% conversion within 15

min (Entry 8) and full conversion in less than 3 hours (Figure S2). A whole-cell reaction with Mb(F43Y,H64V,V68A,I107V)-expressing *E. coli* cells was then performed at a 0.2 mmol scale resulting in the isolation of 36.9 mg of enantiopure **2a** (>99% *ee*) in 90% isolated yield (Entry 7), thus demonstrating the scalability of this biocatalytic method. Additional experiments showed that the Mb variant supports up to 440–450 total turnovers under catalyst-limited conditions (Table S4, entries 7 and 12) with an initial product formation rate of 27 turnovers min⁻¹.

To examine its substrate scope, Mb(F43Y,H64V,V68A,I107V) was then challenged with a diverse panel of allyl α -diazoacetamide derivatives in whole-cell reactions on a semi-preparative (0.2 mmol) scale (Table 2). To our delight, substrates carrying different *N*-substitutions such as methyl (**1b**), methoxy (**1c**) and ethyl (**1d**) groups were all efficiently processed to afford the desired bicyclic products **2b–2d** in high to quantitative yields (90–99%) and with excellent enantioselectivity (>99% *ee*). Allyl diazoacetamides with ‘unprotected’ secondary amide group such as **1e** have represented notoriously difficult substrates for transition metal-catalyzed cyclopropanation due to catalyst poisoning via amide coordination to the metal and/or competition by carbene insertion into the amide N-H bond.^{30–33} In stark contrast, **1e** could be processed by the Mb(F43Y,H64V,V68A,I107V) catalyst to yield the corresponding bicyclic lactam **2e** with excellent enantioselectivity (>99% *ee*) (Table 2, Entry 4). Heterocycle-containing **1f** was also efficiently cyclized to afford **2f** in 94% *ee*.

Encouraged by these results, we extended our studies to diazoacetamides containing unactivated olefinic groups such as **1g–1j**. While Mb(F43Y,H64V,V68A,I107V) was found capable of accepting all these compounds (23%–99% conv.), it showed reduced enantioselectivity for formation of **2h–2j** compared to the aryl-containing **2a** and **2b–2e** (28–90% *ee* vs 90–99% *ee*; Table S5). Importantly, the desired bicyclic products **2h–2j** could be obtained with high degree of enantioselectivity (90–99% *ee*) in good to quantitative yields (54–99%) using Mb(H64V,V68G), an earlier Mb variant identified during the catalyst evolution process (Fig. 2, Table S3). From these results it can be evinced that positions Phe43 and Ile107 are critical for high stereoselection in the presence of large (i.e. Ar) vs. small (-H, -Me) substituents connected to the olefinic group. The results with the *N*-phenyl- and *N*-allyl substituted diazoacetamides **2i** and **2j**, respectively, further demonstrated the broad tolerance of this Mb-based biocatalytic method to substitutions at the level of the amide group.

Finally, the tolerance of Mb(F43Y,H64V,V68A,I107V) toward substitutions of the phenyl group was assessed using **1k–1n**. Notably, *para*, *meta*, and *ortho* substitutions on this group were equally well tolerated by the biocatalyst resulting in the formation of the cyclopropyl- γ -lactams **2k–2n** in good to quantitative yields (71–99%) and with good to excellent enantioselectivity (90–99% *ee*; Entries 10–13). The results with the *ortho*-substituted **2n** highlighted the tolerance of this catalyst to steric hindrance in close proximity to the olefinic bond. Notably, both Mb variants exhibit a consistent (1*R*,5*S*,6*S*) stereoselectivity across this panel of substrates, as determined based on X-ray crystallography (Fig. S4–S5) and comparative chiral SFC/GC analyses.

In addition to enable the asymmetric synthesis of cyclopropyl- γ -lactam core structures for medicinal chemistry (Fig. 1), we envisioned the present method could offer a convenient tool for the chemoenzymatic synthesis of valuable β -cyclopropyl amines and cyclopropyl-fused pyrrolidines, the latter representing key pharmacophores in various drugs and bioactive natural products (Fig. 1). Illustrating this point, arylation of enantiopure **2c** obtained via Mb(F43Y,H64V,V68A,I107V)-catalyzed cyclization afforded the trisubstituted cyclopropane **3** in 83% isolated yield and 99% *ee* (Scheme 2). On the other hand, reduction of enantiopure **2d** in the presence of LiAlH₄ gave the bicyclic pyrrolidine **4** in 82% isolated yield as a single diastereomer (Scheme 2).

In summary, we have reported the development of a highly enantioselective, biocatalytic strategy for the asymmetric construction of bicyclic cyclopropane- γ -lactams from *trans*-allyl diazoacetamide derivatives. These biocatalytic transformations can be performed in whole cells and can be applied to gain stereoselective access to key synthons for medicinal chemistry and natural product synthesis. Based on these results, we foresee that engineered myoglobins can be leveraged to develop iron-based biocatalytic strategies for other types of challenging intramolecular carbene transfer reactions in the future.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

This work was supported by the U.S. National Institute of Health grant GM098628. X. R. acknowledges support from Sunivo LLC (USA). The authors are grateful to Dr. William Brennessel for assistance with crystallographic analyses. MS and X-ray instrumentation are supported by U.S. National Science Foundation grants CHE-0946653 and CHE-1725028.

REFERENCES

- (1). Epstein JW, Brabander HJ, Fanshawe WJ, Hofmann CM, McKenzie TC, Safir SR, Osterberg AC, Cosulich DB, and Lovell FM 1-Aryl-3-azabicyclo[3.1.0]hexanes, a new series of nonnarcotic analgesic agents, *J. Med. Chem.* 1981, 24, 481–490. [PubMed: 7241504]
- (2). Bacon BR, Gordon SC, Lawitz E, Marcellin P, Vierling JM, Zeuzem S, Poordad F, Goodman ZD, Sings HL, Boparai N, Burroughs M, Brass CA, Albrecht JK, Esteban R, and Investigators HR-. Boceprevir for previously treated chronic HCV genotype 1 infection, *N. Engl. J. Med.* 2011, 364, 1207–1217. [PubMed: 21449784]
- (3). Micheli F, Cavanni P, Andreotti D, Arban R, Benedetti R, Bertani B, Bettati M, Bettelini L, Bonanomi G, Braggio S, Carletti R, Checchia A, Corsi M, Fazzolari E, Fontana S, Marchioro C, Merlo-Pich E, Negri M, Oliosi B, Ratti E, Read KD, Roscio M, Sartori I, Spada S, Tedesco G, Tarsi L, Terreni S, Visentini F, Zocchi A, Zonzini L, and Di Fabio R 6-(3,4-dichlorophenyl)-1-[(methoxy)methyl]-3-azabicyclo[4.1.0]heptane: a new potent and selective triple reuptake inhibitor, *J. Med. Chem.* 2010, 53, 4989–5001. [PubMed: 20527970]
- (4). Gootz TD, Zaniewski R, Haskell S, Schmieder B, Tankovic J, Girard D, Courvalin P, and Polzer RJ Activity of the new fluoroquinolone trovafloxacin (CP-99,219) against DNA gyrase and topoisomerase IV mutants of *Streptococcus pneumoniae* selected in vitro, *Antimicrob. Agents Chemother.* 1996, 40, 2691–2697. [PubMed: 9124824]
- (5). Thibodeaux CJ, Chang WC, and Liu HW Enzymatic chemistry of cyclopropane, epoxide, and aziridine biosynthesis, *Chem. Rev.* 2012, 112, 1681–1709. [PubMed: 22017381]

- (6). Wessjohann LA, Brandt W, and Thiemann T Biosynthesis and metabolism of cyclopropane rings in natural compounds, *Chem. Rev.* 2003, 103, 1625–1648. [PubMed: 12683792]
- (7). Furuta T, Koike M, and Abe M Isolation of Cycloclavine from the Culture Broth of *Aspergillus-japonicus* Saito, *Agr. Biol. Chem.* 1982, 46, 1921–1922.
- (8). Gomi S, Ikeda D, Nakamura H, Naganawa H, Yamashita F, Hotta K, Kondo S, Okami Y, Umezawa H, and Iitaka Y Isolation and Structure of a New Antibiotic, Indolizomycin, Produced by a Strain Sk2–52 Obtained by Interspecies Fusion Treatment, *J. Antibiot.* 1984, 37, 1491–1494. [PubMed: 6549003]
- (9). Jakubczyk D, Caputi L, Hatsch A, Nielsen CAF, Diefenbacher M, Klein J, Molt A, Schroder H, Cheng JZ, Naesby M, and O'Connor SE Discovery and reconstitution of the cycloclavine biosynthetic pathway-enzymatic formation of a cyclopropyl group, *Angew. Chem. Int. Ed.* 2015, 54, 5117–5121.
- (10). Jakubczyk D, Caputi L, Stevenson CEM, Lawson DM, and O'Connor SE Structural characterization of EasH (*Aspergillus japonicus*) - an oxidase involved in cycloclavine biosynthesis, *Chem. Commun.* 2016, 52, 14306–14309.
- (11). Yan L, and Liu Y Insights into the mechanism and enantioselectivity in the biosynthesis of ergot alkaloid cycloclavine catalyzed by aj_eash from *Aspergillus japonicus*, *Inorg. Chem.* 2019, 58, 13771–13781. [PubMed: 31560525]
- (12). Bordeaux M, Tyagi V, and Fasan R Highly Diastereoselective and enantioselective olefin cyclopropanation using engineered myoglobin-based catalysts, *Angew. Chem. Int. Ed.* 2015, 54, 1744–1748.
- (13). Bajaj P, Sreenilayam G, Tyagi V, and Fasan R Gram-scale synthesis of chiral cyclopropane-containing drugs and drug precursors with engineered myoglobin catalysts featuring complementary stereoselectivity, *Angew. Chem. Int. Ed.* 2016, 55, 16110–16114.
- (14). Tinoco A, Steck V, Tyagi V, and Fasan R Highly Diastereo- and enantioselective synthesis of trifluoromethyl-substituted cyclopropanes via myoglobin-catalyzed transfer of trifluoromethylcarbene, *J. Am. Chem. Soc.* 2017, 139, 5293–5296. [PubMed: 28366001]
- (15). Chandgude AL, and Fasan R Highly diastereo- and enantioselective synthesis of nitrile-substituted cyclopropanes by myoglobin-mediated carbene transfer catalysis, *Angew. Chem. Int. Ed.* 2018, 57, 15852–15856.
- (16). Vargas D, Khade R, Zhang Y, and Fasan R Biocatalytic strategy for highly diastereo- and enantioselective synthesis of 2,3-dihydrobenzofuran based tricyclic scaffolds., *Angew. Chem. Int. Ed.* 2019, 58, 10148–10152.
- (17). Coelho PS, Brustad EM, Kannan A, and Arnold FH Olefin cyclopropanation via carbene transfer catalyzed by engineered cytochrome P450 enzymes, *Science* 2013, 339, 307–310. [PubMed: 23258409]
- (18). Knight AM, Kan SBJ, Lewis RD, Brandenburg OF, Chen K, and Arnold FH Diverse Engineered heme proteins enable stereodivergent cyclopropanation of unactivated alkenes, *ACS Central Sci.* 2018, 4, 372–377.
- (19). Brandenburg OF, Prier CK, Chen K, Knight AM, Wu Z, and Arnold FH Stereoselective enzymatic synthesis of heteroatom-substituted cyclopropanes, *ACS Catal.* 2018, 8, 2629–2634.
- (20). Chen K, Zhang SQ, Brandenburg OF, Hong X, and Arnold FH Alternate heme ligation steers activity and selectivity in engineered cytochrome P450-catalyzed carbene-transfer reactions, *J. Am. Chem. Soc.* 2018, 140, 16402–16407. [PubMed: 30372623]
- (21). Srivastava P, Yang H, Ellis-Guardiola K, and Lewis JC Engineering a dirhodium artificial metalloenzyme for selective olefin cyclopropanation, *Nat. Commun.* 2015, 6, 7789. [PubMed: 26206238]
- (22). Sreenilayam G, Moore EJ, Steck V, and Fasan R Metal substitution modulates the reactivity and extends the reaction scope of myoglobin carbene transfer catalysts, *Adv. Synth. Cat.* 2017, 359, 2076–2089.
- (23). Sreenilayam G, Moore EJ, Steck V, and Fasan R Stereoselective olefin cyclopropanation under aerobic conditions with an artificial enzyme incorporating an iron-chlorin e6 cofactor, *ACS Catal.* 2017, 7, 7629–7633. [PubMed: 29576911]

- (24). Dydio P, Key HM, Nazarenko A, Rha JYE, Seyedkazemi V, Clark DS, and Hartwig JF An artificial metalloenzyme with the kinetics of native enzymes, *Science* 2016, 354, 102–106. [PubMed: 27846500]
- (25). Ohora K, Meichin H, Zhao LM, Wolf MW, Nakayama A, Hasegawa J, Lehnert N, and Hayashi T Catalytic cyclopropanation by myoglobin reconstituted with iron porphycene: acceleration of catalysis due to rapid formation of the carbene species, *J. Am. Chem. Soc.* 2017, 139, 17265–17268. [PubMed: 29148750]
- (26). Villarino L, Splan KE, Reddem E, Alonso-Cotchico L, de Souza CG, Lledos A, Marechal JD, Thunnissen AMWH, and Roelfes G An artificial heme enzyme for cyclopropanation reactions, *Angew. Chem. Int. Ed.* 2018, 57, 7785–7789.
- (27). Zhao JM, Bachmann DG, Lenz M, Gillingham DG, and Ward TR An artificial metalloenzyme for carbene transfer based on a biotinylated dirhodium anchored within streptavidin, *Catal. Sci. Technol.* 2018, 8, 2294–2298.
- (28). Carminati DM, and Fasan R Stereoselective cyclopropanation of electron-deficient olefins with a cofactor redesigned carbene transferase featuring radical reactivity, *ACS Catal.* 2019, 9, 9683–9697.
- (29). Chandgude AL, Ren X, and Fasan R Stereodivergent intramolecular cyclopropanation enabled by engineered carbene transferases, *J. Am. Chem. Soc.* 2019, 141, 9145–9150. [PubMed: 31099569]
- (30). Doyle MP, Austin RE, Bailey AS, Dwyer MP, Dyatkin AB, Kalinin AV, Kwan MMY, Liras S, Oalman CJ, Pieters RJ, Protopopova MN, Raab CE, Roos GHP, Zhou QL, and Martin SF Enantioselective intramolecular cyclopropanations of allylic and homoallylic diazoacetates and diazoacetamides using chiral dirhodium(ii) carboxamide catalysts, *J. Am. Chem. Soc.* 1995, 117, 5763–5775.
- (31). Doyle MP, and Kalinin AV Highly enantioselective intramolecular cyclopropanation reactions of N-allylic-N-methyldiazoacetamides catalyzed by chiral dirhodium(II) carboxamidates, *J. Org. Chem.* 1996, 61, 2179–2184.
- (32). Abu-Elfotouh AM, Diem PTN, Chanthamath S, Phomkeona K, Shibatomi K, and Iwasa S Water-soluble chiral ruthenium(ii) phenyloxazoline complex: reusable and highly enantioselective catalyst for intramolecular cyclopropanation reactions, *Adv. Synth. Catal.* 2012, 354, 3435–3439.
- (33). Mandour HSA, Nakagawa Y, Tone M, Inoue H, Otag N, Fujisawa I, Chanthamath S, and Iwasa S Reusable and highly enantioselective water-soluble Ru(II)-Amm-Pheox catalyst for intramolecular cyclopropanation of diazo compounds, *Beilstein J. Org. Chem.* 2019, 15, 357–363. [PubMed: 30800184]
- (34). Pedroni J, and Cramer N Chiral gamma-lactams by enantioselective palladium(0)-catalyzed cyclopropane functionalizations, *Angew. Chem. Int. Ed.* 2015, 54, 11826–11829.

Figure 1.
Drugs and bioactive natural products containing cyclopropyl-fused lactam and pyrrolidine moieties

Figure 2.
Overview of biocatalyst evolution process.

a) Cycloclavine biosynthesis: **b) Biocatalytic construction of cyclopropyl-fused lactams and -pyrrolidines (this work):** **Scheme 1.**

Biosynthetic vs. (chemo)biocatalytic synthesis of cyclopropyl-fused lactam/pyrrolidine scaffolds.

Scheme 2. Chemoenzymatic synthesis of β -cyclopropyl amines and cyclopropyl-fused pyrrolidines

Table 1.Intramolecular cyclopropanation of allyl- α -diazoacetamide **1a** with Mb and variants thereof.^a

Entry	Catalyst	OD ₆₀₀	Yield ^b	TON	<i>e.e.</i>
1	Mb	-	13%	15	2%
2	Mb (H64V,I107S)	-	32%	40	45%
3	Mb (H64V,V68A)	-	59%	75	82%
4	Mb (H64V,V68G)	-	57%	70	81%
5	Mb (F43Y,H64V,V68A,I107V)	-	>99%	125	99%
6 ^c	Mb (F43Y,H64V,V68A,I107V)	-	94%	235	98%
7 ^c	Mb (F43Y,H64V,V68A,I107V)	40	>99% (90%) ^e	240	>99%
8 ^d	Mb (F43Y,H64V,V68A,I107V)	40	82%	200	>99%

^aReaction conditions: 2.5 mM (*E*)-2-diazo-*N*-(3-(4-fluorophenyl)allyl)-*N*-methylacetamide (**1a**), 20 μ M Mb variant (or C41(DE3) *E.coli* cells at indicated OD₆₀₀) in KPi buffer (50 mM, pH 7), 10 mM Na₂S₂O₄ (protein only), r.t., 16 hours in anaerobic chamber.

^bGC yield based on the calibration curves with authentic standards.

^cWith 5 mM **1a**.

^dReaction time: 15 min.

^eisolated yield.

Table 2.Substrate scope for Mb(F43Y,H64V,V68A,I107V)-catalyzed cyclization of allyl α -diazacetamides.^a

Entry	Product	Yield ^b	<i>e.e.</i>
1	 2b	93 % (82 %)	>99%
2	 2c	99 % (89 %)	>99%
3	 2d	90 % (82 %)	>99%
4	 2e	31 % (23 %)	>99%
5	 2f	82 % (70 %)	94%
6 ^c	 2g	54 % (45 %)	90%

Entry	Product	Yield ^b	<i>e.e.</i>
7 ^c	 2h	99 % (71 %)	94%
8 ^c	 2i	99 % (83 %)	99%
9 ^c	 2j	57 % (43 %)	93%
10	 2k	75 % (69 %)	>99%
11	 2l	71 % (67 %)	90%
12	 2m	99 % (90 %)	>99%
13	 2n	99 % (81 %)	>99%

^aReaction conditions: 5 mM allyl α -diazoacetamide, Mb(F43Y,H64V,V68A,I107V)-expressing *E.coli*. (OD₆₀₀ = 40) in KPi buffer (50 mM, pH 7), 40 mL-scale, r.t., 16 h.

^bProduct conversion as determined by GC. Yields of isolated products are reported in brackets. Errors are within 10%.

^cUsing Mb(H64V,V68G)-expressing *E.coli*. (OD₆₀₀ = 40).