Published in final edited form as:

J Genet. ; 98: . doi:10.1007/s12041-019-1155-5.

A *de novo* truncating mutation in *ASXL1* associated with segmental overgrowth

Stephanie Efthymiou¹, Vincenzo Salpietro¹, Erica Pironti², Maria Bonsignore², Valentina Ferrazzoli³, Gabriella Di Rosa², Henry Houlden^{1,*}

¹Department of Neuromuscular disorders, Institute of Neurology, University College London, London WC1N 3BG, UK

²Department of Human Pathology of the Adult and Developmental Age 'Gaetano Barresi', Unit of Child Neurology and Psychiatry, University of Messina, Messina 98125, Italy

³Department of Biomedicine and Prevention, University of Tor Vergata, Rome 00133, Italy

Abstract

Mutations in genes involved in chromatin remodelling have been implicated in broad phenotypes of congenital abnormalities and neurodevelopment. However, limited genotype–phenotype correlations are available for some of the rarest genetic disorders that affect chromatin regulation. We hereby describe a 12-year-old girl presented at birth with severe hypotonia, developmental delay, a mid-line capillary malformation and distinctive craniofacial features. During the natural history of her disease, the girl developed severe spasticity and drug-resistant seizures, leading to a diagnosis of Bohring–Opitz syndrome (BOS). We performed whole-exome sequencing (WES) and identified a *de novo* mutation in *ASXL1* (c.2033dupG) which results in the introduction of a premature stop codon (p.R678fs*6). *ASXL1* encodes a polycomb repressive complex protein implicated in chromatin regulation and *de novo* mutations are a known cause of BOS. Phenotypes with segmental craniofacial overgrowth associated to midline capillary malformations enlarge the clinical spectrum of BOS at onset and further expand the differential diagnosis in *ASXL1* mutation carriers.

Keywords

Bohring–Opitz syndrome; *ASXL1* gene; segmental overgrowth; nevus flammeus; macrocephaly capillary malformation

Introduction

Bohring–Opitz syndrome (BOS) is a rare autosomal dominant (AD) syndrome, characterized by the variable combination of severe intrauterine growth restriction, feeding difficulties, hypotonia, profound intellectual disability (ID), trigonocephaly with distinctive facial features, peculiar facial *nevus flammeus* and broad MRI abnormalities (e.g. hypomyelination, hypoplastic corpus callosum, spinal cord abnormalities etc.). BOS has

*For correspondence. h.houlden@ucl.ac.uk.

been reported in literature to be caused in up to 75% of cases by *de novo* truncating variants in *ASXL1*. This gene encodes the additional sex combs-like 1 protein involved in chromatin remodelling. These epigenetic processes are mediated either by histone H2A deubiquitination 1 and/or transcriptional regulation of the polycomb group repressor complex 2 mediated homeobox (*HOX*) gene. *HOX* genes are critical for body patterning and segmental identity during human embryogenesis (Hoischen *et al.* 2011). Bone marrow stromal cells (BMSCs) are an example of such homeotic genes that need to be regulated by *ASXL1* to achieve a homeostatic balance between self-renewal and differentiation. An imbalance in the cell fate or cell lineage commitment of BMSCs can lead to developmental or skeletal defects, as observed in BOS cases (Zhang *et al.* 2016). The Asxl1 protein is composed of an N-terminal DNA binding domain made up of the ASXN and ASXH subdomains, the ASXM1 and ASXM2 domains in the middle region as well as a PHD (zinc finger) domain in the C-terminal region, specifically important for mediating interactions with other proteins (figure 1d). *ASXL1* mutations have been identified as a cause of BOS in 2011 and most individuals reported so far have been found to carry *de novo* truncating mutations (table 1) (Magini *et al.* 2012; Dangiolo *et al.* 2015; Arunachal *et al.* 2016; Urreiziti *et al.* 2016; Carlston *et al.* 2017; Bedoukian *et al.* 2018).

Case report

Here, we report on the clinical characterization of a young girl affected by atypical BOS, whose phenotypic manifestations included global developmental delay with absent speech and severe motor impairment. She was presented with failure to thrive, hirsutism, glabellar *nevus flammeus* (observed since birth) and trigonocephaly with distinctive craniofacial features including a prominent globes, widely set eyes, synophrys and micrognathia (figure 1).

History

Previous neurological or genetic diseases was unremarkable in the family and parents were healthy (figure 1a). Pregnancy was complicated by threatened abortion and an intrauterine growth restriction documented by repeated ultrasound scans. The girl was born at 38 weeks with weight <10th centile for gestational age. At birth, a generalized low muscle tone as well as small hands with contractures of the fingers and a *nevus flammeus* on the face were observed. On the third day of life, she suffered from an epileptic seizure characterized by perioral cyanosis, hypertonia and eye deviation. Axial hypotonia was observed since the neonatal age and a severe impairment of motor milestones was noticed in the first months of life. During the first year of life she started to experience frequent (almost daily) episodes of generalized hypertonic seizures with lateral eye and mouth deviation, eyelid flutter and oral automatisms. Seizures were usually followed by hypotonia and regression of motor developmental milestones. A poor visual interaction was noticed since the first months of life as well as myopia later in life. Severe language impairment was also present, and the girl never attained an intelligible speech. She presented with microcephaly (occipitofrontal circumference <10th) and trigonocephaly as well as distinctive facial features including prominent eyes, micro and retrognathia.

Examination

On her neurological examination at the age of 15 years old, she presented with severe spastic paraparesis and dystonic posturing of the hands, wrists, elbows as well as foot and brisk osteotendinous reflexes. Her shoulders were externally rotated and adducted, elbows and wrists flexed in ulnar deviation, and ulnar deviation of the metacarpophalangeal joints.

Pathological findings

The patient was treated with antiepileptic medication during her first year of life, specifically valproate, and since the second year of life with levetiracetam, with good clinical response, as she was seizure-free until the age of 8 years. Extensive diagnostic and metabolic work-up was reported to be normal. Auditory brain response showed sensorineural hearing loss only in the left ear. The patient also underwent molecular investigation such as karyotype and array comparative genome hybridization (array-CGH), and both were reported as normal.

At the age of 5 years, electroencephalogram (EEG) activity recorded during sleep showed recurrent interictal high-voltage, bilateral, spike-wave discharges, followed by brief sequences of slow delta activities, interspersed with brief tracts of diffuse background slowing according to a fragmented hypsarrhythmia pattern. A progressive organization of background rhythm occurred. At 8 years of age, EEG activity was characterized by interictal high-voltage, bilateral, spike-wave discharges, at 3 Hz, on bilateral postcentral regions. Brain magnetic resonance imaging (MRI) performed at the age of 2 years, revealed brain asymmetry with reduced white matter volume in the left hemisphere, associated with enlargement of the left lateral ventricle and widening of the left hemispheric sulci; the corpus callosum is slightly thinned (figure 1g) These MRI features have been reported in BOS before (Dangiolo *et al.* 2015). The asymmetric appearance of the brain parenchyma appeared of mild degree, cortical abnormalities such as polymicrogyria, frequently associated with this syndrome, hydrocephalus or thickened corpus callosum were also noticed (Russell *et al.* 2015).

Materials and methods

After informed consent, we collected blood samples from the patient and her parents, and extracted DNA using standard procedures. To investigate the genetic cause of the disease, WES was performed in both the affected female and parents (figure 1a: II-1, I-1, I-2). Nextera Rapid Capture Enrichment kit (Illumina) was used according to the manufacturer instructions. Libraries were sequenced in an Illumina HiSeq3000 using a 100-bp paired-end reads protocol. Sequence alignment to the human reference genome (UCSC hg19), and variant calling, and annotation were performed as described elsewhere (Mencacci *et al.* 2016). In total, 55,531,100 (II-1) unique reads were generated. Sequencing data was reviewed for evidence of somatic mosaicism in both the proband and the parent's samples, as the *de novo* variant was present only in 16% of exome sequencing reads and an allele ratio of <35% was present in the proband. After removing all synonymous changes and variants not shared by the patient and the two parents, we filtered single-nucleotide variants (SNVs) and indels, considering only exonic and donor/acceptor splicing variants. In accordance with the pedigree and phenotype, priority was given to rare variants (<1% in

public databases, including 1000 Genomes project, NHLBI Exome Variant Server, Complete Genomics 69, and Exome Aggregation Consortium (ExAC v0.2)) that were fitting a recessive or a *de novo* model. We have not identified rare potentially damaging recessive variants in the girl, and a *de novo* truncating mutation in *ASXL1* (c.2033dupG), resulting in the introduction of a premature stop codon six amino acids downstream (p.R678fs*6), emerged as the most likely explanation for the disease pathogenesis. This is supported by the severe impact of the *ASXL1* mutation on protein function and the existing reports linking loss of function (LoF) variants in this gene to a similar clinical phenotype (Hoischen *et al.* 2011; Magini *et al.* 2012; Dangiolo *et al.* 2015; Arunachal *et al.* 2016; Urreizti *et al.* 2016; Carlston *et al.* 2017; Bedoukian *et al.* 2018). In addition, we screened *ASXL1* in >1000 trios exomes from individuals affected with genetically undiagnosed neurodevelopmental diseases recruited within the SYNAPS Study Group research initiative (neurogenetics.co.uk/synopathies-synaps), without identifying any additional *de novo* variant in this gene. Segregation analysis by traditional Sanger sequencing (primers used: forward (5'-GGTCACCACTGCCATAGAGA-3') and reverse (5'-GAGGATAAGGCGGCAGTAGT-3')) confirmed that mutation occurred *de novo* in the girl (figure 1b). The patient has provided informed consent for publication of the case.

Most *ASXL1* mutations reported so far have been found in the last exon (exon 12) of the gene, that includes ~50% of the whole-coding region of the gene, and where truncating variants are likely to generate transcripts that can escape nonsense mediated decay. The mutation that we identified occurs in exon 12, leading to a premature truncation of the downstream carboxyterminal plant homeodomain finger. Thus, the mutation may affect the binding of Asxl1 to chromatin, altering the activation or silencing of transcription factor genes involved in embryonic development (figure 1d). During the natural history of the disease in the girl, additional progressive craniofacial and neurological features became evident, leading to a clinical diagnosis of atypical BOS.

Discussion

Previous work in mice and *Drosophila* showed a significant variability with the phenotypes observed in BOS individuals carrying *ASXL1* truncating mutations. *Asxl1*^{-/-} mice present multiple developmental abnormalities, including anophthalmia, 30% reduction in body and skull size, resembling the microcephaly and changes in cranial shape observed in BOS individuals, as well as cleft palate and mandibular malformations (Zhang *et al.* 2016). This reinforces the role of Asxl1 as an essential protein needed for multilineage differentiation potential such as osteogenesis, adipogenesis and further skeletal and body development, and potentially explain both the skeletal and brain asymmetries that we observed. The patient was initially evaluated for the combination of her facial *nevus flammeus*, abnormal facial asymmetry (left > right), failure to thrive and global developmental delay and at that point, a segmental overgrowth disorder associated to midline capillary malformation was also suspected as part of the differential diagnosis for her condition.

Segmental overgrowth anomalies associated to midline capillary malformation enlarge the clinical spectrum of BOS at onset and further expand the differential diagnosis in *ASXL1*

mutation carriers, highlighting a possible involvement of chromatin remodelling and epigenetic regulation in the developmental and skeletal defects associated with BOS.

Mutations in other chromatin remodelling genes such as *DNMT3A* have been linked with developmental disorders associated to segmental overgrowth disorders, such as PIK3CA related phenotypes. Future studies are required to fully assess the potential epigenetic changes due to mutations in the above genes and their impact on molecular pathways associated with segmental overgrowth disorders associated with capillary anomalies.

Acknowledgements

We gratefully acknowledge the family for the enthusiastic collaboration to this study. This study was supported in part by The Wellcome Trust in equipment and strategic award (Synaptopathies) funding (WT093205 MA and WT104033AIA).

References

- Arunachal G, Danda S, Omprakash S, Kumar S. A novel de-novo frameshift mutation of the ASXL1 gene in a classic case of Bohring-Opitz syndrome. Clin Dysmorphol. 2016; 25:101–105. [PubMed: 27043953]
- Bedoukian E, Copenhaver D, Bale S, Deardorff M. Bohring-Opitz syndrome caused by an ASXL1 mutation inherited from a germline mosaic mother. Am J Med Genet Part A. 2018; 176:1249–1252. [PubMed: 29681100]
- Carlston CM, O'Donnell-Luria AH, Underhill HR, Cummings BB, Weisburd B, Minikel EV, et al. Pathogenic ASXL1 somatic variants in reference databases complicate germline variant interpretation for Bohring-Opitz Syndrome. Hum Mutat. 2017; 38:517–523. [PubMed: 28229513]
- Dangiolo SB, Wilson A, Jobanputra V, Anyane-Yeboah K. Bohring-Opitz syndrome (BOS) with a new ASXL1 pathogenic variant: Review of the most prevalent molecular and phenotypic features of the syndrome. Am J Med Genet Part A. 2015; 167a:3161–3166. [PubMed: 26364555]
- Hoischen A, van Bon BW, Rodriguez-Santiago B, Gilissen C, Vissers LE, de Vries P, et al. De novo nonsense mutations in ASXL1 cause Bohring-Opitz syndrome. Nat Genet. 2011; 43:729–731. [PubMed: 21706002]
- Magini P, Della Monica M, Uzielli ML, Mongelli P, Scarselli G, Gambineri E, et al. Two novel patients with Bohring-Opitz syndrome caused by de novo ASXL1 mutations. Am J Med Genet Part A. 2012; 158a:917–921. [PubMed: 22419483]
- Mencacci NE, Kamsteeg EJ, Nakashima K, R'Bibo L, Lynch DS, Balint B, et al. De novo mutations in pde10a cause childhood-onset chorea with bilateral striatal lesions. Am J Hum Genet. 2016; 98:763–771. [PubMed: 27058447]
- Russell B, Johnston JJ, Biesecker LG, Kramer N, Pickart A, Rhead W, et al. Clinical management of patients with ASXL1 mutations and Bohring-Opitz syndrome, emphasizing the need for Wilms tumor surveillance. Am J Med Genet Part A. 2015; 167a:2122–2131. [PubMed: 25921057]
- Urreizti R, Roca-Ayats N, Trepas J, Garcia-Garcia F, Aleman A, Orteschi D, et al. Screening of CD96 and ASXL1 in 11 patients with Opitz C or Bohring-Opitz syndromes. Am J Med Genet Part A. 2016; 170a:24–31. [PubMed: 26768331]
- Zhang P, Xing C, Rhodes SD, He Y, Deng K, Li Z, et al. Loss of Asxl1 Alters self-renewal and cell fate of bone marrow stromal cell, leading to bohring-opitz-like syndrome in mice. Stem Cell Rep. 2016; 6:914–925.

Figure 1.

(a) The pedigree diagram of the family with an *ASXL1* mutation. (b) Individual results of Sanger sequencing showing the proband (II.1) carrying the heterozygous *ASXL1* truncating mutation (c.2033dupG, p.R678fs), while being absent parents, I.1 and I.2. (c) Representation of *ASXL1* gene showing the amino acid count. (d) Representation of the protein domains; ASXN, conserved domain at the N-terminus; ASXM, conserved domain in the middle part; NR, nuclear receptor; PHD, plant homeodomain, and the position of the frameshift mutation in *ASXL1* (c.2033dupG, p.R678fs) as indicated. (e) An early and (f) late childhood portrait

photo showing nevus flammeus on the forehead of the proband, and facial dysmorphism typical of BOS syndrome (g) Axial and coronal T2-weighted images (wi) (top left, bottom left) and axial T1-wi (top right) show the asymmetry of the left hemisphere due to reduced white matter volume and consequent enlargement of the ipsilateral lateral ventricle and sulci. Midline sagittal T1-wi (bottom right) shows the thinned appearance of the corpus callosum and no features of hydrocephalus.

Table 1

Previously reported cases of children which have been clinically diagnosed with BOS and also genetically confirmed with *ASXL1* mutations. The three most common mutations are shown in bold.

Author	Cases	Mutations
Urriziti <i>et al.</i> (2018)	1 case	p.G646W/ <i>fs</i> *12
Bedoukian <i>et al.</i> (2018)	1 case	p.R965*
Carlston <i>et al.</i> (2017)	1 case	p.R404*
Arunachal <i>et al.</i> (2016)	1 case	p.G680R/ <i>fs</i> *38
Dangiolo <i>et al.</i> (2015)	1 case	p.F1373/ <i>fs</i>
Russell <i>et al.</i> (2015)	8 cases	p.C672W/ <i>fs</i> *4, p.G1026fs, p.G642*, p.L775*, p.I1919fs, p.Y425Q/ <i>fs</i> *12, p.S846/<i>fs</i>*5
Urriziti <i>et al.</i> (2015)	1 case	p.P701S/ <i>fs</i> *16
Magini <i>et al.</i> (2012)	2 cases	p.E803T/ <i>fs</i> *17, p.R965*
Hoischen <i>et al.</i> (2011)	7 cases	p. R404* , p.Q733*, p.Q778*, p.L823*, p.S846Q/<i>fs</i>*5 , p.Q925*, p.S1028*