Published in final edited form as:

Org Lett. 2020 April 03; 22(7): 2501–2505. doi:10.1021/acs.orglett.9b04347.

Enantioselective Synthesis of γ -Lactams by Lewis-Base Catalyzed Sulfenoamidation of Alkenes

Jesse L. Panger, Scott E. Denmark*

Roger Adams Laboratory, University of Illinois, 600 S. Mathews Ave., Urbana, Illinois 61801

Abstract

A method for the catalytic, enantioselective, intramolecular, 1,2-sulfenoamidation of alkenes is described. Lewis base activation of a suitable sulfur electrophile generates an enantioenriched, thiiranium ion intermediate from a β,γ -unsaturated sulfonyl carboxamide. This intermediate is subsequently intercepted by the sulfonamide nitrogen resulting in cyclization to form γ -lactams. Electron-poor alkenes required the use of a new selenophosphoramidate Lewis base catalyst. Subsequent manipulations of the products harness the latent reactivity of both the amide and thioether functionality.

Graphical Abstract

Alkenes serve as important building blocks for a variety of transformations central to synthetic chemistry.¹ Over the past few decades, enantioselective alkene difunctionalization has received considerable attention as a way to form new, stereochemically defined motifs from simple and abundant starting materials.^{2a-e} Whereas the use of Group 16 and 17 electrophiles to activate alkenes has been extensively studied in lactonization processes (Scheme 1A),^{2c,3} analogous reports for lactamizations are rare owing to the intrinsic preference for amide nucleophiles to react through the oxygen atom.⁴ Furthermore, a long-lived, configurationally stable iranium ion must be generated that allows for *N*-cyclization. In 2014, a preliminary report from these laboratories demonstrated the use of sulfonamides and carbamates as nucleophiles in the catalytic, enantioselective, intramolecular

*Corresponding Author: sdenmark@illinois.edu.

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/jacs.9b07019

Experimental Procedures, characterization data for all new compounds along with copies of spectra and chromatograms (PDF)

Data for **4b** (CIF)

Data for **4q** (CIF)

The authors declare no competing financial interests.

sulfenoamination of unactivated alkenes (Scheme 1B).^{5–6} A single example employed an amide nucleophile to give a δ -lactam, albeit with moderate enantioselectivity and under strongly acidic conditions. In 2015, Yeung and coworkers described an enantioselective halolactamization catalyzed by a cinchonaderived carbamate (Scheme 1C).⁷

As part of longstanding program on the enantioselective difunctionalization of alkenes, our laboratories have developed the Lewis base activation of Group 16 Lewis acid electrophiles to generate stereodefined thiiranium ions from various classes of alkenes followed by diastereospecific capture by a wide range of oxygen, nitrogen and carbon nucleophiles.^{8–9} In 2019, this strategy was exploited for the intermolecular capture of anilines and benzyl amines using hexafluoroisopropyl alcohol (HFIP) as a mildly acidic, activating solvent.¹⁰ With HFIP as both solvent and a weak acid source (pK_a 9.3),¹¹ we sought to re-examine amides as nucleophiles. The pK_a drop from sulfonamides (~ 12.3)¹² to sulfonyl carboxamides (<2)¹³ should enable efficient *N*-cyclization to the desired γ -lactams.

The γ -lactam ring is a privileged scaffold in medicinal chemistry.¹⁴ In addition to expressing a range of biological activities, γ -lactams are useful synthetic intermediates possessing latent reactivity allowing for further diversification.¹⁵ Various catalytic, enantioselective methods have been developed for the preparation of γ -lactams with NHCs^{16a–d} and transition-metal catalysis.¹⁷ Herein we disclose a general, enantioselective construction of γ -lactams using sulfenium ion-initiated amidation, in which the reactivity of both electron-rich and electron-poor alkenes have been successfully engaged (Scheme 1D).

Initial optimization studies (Table 1) were performed with (*E*)-*N*-tosylpent-3-enamide (**1m**), sulfonylating agent **3b**, catalyst (*S*)-**2a** or tetrahydrothiophene (THT), in HFIP as solvent. Without any Lewis base present, no reaction was observed (entry 1). The use of achiral Lewis base THT (entry 2), led to near quantitative conversion to the desired pyrrolidinone. However, using catalyst (*S*)-**2a** afforded much lower conversion after 24 h (entry 3). To increase the reaction rate a second set of experiments were conducted with a less bulky sulfonylating agent and a more nucleophilic styryl alkene (entries 4 and 5). In both cases, the yield of the racemic and enantioenriched products were high and an er of 84:16 was obtained. A further increase to the steric bulk of the sulfonylating agent (entry 6) did not appreciably erode the yield and there was a marked increase in er.

Although yet fully optimized, the scope of the sulfenolactamization with styryl-derived alkenes was explored using catalyst (*S*)-**2a** along with sulfonylating agent **3b** (Table 2). Electron-rich styryl alkenes proved to be competent reaction partners to generate highly enantioenriched γ -lactams. The parent β,γ -unsaturated sulfonamide **1a** gave product **4a** in 92% yield and 97:3 er. Furthermore, electron rich aryl groups (**4b–4d**) led to similarly good yields and high enantioselectivities. The (4*R*,5*S*)-configuration for the major enantiomer of **4b** was established by single crystal X-ray analysis¹⁸ which is in agreement with the stereochemical model from previous studies.^{8g} 2-Naphthyl and 4-tolylsulfonamide **1e** and **1f** afforded cyclized pyrrolidinones **4e** and **4f** in slightly reduced yields but maintained high enantioselectivity. Substitution at the *ortho*-position on the styryl arene with an electron-rich group (**4g**) was efficiently accommodated and indole sulfonamide **1h** was compatible under the reaction conditions to afford lactam **4h** in nearly quantitative yield and 98:2 er.

As was observed with aliphatic substrate **1m**, electron-deficient alkenes were much less reactive leading to low yields even after extended reaction times. To address this problem, two potential solutions were considered: (1) decrease the electron density on the sulfonylating agent to facilitate sulfonyl group transfer to the alkene, and (2) modify the catalyst itself to enhance reactivity of the sulfenium group. On the basis of previous studies, it was found that electronic perturbations of sulfonylating agents had little impact on the rate of sulfenium ion transfer.^{8g} Instead, a catalyst modification was envisioned such that a *decrease* in Lewis basicity was postulated to increase the rate of the sulfenium ion transfer. The justification for this counterintuitive hypothesis was that the attenuated Lewis basicity would increase the electrophilicity of the sulfenium ion so long as it retained sufficient Lewis basicity to form the catalytically active intermediate. To test this hypothesis, the simplest modification was to change from a BINAM-derived catalyst to one derived from BINOL.¹⁹ The synthesis began with enantioenriched (*R*)-BINOL which was deprotonated with NaH and then treated with diisopropylaminochlorophosphine to generate the P(III) intermediate and then was immediately oxidized with elemental selenium to form (*R*)-**2b** (Scheme 2). The preparation of this new Lewis base catalyst proved advantageous in several ways: the starting material is readily available, milder conditions are required for deprotonation, and the catalyst itself could be obtained in nearly quantitative yield after a simple filtration.

To test the viability of Lewis base (*R*)-**2b**, a comparison experiment was performed in which sulfonamide **1b** and (*R*)-**2b** were combined to determine if (*R*)-**2b** would function with a competent substrate. The BINOL-derived catalyst delivered (4*S*,5*R*)-**4b**²⁰ in 93% yield and 97:3 er, demonstrating that the two catalysts behave similarly with this substrate. Next, electron-deficient alkenes **1i**, **1j**, and **1k** were subjected to the standard reaction conditions with (*R*)-**2b** and afforded the corresponding cyclized products (**4i**, **4j**, **4k**) in synthetically acceptable yields and excellent enantioselectivities (Table 3). The presence of an *ortho* substituent on the styryl group resulted in the reduced yield and enantioselectivity (**4l**). If a second *ortho*-methyl group was incorporated, the reaction failed.

Next, catalyst (*R*)-**2b** was evaluated with the previously refractory substrates (*E*)-alkenes bearing alkyl substituents. Sulfenamidation of alkenes **1m** and **1n** still proved sluggish but nevertheless afforded products **4m** and **4n** in 55% and 57% yields, respectively, with a moderate drop in er for **4n**. 3-Butenamide substrate **1o** also exhibited a further drop in yield but maintained a 91:9 er. Prenyl sulfonamide **1p** performed better than its other alkyl counterparts in both yield and enantioselectivity, matching that of aryl substrates. Finally, the cyclic skipped diene **1q** was afforded the desymmetrized product **4q** in 78% yield and 90:10 er.²¹ The respectable enantioselectivity observed for this substrate was unexpected in view of the generally poor selectivity seen previously with (*Z*)-alkenes.^{8g} Furthermore, given that the opening of thiiranium ions is diastereospecifically *anti*, this result demonstrates that (*R*)-**2b** is capable of enantiotopic group differentiation.²²

To illustrate the utility of these products, synthetic manipulations of enantioenriched γ -lactam **4a** were explored (Scheme 3) following recrystallization to 99:1 enantiopurity. Detosylation was carried out with magnesium in methanol in 85% yield to unveil deprotected lactam **5**. A one-pot procedure to generate a mixture of sulfoxides followed by

spontaneous elimination upon addition of aqueous $\text{Na}_2\text{S}_2\text{O}_3$ proceeded to form the unsaturated lactam **6**. Deoxygenation of **4a** proceeded smoothly with $\text{BH}_3\cdot\text{SMe}_2$ to give enantiomerically enriched 2,3-disubstituted pyrrolidine **7** in 95% yield. Treatment with LiAlH_4 resulted in formation of ring-opened amino-alcohol **8** in excellent yield. Conversely, using DIBAL as a weaker reductant yielded a mixture of diastereomeric hemiaminal products which epimerized on silica gel to diastereomerically enriched intermediate **9** in 94% yield. The hemiaminal could undergo further transformations such as stereoselective cyanation with TMSCN in 96% yield (**10**) and allylation with allyltrimethylsilane in 65% yield (**11**) albeit in reduced yield owing to competitive formation of a pyrrole side product. Lastly, treatment with TFAA and lutidine allowed for the facile conversion of lactam **4a** to highly enantioenriched 4,5-disubstituted-2-pyrroline **12** in 98% yield.

In summary, an enantioselective Lewis base-catalyzed sulfenolactamization of β,γ -unsaturated sulfonyl carboxamides has been described. The reaction proceeds under mild conditions through formation of an enantiomerically enriched thiiranium ion followed by a diastereospecific capture with nitrogen to afford highly functionalized γ -lactams. The development of a new Lewis base catalyst has allowed for an expanded substrate scope of less nucleophilic alkenes. Moreover, the lactam products have been shown to undergo a variety of manipulations harnessing both the amide and thioether functionality to reach valuable synthetic targets.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We are grateful to the National Institutes of Health (GM R35127010) for generous financial support. We also thank the UIUC SCS support facilities (microanalysis, mass spectrometry, X-ray and NMR spectroscopy) for their assistance.

REFERENCES

- (1). Patai S The Chemistry of Double-Bonded Functional Groups; John Wiley & Sons: West Sussex, 1989.
- (2). (a) Kolb HC; VanNieuwenhze MS; Sharpless KB Catalytic Asymmetric Dihydroxylation. Chem. Rev 1994, 94 (8), 2483–2547. (b) Bodkin JA; McLeod D The Sharpless Asymmetric Aminohydroxylation. J. Chem. Soc. Perkin Trans 2002, 1 (24), 2733–2746. (c) Denmark SE; Kuester WE; Burk MT Catalytic, Asymmetric Halofunctionalization of Alkenes-A Critical Perspective. Angew. Chem., Int. Ed 2012, 51 (44), 10938–10953. (d) Landry ML; Burns NZ Catalytic Enantioselective Dihalogenation in Total Synthesis. Acc. Chem. Res 2018, 51 (5), 1260–1271. [PubMed: 29664281] (e) Xia Q-H; Ge H-Q; Ye C-P; Liu Z-M; Su K-X Advances in Homogeneous and Heterogeneous Catalytic Asymmetric Epoxidation. Chem. Rev 2005, 105 (5), 1603–1662. [PubMed: 15884785]
- (3). For reviews, see: (a) Tan CK; Zhou L; Yeung YY Organocatalytic Enantioselective Halolactonizations: Strategies of Halogen Activation. Synlett 2011, 1335–1339. (b) Murai K; Fujioka H Recent Progress in Organocatalytic Asymmetric Halocyclization. Heterocycles 2013, 87, 763–805. (c) Nolsøe JMJ; Hansen TV Asymmetric Iodolactonization: An Evolutionary Account. Eur. J. Org. Chem 2014, 3051–3065. (d) Kalyani D; Kornfilt DJ-P; Burk MT; Denmark SE Lewis Base Catalysis: A Platform for Enantioselective Addition to Alkenes Using Group 16 and 17 Lewis Acids ($\text{N} \rightarrow \Sigma^*$) In Lewis Base Catalysis in Organic Synthesis; Wiley-VCH Verlag

GmbH & Co. KGaA: Weinheim, Germany, 2016; pp 1153–1212.(e)Gieuw MH; Ke Z; Yeung YY Lewis Base Catalyzed Stereo- and Regioselective Bromocyclization. *Chem. Rec* 2017, 17, 287–311. [PubMed: 27701807]

- (4). (a)Alanine AID; Fishwick CWG; Szantay C Jr. Facile Preparation of 2-Imino Tetrahydrofurans, Pyrans, and Oxepans. *Tetrahedron Lett.* 1989, 30 (47), 6571–6572.(b)Cardillo G; Orena M Stereocontrolled Cyclofunctionalizations of Double Bonds Through Heterocyclic Intermediates. *Tetrahedron* 1990, 46 (10), 3321–3408.(c)Jaganathan A; Garzan A; Whitehead DC; Staples RJ; Borhan B A Catalytic Asymmetric Chlorocyclization of Unsaturated Amides. *Angew. Chem. Int. Ed* 2011, 50, 2593–2596.(d)Yin Q; You S-L Asymmetric Chlorocyclization of Indole-3-yl-benzamides for the Construction of Fused Indolines. *Org. Lett* 2014, 16 (9) 2426–2429. [PubMed: 24730758] (e)Yin Q; You S-L Enantioselective Chlorocyclization of Indole Derived Benzamides for the Synthesis of Spiro-indolines. *Org. Lett* 2013, 15 (16) 4266–4269. [PubMed: 23909796] (f)Wang YM; Wu J; Hoong C; Rauniyar V; Toste FD Enantioselective Halocyclization Using Reagents Tailored for Chiral Anion Phase-Transfer Catalysis. *J. Am. Chem. Soc* 2012, 134 (31), 12928–12931. [PubMed: 22830953] (g)Rauniyar V; Lackner AD; Hamilton GL; Toste FD Asymmetric Electrophilic Fluorination Using an Anionic Chiral Phase-Transfer Catalyst. *Science* 2011, 334, 1681–1684. [PubMed: 22194571] (h)Kawato Y; Kubota A; Ono H; Egami H; Hamashima Y Enantioselective Bromocyclization of Allylic Amides Catalyzed by BINAP Derivatives. *Org. Lett* 2015, 17 (5) 1244–1247. [PubMed: 25699577]
- (5). Denmark SE; Chi HM Lewis Base Catalyzed, Enantioselective, Intramolecular Sulfenoamination of Olefins. *J. Am. Chem. Soc* 2014, 136 (25), 8915–8918. [PubMed: 24926794]
- (6). Denmark SE; Chi HM Catalytic, Enantioselective, Intramolecular Sulfenoamination of Alkenes with Anilines. *J. Org. Chem* 2017, 82 (7), 3826–3843. [PubMed: 28294614]
- (7). Cheng YA; Yu WZ; Yeung YY Carbamate-Catalyzed Enantioselective Bromolactamization. *Angew. Chem. Int. Ed* 2015, 54, 12102–12106.
- (8). (a)Tao Z; Robb KA; Panger JL; Denmark SE Enantioselective, Lewis Base-Catalyzed Carbosulfonylation of Alkenylboronates by 1,2-Boronate Migration. *J. Am. Chem. Soc* 2018, 140 (46), 15621–15625. [PubMed: 30411878] (b)Matviitsuk A; Denmark SE Enantio- and Diastereoselective, Lewis Base Catalyzed, Cascade Sulfenoacetalization of Alkenyl Aldehydes. *Angew. Chem. Int. Ed* 2019, 58, 12486–12490.(c)Tao Z; Robb KA; Zhao K; Denmark SE Enantioselective, Lewis Base-Catalyzed Sulfenocyclization of Polyenes. *J. Am. Chem. Soc* 2018, 140 (10), 3569–3573. [PubMed: 29509003] (d)Denmark SE; Kornfilt DJP Catalytic, Enantioselective, Intramolecular Sulfenofunctionalization of Alkenes with Phenols. *J. Org. Chem* 2017, 82 (6), 3192–3222. [PubMed: 28257203] (e)Denmark SE; Rossi S; Webster MP; Wang H Catalytic, Enantioselective Sulfonylation of Ketone-Derived Enoxysilanes. *J. Am. Chem. Soc* 2014, 136 (37), 13016–13028. [PubMed: 25192220] (f)Denmark SE; Jaunet A Catalytic, Enantioselective, Intramolecular Carbosulfonylation of Olefins. *Preparative and Stereochemical Aspects. J. Org. Chem* 2014, 79 (1), 140–171. [PubMed: 24328051] (g)Denmark SE; Hartmann E; Kornfilt DJP; Wang H Mechanistic, Crystallographic, and Computational Studies on the Catalytic, Enantioselective Sulfenofunctionalization of Alkenes. *Nat. Chem* 2014, 6, 1056–1064. [PubMed: 25411883] (h)Hartmann E; Denmark SE Structural, Mechanistic, Spectroscopic, and Preparative Studies on the Lewis Base Catalyzed, Enantioselective Sulfenofunctionalization of Alkenes *Helv. Chim. Acta* 2017, 100, e1700158. [PubMed: 29311750] (i)Denmark SE; Kornfilt DJP; Vogler T Catalytic Asymmetric Thiofunctionalization of Unactivated Alkenes. *J. Am. Chem. Soc* 2011, 133 (39), 15308–15311. [PubMed: 21859086] (j)Denmark SE; Jaunet A Catalytic, Enantioselective, Intramolecular Carbosulfonylation of Olefins. *J. Am. Chem. Soc* 2013, 135 (17), 6419–6422. [PubMed: 23597174]
- (9). For examples of other enantioselective thiofunctionalizations, see:(a)Lucchini V; Modena G; Pasquato L Enantiopure Thiosulfonium Salts in Asymmetric Synthesis. *Face Selectivity in Electrophile Additions to Unfunctionalised Olefins. J. Chem. Soc., Chem. Commun* 1994, 1565–1566.(b)Liang Y; Zhao X Enantioselective Construction of Chiral Sulfides via Catalytic Electrophilic Azidothiolation and Oxythiolation of N-Allyl Sulfonamides. *ACS Catal.* 2019, 9, 6896–6902.(c)Xu J; Zhang Y; Qin.; Zhao X Catalytic, Regio-, and Enantioselective Oxytrifluoromethylthiolation of Aliphatic Internal Alkenes by Neighboring Group Assistance. *Org. Lett* 2018, 20, 6384–6388. [PubMed: 30265554] (d)Luo H-Y; Dong J-W; Xie Y-Y; Song X-F; Zhu D; Ding T; Liu Y; Chen Z-M Lewis Base/Brønsted Acid Co-Catalyzed Asymmetric

Thiolation of Alkenes with Acid-Controlled Divergent Regioselectivity. *Chem. Eur. J* 2019, 25, 15411–15418. [PubMed: 31489999] For an example of racemic thiofunctionalization of γ -lactams, see: Danilyuk IY; Vas'kevich RI; Vas'kevich AI; Vovk MV Electrophilic Intramolecular Cyclization of Functional Derivative of Unsaturated Compounds: VIII. Cyclization of 4-Aryl-N-(thiophen-3-yl)but-3-enamides by the Action of Polyphosphoric Acid and Chlorosulfanilarenes. *Russ. J. Org. Chem* 2016, 52, 987–992.

- (10). Roth A; Denmark SE Enantioselective, Lewis Base-Catalyzed, Intermolecular Sulfenaoamination of Alkenes. *J. Am. Chem. Soc* 2019, 141 (35), 13767–13771. [PubMed: 31433174]
- (11). Filler R; Schure RM Highly Acidic Perhalogenated Alcohols. A New Synthesis of Perfluoro-*t*-butyl Alcohol. *J. Org. Chem* 1967, 32 (4), 1217–1219.
- (12). Bordwell FG; Fried HE; Hughes DL; Lynch TY; Satish AV; Whang YE Acidities of Carboxamides, Hydroxamic Acids, Carbohydrazides, Benzenesulfonamides, and Benzenesulfonohydrazides in DMSO Solution. *J. Org. Chem* 1990, 55 (10), 3330–3336.
- (13). Serjeant EP, Dempsey B Ionisation Constants of Organic Acids in Aqueous Solution. International Union of Pure and Applied Chemistry (IUPAC). Pergamon Press: New York, New York, 1979.
- (14). Saldívar-González FI; Lenci E; Trabocchi A; Medina-Franco JL Exploring the Chemical Space and the Bioactivity Profile of Lactams: a Chemoinformatic Study. *RSC. Adv* 2019, 9, 27105–27116.
- (15). Caruano J; Muccioli GG; Robiette R Biologically active γ -lactams: Synthesis and Natural Sources. *Org. Biomol. Chem* 2016, 14, 10134–10156. [PubMed: 27748489]
- (16). (a) Dong S; Frings M; Zhang D; Guo Q; Daniliuc CG; Cheng H; Bolm C Organocatalytic Asymmetric Synthesis of trans- γ -Lactams. *Chem. Eur. J* 2017, 23, 13888–13892. [PubMed: 28833642] (b) Rommel M; Fukuzumi T; Bode JW Cyclic Ketimines as Superior Electrophiles for NHC-Catalyzed Homoenolate Additions with Broad Scope and Low Catalyst Loadings. *J. Am. Chem. Soc* 2008, 130 (51), 17266–17267. [PubMed: 19053399] (c) Raup DEA; Cardinal-David B; Holte D; Scheidt KA Cooperative Catalysis by Carbenes and Lewis Acids in a Highly Stereoselective Route to γ -Lactams. *Nat. Chem* 2010, 2, 766–771. [PubMed: 20729898] (d) Zhao X; DiRocco DA; Rovis T N-Heterocyclic Carbene and Brønsted Acid Cooperative Catalysis: Asymmetric synthesis of trans- γ -Lactams. *J. Am. Chem. Soc* 2011, 133 (32), 12466–12469. [PubMed: 21780842]
- (17). Ye LW; Shu C; Gagosz F Recent Progress Towards Transition Metal-Catalyzed Synthesis of γ -Lactams. *Org. Biomol. Chem* 2014, 12, 1833–1845. [PubMed: 24473105]
- (18). CCDC 1952721 contains the supplementary crystallographic data for (4R,5S)-4b. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
- (19). BINOL derived catalysts had been prepared previously in the context of selenofunctionalization and were inferior Postdoctoral report from Thomas Vogler, University of Illinois at Urbana-Champaign, 2009.
- (20). Determined by CSP-HPLC comparison with its enantiomer.
- (21). CCDC 1952722 contains the supplementary crystallographic data for (S,S,S)002D4q. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
- (22). Izumi Y; Tai A Stereo-Differentiating Reactions, Academic Press, New York, 1977.

A) Well studied halogen and chalcogen initiated lactonizations **B) Previous work (2014) - 1 example** **C) Yeung (2015) - single report** **D) This work**

Scheme 1.
Enantioselective Group 16 and 17 Initiated Cyclizations

Scheme 2.
Synthesis of BINOL-derived Lewis Base Catalyst (*R*)-2b.

Scheme 3. Product Manipulations.

^aAll reactions performed on 1.00 mmol scale. Diastereomeric ratio determined by ¹H NMR spectroscopic analysis of the crude reaction mixture. Enantiomeric ratio determined by CSP-HPLC analysis. ^bProduct epimerized at C(5) upon silica gel chromatography. ^cProduct epimerized at C(5) upon CSP-HPLC analysis.

Table 1.

Reaction Optimization.^a

<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;"> <p>3a: Aryl = Ph</p> <p>3b: Aryl = 2,6-(<i>i</i>-Pr)₂C₆H₃</p> </div> <div style="text-align: center;"> <p>(S)-2a</p> </div> <div style="text-align: center;"> <p>THT</p> </div> </div>						
entry	1	Sulfonylating agent	catalyst	conc. (M)	Yield (%) ^b	er ^c
1	1m	3b	-	0.2	0	-
2	1m	3b	THT	0.2	85 (93)	-
3	1m	3b	(S)-2a	0.2	(12)	Nd
4	1a	3a	THT	0.5	85 (88)	-
5	1a	3a	(S)-2a	0.5	83 (83)	84:16
6	1a	3b	(S)-2a	0.5	76 (85)	98:2

^aReactions performed on 0.1 mmol scale.^byield in parentheses determined by ¹H NMR spectroscopic analysis using 1,1,2,2-tetrachloroethane as an internal standard.^cdetermined by CSP-HPLC analysis.

Table 2.

Sulfenoamidation of Electron-Rich Styrenes.^{a-c}

92%
97:3 er

94%
97:3 er

88%
98:2 er

80%
96:4 er

82%
95:5 er

77%
96:4 er

95%
95:5 er

99%
98:2 er

^a All reactions performed on a 1.00 mmol scale.^b Yield of isolated, analytically pure product.

^cEnantiomeric ratio determined by CSP-HPLC analysis.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3.

Sulfenoamidation of Electron-Deficient Styrenes and Alkenes.^{a-c} ^a All reactions performed on a 1.00 mmol scale.^b Yield of isolated, analytically pure product.

^cEnantiomeric ratio determined by CSP-HPLC analysis.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript