Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Reference to Preferences for Cardiopulmonary Resuscitation

To the Editor:

Our comment is directed to the recommendation made in the recent Letters to the Editor by Dr. Bowers¹ stating that all hospitals should require informed consent for patients requesting cardiopulmonary resuscitation (CPR). We agree that the meaning of cardiopulmonary arrest and cardiopulmonary resuscitative measures, including survival data, needs to be explained in detail to the patient or next of kin/healthcare proxy. We also fully agree that if CPR were explained in detail, most, if not all, of the octogenarian patients and their families would choose a do-not-resuscitate (DNR) order. In our experience, when CPR is explained through informed consent, then the refusal rate is more than 90%. However, the assertion by Dr. Bowers that all hospitals should require informed consent for patients requesting CPR goes to the extreme and negates the basics of medicine: the sanctity of life, and changes the focus from saving life to that of withholding treatment. By making DNR the defaulted choice and CPR requiring informed consent would imply that life would not be maintained unless it is requested. This recommendation is not only entirely opposite to current practice, but contrary to the Hippocratic oath that is pledged by physicians² and also in opposition to the three Abrahamic monotheistic religions.³ In addition, it is in contradiction to current existing medical, ethical, and legal standards.

We suggest that all physicians continue to be more vigilant toward the discussion of code status and encouragement of formulation of advance directives (healthcare proxy and treatment preferences) on admission and throughout the course of hospitalization. We also recommend that physicians keep themselves engaged in discussion with the patient and next of kin/healthcare proxy so that if the time comes, the patient and/or healthcare proxy can make more informed decisions about withholding of treatment and DNR.

Abid Iraqi, MD, FACP, CMD

Terry Lynn Hughes, MS, RN, CS
Syracuse VA Medical Center
Syracuse, New York

REFERENCES

1. Bowers WR. Preferences for cardiopulmonary resuscitation [Letter]. *JAMDA* 2003;4:283.
2. Graham D. Revisiting Hippocrates: Does an oath really matter? *JAMA* 2000;284:2841–2842.
3. Clarfield AM, Gordon M, Markwell H, et al. Ethical issues in end-of-life geriatric care: The approach of three monotheistic religions—Judaism, Catholicism and Islam. *J Am Geriatr Soc* 2003;51:1149–1154.

DOI: 10.1097/01.JAM.0000126428.89797.38

Author's Reply

To the Editor:

Initially, my concerns were about cardiopulmonary resuscitation (CPR) for octogenarians; the median between life expectancy (in the United States) and lifespan (in the United States), 75–85 years. CPR has proven useless in this age group.

Informed consent for octogenarians requesting CPR does not negate the basics of medicine. It affirms our current legal standard of providing informed consent for invasive procedures.

Defaulting to do-not-resuscitate status simply recognizes a natural end-of-life phenomenon and is not in opposition to the three Abrahamic monotheistic religions, which, according to the reference article of Dr. Iraqi, supports a respect for the dying process when it is clearly imminent and irreversible!

Attending to the physical distress of an octogenarian is a physician's duty—comfort care. Trying to revive an octogenarian after cardiopulmonary arrest is simply paying homage to the false idol of medical technology, which is unrealistically promoted by the media.

The Hippocratic oath is a pledge that we physicians will not use our knowledge contrary to the laws of humanity. The laws of humanity have always respected the natural course of life and inevitable death, hopefully without suffering.

I agree with Dr. Iraqi's recommendation to keep ourselves engaged in discussion with our octogenarian patients and kin. I am very skeptical when an octogenarian is admitted to the nursing home as a "full code." These new residents and families are given further information and informed consent with the same refusal rate as Dr. Iraqi has experienced.

"Do unto others. . .," the Golden Rule, is also preached by the monotheistic religions. I do not know any physicians who would want their octogenarian parents or their octogenarian selves to be resuscitated after cardiopulmonary arrest. We need to preach what we practice. I have found that when I tell patients what I would do for my parents or for myself, refusal of CPR is approximately 100%.

William R. Bowers, MD, CMD
Athens Regional Medical Center
Life Care of Athens, TN
Brookwood NH of Decatur, TN
Athens, Tennessee

DOI: 10.1097/01.JAM.0000126429.97420.3A

What Has SARS Taught Us About Infection Control in Nursing Homes?

To the Editor:

Following submission of my editorial, "What Has SARS Taught Us About Infection Control in Nursing Homes?,"¹ the Centers for Disease Control and Prevention (CDC) has made new recommendations based on what they learned during the worldwide SARS outbreak. The CDC has recommended a

"Respiratory Hygiene/Cough Etiquette Strategy for Healthcare Facilities," including specific "Recommendations for Long-Term Care Facilities."² My editorial mentions the lack of sensitive, real-time diagnostic tests to determine which organism is responsible for cough and fever in a resident with an acute change in status. Even the rapid influenza tests have problems with sensitivity. The editorial also mentions that familiar respiratory viruses such as influenza, parainfluenza, and respiratory syncytial virus (RSV), especially complicated by pneumonia, have mortality rates in nursing home residents greater than the 9% figure cited for SARS cases in the community. Because clinicians seldom know the precise etiology of an infectious respiratory syndrome, the CDC now recommends universal respiratory hygiene for *all* patients who present with an infectious respiratory syndrome, especially during periods of increased respiratory activity in the community (ie, evidence of ongoing transmission). The approach includes droplet secretion precautions. Given the fact that influenza, SARS-CoV, parainfluenza, and RSV are all transmissible and potentially lethal in nursing home residents, and could be clinically indistinguishable, it seems prudent to initially approach all residents with symptoms of acute respiratory infection with droplet secretion precautions modified to the reality of your facility.³⁻⁵

Paul J. Drinka, MD, CMD
Wisconsin Veterans Home
King, Wisconsin
Department of Internal
Medicine/Geriatrics
University of
Wisconsin-Madison
Madison, Wisconsin
Medical College of
Wisconsin-Milwaukee
Milwaukee, Wisconsin

REFERENCES

1. Drinka PJ. Editorial: What has SARS taught us about infection control in nursing homes? *JAMDA* 2004;5:59-60.
2. Public Health Guidance for Community-Level Preparedness and Response to Severe Acute Respiratory Syndrome (SARS)-version 2, January 8, 2004. Available at: <http://www.cdc.gov/ncidod/sars/guidance/> (See Supplement C for Preparedness and Response in Healthcare Facilities). Accessed January 12, 2004.
3. Garner JS, Hospital Infection Control Practices Advisory Committee for CDC. 1996 guideline for isolation precautions in hospitals. *Infect Control Hosp Epidemiol* 1996;17:63-80.
4. Drinka P, Gravenstein S, Krause P, et al. Non-influenza respiratory viruses may overlap and obscure influenza activity. *J Am Geriatr Soc* 1999;47:1087-1093.
5. Drinka PJ, Langer L, Krause P, et al. Mortality following isolation of various respiratory viruses in nursing home residents. *Infect Control Hosp Epidemiol* 1999;20:812-815.

DOI: 10.1097/01.JAM.0000126430.97420.EC

Nurse Practitioner/Physician Collaborative Models of Care

To the Editor:

The comparative study by Aigner et al.¹ raises interesting issues regarding nurse practitioner/physician collaborative

models of care for nursing facility residents. Previous authors have documented differences in outcomes when advanced practice nurses (APNs) and physicians work as a team to enhance nursing services and provide medical care.²⁻⁴ A collegial review of the model is offered here in an effort to further explain the results.

We believe that comprehensive, consistent care is critical for frail institutionalized older adults who have complex medical, psychosocial, and behavioral issues. Understanding the interaction between these factors requires significant training in geriatric syndromes, as well as an understanding of the nursing process and systems unique to long-term care. The APNs in this study were placed on a rotating basis, which could have inhibited the opportunity to build consistent, stable relationships with patients, families, or staff. By rotating the APNs every 3 months, comprehensive primary care could have been compromised.

In addition, the APNs in this study were not geriatric nurse practitioners (GNPs). The APNs most likely had received training in the management of diseases of adulthood, but may have had less education and training than GNPs in the evaluation and management of geriatric syndromes such as falls, incontinence, geriatric depression, dementia behaviors, polypharmacy, involuntary weight loss, and pressure ulcers. Many GNP educational programs include content on nursing facility culture and processes, which can affect patient care. Geriatric nurse practitioners have had a positive impact on individual patients with attention to the geriatric syndromes,⁶ and geriatric clinical nurse specialists have been adept at influencing the system of care delivery.⁷ It appears the model in the study was primarily a physician substitution model with less attention to augmenting nursing services.

The APNs provided services common in a fee-for-service environment, including acute visits, mandated progress visits, and annual comprehensive examinations. Progress visits were to be made every 60 days, alternating with the physician. This goal was not met because residents were seen only 4.6 times per year for progress visits when six would have been expected. The APNs managed approximately 14 paged or voice-mail messages annually per patient but made only three acute visits per patient per year as a team. The APNs could have missed opportunities to intervene early in acute illness and prevent emergency department visits or hospitalizations. The American Medical Directors' Association (AMDA) has suggested one visit per month could be appropriate for a person who requires nursing care around the clock and one visit a week for persons receiving skilled nursing facility (SNF) services under Medicare part A.⁵

The physician-only arm consisted of physicians who would continue to follow their own patients as they entered nursing facilities in the community. Thus, patients were followed by a provider who presumably knew them and their families very well. The patients in this study were all in the same eight nursing facilities. Given the mix of the team-patients and physician-only patients in the same facilities, this leaves open the possibility that any benefit from the APN presence was also seen with the physician-only group, resulting in a halo effect. The same nurses and