Published in final edited form as:

J Abnorm Psychol. 2020 January ; 129(1): 38–48. doi:10.1037/abn0000465.

Levers and Barriers to Success in the Use of Translational Neuroscience for the Prevention and Treatment of Mental Health and Promotion of Well-Being Across the Lifespan

Sarah R. Horn, Philip A. Fisher, Jennifer H. Pfeifer, Nicholas B. Allen, Elliot T. Berkman

Department of Psychology and Center for Translational Neuroscience, University of Oregon.

Abstract

Neuroscientific tools and approaches such as neuroimaging, measures of neuroendocrine and psychoneuroimmune activity, and peripheral physiology are increasingly used in clinical science and health psychology research. We define *translational neuroscience* (TN) as a systematic, theory-driven approach that aims to develop and leverage basic and clinical neuroscientific knowledge to aid the development and optimization of clinical and public health interventions. There is considerable potential across basic and clinical science fields for this approach to provide insights into mental and physical health pathology that had previously been inaccessible. For example, TN might hold the potential to enhance diagnostic specificity, better recognize increased vulnerability in at-risk populations, and augment intervention efficacy. Despite this potential, there has been limited consideration of the advantages and limitations of such an approach. In this article, we articulate extant challenges in defining TN and propose a unifying conceptualization. We illustrate how TN can inform the application of neuroscientific tools to realistically guide clinical research and inform intervention design. We outline specific leverage points of the TN approach and barriers to progress. Ten principles of TN are presented to guide and shape the emerging field. We close by articulating ongoing issues facing TN research.

General Scientific Summary

We present an overview of translational neuroscience (TN) as a systematic and theory-driven approach with promise to develop and leverage basic and clinical neuroscientific knowledge to aid the development and optimization of clinical and public health interventions. In this article, we outline specific leverage points of the TN approach and barriers to progress. Finally, 10 principles of TN (with corresponding examples) are presented to guide and shape the emerging field. We close by articulating outstanding issues facing TN research.

Keywords

translational neuroscience; neuroimaging; psychoneuroimmune; physiology

Basic psychological and clinical intervention science have historically operated on somewhat independent tracks. These tracks must meet to address pressing gaps in the field (Allen & Dahl, 2015; Fisher & Berkman, 2015; Roos, Horn, Berkman, Pears, & Fisher, 2018). Researchers have made progress in delineating the basic neurobiological mechanisms underlying mental and physical health outcomes (e.g., depression, obesity), and advances have been made in using basic psychological processes to improve therapeutic outcomes (e.g., cognitive principles with cognitive-behavioral therapy [CBT]). However, despite promising translational efforts to directly alter brain function (e.g., deep brain stimulation and depression; Mayberg et al., 2005), we have yet to witness targeted and readily scalable psychosocial programs that were designed with an advancing understanding of neurobiological mechanisms (Shonkoff et al., 2012). Translational neuroscience (TN) is a systematic research approach for using basic and clinical neuroscientific knowledge to aid the design and optimization of clinical and public health interventions (Fisher & Berkman, 2015). Here, we describe TN as it has been used in developmental and clinical psychology, affective and social neuroscience, and prevention and intervention science, with a focus on demonstrating how TN can inform scalable, noninvasive intervention strategies. Our purpose is to describe the specific ways TN can be useful in advancing this agenda and its current limitations.

In the section that follows, we present two key issues in which TN may be useful in advancing the field, accompanied by examples. We then propose 10 guiding TN principles imperative to future work in the field. Next, we highlight five “levers” or strengths inherent in a TN approach and five “barriers” to be addressed. We conclude with a synopsis and future directions.

TN to Address Variability and Individual Differences

Heterogeneity is a pervasive problem in clinical psychological research. This is true in terms of variation in responses to environmental influences and in the definition and measurement of psychopathology.

Heterogeneity in Response to Environmental Influences

Clinical science has accumulated a strong evidence base about the psychosocial factors that predispose individuals to mental and physical health disorders (Bruce, 2002; Donovan, 2004). For example, early adversity predicts poor outcomes across the lifespan (Albott, Forbes, & Anker, 2018; Felitti et al., 1998; Shonkoff et al., 2012). However, prospective prediction of illness is complicated by significant multifinality (i.e., early adversity predicts a wide range of outcomes; Albott et al., 2018) and heterogeneity (i.e., many children are resilient to the impact of adversity; Masten, 2001). A full account of the biological pathways linking early adversity to later outcomes, as well as individual differences that might predispose individuals toward one outcome versus another, is not yet available.

As such, there has been a significant push to identify biomarkers that signal high risk for pathology (Jaffee, 2018). Valid biomarkers could identify risk before behavioral indicators emerge, particularly in developmental periods (e.g., birth to 3) when behavioral measures are less reliable (Bauer, Wiebe, Carver, Waters, & Nelson, 2003) and significant heterogeneity

exists (e.g., in early presentation of autism spectrum disorder [ASD]; Pelphey, Shultz, Hudac, & Vander Wyk, 2011). Neurobiological processes can help elucidate behavioral phenotypes by providing greater specificity about underlying mechanisms. For example, adversity-exposed children, who often exhibit early signs of oppositional defiant symptomatology, have been observed to have a neural deficit in processing corrective feedback as indexed with electroencephalography (Bruce, McDermott, Fisher, & Fox, 2009; McDermott et al., 2018; Roos, Pears, Bruce, Kim, & Fisher, 2015), and this index has been linked to impulsivity (Roos et al., 2015). One implication is that adversity-exposed children may not be purposely defiant but rather possess a deficiency in neural processes contributing to blunted receptiveness to external feedback, which may underlie important behavioral phenotypes.

Biomarkers are not the only way to identify risk. There are instances when lower-cost, noninvasive methods (e.g., surveys, eye-tracking) perform well on their own. We advocate a transdisciplinary approach to TN wherein the goal is to determine an individual's comprehensive risk profile across domains and in which biomarkers complement behavioral assessments. How well neurobiological markers can enhance behavioral assessments for early detection of risk is an open question.

Heterogeneity in Psychopathology Measurement

There is substantial diagnostic overlap as well as heterogeneity within individual disorder presentations. Heterogeneity in presentation is ubiquitous, with upward of 1,000 unique possible symptom profiles for major depressive disorder (MDD) alone (Fried & Nesse, 2015). Neurobiological phenotypes have also not linked robustly onto clinical diagnostic categories, as indicated by the low replicability of candidate gene studies for MDD (Bosker et al., 2011). Diagnostic heterogeneity hinders our ability to accurately examine neurobiological underpinnings and is an obstacle to successfully translating neuroscientific findings to inform treatment. The National Institute of Mental Health released the Research Domain Criteria (RDoC) project in recognition of this problem (Insel et al., 2010). The goals of the TN approach are complementary to those of RDoC, but we view TN as superseding the RDoC by presenting a means of conducting integrative research rather than organizing knowledge from separate domains.

TN Example: Leveraging Psychoneuroimmunology to Enhance Etiology and Diagnostic Specificity of MDD

Early life adversity is a potent risk factor for the development of MDD (Heim & Nemeroff, 2001; Horesh, Klomek, & Apter, 2008). However, not everyone with a history of early adversity exposure develops MDD, indicating moderators of this relationship (Wingo et al., 2010). Immune functioning is one candidate pathway implicated in both early life adversity (Baumeister, Akhtar, Ciufolini, Pariante, & Mondelli, 2016) and MDD (Dantzer, O'Connor, Freund, Johnson, & Kelley, 2008). In a prospective study, children exposed to early adversity had elevated risk for both depression and high inflammation in adulthood (Danese et al., 2009). In a community sample, levels of inflammation in 9-year-olds predicted both depression and psychosis at age 18, even after adjustment for several covariates (Khandaker, Pearson, Zammit, Lewis, & Jones, 2014).

Tools from psychoneuroimmunology can enhance diagnostic specificity by identifying transdiagnostic symptom clusters. For example, inflammation has been associated with MDD (Howren, Lamkin, & Suls, 2009), albeit somewhat inconsistently (Horn et al., 2018). Researchers have begun to test the notion of an “inflammatory subtype” of depression whereby specific (sub)sets of symptoms are more robustly associated with hyperinflammation than others. Emerging evidence supports that inflammation may be most linked to somatic symptoms of depression, such as fatigue, sleep disturbance, and appetite changes (Bai, Chiou, Su, Li, & Chen, 2014; Jokela, Virtanen, Batty, & Kivimäki, 2016; White, Kivimäki, Jokela, & Batty, 2017). Differentiating the neurobiological underpinnings of symptom profiles in this way can help clarify the etiology and diagnosis of disorders such as MDD.

A next step is to identify transdiagnostic neurobiological systems implicated across adverse outcomes such as those related to threat and reward brain systems (Nelson et al., 2013), anhedonia (Christensen, Bisgaard, & Wiborg, 2011), insomnia (Gehrman et al., 2018), and addiction (Kwako, Bickel, & Goldman, 2018). These are promising targets for research because they might function as intermediary links between clinical disorders and associated propensities and past experiences.

TN for Indices of Treatment Response

Another promising direction is the use of neuroscientific markers to predict treatment response (Fishbein & Dariotis, 2019). For instance, despite significant movement in delineating neurobiological mechanisms associated with MDD (Otte et al., 2016), the majority of patients are still treated first with antidepressants that work only for a minority of individuals, whose underlying mechanisms are poorly understood, and which only slightly outperform placebo (Cipriani et al., 2018). Clinical science is moving past the “one-size-fits-all” mentality toward one that accounts for individual difference moderators of treatment efficacy.

Neuroscientific measures might also afford theory-driven hypotheses about who is likely to respond (i.e., does this neuroscientific index predict who will respond?) and how certain therapeutic strategies are working (i.e., does a neuroscientific index change pre/post treatment?). This information can inform individualized approaches to treatment planning and provide guidance on optimizing treatment designs. Treatment response might be apparent more quickly on neuroscientific indices than on behavioral measures when the former measure processes are closer to the underlying targets of change.

Example: Moderators of Treatment Response With Resting-State Functional Connectivity MRI

Resting-state functional connectivity MRI (rs-fcMRI) is a neuroscientific method with significant potential to predict and detect moderators of treatment response. Rs-fcMRI measures correlations in spontaneous neural activity when participants are not engaged in an explicit task, believed to reflect more trait-like, “intrinsic” brain networks. A systematic review found increased connectivity between frontal and limbic brain regions predicted response to antidepressant treatment, potentially indicating better functioning of inhibitory

control in circuits that process emotions (Dichter, Gibbs, & Smoski, 2015). The clinical translatability of this line of work can be enhanced by going beyond associations with treatment response to identify measures that prospectively predict treatment response. Recent work has used predictive models to take this next step. For example, a single-subject pretreatment functional MRI (fMRI) study identified brain activation that predicted response to CBT treatment response for individuals with generalized anxiety or panic disorder. The neuroimaging measures outperformed clinical or demographic variables in predicting individual patient treatment response (Ball, Stein, Ramsawh, Campbell-Sills, & Paulus, 2014). This approach could predict who is more likely to respond to therapeutic approaches and help to delineate how a treatment might work.

These examples illustrate only some of the ways that the TN approach might inform clinical science. Creative researchers will undoubtedly identify more strategies for deploying the TN approach as the field accelerates. As they do so, a careful analysis of not only the strengths but also the limitations of TN will be critical. A skeptical view is that the attention that TN has received outstrips the advances it has brought to clinical science. The field has yet to develop a systematic approach to articulate how neuroscience tools can tangibly and realistically inform, guide, and be integrated into clinical research, nor rigorously mapped out the barriers to progress. Despite the rapid accumulation of knowledge about the neural alterations associated with mental illness, there is little concrete evidence that this knowledge has facilitated improved treatments.

TN Principles, Leverage Points, and Barriers

We have identified 10 principles of TN as a systematic framework to guide future work in the field toward a resolution of these challenges. These principles are intended to enable TN research to efficiently and realistically inform and improve clinical science interventions. Each principle is paired with an example to illustrate these principles in practice. We describe the principles below in terms of how they pertain to TN specifically; however, we also acknowledge the principles may be more broadly applicable to other domains of psychology.

Alongside the principles and examples, we integrate discussion of five leverage points (“levers”) and five barriers of TN (“barriers”; see Table 1). The leverage points highlight strengths of the TN approach that can fuel progress in basic and clinical sciences and inform public policy. Specifically, neuroscientific knowledge can improve each step of the translational process from identification of neurobiological mechanistic pathways (Lever 1) to treatment development and optimization (Lever 2) to addressing public policy (Lever 3). TN also can help spur new basic neuroscience advances, including establishing a bridge between superficially dissimilar topics (Lever 4) and discriminating between superficially similar processes (Lever 5).

However, there are also significant barriers for the field of TN. Notably, these barriers are potentially germane to many methodologies but are particularly salient for TN. These include the “So what?” (Barrier 1) and “Now what?” (Barrier 2) problems, which highlight the lack of relevant progress in the clinical applicability of TN. These barriers arise when a

novel “biomarker” differentiates between subgroups but is not clearly implicated as a causal factor (“So what?”), or a neurological pathway offers no path to intervention (“Now what?”). Scalability (Barrier 3) and costliness (Barrier 4) are additional significant obstacles for TN to achieve meaningful, real-world impacts. A related barrier is the need for a diverse set of skills required for the entire translational process (Barrier 5). Overcoming these barriers requires a combination of high-quality data and methods; collaboration among scientists across the translational spectrum; changes in the funding model to support rapid-scale, iterative clinical trials; and new ways of training the next generation of translational neuroscientists who will be well versed in clinical science and practice, neurobiology and neuroscientific research tools, and transdisciplinary collaboration.

1. Identify Environmentally Malleable Neurobiological Functions, Circuits, and Systems That Underlie Behavioral Risk and Resilience

Identifying *malleable* neurobiological pathways is critical for preventative and intervention efforts. The relationship between neurobiological and environmental processes is reciprocal and dynamic (Fishbein, 2000). Environmental conditions such as early adversity, poverty, parenting, and neighborhood conditions can influence neurobiological development and functioning across the lifespan. Neurobiological functioning is also malleable and sensitive to environmental influences. Translational neuroscience must identify relevant and malleable environmental, psychosocial, and neurobiological systems spanning across genetic, epigenetic, peripheral (e.g., neuroendocrine, immune) and brain functioning, to inform the development of prevention and intervention efforts.

Example. As discussed previously, the immune system is an environmentally malleable neurobiological system and thus highly germane to TN. Several environmental factors can influence inflammation, such as parenting (Byrne et al., 2017), neighborhood violence (Janusek, Tell, Gaylord-Harden, & Mathews, 2017), early adversity (Miller & Cole, 2012), and physical activity (Ford, 2002), and some are easier to target with intervention than others (see Barrier 2). Moreover, inflammatory mechanisms may be malleable by treatments, such as CBT (Lopresti, 2017), antidepressant medication (Strawbridge et al., 2015), and a family-oriented psychosocial intervention (Miller, Brody, Yu, & Chen, 2014).

Inflammation has also been shown to underlie specific behavioral indices of risk and resilience. For example, inflammatory markers may relate to cognitive vulnerabilities, such as rumination, in adolescents (Moriarty et al., 2018). Rumination is a behavioral risk factor implicated as a target to improve cognitive-behavioral treatments (Watkins, 2009). In a longitudinal study with adolescents, Moriarty et al. (2018) found that the inflammatory cytokine interleukin-6 (IL-6) mediated the association between baseline anxiety symptoms and later-onset depressive symptoms, such that as rumination increased, anxiety symptoms predicted greater inflammation and depressive symptoms. This study illustrates Lever 1 by elucidating how inflammation may be a neurobiological mechanism underlying rumination. This research may help to distinguish between processes that appear similar (i.e., if inflammation is associated with rumination above and beyond other cognitive symptoms; Lever 5).

2. Focus on Central Neurobiological Processes Broadly Implicated in Well-Being and Maladjustment

A multilevel focus on neurobiological systems implicated across well-being and maladjustment is necessary for translating neuroscientific knowledge into clinical applicability. Central nervous system processes play a critical role in psychological functioning and coordinating peripheral processes (e.g., the hypothalamic-pituitary-adrenal [HPA] axis is associated with several disorders; Tsigos & Chrousos, 2002). The dynamic and interactive pathways between peripheral systems (e.g., coregulation of HPA axis and immune system; Kuhlman, Chiang, Horn, & Bower, 2017) are also significant. A comprehensive account requires examination of common transdiagnostic processes at multiple levels, including broad central level (e.g., functional connectivity of neural regions), focal brain region levels (e.g., areas involved in reward processing and sensitivity), at the peripheral biological level (e.g., inflammation), and even at the cellular level (e.g., mitochondria).

Example. Reward processing and sensitivity is an example of a core transdiagnostic process implicated in a range of psychiatric disorders (Nusslock & Alloy, 2017). Reward processing has been implicated as a mechanism underlying important transdiagnostic symptoms, such as anhedonia (Corral-Frías et al., 2015; Treadway & Zald, 2013). Both the activation of reward-related brain structures and their pattern of connectivity with other regions and peripheral systems have been identified as a target for treatments across a broad range of maladaptive behavior and outcomes, including alcohol addiction (Becker et al., 2018), smoking (Cinciripini et al., 2017), depression (Allen et al., 2019), and binge eating (Balodis et al., 2014).

Reduced reward responsivity is linked to dissociable patterns of connectivity between the nucleus accumbens, the default mode network, and the cingulo-opercular network (Sharma et al., 2017) in adults with mood and/or psychotic disorders. This study highlights the principle by examining a central process and its relation to mental health outcomes and Lever 4 by establishing a bridge between superficially dissimilar topics (mood and psychotic-based disorders).

3. Employ Developmentally Sensitive, Neurobiologically Informed Intervention Strategies

Developmental processes are crucial to the onset and maintenance of psychological disorders and to designing effective prevention and intervention programs. Infancy (Kieling et al., 2011) and adolescence (Allen & Dahl, 2015) are particularly critical periods of interest. Yet, developmental research often does not inform intervention strategies. Gold-standard interventions designed for and tested originally with an adult population (e.g., CBT) are often adapted for younger populations (e.g., Cohen, Mannarino, Berliner, & Deblinger, 2000) with less consideration of the relevant developmental processes. Neuroscientific insights can inform developmentally sensitive treatments (Cicchetti & Toth, 2009).

Example. Infancy is a highly sensitive period characterized by dynamic brain growth associated with hormonal changes, synaptic pruning, and metabolic shifts (Gee & Casey,

2015). Attachment patterns form in early life, and these have strong implications for children's health and development (Meins, Bureau, & Fernyhough, 2018). A secure, reciprocal, and warm attachment between child and caregiver during infancy is a moderator of developmental processes and psychological well-being (Mikulincer & Shaver, 2007).

The Attachment and Biobehavioral Catch-up (ABC) intervention illustrates how an intervention can be tailored to a developmental period. ABC was developed for foster infants and infants living with potentially neglecting parents (Dozier, Roben, Caron, Hoyer, & Bernard, 2018). In building the program, the developers of ABC leveraged knowledge from developmental psychology about the sensitive nature of infancy; public policy regarding elevated rates of neglect with foster care and child welfare involved children (Lever 3), attachment theory with respect to the links between neglect and disorganized attachment styles, and neurobiology about the impact of neglect on neuroendocrine dysfunction (Dozier et al., 2018). Several studies have demonstrated the efficacy of the ABC program compared to a control intervention on child and caregiver outcomes (Dozier et al., 2018).

4. Prioritize Those Neurobiological Models That Parsimoniously Add Explanatory Power to Behavioral Theories and Evidence and to Inform Intervention

In addition to knowledge about developmental periods, neuroscientific knowledge can also inform prevention and intervention design (Shonkoff & Fisher, 2013). For example, prolonged and/or disrupted activation of stress response systems impact neurobiological and neural development (Lupien, McEwen, Gunnar, & Heim, 2009). The specificity provided by such studies can sharpen the hypotheses and assessment strategies in testing the hypothesized mechanisms underlying intervention effects (Shonkoff & Fisher, 2013). Neurobiological measures are not without their costs (see Barrier 4), but a clear theoretical rationale can justify their inclusion in intervention trials. The more precision with which an intervention target is described, the more compelling such rationales become.

Example. The Filming Interactions to Nurture Development (FIND) video coaching program is a brief video coaching program developed for parents and caregivers with infants and young children (Fisher, Frenkel, Noll, Berry, & Yockelson, 2016). The FIND intervention draws on knowledge about the neurobiological effects of early adversity, particularly its disruptive effects on the stress, immune, and metabolic systems (Lupien et al., 2009; Maccari, Krugers, Morley-Fletcher, Szyf, & Brunton, 2014). The FIND targets specifically the interactions between caregivers and children when children's brain circuitry is most sensitive to alter emerging language, socioemotional, cognitive, and self-regulatory capacities (Lever 1; Fisher et al., 2016). The FIND program also leverages technology (i.e., video editing) to maximize scalability (Barrier 3); the program is under evaluation worldwide.

5. Use Intervention Trials to Test Theories That Connect Causes to Outcomes via Underlying Neurobiological Mediators

A TN approach draws on neuroscientific knowledge to examine neurobiological mediators of intervention outcomes. Researchers can advance the knowledge of neurobiological pathways implicated in health outcomes and begin to delineate how certain treatments work

and *for whom* they are most effective (Lever 2). This requires pinpointing the neural and/or biological targets that correspond to the mechanisms of the intervention action. Neurobiological moderators may indicate relevant individual differences that enable the development of increasingly refined hypotheses, specify a nuanced understanding of how and for whom an intervention is working, and can help increase intervention efficacy (Fisher & Berkman, 2015).

Example. Multidimensional Treatment Foster Care for Preschoolers (MTFC-P) is an intervention for preschool-aged foster care children with a history of maltreatment (Fisher, Burraston, & Pears, 2005; Fisher, Ellis, & Chamberlain, 1999). The theory behind MTFC-P is that neglect exerted its effects via chronic stress and HPA axis dysfunction as indicated by a blunted diurnal cortisol slope. A mechanistic efficacy trial tested not only the intervention's effects on outcomes but also whether it would normalize children's diurnal cortisol slopes (Fisher, Stoolmiller, Gunnar, & Burraston, 2007). Foster children in the MTFC-P group showed more typical diurnal cortisol slopes comparable to a community control group (Fisher et al., 2007), while foster care children receiving routine services demonstrated a blunted morning-to-evening cortisol slope. Additionally, foster parents in the routine services group but not the other groups had elevated levels of caregiver stress that were associated with atypical cortisol activity in children (Fisher & Stoolmiller, 2008). This line of work illustrates how TN can test not only the efficacy of an intervention but also its hypothesized neurobiological mechanisms (Lever 2).

6. Include Rigorous Measurement of Potentially Moderating Contextual Factors (e.g., Poverty, Neighborhood, Culture) That Confer Risk or Resiliency

Contextual environmental factors are related to maladjustment and well-being and also to neurobiological underpinnings of health and resilience (Rutten et al., 2013). A comprehensive understanding of risk requires uncovering the interactive and additive impacts of contextual factors in relation to neurobiological pathways and psychosocial development. For example, race and ethnicity are often regarded as “nuisance” variables or obstacles garnering little attention beyond their perfunctory role in statistical control and analysis (Bernal, Cumba-Avilés, & Rodríguez-Quintana, 2014; Hall, Yip, & Zárate, 2016). However, there is a rapidly growing field that has examined how challenges unique to ethnic minority groups (e.g., racial discrimination) not only confer risk for poor health outcomes but are clearly linked to stress neurobiology, including impacts on neural and hormonal pathways (Berger & Sarnyai, 2015).

Example. Contextual knowledge about ethnic differences in sleep quality and immune functioning informed a recent study testing the idea that sleep disturbances in the postpartum period might confer risk for mood symptoms in African American and White women (Christian, Kowalsky, Mitchell, & Porter, 2018). The study showed that poor sleep and stress predicted greater stimulated cytokine production, but only in African American women. A thorough understanding of mechanisms underlying racial differences in immunological disruption and depression will be essential to delineating the etiology of depression and informing novel treatment strategies.

7. Apply Precision Interventions That Are Tailored to Individuals Based on Biobehavioral Characteristics and Grounded in Robust Neuroscience

A long-term aim of clinical science is to provide precision treatment by integrating person-level data to inform the selection of treatment (Prendes-Alvarez & Nemeroff, 2018). A single biomarker is unlikely to predict a diagnosis or treatment, but a panel of independent biomarkers, including genetic and epigenetic factors and biomarkers of neuroendocrine, immune, and metabolic function, might have greater predictive validity. Along with clinical presentation and environmental factors (e.g., culture; Matheson, Bombay, & Anisman, 2018), a set of valid, reliable biomarkers might help optimize existing treatments, design new interventions for treatment-resistant or complex presentations, and predict who is most likely to response to an intervention. A challenge lies in establishing reliable and valid individual difference measures of intervention efficacy, requiring substantial measurement expertise (Barrier 5). There is scant empirical attention to the basic psychometric features, such as reliability and validity of neurobiological and neural measures (DuBois & Adolphs, 2016), and reliability is often found to be low when examined, such as with fMRI (Bennett & Miller, 2010; Herting, Gautam, Chen, Mezher, & Vetter, 2018). The scarcity of psychometric data for many measures is an outstanding obstacle for TN, which cannot be useful without reliable measures.

Example. The drug sirukumab modulates the immune response by acting against IL-6 and reduces depressive symptoms in patients with rheumatoid arthritis (Sun et al., 2017) and is a candidate adjunctive treatment for MDD. An RCT should be changed to randomized controlled trial is under way with depressed patients on monoaminergic antidepressants testing a precise hypothesis grounded in neuroscientific theory. The model predicts that the medication would work only for depressed individuals with elevated inflammation; therefore, only participants with elevated systemic inflammation are eligible (NCT02473289; Lever 2). Translating neuroscientific knowledge about inflammation and depression to test a new treatment is also an example of overcoming the “So what?” and “Now what?” barriers (Barriers 1 and 2).

8. Emphasize Applications of Neuroscience to Interpersonal Processes Rather Than Solely Individual Processes

Interpersonal processes are intrinsic to nearly all health outcomes. Social neuroscience research uncovering the ways that “social emotion regulation” is different from other forms of self-regulation can indicate ways to facilitate emotion regulation in relationships (Levers 4 and 5; Reeck, Ames, & Ochsner, 2016). Coregulation is central to a range of close relationships (e.g., parenting, romantic relationships). For example, parenting quality is related to children’s emotion regulation (e.g., Chang, Schwartz, Dodge, & McBride-Chang, 2003), and the family environment is a key contextual factor (see Principle 6) in the development of emotion regulation capabilities (Morris, Silk, Steinberg, Myers, & Robinson, 2007). Further, the neurobiology of caregiving is an essential component of child development and outcomes (Abraham & Feldman, 2018). Future directions are to understand the neurobiological underpinnings of these interpersonal processes and examine coregulation of neurobiological processes in dyads (e.g., parent–child, couples).

Example. Parental executive function (EF) is critical to children's EF development (Cipriano-Essel, Skowron, Stifter, & Teti, 2013). For instance, a recent study found that maternal working memory capacity is positively related to preschool-aged children's inhibitory control, suggesting that maternal influences play a significant role in the development of children's EF in high-risk families (Kim, Shimomada, Giuliano, & Skowron, 2017). This study illustrates Principle 6 by accounting for the relation of EF to socioeconomic diversity, an important contextual factor, and Lever 4 by establishing a bridge between processes that may appear unrelated (i.e., maternal working memory and children's inhibitory control).

9. Elucidate Linkages Between Mental Health and Physical Systems (e.g., Autonomic, Neuroendocrine, Immune, Metabolic, Gut Microbiome)

The origins of many adulthood physical and psychological disorders have roots in early development and neurobiological disruptions (Shonkoff, Boyce, & McEwen, 2009). Cumulative damage (i.e., allostatic load; Juster, McEwen, & Lupien, 2010) and biological embedding of adversity within sensitive windows (Miller, Chen, & Parker, 2011) can influence health trajectories. Children exposed to early adversity are vulnerable to mental and physical health illnesses (Johnson & Schoeni, 2011), suggesting the need to examine mechanistic pathways that confer risk for a range of health outcomes (Nusslock & Miller, 2016). TN can examine the shared neurobiological processes across physical and mental health outcomes as a way to increase scientific understanding of the etiology and developmental trajectory of pathology. This approach also helps to connect unknown disease-relevant processes (e.g., asthma and psychological health; Goldbeck, Koffmane, Lecheler, Thiessen, & Fegert, 2007) by emphasizing common neurobiological pathways (e.g., immune functioning; Chen, Fisher, Bacharier, & Strunk, 2003).

Example. Physical and mental health share several mechanistic pathways (Scott et al., 2016). For example, there is a high prevalence of gastrointestinal (GI) disorders and symptoms observed in children with ASD (McElhanon, McCracken, Karpen, & Sharp, 2014), with increasing attention being paid to potential shared neurobiological underpinnings (e.g., immunometabolic pathways, the gut microbiome). A novel study contrasted children with ASD with and without GI symptoms with typically developing children with and without GI symptoms. Findings indicated that children with ASD and GI symptomatology produced increased levels of mucosa-relevant inflammatory cytokines to an inflammatory stimulation challenge in contrast to children with ASD but no GI symptoms. Further, the researchers found family-level differences in gut microbiome composition between ASD and typically developing children with GI symptoms (Rose et al., 2018). Such results have several implications, such as addressing heterogeneity in ASD symptom profiles (e.g., why not all children with ASD have comorbid GI problems) and advancing new treatment possibilities. This example also highlights Lever 4 by drawing a connection between unknown disease-relevant processes (e.g., GI and ASD symptomatology).

10. Select Neurobiological Methodologies and Measures Based on Their Suitability to the Hypotheses Being Tested

Translational neuroscience encourages a thoughtful selection of methods with respect to the focal hypothesis. Simply put, the theoretical models should drive the selection of the measurement tools. Assessing the theoretical model beforehand enables selecting valid, reliable measures of relevant systems that are sensitive to change and can detect individual differences (Lever 2). Especially careful consideration is warranted within certain methodologies, such as neuroimaging (Barriers 3 and 4; Pfeifer, Allen, Byrne, & Mills, 2018).

Example. A study of neural responses to peer emotional expressions in adolescent girls leveraged prior knowledge to make reasoned, a priori decisions about the optimal neuroimaging paradigm (Flannery, Giuliani, Flournoy, & Pfeifer, 2017). The study tested a neurodevelopmental model stating that adolescent risk-taking results from an imbalance in the maturation of networks related to self-regulation versus affect and reward. Prior research relied on static, adult face stimuli, which are particularly inappropriate given the increased salience of peers during adolescence. So, in this case, the researchers enhanced the relevance of the stimuli by using videos of children and adolescents making a range of emotional expressions. Doing so revealed very different age-based trajectories than those predicted by previous models, suggesting possible salience effects in previous work (Flannery et al., 2017). Theory-driven decision making can also combat problems associated with costliness (Barrier 4) and participant burden (Barrier 3).

Conclusions

We highlighted areas where progress in TN has been made and where growth is needed. We see several directions for TN in the coming years. Elucidating the neurobiological mechanisms of psychological processes with ever-increasing precision (Lever 1) will be critical, whether they be individual (e.g., rumination) or interpersonal processes (e.g., dyadic coregulation and EF). A larger task, which requires additional high-quality evidence, is to identify connections between seemingly dissimilar topics as well as establishing dissociations between processes that are believed to be similar (Levers 4 and 5). A downstream goal is to then apply that accumulated knowledge to prevention and intervention evaluation (Lever 2). However, as highlighted in Barrier 1, this has yet to widely occur. Interventions such as ABC and FIND are promising examples of overcoming this barrier.

The TN approach also embraces transdiagnostic constructs broadly implicated in health. Delineating the neurobiological mechanisms of symptomatology spanning physical and mental health is critical to advancing scientific knowledge of the etiology and trajectory of disease. The field can formulate integrated theoretical models crossing behavioral, psychological, and neurobiological levels only once this knowledge is obtained and then test the models in translational interventions (Lever 2). The existential questions presented in Barriers 1 and 2 (“So what?” and “Now what?”) can be answered only when the field can design, evaluate, and implement psychosocial interventions that precisely alter neurobiological function and demonstrate that their effects are mediated by the specified neurobiological mechanisms. That is admittedly a lofty goal, and significant challenges

persist (Barriers 3–5). Ultimately, TN is a collaborative, transdisciplinary field whose scope is deliberately ambitious to be commensurate with the magnitude of the goal of transforming the scientific approach to alleviating the burden of mental illness.

References

- Abraham E, & Feldman R (2018). The neurobiology of human allomaternal care; implications for fathering, coparenting, and children's social development. *Physiology & Behavior*, 193, 25–34. 10.1016/j.physbeh.2017.12.034 [PubMed: 29730038]
- Albott CS, Forbes MK, & Anker JJ (2018). Association of childhood adversity with differential susceptibility of transdiagnostic psychopathology to environmental stress in adulthood. *Journal of the American Medical Association Network Open*, 1, e185354 10.1001/jamanetworkopen.2018.5354
- Allen NB, & Dahl RE (2015). Multi-level models of internalizing disorders and translational developmental science: Seeking etiological insights that can inform early intervention strategies. *Journal of Abnormal Child Psychology*, 43, 875–883. 10.1007/s10802-015-0024-9 [PubMed: 25947071]
- Allen TA, Lam RW, Milev R, Rizvi SJ, Frey BN, MacQueen GM, ... Quilty LC (2019). Early change in reward and punishment sensitivity as a predictor of response to antidepressant treatment for major depressive disorder: A CAN-BIND-1 report. *Psychological Medicine*, 49, 1629–1638. [PubMed: 30220263]
- Bai Y-M, Chiou WF, Su TP, Li CT, & Chen MH (2014). Pro-inflammatory cytokine associated with somatic and pain symptoms in depression. *Journal of Affective Disorders*, 155, 28–34. 10.1016/j.jad.2013.10.019 [PubMed: 24176538]
- Ball TM, Stein MB, Ramsawh HJ, Campbell-Sills L, & Paulus MP (2014). Single-subject anxiety treatment outcome prediction using functional neuroimaging. *Neuropsychopharmacology*, 39, 1254–1261 [PubMed: 24270731]
- Balodis IM, Grilo CM, Kober H, Worhunsky PD, White MA, Stevens MC, ... Potenza MN (2014). A pilot study linking reduced fronto-striatal recruitment during reward processing to persistent bingeing following treatment for binge-eating disorder. *International Journal of Eating Disorders*, 47, 376–384. 10.1002/eat.22204 [PubMed: 24729034]
- Bauer PJ, Wiebe SA, Carver LJ, Waters JM, & Nelson CA (2003). Developments in long-term explicit memory late in the first year of life: Behavioral and electrophysiological indices. *Psychological Science*, 14, 629–635. 10.1046/j.0956-7976.2003.psci_1476.x [PubMed: 14629697]
- Baumeister D, Akhtar R, Ciufofini S, Pariante CM, & Mondelli V (2016). Childhood trauma and adulthood inflammation: A meta-analysis of peripheral C-reactive protein, interleukin-6 and tumour necrosis factor- α . *Molecular Psychiatry*, 21, 642–649. 10.1038/mp.2015.67 [PubMed: 26033244]
- Becker A, Gerchen MF, Kirsch M, Hoffmann S, Kiefer F, & Kirsch P (2018). Striatal reward sensitivity predicts therapy-related neural changes in alcohol addiction. *European Archives of Psychiatry and Clinical Neuroscience*, 268, 231–242. 10.1007/s00406-017-0805-y [PubMed: 28488021]
- Bemal G, Cumba-Avilés E, & Rodriguez-Quintana N (2014). *APA handbook of multicultural psychology: Vol. 1. Methodological challenges in research with ethnic, racial, and ethnocultural groups*. Washington, DC: American Psychological Association.
- Bennett CM, & Miller MB (2010). How reliable are the results from functional magnetic resonance imaging? *Annals of the New York Academy of Sciences*, 1191, 133–155. 10.1111/j.1749-6632.2010.05446.x [PubMed: 20392279]
- Berger M, & Sarnyai Z (2015). "More than skin deep": Stress neurobiology and mental health consequences of racial discrimination. *Stress: The International Journal on the Biology of Stress*, 18, 1–10. 10.3109/10253890.2014.989204
- Bosker FJ, Hartman CA, Nolte IM, Prins BP, Terpstra P, Posthuma D, ... Nolen WA (2011). Poor replication of candidate genes for major depressive disorder using genome-wide association data. *Molecular Psychiatry*, 16, 516–532. 10.1038/mp.2010.38 [PubMed: 20351714]

- Bruce J, McDermott JM, Fisher PA, & Fox NA (2009). Using behavioral and electrophysiological measures to assess the effects of a preventive intervention: A preliminary study with preschool-aged foster children. *Prevention Science*, 10, 129–140. 10.1007/s11121-008-0115-8 [PubMed: 19030992]
- Bruce ML (2002). Psychosocial risk factors for depressive disorders in late life. *Biological Psychiatry*, 52, 175–184. 10.1016/S0006-3223(02)01410-5 [PubMed: 12182924]
- Byrne ML, Badcock PB, Simmons JG, Whittle S, Pettitt A, Olsson CA, ... Allen NB (2017). Self-reported parenting style is associated with children's inflammation and immune activation. *Journal of Family Psychology*, 31, 374–380. 10.1037/fam0000254 [PubMed: 27819440]
- Chang L, Schwartz D, Dodge KA, & McBride-Chang C (2003). Harsh parenting in relation to child emotion regulation and aggression. *Journal of Family Psychology*, 17, 598–606. 10.1037/0893-3200.17.4.598 [PubMed: 14640808]
- Chen E, Fisher EB, Bacharier LB, & Strunk RC (2003). Socioeconomic status, stress, and immune markers in adolescents with asthma. *Psychosomatic Medicine*, 65, 984–992. 10.1097/01.PSY.0000097340.54195.3C [PubMed: 14645776]
- Christensen T, Bisgaard CF, & Wiborg O (2011). Biomarkers of anhedonic-like behavior, antidepressant drug refractoriness, and stress resilience in a rat model of depression. *Neuroscience*, 196, 66–79. 10.1016/j.neuroscience.2011.08.024 [PubMed: 21889970]
- Christian LM, Kowalsky JM, Mitchell AM, & Porter K (2018). Associations of postpartum sleep, stress, and depressive symptoms with LPS-stimulated cytokine production among African American and White women. *Journal of Neuroimmunology*, 316, 98–106. 10.1016/j.jneuroim.2017.12.020 [PubMed: 29406850]
- Cicchetti D, & Toth SL (2009). The past achievements and future promises of developmental psychopathology: The coming of age of a discipline. *Journal of Child Psychology and Psychiatry*, 50, 16–25. 10.1111/j.1469-7610.2008.01979.x [PubMed: 19175810]
- Cinciripini PM, Green CE, Robinson JD, Karam-Hage M, Engelman JM, Minnix JA, ... Versace F (2017). Benefits of varenicline vs. bupropion for smoking cessation: A Bayesian analysis of the interaction of reward sensitivity and treatment. *Psychopharmacology*, 234, 1769–1779. 10.1007/s00213-017-4580-2 [PubMed: 28275830]
- Cipriani A, Furukawa TA, Salanti G, Chaimani A, Atkinson LZ, Ogawa Y, ... Geddes JR (2018). Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: A systematic review and network meta-analysis. *The Lancet*, 391, 1357–1366. 10.1016/S0140-6736(17)32802-7
- Cipriano-Essel E, Skowron EA, Stifter CA, & Teti DM (2013). Heterogeneity in maltreated and non-maltreated preschool children's inhibitory control: The interplay between parenting quality and child temperament. *Infant and Child Development*, 22, 501–522. 10.1002/icd.1801 [PubMed: 24729743]
- Cohen JA, Mannarino AP, Berliner L, & Deblinger E (2000). Trauma-focused cognitive behavioral therapy for children and adolescents: An empirical update. *Journal of Interpersonal Violence*, 15, 1202–1223. 10.1177/088626000015011007
- Corral-Frías NS, Nikolova YS, Michalski LJ, Baranger DA, Hariri AR, & Bogdan R (2015). Stress-related anhedonia is associated with ventral striatum reactivity to reward and transdiagnostic psychiatric symptomatology. *Psychological Medicine*, 45, 2605–2617. 10.1017/S0033291715000525 [PubMed: 25853627]
- Danese A, Moffitt TE, Harrington H, Milne BJ, Polanczyk G, Pariante CM, ... Caspi A (2009). Adverse childhood experiences and adult risk factors for age-related disease: Depression, inflammation, and clustering of metabolic risk markers. *Archives of Pediatrics & Adolescent Medicine*, 163, 1135–1143. 10.1001/archpediatrics.2009.214 [PubMed: 19996051]
- Dantzer R, O'Connor JC, Freund GG, Johnson RW, & Kelley KW (2008). From inflammation to sickness and depression: When the immune system subjugates the brain. *Nature Reviews Neuroscience*, 9, 46–56. 10.1038/nrn2297 [PubMed: 18073775]
- Dichter GS, Gibbs D, & Smoski MJ (2015). A systematic review of relations between resting-state functional-MRI and treatment response in major depressive disorder. *Journal of Affective Disorders*, 172, 8–17. 10.1016/j.jad.2014.09.028 [PubMed: 25451389]

- Donovan JE (2004). Adolescent alcohol initiation: A review of psychosocial risk factors. *Journal of Adolescent Health*, 35, 529.e7–529.e18.
- Dozier M, Roben CKP, Caron E, Hoyer J, & Bernard K (2018). Attachment and biobehavioral catch-up: An evidence-based intervention for vulnerable infants and their families. *Psychotherapy Research*, 28, 18–29. 10.1080/10503307.2016.1229873 [PubMed: 27729003]
- Dubois J, & Adolphs R (2016). Building a science of individual differences from fMRI. *Trends in Cognitive Sciences*, 20, 425–443. 10.1016/j.tics.2016.03.014 [PubMed: 27138646]
- Felitti VJ, Anda RF, Nordenberg D, Williamson DF, Spitz AM, Edwards V, ... Marks JS (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, 14, 245–258. 10.1016/S0749-3797(98)00017-8 [PubMed: 9635069]
- Fishbein D (2000). The importance of neurobiological research to the prevention of psychopathology. *Prevention Science*, 1, 89–106. 10.1023/A:1010090114858 [PubMed: 11521962]
- Fishbein DH, & Dariotis JK (2019). Personalizing and optimizing preventive intervention models via a translational neuroscience framework. *Prevention Science*, 20, 10–20. 10.1007/s1121-017-0851-8 [PubMed: 29101644]
- Fisher PA, & Berkman ET (2015). Designing interventions informed by scientific knowledge about effects of early adversity: A translational neuroscience agenda for next generation addictions research. *Current Addiction Reports*, 2, 347–353. 10.1007/s40429-015-0071-x [PubMed: 26985399]
- Fisher PA, Burraston B, & Pears K (2005). The early intervention foster care program: Permanent placement outcomes from a randomized trial. *Child Maltreatment*, 10, 61–71. 10.1177/1077559504271561 [PubMed: 15611327]
- Fisher PA, Ellis BH, & Chamberlain P (1999). Early intervention foster care: A model for preventing risk in young children who have been maltreated. *Children's Services: Social Policy, Research, & Practice*, 2, 159–182. 10.1207/s15326918cs0203_3
- Fisher PA, Frenkel TI, Noll LK, Berry M, & Yockelson M (2016). Promoting healthy child development via a two-generation translational neuroscience framework: The Filming Interactions to Nurture Development video coaching program. *Child Development Perspectives*, 10, 251–256. 10.1111/cdep.12195 [PubMed: 28936231]
- Fisher PA, & Stoolmiller M (2008). Intervention effects on foster parent stress: Associations with child cortisol levels. *Development and Psychopathology*, 20, 1003–1021. 10.1017/S0954579408000473 [PubMed: 18606041]
- Fisher PA, Stoolmiller M, Gunnar MR, & Burraston BO (2007). Effects of a therapeutic intervention for foster preschoolers on diurnal cortisol activity. *Psychoneuroendocrinology*, 32, 892–905. 10.1016/j.psyneuen.2007.06.008 [PubMed: 17656028]
- Flannery JE, Giuliani NR, Flournoy JC, & Pfeifer JH (2017). Neurodevelopmental changes across adolescence in viewing and labeling dynamic peer emotions. *Developmental Cognitive Neuroscience*, 25, 113–127. 10.1016/j.dcn.2017.02.003 [PubMed: 28262423]
- Ford ES (2002). Does exercise reduce inflammation? Physical activity and C-reactive protein among U.S. adults. *Epidemiology*, 13, 561–568. 10.1097/00001648-200209000-00012 [PubMed: 12192226]
- Fried EI, & Nesse RM (2015). Depression is not a consistent syndrome: An investigation of unique symptom patterns in the STAR*D study. *Journal of Affective Disorders*, 172, 96–102. 10.1016/j.jad.2014.10.010 [PubMed: 25451401]
- Gee DG, & Casey BJ (2015). The impact of developmental timing for stress and recovery. *Neurobiology of Stress*, 1, 184–194. 10.1016/j.ynstr.2015.02.001 [PubMed: 25798454]
- Gehrman P, Sengupta A, Harders E, Ubeydullah E, Pack AI, & Weljie A (2018). Altered diurnal states in insomnia reflect peripheral hyperarousal and metabolic desynchrony: A preliminary study. *Sleep: Journal of Sleep and Sleep Disorders Research*, 41, zsy043 10.1093/sleep/zsy043
- Goldbeck L, Koffmane K, Lecheler J, Thiessen K, & Fegert JM (2007). Disease severity, mental health, and quality of life of children and adolescents with asthma. *Pediatric Pulmonology*, 42, 15–22. 10.1002/ppul.20509 [PubMed: 17106900]

- Hall GCN, Yip T, & Zárate MA (2016). On becoming multicultural in a monocultural research world: A conceptual approach to studying ethnocultural diversity. *American Psychologist*, 71, 40–51. 10.1037/a0039734 [PubMed: 26766764]
- Heim C, & Nemeroff CB (2001). The role of childhood trauma in the neurobiology of mood and anxiety disorders: Preclinical and clinical studies. *Biological Psychiatry*, 49, 1023–1039. 10.1016/S0006-3223(01)01157-X [PubMed: 11430844]
- Herting MM, Gautam P, Chen Z, Mezher A, & Vetter NC (2018). Test-retest reliability of longitudinal task-based fMRI: Implications for developmental studies. *Developmental Cognitive Neuroscience*, 33, 17–26. 10.1016/j.dcn.2017.07.001 [PubMed: 29158072]
- Horesh N, Klomek AB, & Apter A (2008). Stressful life events and major depressive disorders. *Psychiatry Research*, 160, 192–199. 10.1016/j.psychres.2007.06.008 [PubMed: 18582951]
- Horn SR, Long MM, Nelson BW, Allen NB, Fisher PA, & Byrne ML (2018). Replication and reproducibility issues in the relationship between C-reactive protein and depression: A systematic review and focused meta-analysis. *Brain, Behavior, and Immunity*, 73, 85–114. 10.1016/j.bbi.2018.06.016
- Howren MB, Lamkin DM, & Suls J (2009). Associations of depression with C-reactive protein, IL-1, and IL-6: A meta-analysis. *Psychosomatic Medicine*, 71, 171–186. 10.1097/PSY.0b013e3181907c1b [PubMed: 19188531]
- Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, ... Wang P (2010). Research domain criteria (RDoC): Toward a new classification framework for research on mental disorders. *American Journal of Psychiatry*, 167, 748–751. 10.1176/appi.ajp.2010.09091379 [PubMed: 20595427]
- Jaffee S (2018). Promises and pitfalls in the development of biomarkers that can promote early intervention in children at risk. *Journal of Child Psychology and Psychiatry*, 59, 97–98. 10.1111/jcpp.12869 [PubMed: 29349808]
- Janusek LW, Tell D, Gaylord-Harden N, & Mathews HL (2017). Relationship of childhood adversity and neighborhood violence to a proinflammatory phenotype in emerging adult African American men: An epigenetic link. *Brain, Behavior, and Immunity*, 60, 126–135. 10.1016/j.bbi.2016.10.006
- Johnson RC, & Schoeni RF (2011). Early-life origins of adult disease: National longitudinal population-based study of the United States. *American Journal of Public Health*, 101, 2317–2324. 10.2105/AJPH.2011.300252 [PubMed: 22021306]
- Jokela M, Virtanen M, Batty GD, & Kivimäki M (2016). Inflammation and specific symptoms of depression. *Journal of the American Medical Association Psychiatry*, 73, 87–88. 10.1001/jamapsychiatry.2015.1977 [PubMed: 26579988]
- Juster R-P, McEwen BS, & Lupien SJ (2010). Allostatic load biomarkers of chronic stress and impact on health and cognition. *Neuroscience and Biobehavioral Reviews*, 35, 2–16. 10.1016/j.neubiorev.2009.10.002 [PubMed: 19822172]
- Khandaker GM, Pearson RM, Zammit S, Lewis G, & Jones PB (2014). Association of serum interleukin 6 and C-reactive protein in childhood with depression and psychosis in young adult life: A population-based longitudinal study. *Journal of the American Medical Association Psychiatry*, 71, 1121–1128. 10.1001/jamapsychiatry.2014.1332 [PubMed: 25133871]
- Kieling C, Baker-Henningham H, Belfer M, Conti G, Ertem I, Omigbodun O, ... Rahman A (2011). Child and adolescent mental health worldwide: Evidence for action. *The Lancet*, 378, 1515–1525. 10.1016/S0140-6736(11)60827-1
- Kim MH, Shimomaeda L, Giuliano RJ, & Skowron EA (2017). Intergenerational associations in executive function between mothers and children in the context of risk. *Journal of Experimental Child Psychology*, 164, 1–15. 10.1016/j.jecp.2017.07.002 [PubMed: 28759782]
- Kuhlman KR, Chiang JJ, Horn S, & Bower JE (2017). Developmental psychoneuroendocrine and psychoneuroimmune pathways from childhood adversity to disease. *Neuroscience and Biobehavioral Reviews*, 80, 166–184. 10.1016/j.neubiorev.2017.05.020 [PubMed: 28577879]
- Kwako LE, Bickel WK, & Goldman D (2018). Addiction biomarkers: Dimensional approaches to understanding addiction. *Trends in Molecular Medicine*, 24, 121–128. 10.1016/j.molmed.2017.12.007 [PubMed: 29307501]

- Lopresti AL (2017). Cognitive behaviour therapy and inflammation: A systematic review of its relationship and the potential implications for the treatment of depression. *Australian and New Zealand Journal of Psychiatry*, 51, 565–582. 10.1177/0004867417701996 [PubMed: 28382827]
- Lupien SJ, McEwen BS, Gunnar MR, & Heim C (2009). Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature Reviews Neuroscience*, 10, 434–445. 10.1038/nrn2639 [PubMed: 19401723]
- Maccari S, Krugers HJ, Morley-Fletcher S, Szyf M, & Brunton PJ (2014). The consequences of early-life adversity: Neurobiological, behavioural and epigenetic adaptations. *Journal of Neuroendocrinology*, 26, 707–723. 10.1111/jne.12175 [PubMed: 25039443]
- Masten AS (2001). Ordinary magic: Resilience processes in development. *American Psychologist*, 56, 227–238. 10.1037/0003-066X.56.3.227 [PubMed: 11315249]
- Matheson K, Bombay A, & Anisman H (2018). Culture as an ingredient of personalized medicine. *Journal of Psychiatry & Neuroscience*, 43, 3–6. 10.1503/jpn.170234 [PubMed: 29252161]
- Mayberg HS, Lozano AM, Voon V, McNeely HE, Seminowicz D, Hamani C, ... Kennedy SH (2005). Deep brain stimulation for treatment-resistant depression. *Neuron*, 45, 651–660. 10.1016/j.neuron.2005.02.014 [PubMed: 15748841]
- McDermott JM, Pears KC, Bruce J, Kim HK, Roos L, Yoerger KL, & Fisher PA (2018). Improving kindergarten readiness in children with developmental disabilities: Changes in neural correlates of response monitoring. *Applied Neuropsychology: Child*, 7, 187–199. 10.1080/21622965.2017.1286239 [PubMed: 28631968]
- McElhanon BO, McCracken C, Karpen S, & Sharp WG (2014). Gastrointestinal symptoms in autism spectrum disorder: A meta-analysis. *Pediatrics*, 133, 872–883. 10.1542/peds.2013-3995 [PubMed: 24777214]
- Meins E, Bureau JF, & Fernyhough C (2018). Mother–child attachment from infancy to the preschool years: Predicting security and stability. *Child Development*, 89, 1022–1038. 10.1111/cdev.12778 [PubMed: 28294294]
- Mikulincer M, & Shaver PR (2007). Boosting attachment security to promote mental health, prosocial values, and inter-group tolerance. *Psychological Inquiry*, 18, 139–156. 10.1080/10478400701512646
- Miller GE, Brody GH, Yu T, & Chen E (2014). A family-oriented psychosocial intervention reduces inflammation in low-SES African American youth. *Proceedings of the National Academy of Sciences of the United States of America*, 111, 11287–11292. 10.1073/pnas.1406578111 [PubMed: 25049403]
- Miller GE, Chen E, & Parker KJ (2011). Psychological stress in childhood and susceptibility to the chronic diseases of aging: Moving toward a model of behavioral and biological mechanisms. *Psychological Bulletin*, 137, 959–997. 10.1037/a0024768 [PubMed: 21787044]
- Miller GE, & Cole SW (2012). Clustering of depression and inflammation in adolescents previously exposed to childhood adversity. *Biological Psychiatry*, 72, 34–40. 10.1016/j.biopsych.2012.02.034 [PubMed: 22494534]
- Moriarty DP, McArthur BA, Ellman LM, Coe CL, Abramson LY, & Alloy LB (2018). Immunocognitive model of depression secondary to anxiety in adolescents. *Journal of Youth and Adolescence*, 47, 2625–2636. 10.1007/s10964-018-0905-7 [PubMed: 30088131]
- Morris AS, Silk JS, Steinberg L, Myers SS, & Robinson LR (2007). The role of the family context in the development of emotion regulation. *Social Development*, 16, 361–388. 10.1111/j.1467-9507.2007.00389.x [PubMed: 19756175]
- Nelson BD, McGowan SK, Sarapas C, Robison-Andrew EJ, Altman SE, Campbell ML, ... Shankman SA (2013). Biomarkers of threat and reward sensitivity demonstrate unique associations with risk for psychopathology. *Journal of Abnormal Psychology*, 122, 662–671. 10.1037/a0033982 [PubMed: 24016008]
- Nusslock R, & Alloy LB (2017). Reward processing and mood-related symptoms: An RDoC and translational neuroscience perspective. *Journal of Affective Disorders*, 216, 3–16. 10.1016/j.jad.2017.02.001 [PubMed: 28237133]

- Nusslock R, & Miller GE (2016). Early-life adversity and physical and emotional health across the lifespan: A neuroimmune network hypothesis. *Biological Psychiatry*, 80, 23–32. 10.1016/j.biopsych.2015.05.017 [PubMed: 26166230]
- Otte C, Gold SM, Penninx BW, Pariante CM, Etkin A, Fava M, ... Schatzberg AF (2016). Major depressive disorder. *Nature Reviews Disease Primers*, 2, 16065 10.1038/nrdp.2016.65
- Pelphrey KA, Shultz S, Hudac CM, & Vander Wyk BC (2011). Research review: Constraining heterogeneity: The social brain and its development in autism spectrum disorder. *Journal of Child Psychology and Psychiatry*, 52, 631–644. 10.1111/j.1469-7610.2010.02349.x [PubMed: 21244421]
- Pfeifer JH, Allen NB, Byrne ML, & Mills KL (2018). Modeling developmental change: Contemporary approaches to key methodological challenges in developmental neuroimaging. *Developmental Cognitive Neuroscience*, 33, 1–4. 10.1016/j.dcn.2018.10.001 [PubMed: 30384956]
- Prendes-Alvarez S, & Nemeroff CB (2018). Personalized medicine: Prediction of disease vulnerability in mood disorders. *Neuroscience Letters*, 669, 10–13. 10.1016/j.neulet.2016.09.049 [PubMed: 27746310]
- Reeck C, Ames DR, & Ochsner KN (2016). The social regulation of emotion: An integrative, cross-disciplinary model. *Trends in Cognitive Sciences*, 20, 47–63. 10.1016/j.tics.2015.09.003 [PubMed: 26564248]
- Roos LE, Horn S, Berkman ET, Pears K, & Fisher PA (2018). Leveraging translational neuroscience to inform early intervention and addiction prevention for children exposed to early life stress. *Neurobiology of Stress*, 9, 231–240. 10.1016/j.ynstr.2018.10.004 [PubMed: 30450387]
- Roos LE, Pears K, Bruce J, Kim HK, & Fisher PA (2015). Impulsivity and the association between the feedback-related negativity and performance on an inhibitory control task in young at-risk children. *Psychophysiology*, 52, 704–713. 10.1111/psyp.12389 [PubMed: 25424570]
- Rose DR, Yang H, Serena G, Sturgeon C, Ma B, Careaga M, ... Ashwood P (2018). Differential immune responses and microbiota profiles in children with autism spectrum disorders and co-morbid gastrointestinal symptoms. *Brain, Behavior, and Immunity*, 70, 354–368. 10.1016/j.bbi.2018.03.025
- Rutten BP, Hammels C, Geschwind N, Menne-Lothmann C, Pishva E, Schruers K, ... Wichers M (2013). Resilience in mental health: Linking psychological and neurobiological perspectives. *Acta Psychiatrica Scandinavica*, 128, 3–20. 10.1111/acps.12095 [PubMed: 23488807]
- Scott KM, Lim C, Al-Hamzawi A, Alonso J, Bruffaerts R, Caldasde-Almeida JM, ... Kessler RC (2016). Association of mental disorders with subsequent chronic physical conditions: World mental health surveys from 17 countries. *Journal of the American Medical Association Psychiatry*, 73, 150–158. 10.1001/jamapsychiatry.2015.2688 [PubMed: 26719969]
- Sharma A, Wolf DH, Ciric R, Kable JW, Moore TM, Vandekar SN, ... Satterthwaite TD (2017). Common dimensional reward deficits across mood and psychotic disorders: A connectome-wide association study. *American Journal of Psychiatry*, 174, 657–666. 10.1176/appi.ajp.2016.16070774 [PubMed: 28135847]
- Shonkoff JP, Boyce WT, & McEwen BS (2009). Neuroscience, molecular biology, and the childhood roots of health disparities: Building a new framework for health promotion and disease prevention. *Journal of the American Medical Association*, 301, 2252–2259. 10.1001/jama.2009.754 [PubMed: 19491187]
- Shonkoff JP, & Fisher PA (2013). Rethinking evidence-based practice and two-generation programs to create the future of early childhood policy. *Development and Psychopathology*, 25, 1635–1653. 10.1017/S0954579413000813 [PubMed: 24342860]
- Shonkoff JP, Garner AS, Siegel BS, Dobbins MI, Earls MF, McGuinn L, ... the Committee on Psychosocial Aspects of Child and Family Health, the Committee on Early Childhood, Adoption, and Dependent Care, & the Section on Developmental and Behavioral Pediatrics. (2012). The lifelong effects of early childhood adversity and toxic stress. *Pediatrics*, 129, e232–e246. 10.1542/peds.2011-2663 [PubMed: 22201156]
- Strawbridge R, Arnone D, Danese A, Papadopoulos A, Herane Vives A, & Cleare AJ (2015). Inflammation and clinical response to treatment in depression: A meta-analysis. *European Neuropsychopharmacology*, 25, 1532–1543. 10.1016/j.euroneuro.2015.06.007 [PubMed: 26169573]

- Sun Y, Wittenberg G, Chen G, Drevets W, Hsu B, & Curran M (2017). Improvement in measures of depressed mood and anhedonia in two randomized, placebo-controlled Phase III studies of sirukumab, a human anti-interleukin-6 antibody, in patients with rheumatoid arthritis. *Biological Psychiatry*, 81, S123 10.1016/j.biopsych.2017.02.314
- Treadway MT, & Zald DH (2013). Parsing anhedonia: Translational models of reward-processing deficits in psychopathology. *Current Directions in Psychological Science*, 22, 244–249. 10.1177/0963721412474460 [PubMed: 24748727]
- Tsigos C, & Chrousos GP (2002). Hypothalamic-pituitary-adrenal axis, neuroendocrine factors and stress. *Journal of Psychosomatic Research*, 53, 865–871. 10.1016/S0022-3999(02)00429-4 [PubMed: 12377295]
- Watkins ER (2009). Depressive rumination: Investigating mechanisms to improve cognitive behavioural treatments. *Cognitive Behaviour Therapy*, 38(Suppl. 1), 8–14. 10.1080/16506070902980695 [PubMed: 19697180]
- White J, Kivimäki M, Jokela M, & Batty GD (2017). Association of inflammation with specific symptoms of depression in a general population of older people: The English Longitudinal Study of Ageing. *Brain, Behavior, and Immunity*, 61, 27–30. 10.1016/j.bbi.2016.08.012
- Wingo AP, Wrenn G, Pelletier T, Gutman AR, Bradley B, & Ressler KJ (2010). Moderating effects of resilience on depression in individuals with a history of childhood abuse or trauma exposure. *Journal of Affective Disorders*, 126, 411–414. 10.1016/j.jad.2010.04.009 [PubMed: 20488545]

Table 1

Five Levers and Five Barriers of Translational Neuroscience

Levers	Barriers
<p>1. Elucidate underlying neural and neurobiological mechanisms of behavior and health</p> <p><i>Moving “down” from behavior to biological level of analysis to facilitate creative insights and inform treatment design.</i></p>	<p>1. The “So what?” problem</p> <p><i>Successfully pinpointing neurobiological mechanisms of disease-relevant processes has yet to inform treatment now.</i></p> <p><i>Overcome: Higher-quality neuroscientific data and greater acknowledgment within the field of limitations of inferences based on brain and biological-level data.</i></p>
<p>2. Understand individual differences in treatment response and outcomes at a neurobiological level to increase precision of targets</p> <p><i>Identifying neurobiological moderators of treatment by identifying points of interface between individual processes and treatment mechanisms.</i></p>	<p>2. The “Now what?” problem</p> <p><i>There is a limited quantity of tools available to directly intervene on a system once it is known to be causally involved in the disease.</i></p> <p><i>Overcome: A more nuanced mapping between neural, biological, and mental processes is necessary as well as rigorous examination of biomarkers within intervention trials.</i></p>
<p>3. Catalogue effects of context or disorder on neurobiological processes</p> <p><i>Insights about biological foundations of illness can inform public discourse and policy.</i></p>	<p>3. Scalability</p> <p><i>Complex neuroscientific tools are difficult to scale in community settings (e.g., primary care, school).</i></p> <p><i>Overcome: Transdisciplinary teams of neuroscientists, interventionists, implementation scientists, and community partners at all stages of TN.</i></p>
<p>4. Compare neurobiological systems to make connections between unknown disease-relevant processes</p> <p><i>Identify overlap in neurobiological processes to establish a bridge between superficially dissimilar topics and spur new advances.</i></p>	<p>4. Costliness</p> <p><i>Neuroscientific tools (e.g., neuroimaging) are costly in terms of money and time.</i></p> <p><i>Overcome: Develop new funding models to allow for more rapid progress in the translational cycle.</i></p>
<p>5. Compare neurobiological systems to discriminate between processes that appear similar</p> <p><i>Identify unique neurobiological profiles to refine scientific concepts to present new pathways for targeted interventions.</i></p>	<p>5. Need for diverse skills</p> <p><i>A single researcher cannot possess the set of skills necessary to fully implement translational neuroscience.</i></p> <p><i>Overcome: Collaboration and the revision of prevailing models of training to support transdisciplinary work.</i></p>