Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Short Communication

COVID-19 pandemic and environmental pollution: A blessing in disguise?

Sulaman Muhammad^{*}, Xingle Long^{*}, Muhammad Salman

School of Management, Jiangsu University, Zhenjiang 212013, China

HIGHLIGHTS

- Half of the world population is under some form of lockdown due to COVID-19.
- Environmental pollution is reduced up to 30%.
- Mobility is reduced up to 90%.

GRAPHICAL ABSTRACT

Before Lockdown

After Lockdown

ARTICLE INFO

Article history:

Received 13 April 2020

Received in revised form 16 April 2020

Accepted 17 April 2020

Available online 20 April 2020

Editor: Jianmin Chen

Keywords:

COVID-19

Environmental pollution

NO₂ emissions

ABSTRACT

In late 2019, a novel infectious disease with human to human transmission (COVID-19) was identified in Wuhan China, which now has turned into a global pandemic. Countries all over the world have implemented some sort of lockdown to slow down its infection and mitigate it. Lockdown due to COVID-19 has drastic effects on social and economic fronts. However, this lockdown also has some positive effect on natural environment. Recent data released by NASA (National Aeronautics and Space Administration) and ESA (European Space Agency) indicates that pollution in some of the epicenters of COVID-19 such as Wuhan, Italy, Spain and USA etc. has reduced up to 30%. This study compiled the environmental data released by NASA and ESA before and after the coronavirus pandemic and discusses its impact on environmental quality.

© 2020 Elsevier B.V. All rights reserved.

1. Introduction

A novel infectious disease of coronavirus family was identified in Wuhan, China in late December 2019, which was later named as COVID-19 (Chen et al., 2020). In January 2020, WHO (World Health Organization) confirmed human to human transmission of COVID-19 through respiratory droplets (WHO, 2020). Later on the same month, authorities confirmed a cluster of COVID-19 cases in Wuhan, which increased rapidly not only in surrounding areas but also spread in the whole country and the outbreak turned into epidemic (Dutheil et al., 2020). On January 23rd, Wuhan was placed under quarantine, while

Hubei province follows within few days. Subsequently, Chinese government placed the whole country on lockdown to slow down the spread of infection and ease burden on health facilities (Wilder-Smith and Freedman, 2020). Chinese authorities shut down public transport, educational institutes, businesses centers, parks and other social interaction points to curtail the transmission of COVID-19. On January 30th, WHO declared worldwide public health emergency. In February, outbreaks begin in Iran, Italy and other countries around the globe. Subsequently, the epidemic turns into pandemic and by end of March half of the world population was under some form of lockdown (Tosepu et al., 2020). As of April 16, 2020, the total no of COVID-19 cases surpassed 2.1 million worldwide, with no of total deaths more than 135,000 (WHO, 2020).

As countries went into lockdown the industrial activities shut down globally. Among many other sectors, transport is the most hard hit sector due to lockdown. Road and air transport came to halt as people are

^{*} Corresponding authors.

E-mail addresses: Sulaman_muhammad@yahoo.com (S. Muhammad), longxingle@163.com (X. Long).

Table 1
NO₂ emissions data acquisition across different regions.

Location	Agency	Satellite	Time	% Reduction	Source
Wuhan	NASA and ESA	Aura and Sentinel-5P	Jan–Feb (2019 and 2020)	30%	(NASA, 2020)
China	ESA	Sentinel-5P	Jan and Feb 2020	20–30%	(ESA, 2020)
Europe	ESA	Sentinel-5P	Mar 2019 and Mar 2020	20–30%	(ESA, 2020)
Italy	ESA	Sentinel-5P	Mar 2019 and Mar 2020	20–30%	(ESA, 2020)
France	ESA	Sentinel-5P	Mar 2019 and Mar 2020	20–30%	(ESA, 2020)
Spain	ESA	Sentinel-5P	Mar 2019 and Mar 2020	20–30%	(ESA, 2020)
USA	NASA	Aura	Mar 2015–19 and Mar 2020	30%	(NASA, 2020)

Table 2
Mobility index report based on google tracking.

Location	Transport	Grocery and pharmacy	Retail and recreation	Work place	Parks and outing	Residential
USA	–54%	–20%	–49%	–40%	–20%	+13%
Spain	–89%	–77%	–94%	–68%	–90%	+23%
Italy	–86%	–82%	–95%	–62%	–90%	+24%
France	–82%	–62%	–85%	–53%	–73%	+17%
Germany	–47%	–13%	–58%	–30%	+61%	+8%
UK	–70%	–41%	–82%	–54%	–29%	+15%

not allowed or hesitate to travel. According to report, air travel dropped by 96% due to COVID-19, lowest in 75 years (CNN, 2020). Furthermore, not only transport sector but also industrial and manufacturing sector is heavily affected by pandemic. Global oil demand declined drastically and prices cut down sharply, as industrial and transport sectors came to halt worldwide. COVID-19 has severe negative impact on human health and world economy, however it also results in pollution reduction due to limited social and economic activities (Dutheil et al., 2020).

NO₂ (nitrogen dioxide) is a highly reactive pollutant and emitted especially from the combustion of fossil fuels. Traffic pollution is considered as the major source of NO₂ emissions (He et al., 2020a, 2020b). NO₂ is considered highly lethal to human health as studies shows that both short term and long term exposure to NO₂ can increase mortality rate (Faustini et al., 2014). NO₂ may cause bronchial hyper responsiveness, cellular inflammation and respiratory problems (He et al., 2020a, 2020b). Annually 4.6 million people die worldwide due to poor air quality. Air pollution is a global problem and its effects can be seen even across developed nations such as Europe where 193,000 people died due to air pollution in 2012 (Cohen et al., 2017).

2. Pollution assessment during COVID-19

Lockdown due to COVID-19 reduced transport activities which results in less energy consumption and lower oil demand. These changes in transport activities and oil demand exert a significant impact on the environmental quality. NASA (National Aeronautics and Space Administration) and ESA (European Space Agency) released fresh evidence which suggests that environmental quality improved and the emission of NO₂ reduced up to 30%. NASA collect the data using OMI (Ozone

Fig. 1. NO₂ emissions in Wuhan during 2019 and 2020. (NASA, 2020)

Fig. 2. NO₂ emissions in China before and after lockdown. (ESA, 2020)

Monitoring Instruments) on its AURA satellite. While, ESA collect the data through Sentinel-5P satellite using TROPOMI (TROPOspheric Monitoring Instrument). NASA and ESA release satellite images of various countries before and after lockdown (Table 1).

In addition, Google release data of mobility index report from February 23 to April 05 (see Table 2). The data shows that mobility has reduced up to 90%. Particularly across Spain, Italy and France, while, mobility has reduced the least across USA.

2.1. Wuhan

Fig. 1 shows the NO₂ concentration in Wuhan during 2019 and 2020. Where NO₂ emissions is reduced up to 30% (NASA, 2020). NO₂ emissions is measured using TROPOMI instruments through Sentinel-5P satellite. The image provide comparison between Wuhan 2019 (Jan and Feb) and 2020 (Jan and Feb), NO₂ emissions. Where it clearly indicates that NO₂ emissions reduced significantly due to COVID-19.

Fig. 3. NO₂ emissions in Spain before and after lockdown. (ESA, 2020)

Fig. 4. NO₂ emissions in France before and after lockdown. (ESA, 2020).

2.2. China

Fig. 2 represents NO₂ emissions sequence of China before and after lockdown. Where NO₂ emissions is reduced up to 20–30% from February 10 to 25 after lockdown was implemented (ESA, 2020). The satellite image was captured by ESA satellite Sentinel-5P using TROPOMI Instrument.

2.3. Spain

Fig. 3 represents NO₂ emissions concentration in Spain during March 2019 and March 2020. According to (ESA, 2020), the NO₂ emissions reduced up to 20 to 30% in Spain due to lockdown, especially across the major cities such as Madrid, Barcelona and Seville. The satellite image was captured by ESA satellite Sentinel-5P using TROPOMI Instrument.

2.4. France

Fig. 4 represents NO₂ emissions concentration in France during March 2019 and March 2020. NO₂ emissions is reduced up to 20 to 30% in France (ESA, 2020). The satellite image was captured by ESA satellite Sentinel-5P using TROPOMI Instrument. NO₂ emissions reduced

significantly during lockdown across Paris and other major cities due to transportation shutdown.

2.5. Italy

Fig. 5 represents NO₂ emissions concentration in France during March 2019 and March 2020. The image indicate that NO₂ concentration reduced significantly during lockdown due to transport shutdown and low mobility. The satellite image was captured by ESA satellite Sentinel-5P using TROPOMI Instrument. NO₂ emissions across Italy is reduced up to 20 to 30% (ESA, 2020).

2.6. USA

Fig. 6 represents NO₂ emissions concentration in northeastern part of United States during March 2015 to 2019 and March 2020. Satellite image was captured by NASA through AURA satellite using OMI instrument. Where NO₂ emissions is reduced up to 30% due to lockdown in northeastern part of USA (NASA, 2020).

3. Conclusion

COVID-19 is a global pandemic and serious threat to human health which halt the economic activities, however it is also considered as a

Fig. 5. NO₂ emissions in Italy before and after lockdown. (ESA, 2020)

Fig. 6. NO₂ emissions in northeastern USA before and after lockdown. (NASA, 2020).

“Blessing in Disguise”, where pollution is reducing and nature is reclaiming itself. This positive impact on environment maybe temporary but governments and individuals should learn from this lockdown on how to reduce pollution on long term basis.

CRedit authorship contribution statement

Sulaman Muhammad: Conceptualization, Investigation, Writing - review & editing. **Xingle Long:** Conceptualization, Supervision, Writing - review & editing. **Muhammad Salman:** Conceptualization, Investigation.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

This study is funded by National Natural Science Foundation of China (Nos. 71911540483–71603105). Ministry of Educations and the National Research Foundation of Korea (No. NRF - 2018S1A5A2A03036952).

References

Chen, H., Guo, J., Wang, C., Luo, F., Yu, X., Zhang, W., Li, J., Zhao, D., Xu, D., Gong, Q., Liao, J., Yang, H., Hou, W., Zhang, Y., 2020. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet* [https://doi.org/10.1016/S0140-6736\(20\)30360-3](https://doi.org/10.1016/S0140-6736(20)30360-3).

CNN, 2020. . CNN, 2020. doi. <https://edition.cnn.com/2020/04/09/politics/airline-passengers-decline/index.html>.

Cohen, A.J., Brauer, M., Burnett, R., Anderson, H.R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef, B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling, A., Kan, H., Knibbs, L., Liu, Y., Martin, R., Morawska, L., Pope, C.A., Shin, H., Straif, K., Shaddick, G., Thomas, M., van Dingenen, R., van Donkelaar, A., Vos, T., Murray, C.J.L., Forouzanfar, M.H., 2017. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet* [https://doi.org/10.1016/S0140-6736\(17\)30505-6](https://doi.org/10.1016/S0140-6736(17)30505-6).

Dutheil, F., Baker, S.J., Navel, V., 2020. COVID-19 as a factor influencing air pollution? *Environ. Pollut.* <https://doi.org/10.1016/j.envpol.2020.114466>.

ESA, 2020. . ESA, 2020. doi. https://www.esa.int/Applications/Observing_the_Earth/Copernicus/Sentinel-5P.

Faustini, A., Rapp, R., Forastiere, F., 2014. Nitrogen dioxide and mortality: review and meta-analysis of long-term studies. *Eur. Respir. J.* <https://doi.org/10.1183/09031936.00114713>.

He, M.Z., Kinney, P.L., Li, T., Chen, C., Sun, Q., Ban, J., Wang, J., Liu, S., Goldsmith, J., Kioumourtzoglou, M.A., 2020a. Short- and intermediate-term exposure to NO₂ and mortality: a multi-county analysis in China. *Environ. Pollut.* <https://doi.org/10.1016/j.envpol.2020.114165>.

He, L., Zhang, S., Hu, J., Li, Z., Zheng, X., Cao, Y., Xu, G., Yan, M., Wu, Y., 2020b. On-road emission measurements of reactive nitrogen compounds from heavy-duty diesel trucks in China. *Environ. Pollut.* <https://doi.org/10.1016/j.envpol.2020.114280>.

NASA, 2020. NASA, 2020. <https://earthobservatory.nasa.gov/images>.

Tosepu, R., Gunawan, J., Effendy, S.D., Ahmad, A.I., Lestari, H., Bahar, H., Asfian, P., 2020. Correlation between weather and Covid-19 pandemic in Jakarta, Indonesia. *Sci. Total Environ.* <https://doi.org/10.1016/j.scitotenv.2020.138436>.

WHO, 2020. WHO. <http://www.who.int>.

Wilder-Smith, A., Freedman, D.O., 2020. Isolation, quarantine, social distancing and community containment: pivotal role for old-style public health measures in the novel coronavirus (2019-nCoV) outbreak. *J. Travel Med.* <https://doi.org/10.1093/jtm/taaa020>.