Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Occasional Survey

CLINICAL VIRAL INFECTIONS AND MULTIPLE SCLEROSIS

WILLIAM A. SIBLEY COLIN R. BAMFORD
KATHERINE CLARK

Department of Neurology, University of Arizona, Tucson, AZ, USA

Summary Over an 8 year period, 170 patients with multiple sclerosis (MS) and 134 healthy controls were assessed at monthly intervals in order to ascertain environmental factors which might be important in producing exacerbation or progression of the illness, and to compare the frequency of common viral infections in the two groups. During cumulative periods designated "at risk" (2 weeks before the onset of infection until 5 weeks afterwards) annual exacerbation rates were almost 3-fold greater than those during periods not at risk. Approximately 9% of infections were temporally related to exacerbations, whereas 27% of exacerbations were related to infections. Frequency of common infections was approximately 20–50% less in MS patients than controls; it was progressively less in those with greater disability. Even in minimally disabled patients with similar potential for infectious contacts, the infection rate was significantly less than in controls, suggesting that MS patients could have superior immune defences against common viruses.

INTRODUCTION

In 1976 we began a prospective study of patients with multiple sclerosis (MS) and healthy controls in order to ascertain factors which might trigger exacerbations. The effects of physical trauma,¹ anaesthesia,² influenza immunisation,³ and season⁴ have been published; we now report our findings on the frequency of clinical viral infections and the effects of such infections on annual exacerbation rates.

PATIENTS AND METHODS

Patients

Patients with MS, as defined by Schumacher Committee criteria⁵ were invited to participate in the study. Over an 8 year period (Jan 1, 1976–Dec 31, 1983) 170 patients (67 males and 103 females) were followed for at least 1 year; average duration of observation was 5.3 years. Each patient was contacted monthly to complete a questionnaire about the occurrence of various events during the preceding 30 days, and was examined routinely every 3 months; additional examinations were done whenever exacerbations occurred. At the time of each neurological examination a disability status scale (DSS) rating was given.⁶ Between clinic visits there was

TABLE I—COMPARISON OF MS PATIENTS WITH MINIMAL DISABILITY (DSS 0–2) WITH CONTROLS OF SIMILAR AGE

	Patients (n = 36)	Controls (n = 105)
Mean age	34.7 (±10.8 SD)	34.4 (±9.9 SD)
Mean household size	2.9	3.0
Employed outside home	61%	68%
Children under 18 yr at home	53%	38%
Children under 10 yr at home	33%	28%
Months involved with viral infections*	10.5%	13.3%

*See table II.

frequent telephone contact with the clinic nurse (K. C.) to report new symptoms and to assure completion of the questionnaire on time. Mildly affected patients (DSS 0–2) were mostly employed outside the home in occupations similar to the controls (table I).

Controls

134 age-matched healthy adults (49 males and 85 females) were followed for a minimum of 1 year; mean period of observation was 4.6 years. 42 controls lived in the same household as a patient (38 spouses and 4 blood relatives). 17 additional controls were blood relatives of MS patients, but the remaining 113 were not related. Control subjects completed the same monthly questionnaire as the patients and telephone interviews were also used in this group to avoid a lapse of more than 30 days between reporting periods.

Exacerbation

An exacerbation was defined as a new neurological symptom, associated with an appropriate change in neurological examination, which lasted more than 48 h and was not associated with fever. The term implied a new lesion in the nervous system, or worsening of an existing lesion.

Viral Infections

Viral infections were identified by the symptoms commonly associated with them. Thus complaints such as coryza, sore throat, cough, and malaise, with or without fever, characterised most of the respiratory infections. Infections without fever or with fever <38°C were called "colds"; those with fever >38°C were designated as "flu", with full recognition that many of the illnesses were not due to influenza viruses. When sore throat, malaise, cramps, and diarrhoea were prominent, the infection was called enteric. Herpetic eruptions (genital or oral) were usually characteristic but could not always be distinguished from aphthous ulcers. Viral cultures of patients, controls, and family were not done because this was beyond the scope of our exploratory study.

Periods at Risk and Not at Risk

The at risk (AR) period extended from 2 weeks before the onset of symptoms of infection until 5 weeks afterwards (a total of 7 weeks). All other periods in the study were not at risk (NAR). Annual exacerbation risks were calculated separately for each period. The AR period was defined at the beginning of the study. In a previous study⁷ we used an AR period of 1 week before onset of symptoms

TABLE II—FREQUENCY OF COMMON VIRAL INFECTIONS IN MS PATIENTS WITH VARIOUS DISABILITY RATINGS AND CONTROLS OF SAME AGE

Patients						Controls						
Mean* DSS (range)	No	Mean age (yr)	Infections	Months	Mo involved (%)	No	Mean Age (yr)	Infections	Mo	Mo involved (%)	χ ²	p
0–2	36	34.7	236	2331	10.5	105	34.4	744	5574	13.3	9.45	<0.005
2–4	32	37.8	143	1728	8.3	107	37.8	739	5862	12.6	21.53	<0.001
4–6	48	48.2	214	3185	6.7	51	48.4	296	2896	10.2	22.56	<0.001
6–8	44	47.7	163	2962	5.5	61	47.6	359	3436	10.4	47.53	<0.001
8–10	10	47.1	23	730	3.1	73	47.3	426	4312	9.9	31.76	<0.001

*Mean DSS = average of entry and exit DSS scores.

TABLE III—FREQUENCY OF VARIOUS INFECTIONS IN MS PATIENTS AND CONTROLS

—	Mo in study	Colds (no/%mo)	Flu (no/%mo)	Enteric (no/%mo)	Herpes (no/%mo)
Controls	7418	590/7.9%	141/1.9%	152/2.0%	25/0.3%
Patients	11 486	523/4.5%	143/1.2%	101/0.8%	4/0.03%

p<0.001 for comparison between patients and controls for all infections.

until 3 weeks afterwards; it was subsequently realised that these limits were too narrow and therefore they were extended for the present study. The rationale for an AR period of this duration stems from the fact that most cases of encephalomyelitis due to rabies vaccine—perhaps the best human model of demyelinating disease resulting from antigenic sensitisation—occur within a latent period of 35 days.⁸ Extending the AR period to 2 weeks before the onset of the infection seemed wise because of difficulties in dating the onset of infections with precision. We were also influenced by the studies of Fox et al⁹ who noted that virus shedding sometimes occurred as early as 13 days before the onset of clinical symptoms.

RESULTS

Frequency of Total Viral Infections in MS Patients and Controls

Viral-like infections were significantly less frequent in patients than in controls; the difference was most pronounced in patients with severe disability. Table II shows that there was a linear drop in the frequency of such infections with advancing disability. Although much of this difference might be attributed to the patients' sheltered lives, giving rise to fewer infectious contacts, this may be only a partial explanation. The frequency of viral infections was also significantly lower in patients with trivial disability (DSS 0–2) whose daily activities were comparable to controls. Table I shows that patients with mild MS resembled controls with respect to important determinants of infection frequency such as age, household size, employment outside the home, and contacts with children under the age of 10 years.¹⁰ Moreover, although the differences in infection rates were evident for all categories of viral infection, they were most pronounced for herpes simplex-like infections (table III), in which clinical symptoms in adults are usually due to re-emergence of latent virus rather than person-to-person contact. Only 4 cases of herpetic eruption occurred in 170 MS patients during 5.3 years, whereas 25 cases occurred in 134 controls during 4.6 years.

Viral-like Infections, Exacerbation Rates, and Progression of MS

Exacerbation rates in AR periods were almost 3-fold greater than those in NAR periods (table IV). This effect was apparent not only in patients with frequent exacerbations but also in those with moderate and low rates (table V). The same accentuation of exacerbation rates occurred in patients of all levels of disability, except that the numbers of exacerbations

TABLE IV—RELATION BETWEEN MS EXACERBATIONS AND 779 VIRAL INFECTIONS IN 170 PATIENTS IN 5.3 YR

Period	Wk	Exacerbations	Exacerbation rate/yr
AR	5453 (12%)	67 (27%)	0.64* (0.49–0.79)†
NAR	41 150 (88%)	179 (73%)	0.23* (0.19–0.26)†
Total	46 603	246	..

* $\chi^2 = 56.3$; p<0.001.
†95% confidence limits.

in severely disabled patients (DSS 7–9) were so few that the figures did not reach significance for this group (table VI). 75 of the 170 patients had no exacerbations during the course of the study. 50 of 95 patients had at least 1 AR exacerbation, while 17 had more than 1 (colds 44, enteric infections 11, flu 10, and herpes simplex-like 2).

The effect of frequency of infection on exacerbation was studied by dividing patients into two groups, one with <1 infection per year and the other with more (table VII). The exacerbation rate during AR periods in mildly disabled patients with frequent infections was 5-fold greater than that of similarly disabled patients with infrequent infections. Nevertheless, the patients with infrequent infections were 10 years older and had a significantly more rapid progression of MS, the reasons for which are not apparent.

Median time between the onset of infection and occurrence of MS exacerbation was plus 8 days, and this was also the most common time; the mean time was plus 10 days. In 30 instances the interval between infection and a new attack varied from 9–35 days; in 18 instances infection followed MS exacerbation, usually by a few days, but in 5 by as long as 10–14 days. It seems most unlikely that MS exacerbations could be related to clinical infection that occurred 2 weeks later, and we suspect that in such cases onset of infection was dated erroneously. Even with a 30-day reporting period, it was often difficult for patients and controls to date the onset of infections precisely; an error of ± 7 days would probably be a reasonable assumption.

DISCUSSION

Modern imaging methods confirm that new symptoms and signs in MS are usually associated with one or more new lesions in the white matter of the central nervous system;¹¹ sometimes this represents renewed activity at the margin of an old plaque.¹² In either event, it would appear important to identify factors associated with new demyelination, since this determines either a step-wise or slow increase in disability.

The present study strongly suggests that common viral-like infections are temporally associated with exacerbations. Most earlier studies of the relation between infection and MS have been retrospective, relying either on medical record reviews or patients' memories to recall events before the onset of MS. Such studies, reviewed by McAlpine et al,¹³ showed frequencies of infection before the first attack of MS of

TABLE V—ASSOCIATION OF MS EXACERBATIONS AND VIRAL INFECTIONS BY FREQUENCY OF EXACERBATION*

Frequency† of exacerbation	Mean rate	Patients	Wk AR/ exacerbations	Wk NAR/ exacerbations	Annual infection rate	DSS (entry)	DSS (exit)	Exacerbation rate AR	Exacerbation rate NAR
High	0.70	36	1505/40	7610/83	1.22	3.0	3.9	1.38‡	0.57
Moderate	0.31	30	1260/19	8754/42	0.93	3.6	4.6	0.78‡	0.25
Low	0.04	89	2681/8	22 330/12	0.79	5.0	5.84	0.16‡	0.03

*15 patients who had no infections are excluded.
†High=0.5 or more exacerbations/yr; moderate=0.25–0.49/yr; low=0–0.24/yr.
‡p<0.001 (χ^2) by comparison with NAR.

TABLE VI—MEAN MS EXACERBATION RATE IN AR AND NAR PERIODS BY MEAN DISABILITY

Mean DSS DSS rating	Patients	Mean age (entry)	Infections/yr	Exacerbations/AR	Exacerbations/NAR	Wk AR	Wk NAR	Exacerbation rate/yr AR	Exacerbation rate/yr NAR
0-2	36	34.7	1.28	28	47	1652	7941	0.88*	0.31
2-4	32	37.8	1.00	19	51	1001	6430	0.99*	0.41
4-6	48	48.2	0.81	14	47	1498	12207	0.49*	0.20
6-8	44	47.7	0.66	6	31	1141	11570	0.27	0.14
8-10	10	47.1	0.38	0	3	161	3002	0.00	0.06

* $p < 0.001$ by comparison with NAR.

10-41%. The present results are similar to those of Sibley and Foley who did a prospective study of 34 patients over a period of 3 years and found that 33 of 69 exacerbations (48%) were temporally associated with a common infection—ie, the infection occurred within a period from 3 weeks before to 1 week after the exacerbation.⁷ Concurrent controls were not used in that study but it was noted that the infection rate was lower than that in a large study of infections in healthy families in the same area;¹⁰ at the time, under-reporting of infections not associated with MS exacerbations was suspected, but in light of the present results this conclusion was probably incorrect. In the previous study, patients were aware that the relation of infection to MS was the subject of the investigation; in the present study, infection was only one of many factors being investigated, and care was taken to avoid any patient preconception.

The apparent triggering of MS attacks by infections is probably not a nonspecific effect of any infection. By the same methods of analysis, for example, we have found no relation between bacterial infections (mostly of the urinary tract) and worsening of MS. Moreover, although the annual exacerbation rate almost trebled during AR periods, only 8.6% of infections were associated with worsening of MS. Such a low rate of triggering might be explained, in part, by the fact that many new lesions find no clinical expression. However, specific triggering only by a certain virus or class of viruses might be another reason.

If viral triggering is important to formation of MS lesions, why are only 27% of MS exacerbations associated with evidence of infection? One possible explanation is the known frequency of inapparent infection—eg, as many as 80% of primary enteroviral infections are symptomless. Recurrent infections by the same virus are often associated with partial immunity and inapparent infection; reinfections with rhinoviruses, parainfluenza and respiratory syncytial viruses, and other important organisms in respiratory illness, are usually symptomless.¹⁴

Since seasonal studies in Arizona,⁵ Cleveland,⁷ Canada,¹⁵ and Newcastle upon Tyne¹⁶ have shown spring and summer peaks in exacerbation rates; this clustering may be attributable to certain viruses with a similar seasonal incidence. Coronaviruses show this pattern,¹⁷ but, because

there was a 4-5 week latency between infection and exacerbation in some of our cases, other viruses with slightly earlier seasonal peaks must also be considered. Isolation and identification of viruses during clinical infections in MS patients and family members should clarify the precise role of these infections in the pathogenesis of MS, and establish which agents may be most important.

While MS patients have fewer common infections than either family members or controls, it is difficult to know whether sheltering, immunity, or both, are responsible. The fact that a difference from controls can be shown even in MS patients with insignificant disability and the same number of contacts—especially with school children—suggests that immune defences may play a role. It is possible that they have more inapparent infections, perhaps as the result of more efficient natural defences against these common viruses. The finding of a much more rapid progression rate in mildly affected MS patients with infrequent symptoms of clinical infection suggests that overactive immune mechanisms determine not only a greater frequency of inapparent infection, but also swifter progress of disease.

This work was supported by research grant 1048-C-13 from the National Multiple Sclerosis Society, New York.

Correspondence should be addressed to W. A. S., University Hospital, Tucson, AZ 85724, USA.

REFERENCES

- Bamford CR, Sibley WA, Thies C, Laguna JF, Smith MS, Clark K. Trauma as an etiologic and aggravating factor in multiple sclerosis. *Neurology (NY)* 1981; **31**: 1229-34.
- Bamford CR, Sibley WA, Laguna JF. Anesthesia in Multiple sclerosis. *Can J Neurol Sci* 1978; **5**: 41-44.
- Bamford CR, Sibley WA, Laguna JF. Swine influenza vaccination in patients with multiple sclerosis. *Arch Neurol* 1978; **35**: 242-43.
- Bamford CR, Sibley WA, Thies C. Seasonal variation of multiple sclerosis exacerbations in Arizona. *Neurology (NY)* 1983; **33**: 697-701.
- Schumacher GA, Beebe G, Kibler RF et al. Problems of experimental trials of therapy in multiple sclerosis: Report by the panel on evaluation of experimental trials of therapy in multiple sclerosis. *Ann NY Acad Sci* 1965; **122**: 552-68.
- Kurtzke JF. On the evaluation of disability in multiple sclerosis. *Neurology (NY)* 1961; **11**: 686-94.
- Sibley WA, Foley JM. Infection and immunization in multiple sclerosis. *Ann NY Acad Sci* 1965; **122**: 457-68.
- Shiraki H, Otani S. Clinical and pathological features of rabies post-vaccinal encephalomyelitis in man. In: Kies MW, Alvord EC, eds. 'Allergic' encephalomyelitis. Springfield: Charles C Thomas, 1959: 58-129.
- Fox JP, Hall CE. Viruses in families. Littleton: PSG Publ Co, 1980: 82.
- Dingle JH, Badger GF, Jordan WS. Illness in the home—a study of 25 000 illnesses in a group of Cleveland families. Cleveland: Western Reserve University Press, 1964: 19-96.
- Ebers GC, Paty DW, Sears ES. Imaging in multiple sclerosis. In: Poser CM, Paty DW, Scheinberg L, McDonald WI, Ebers GC, eds. The diagnosis of multiple sclerosis. New York and Stuttgart: Thieme-Stratton, 1984: 186.
- Adams CWM. The general pathology of multiple sclerosis: morphological and chemical aspects of the lesions. In: Hallpike JF, Adams CWM, Tourtellotte WW, eds. Multiple sclerosis. Baltimore: Williams and Wilkins, 1983: 203-40.
- McAlpine D, Lumsden CE, Acheson ED. Multiple sclerosis. A reappraisal. Edinburgh: Churchill Livingstone, 1972: 99-100.
- Chang T. Recurrent viral infection (reinfection). *New Engl J Med* 1971; **284**: 765-73.
- Sibley WA, Paty DW. A comparison of multiple sclerosis in Ontario, Canada and Arizona, USA. *Acta Neurol Scand* 1981; **64** (suppl 87): 60-68.
- Prineas JW. The etiology and pathogenesis of multiple sclerosis. In: Vinken PJ, Bruyn GW, eds. Vol 9 Handbook of clinical neurology. Multiple sclerosis and other demyelinating diseases. Amsterdam: North Holland, 1970: 119.
- Monto AS. Coronaviruses. In: Evans AS, ed. Viral infections of humans. New York: Plenum, 1982: 157.

TABLE VII—FREQUENCY OF CLINICAL VIRAL INFECTION, AND PROGRESSION OF MS DISABILITY IN 170 PATIENTS OVER 5.3 YR

Entry DSS rating	Infections	Patients	Mean age	Mean infections /yr	Mean progression of DSS/yr	Exacer- bation rate/yr AR	Exacer- bation rate/yr NAR
0-4	<1/yr	37	43	0.53	0.43*	0.19†	0.42
	≥1/yr	46	33	1.68	0.19*	1.02†	0.37
5-9	<1/yr	64	50	0.41	0.13	0.39	0.16
	≥1/yr	23	44	1.55	0.15	0.22	0.21

*SE of difference in means = 0.08. $p = 0.0027$.

†SE of difference in means = 0.16; $p < 0.001$.