Cochrane
Library

Cochrane Database of Systematic Reviews

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Chan EY, Fransen M, Parker DA, Assam PN, Chua N

Chan EY, Fransen M, Parker DA, Assam PN, Chua N.
Femoral nerve blocks for acute postoperative pain after knee replacement surgery.
Cochrane Database of Systematic Reviews 2014, Issue 5. Art. No.: CD009941.
DOI: [10.1002/14651858.CD009941.pub2](https://doi.org/10.1002/14651858.CD009941.pub2).

www.cochranelibrary.com

TABLE OF CONTENTS

HEADER	1
ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
SUMMARY OF FINDINGS	4
BACKGROUND	15
OBJECTIVES	16
METHODS	16
RESULTS	20
Figure 1.	21
Figure 2.	23
Figure 3.	25
Figure 4.	26
Figure 5.	29
Figure 6.	32
DISCUSSION	33
AUTHORS' CONCLUSIONS	35
ACKNOWLEDGEMENTS	35
REFERENCES	36
CHARACTERISTICS OF STUDIES	43
DATA AND ANALYSES	109
Analysis 1.1. Comparison 1 FNB versus PCA opioid, Outcome 1 Pain at rest first 2 hours.	114
Analysis 1.2. Comparison 1 FNB versus PCA opioid, Outcome 2 Pain at rest 3 to 12 hours.	115
Analysis 1.3. Comparison 1 FNB versus PCA opioid, Outcome 3 Pain at rest 24 hours.	116
Analysis 1.4. Comparison 1 FNB versus PCA opioid, Outcome 4 Pain at rest 48 hours.	117
Analysis 1.5. Comparison 1 FNB versus PCA opioid, Outcome 5 Pain at rest 72 hours.	118
Analysis 1.6. Comparison 1 FNB versus PCA opioid, Outcome 6 Pain on movement first 2 hours.	119
Analysis 1.7. Comparison 1 FNB versus PCA opioid, Outcome 7 Pain on movement 3 to 12 hours.	119
Analysis 1.8. Comparison 1 FNB versus PCA opioid, Outcome 8 Pain on movement 24 hours.	120
Analysis 1.9. Comparison 1 FNB versus PCA opioid, Outcome 9 Pain on movement 48 hours.	121
Analysis 1.10. Comparison 1 FNB versus PCA opioid, Outcome 10 Pain on movement 72 hours.	122
Analysis 1.11. Comparison 1 FNB versus PCA opioid, Outcome 11 Pain at rest 24 hours—subgrouped by FNB with and without concurrent PCA.	123
Analysis 1.12. Comparison 1 FNB versus PCA opioid, Outcome 12 Pain on movement 24 hours—subgrouped by FNB with and without concurrent PCA.	124
Analysis 1.13. Comparison 1 FNB versus PCA opioid, Outcome 13 Pain at rest 24 hours—subgrouped by FNB ropivacaine vs bupivacaine.	124
Analysis 1.14. Comparison 1 FNB versus PCA opioid, Outcome 14 Pain on movement 24 hours—subgrouped by FNB ropivacaine vs bupivacaine.	125
Analysis 1.15. Comparison 1 FNB versus PCA opioid, Outcome 15 Pain at rest 24 hours—sensitivity analysis by low bias for allocation concealment.	126
Analysis 1.16. Comparison 1 FNB versus PCA opioid, Outcome 16 Pain on movement 24 hours—sensitivity analysis by low bias for allocation concealment.	126
Analysis 1.17. Comparison 1 FNB versus PCA opioid, Outcome 17 Pain on rest 24 hours—sensitivity analysis by blinding of participants, personnel and outcome assessor.	127
Analysis 1.18. Comparison 1 FNB versus PCA opioid, Outcome 18 Pain on movement 24 hours—sensitivity analysis by blinding of participants, personnel and outcome assessor.	127
Analysis 1.19. Comparison 1 FNB versus PCA opioid, Outcome 19 Opioid consumption 24 hours (mg).	128
Analysis 1.20. Comparison 1 FNB versus PCA opioid, Outcome 20 Opioid consumption 48 hours (mg).	129
Analysis 1.21. Comparison 1 FNB versus PCA opioid, Outcome 21 Nausea and/or vomiting.	130
Analysis 1.22. Comparison 1 FNB versus PCA opioid, Outcome 22 Sedation.	131
Analysis 1.23. Comparison 1 FNB versus PCA opioid, Outcome 23 Urinary retention.	132
Analysis 1.24. Comparison 1 FNB versus PCA opioid, Outcome 24 Knee flexion range of motion (postoperative day 2 to 4).	133

Analysis 1.25. Comparison 1 FNB versus PCA opioid, Outcome 25 Participant satisfaction with analgesia during hospital stay. ..	133
Analysis 2.1. Comparison 2 FNB versus epidural, Outcome 1 Pain at rest first 2 hours.	136
Analysis 2.2. Comparison 2 FNB versus epidural, Outcome 2 Pain at rest 3 to 12 hours.	137
Analysis 2.3. Comparison 2 FNB versus epidural, Outcome 3 Pain at rest 24 hours.	137
Analysis 2.4. Comparison 2 FNB versus epidural, Outcome 4 Pain at rest 48 hours.	138
Analysis 2.5. Comparison 2 FNB versus epidural, Outcome 5 Pain on movement 24 hours.	138
Analysis 2.6. Comparison 2 FNB versus epidural, Outcome 6 Pain on movement 48 hours.	139
Analysis 2.7. Comparison 2 FNB versus epidural, Outcome 7 Pain at rest 24 hours—subgrouped by FNB ropivacaine vs bupivacaine.	140
Analysis 2.8. Comparison 2 FNB versus epidural, Outcome 8 Pain on movement 24 hours—subgrouped by FNB ropivacaine vs bupivacaine.	140
Analysis 2.9. Comparison 2 FNB versus epidural, Outcome 9 Pain at rest 24 hours—sensitivity analysis by low bias for allocation concealment.	141
Analysis 2.10. Comparison 2 FNB versus epidural, Outcome 10 Pain on movement 24 hours—sensitivity analysis by low bias for allocation concealment.	141
Analysis 2.11. Comparison 2 FNB versus epidural, Outcome 11 Opioid consumption 24 hours (mg).	142
Analysis 2.12. Comparison 2 FNB versus epidural, Outcome 12 Opioid consumption 48 hours (mg).	142
Analysis 2.13. Comparison 2 FNB versus epidural, Outcome 13 Nausea and/or vomiting.	143
Analysis 3.1. Comparison 3 FNB versus local infiltration analgesia, Outcome 1 Pain at rest 24 hours.	144
Analysis 3.2. Comparison 3 FNB versus local infiltration analgesia, Outcome 2 Pain at rest 48 hours.	145
Analysis 3.3. Comparison 3 FNB versus local infiltration analgesia, Outcome 3 Pain on movement 24 hours.	145
Analysis 3.4. Comparison 3 FNB versus local infiltration analgesia, Outcome 4 Pain on movement 48 hours.	145
Analysis 3.5. Comparison 3 FNB versus local infiltration analgesia, Outcome 5 Nausea and/or vomiting.	146
Analysis 4.1. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 1 Pain at rest first 2 hours.	147
Analysis 4.2. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 2 Pain at rest 3 to 12 hours.	147
Analysis 4.3. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 3 Pain at rest 24 hours.	148
Analysis 4.4. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 4 Pain at rest 48 hours.	148
Analysis 4.5. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 5 Pain on movement 24 hours.	148
Analysis 4.6. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 6 Pain on movement 48 hours.	149
Analysis 4.7. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 7 Pain on rest 24 hours—subgrouped by continuous FNB with or without concurrent PCA.	149
Analysis 4.8. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 8 Pain on movement 24 hours—subgrouped by continuous FNB with or without concurrent PCA.	149
Analysis 4.9. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 9 Pain at rest 24 hours—sensitivity analysis by low bias for allocation concealment.	150
Analysis 4.10. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 10 Pain on movement 24 hours—sensitivity analysis by low bias for allocation concealment.	150
Analysis 4.11. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 11 Opioid consumption 24 hours (mg).	150
Analysis 4.12. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 12 Opioid consumption 48 hours (mg).	151
APPENDICES	151
WHAT'S NEW	162
CONTRIBUTIONS OF AUTHORS	162
DECLARATIONS OF INTEREST	163
SOURCES OF SUPPORT	163
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	163
INDEX TERMS	164

[Intervention Review]

Femoral nerve blocks for acute postoperative pain after knee replacement surgery

Ee-Yuee Chan^{1,2}, Marlene Fransen¹, David A Parker³, Pryseley N Assam^{4,5}, Nelson Chua⁶¹Faculty of Health Sciences, University of Sydney, Sydney, Australia. ²Nursing Service, Tan Tock Seng Hospital, Singapore, Singapore.³Sydney Orthopaedic Research Institute, Sydney, Australia. ⁴Centre for Quantitative Medicine, Office of Clinical Sciences, Duke-NUS Graduate Medical School, Singapore, Singapore. ⁵Department of Biostatistics, Singapore Clinical Research Institute Pte Ltd, Singapore, Singapore. ⁶Department of Anaesthesiology, Tan Tock Seng Hospital, Singapore, Singapore**Contact address:** Ee-Yuee Chan, Faculty of Health Sciences, University of Sydney, Cumberland Campus C42, Room 205, O Block,, 75 East Street, Sydney, NSW, 1825, Australia. eeuyee@gmail.com.**Editorial group:** Cochrane Anaesthesia Group.**Publication status and date:** Edited (no change to conclusions), published in Issue 1, 2016.**Citation:** Chan EY, Fransen M, Parker DA, Assam PN, Chua N. Femoral nerve blocks for acute postoperative pain after knee replacement surgery. *Cochrane Database of Systematic Reviews* 2014, Issue 5. Art. No.: CD009941. DOI: [10.1002/14651858.CD009941.pub2](https://doi.org/10.1002/14651858.CD009941.pub2).

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

Total knee replacement (TKR) is a common and often painful operation. Femoral nerve block (FNB) is frequently used for postoperative analgesia.

Objectives

To evaluate the benefits and risks of FNB used as a postoperative analgesic technique relative to other analgesic techniques among adults undergoing TKR.

Search methods

We searched the Cochrane Central Register of Controlled Trials (CENTRAL) 2013, Issue 1, MEDLINE, EMBASE, CINAHL, Web of Science, dissertation abstracts and reference lists of included studies. The date of the last search was 31 January 2013.

Selection criteria

We included randomized controlled trials (RCTs) comparing FNB with no FNB (intravenous patient-controlled analgesia (PCA) opioid, epidural analgesia, local infiltration analgesia, and oral analgesia) in adults after TKR. We also included RCTs that compared continuous versus single-shot FNB.

Data collection and analysis

Two review authors independently performed study selection and data extraction. We undertook meta-analysis (random-effects model) and used relative risk ratios (RRs) for dichotomous outcomes and mean differences (MDs) or standardized mean differences (SMDs) for continuous outcomes. We interpreted SMDs according to rule of thumb where 0.2 or smaller represents a small effect, 0.5 a moderate effect and 0.8 or larger, a large effect.

Main results

We included 45 eligible RCTs (2710 participants) from 47 publications; 20 RCTs had more than two allocation groups. A total of 29 RCTs compared FNB (with or without concurrent treatments including PCA opioid) versus PCA opioid, 10 RCTs compared FNB versus epidural, five RCTs compared FNB versus local infiltration analgesia, one RCT compared FNB versus oral analgesia and four RCTs compared continuous versus single-shot FNB. Most included RCTs were rated as low or unclear risk of bias for the aspects rated in the risk of bias

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

assessment tool, except for the aspect of blinding. We rated 14 (31%) RCTs at high risk for both participant and assessor blinding and rated eight (18%) RCTs at high risk for one blinding aspect.

Pain at rest and pain on movement were less for FNB (of any type) with or without a concurrent PCA opioid compared with PCA opioid alone during the first 72 hours post operation. Pooled results demonstrated a moderate effect of FNB for pain at rest at 24 hours (19 RCTs, 1066 participants, SMD -0.72, 95% CI -0.93 to -0.51, moderate-quality evidence) and a moderate to large effect for pain on movement at 24 hours (17 RCTs, 1017 participants, SMD -0.94, 95% CI -1.32 to -0.55, moderate-quality evidence). Pain was also less in each FNB subgroup: single-shot FNB, continuous FNB and continuous FNB + sciatic block, compared with PCA. FNB also was associated with lower opioid consumption (IV morphine equivalent) at 24 hours (20 RCTs, 1156 participants, MD -14.74 mg, 95% CI -18.68 to -10.81 mg, high-quality evidence) and at 48 hours (MD -14.53 mg, 95% CI -20.03 to -9.02 mg), lower risk of nausea and/or vomiting (RR 0.47, 95% CI 0.33 to 0.68, number needed to treat for an additional harmful outcome (NNTH) four, high-quality evidence), greater knee flexion (11 RCTs, 596 participants, MD 6.48 degrees, 95% CI 4.27 to 8.69 degrees, moderate-quality evidence) and greater patient satisfaction (four RCTs, 180 participants, SMD 1.06, 95% CI 0.74 to 1.38, low-quality evidence) compared with PCA.

We could not demonstrate a difference in pain between FNB (any type) and epidural analgesia in the first 72 hours post operation, including pain at 24 hours at rest (six RCTs, 328 participants, SMD -0.05, 95% CI -0.43 to 0.32, moderate-quality evidence) and on movement (six RCTs, 317 participants, SMD 0.01, 95% CI -0.21 to 0.24, high-quality evidence). No difference was noted at 24 hours for opioid consumption (five RCTs, 341 participants, MD -4.35 mg, 95% CI -9.95 to 1.26 mg, high-quality evidence) or knee flexion (six RCTs, 328 participants, MD -1.65, 95% CI -5.14 to 1.84, high-quality evidence). However, FNB demonstrated lower risk of nausea/vomiting (four RCTs, 183 participants, RR 0.63, 95% CI 0.41 to 0.97, NNTH 8, moderate-quality evidence) and higher patient satisfaction (two RCTs, 120 participants, SMD 0.60, 95% CI 0.23 to 0.97, low-quality evidence), compared with epidural analgesia.

Pooled results of four studies (216 participants) comparing FNB with local infiltration analgesia detected no difference in analgesic effects between the groups at 24 hours for pain at rest (SMD 0.06, 95% CI -0.61 to 0.72, moderate-quality evidence) or pain on movement (SMD 0.38, 95% CI -0.10 to 0.86, low-quality evidence). Only one included RCT compared FNB with oral analgesia. We considered this evidence insufficient to allow judgement of the effects of FNB compared with oral analgesia.

Continuous FNB provided less pain compared with single-shot FNB (four RCTs, 272 participants) at 24 hours at rest (SMD -0.62, 95% CI -1.17 to -0.07, moderate-quality evidence) and on movement (SMD -0.42, 95% CI -0.67 to -0.17, high-quality evidence). Continuous FNB also demonstrated lower opioid consumption compared with single-shot FNB at 24 hours (three RCTs, 236 participants, MD -13.81 mg, 95% CI -23.27 to -4.35 mg, moderate-quality evidence).

Generally, the meta-analyses demonstrated considerable statistical heterogeneity, with type of FNB, allocation concealment and blinding of participants, personnel and outcome assessors reducing heterogeneity in the analyses. Available evidence was insufficient to allow determination of the comparative safety of the various analgesic techniques. Few RCTs reported on serious adverse effects such as neurological injury, postoperative falls or thrombotic events.

Authors' conclusions

Following TKR, FNB (with or without concurrent treatments including PCA opioid) provided more effective analgesia than PCA opioid alone, similar analgesia to epidural analgesia and less nausea/vomiting compared with PCA alone or epidural analgesia. The review also found that continuous FNB provided better analgesia compared with single-shot FNB. RCTs were insufficient to allow definitive conclusions on the comparison between FNB and local infiltration analgesia or oral analgesia.

PLAIN LANGUAGE SUMMARY

Femoral nerve blocks for acute postoperative pain after knee replacement surgery

Total knee replacement is a common and often painful orthopaedic operation. Femoral nerve block (FNB) is an analgesic technique that blocks sensation to the knee to reduce pain following surgery. FNB is given as a single injection or as continuous infusion of numbing medication in the groin area.

This review compared FNB with other common analgesic techniques (patient-controlled analgesia (PCA) with opioids, epidural analgesia, local infiltration analgesia and oral analgesia). A total of 45 randomized trials with 2710 participants were included. Among the included studies, 30 treatment groups compared FNB with PCA opioids, 10 compared FNB with epidural analgesia, five compared FNB with local infiltration analgesia, one compared FNB with oral analgesia and four compared continuous FNB with single-injection FNB. The average methodological quality of the included studies was moderate.

FNB (any type) resulted in less pain at rest and on movement during the first 72 hours after surgery, compared with PCA opioid alone. FNB also resulted in lower opioid consumption, fewer patients with nausea and vomiting, greater knee flexion and higher patient satisfaction, compared with PCA opioid. Additionally, continuous FNB provided better pain relief than was attained with single-injection FNB.

No differences in postoperative pain and opioid consumption at 24 hours were noted between FNB and epidural analgesia, although the former resulted in fewer patients with nausea and vomiting, and higher patient satisfaction was reported with analgesia. Similarly, we

found no difference in postoperative pain between participants given FNB as opposed to local infiltration analgesia. Information was insufficient for review authors to conclude on the comparison of FNB with local infiltration analgesia or oral analgesia, or on the safety of the various analgesic techniques.

SUMMARY OF FINDINGS

Summary of findings for the main comparison. FNB (any type) compared with PCA opioid for knee replacement surgery

FNB (any type) compared with PCA opioid for knee replacement surgery

Patient or population: patients with knee replacement surgery

Settings: hospital

Intervention: FNB (any type)

Comparison: PCA opioid

Out-comes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	PCA opioid	FNB (any type)				
Pain at rest at 24 hours Visual analogue scale Scale from zero to 10 Follow-up: median two days	Mean pain at rest at 24 hours ranged across control groups from 0.7 to 5.5 points	Mean pain at rest at 24 hours in the intervention groups was 1.20 lower (1.60 to 0.87 lower)		1066 (19 studies)	⊕⊕⊕⊖ moderate ¹	SMD -0.72 (-0.93 to -0.51), representing moderate effects between groups. Lower score indicates less pain ²
Pain on movement at 24 hours Visual analogue scale Scale from zero to 10	Mean pain on movement at 24 hours ranged across control groups from 2.8 to 8 points	Mean pain on movement at 24 hours in the intervention groups was 1.66 lower (2.32 to 0.97 lower)		1017 (17 studies)	⊕⊕⊕⊖ moderate ¹	SMD -0.94 (-1.32 to -0.55), representing moderate to large effect between groups. Lower score indicates less pain ²

Fol- low-up: median two days						
Neuro- logical injury Fol- low-up: during hospital- ization	See comment	See comment	Not estimable	390 (4 studies)	See comment	Effect is uncertain as neurolog- ical injuries are extremely rare. No events were reported in four studies
Opioid con- sump- tion at 24 hours using mor- phine equiv- alents Scale from ze- ro to 150 mg Fol- low-up: 24 hours	Mean opioid consumption 24 hours ranged across control groups from 19.3 to 99.8 mg	Mean opioid consumption at 24 hours in the intervention groups was 14.17 lower (18.10 to 10.23 lower)		1156 (20 studies)	⊕⊕⊕⊕ high ³	MD -14.74 mg (-18.68 to -10.81 mg)
Nausea and/or vomit- ing Fol- low-up: median two days	Study population ⁴		RR 0.47 (0.33 to 0.68)	1100 (16 studies)	⊕⊕⊕⊕ high ³	
	47 per 100	22 per 100 (15 to 32)				
	Low ⁴					
	10 per 100	5 per 100 (3 to 7)				
	High ⁴					

	93 per 100	44 per 100 (31 to 63)			
Knee flexion range of motion (degree) ⁵ Follow-up: median three days	Mean knee flexion range of motion ranged across control groups from 55 to 85 degrees	Mean knee flexion in the intervention group was 6.48 higher (4.27 to 8.69 higher)	596 (11 studies)	⊕⊕⊕⊖ moderate ¹	MD 6.48 degrees (4.27 to 8.69 degrees)
Participant satisfaction with analgesia (point) Scale from zero to 10 Follow-up: during hospitalization	Mean participant satisfaction ranged across control groups from 6.3 to 8.7 points	Mean participant satisfaction was 1.79 higher (1.25 to 2.32 higher)	180 (4 studies)	⊕⊕⊕⊖ low ⁶	SMD 1.06 (0.74 to 1.38), representing large effects. Higher score indicates greater satisfaction ²

*The basis for the **assumed risk** (e.g. median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).
CI: Confidence interval; **RR:** Risk ratio.

GRADE Working Group grades of evidence.

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹Downgraded for significant heterogeneity.

²Rule of thumb for interpreting SMD: 0.2 small effect, 0.5 moderate effect, 0.8 large effect (Cohen 1988).

³Downgrading for significant heterogeneity balanced by upgrading for large effect based on consistent evidence from more than two studies.

⁴Assumed risk was based on control group risk in the included studies.

⁵Knee flexion of 90 degrees is usually required for patients to be able to climb stairs.

⁶Downgraded for relatively few participants and for lack of blinding of participants, personnel and/or outcome assessors in some trials.

Summary of findings 2. FNB (any type) compared with epidural analgesia for knee replacement surgery

FNB (any type) compared with epidural analgesia for knee replacement surgery

Patient or population: patients with knee replacement surgery

Settings: hospital

Intervention: FNB (any type)

Comparison: epidural

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Epidural	FNB (any type)				
Pain at rest at 24 hours Visual analogue scale Scale from zero to 10 Follow-up: median two days	Mean pain at rest at 24 hours ranged across control groups from 0.2 to 3.9 points	Mean pain at rest at 24 hours in the intervention groups was 0.10 lower (0.70 lower to 0.67 higher)		328 (6 studies)	⊕⊕⊕⊖ moderate ¹	SMD -0.05 (-0.43 to 0.32), representing small and non-significant effects. Lower score indicates less pain ²
Pain on movement at 24 hours Visual analogue scale Scale from zero to 10 Follow-up: median two days	Mean pain on movement at 24 hours ranged across control groups from 1.8 to 5.5 points	Mean pain on movement at 24 hours		317 (6 studies)	⊕⊕⊕⊕ high	SMD 0.01 (-0.21 to 0.24), representing small and non-significant effects. Lower score indicates less pain ²

		in the inter- vention groups was 0.02 lower (0.48 lower to 0.54 higher)				
Neurological injury Follow-up: during hospitalization	See comment	See com- ment	Not estimable	306 (4 studies)	See comment	Effect is uncertain as neurolog- ical injuries are extremely rare. Only one event was reported in the epidural group and none in the FNB group
Opioid consumption at 24 hours using morphine equivalents Scale from zero to 150 mg Follow-up: 24 hours	Mean opioid consumption 24 hours ranged across control groups from 8 to 50.7 mg	Mean opioid con- sump- tion at 24 hours in the inter- vention groups was 4.35 lower (9.95 lower to 1.26 higher)		341 (5 studies)	⊕⊕⊕⊕ high	MD -4.35 (-9.95 to 1.26)
Nausea and/or vomiting Follow-up: median two days	Study population ³ 38 per 100 Low ³	24 per 100 (15 to 37)	RR 0.63 (0.41 to 0.97)	183 (4 studies)	⊕⊕⊕⊖ moderate ⁴	

	13 per 100	8 per 100 (5 to 13)			
	High ³				
	67 per 100	42 per 100 (27 to 65)			
Knee flexion range of motion (degrees) ⁵ Follow-up: median three days	Mean knee flexion range of motion ranged across control groups from 48 to 84 degrees	Mean knee flexion in the intervention groups was 1.65 lower (5.14 lower to 1.84 higher)	328 (6 studies)	⊕⊕⊕⊕ high	MD -1.65 (-5.14 to 1.84).
Participant satisfaction with analgesia (point) Scale from zero to 10 Follow-up: during hospitalization	Mean participant satisfaction ranged across control groups from 8.2 to 8.7 points	Mean participant satisfaction in the intervention groups was 1.03 higher (0.39 to 1.66 higher)	120 (2 studies)	⊕⊕⊕⊖ low ⁶	SMD 0.60 (0.23 to 0.97), representing small to moderate effects. Higher score indicates greater satisfaction ²

*The basis for the **assumed risk** (e.g. median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio.

GRADE Working Group grades of evidence.

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.
Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.
Very low quality: We are very uncertain about the estimate.

- ¹Downgraded for significant heterogeneity attributable to inadequate allocation concealment.
²Rule of thumb for interpreting SMD: 0.2 small effect, 0.5 moderate effect, 0.8 large effect (Cohen 1988).
³Assumed risk was based on control group risks in the included studies.
⁴Downgraded for relatively few participants.
⁵Knee flexion of 90 degrees is usually required for mobility independence.
⁶Downgraded for relatively few participants and for lack of blinding of participants, personnel and/or outcome assessors.

Summary of findings 3. FNB (any type) compared with local infiltration analgesia for knee replacement surgery

FNB (any type) compared with local infiltration analgesia for knee replacement surgery

Patient or population: patients with knee replacement surgery
Settings: hospital
Intervention: FNB (any type)
Comparison: local infiltration analgesia

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	Local infiltration analgesia	FNB (any type)				
Pain at rest at 24 hours Visual analogue scale Scale from zero to 10 Follow-up: median two days	Mean pain at rest at 24 hours ranged across control groups from 1.6 to 5 points	Mean pain at rest at 24 hours in the intervention groups was 0.10 higher (1.04 lower to 1.22 higher)		216 (4 studies)	⊕⊕⊕⊖ moderate ¹	SMD 0.06 (-0.61 to 0.72), representing small and non-significant effects. Lower score indicate less pain ³
Pain on movement at 24 hours Visual analogue scale Scale from zero to 10 Follow-up: median two days	Mean pain on movement at 24 hours ranged across control groups from 2.4 to 7 points	Mean pain on movement at 24 hours in the intervention groups was 0.63 higher (0.17 lower to 1.43 higher)		153 (3 studies)	⊕⊕⊖⊖ low ^{1,2}	SMD 0.38 (-0.10 to 0.86), representing

					small and non-significant effects. Lower score indicates less pain ³	
Neurological injury: not reported	See comment	See comment	Not estimable	-	See comment	Effect is uncertain as neurological injuries are extremely rare. No study reported on neurological injury
Opioid consumption at 24 hours using morphine equivalents Scale from zero to 150 mg Follow-up: 24 hours	Mean opioid consumption at 24 hours in the control groups was 26 mg	Mean opioid consumption at 24 hours in the intervention group was 11.50 lower (24.08 lower to 1.08 higher)		40 (1 study)	⊕⊕⊕⊖ moderate ²	MD -11.50 (-24.08 to 1.08)
Nausea and/or vomiting Follow-up: median two days	Study population ⁴		RR 1.71 (0.64 to 4.62)	177 (3 studies)	⊕⊕⊕⊖ moderate ²	
	27 per 100	47 per 100 (18 to 100)				
	Low ⁴					
	6 per 100	10 per 100 (4 to 28)				
	High ⁴					
	55 per 100	94 per 100 (35 to 100)				
Participant satisfaction with analgesia Scale from zero to 10	See comment	See comment	Not estimable	-	See comment	Mean (SD) = 7.8 (0) in FNB and 9

(0) in local infiltration reported in 1 RCT. Unable to generate effect estimate, as SD = 0 for both interventions. Higher scores indicate greater satisfaction

Follow-up not reported

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio.

GRADE Working Group grades of evidence.

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹Downgraded for significant heterogeneity.

²Downgraded for relatively few participants.

³Rule of thumb for interpreting SMD: 0.2 small effect, 0.5 moderate effect, 0.8 large effect (Cohen 1988).

⁴Assumed risk was based on control group risks in the included studies.

Summary of findings 4. Continuous FNB compared with single-shot FNB for knee replacement surgery

Continuous FNB compared with single-shot FNB for knee replacement surgery

Patient or population: patients with knee replacement surgery

Settings: hospital

Intervention: continuous FNB

Comparison: single-shot FNB

Outcomes	Illustrative comparative risks* (95% CI)	Relative effect	No. of participants	Quality of the evidence	Comments
----------	--	-----------------	---------------------	-------------------------	----------

	Assumed risk	Corresponding risk	(95% CI)	(studies)	(GRADE)	
	Single-shotFNB	Continuous FNB				
Pain at rest at 24 hours Visual analogue scale Scale from zero to 10 Follow-up: median two days	Mean pain at rest at 24 hours ranged across control groups from 0.8 to 3.9 points	Mean pain at rest at 24 hours in the intervention group was 0.95 lower (1.78 to 0.11 lower)		272 (4 studies)	⊕⊕⊕⊖ moderate ¹	SMD -0.62 (-1.17 to -0.07), representing small to moderate effects. Lower score indicates less pain ²
Pain on movement at 24 hours Visual analogue scale Scale from zero to 10 Follow-up: median two days	Mean pain on movement at 24 hours ranged across control groups from 4.5 to 7.2 points	Mean pain on movement at 24 hours was 0.74 lower (1.18 to 0.30 lower)		272 (4 studies)	⊕⊕⊕⊕ high	SMD -0.42 (-0.67 to -0.17), representing small effects. Lower score indicates less pain ²
Neurological injury: not reported	See comment	See comment	Not estimable	-	See comment	Effect is uncertain as neurological injuries are extremely rare. No events were reported in two trials
Opioid consumption at 24 hours using morphine equivalents Scale from 0 to 150 mg Follow-up: 24 hours	Mean opioid consumption at 24 hours ranged across control groups from 14 to 42 mg	Mean opioid consumption at 24 hours in the intervention group was 13.81 lower (23.27 to 4.35 lower)		236 (3 studies)	⊕⊕⊕⊖ moderate ¹	MD -13.81 (-23.27 to -4.35)
Nausea and/or vomiting	Study population ³		RR 0.55 (0.2 to 1.56)	214 (2 studies)	⊕⊕⊕⊖ moderate ¹	
	36 per 100	20 per 100				

Follow-up: median two days	(7 to 56)					
	Low ³					
	33 per 100	18 per 100	(7 to 51)			
	High ³					
	45 per 100	25 per 100	(9 to 70)			
Participant satisfaction with analgesia: not reported	See comment	See comment	Not estimable	-	See comment	Effect is uncertain, as no study reported on this outcome

*The basis for the **assumed risk** (e.g. median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio.

GRADE Working Group grades of evidence.

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹Downgraded for significant heterogeneity.

²Rule of thumb for interpreting SMD: 0.2 small effect, 0.5 moderate effect, 0.8 large effect (Cohen 1988).

³Assumed risk was based on control group risks in the included studies.

BACKGROUND

Description of the condition

Total knee replacement (TKR) is a common orthopaedic operation, consisting of replacing diseased or damaged knee joint surfaces to relieve the pain and disability of osteoarthritis. The significant public health burden of osteoarthritis is reflected in the findings that nearly half of all adults in the United States are at risk of developing symptomatic knee osteoarthritis by 85 years of age (Murphy 2008). The number of knee replacements undertaken annually around the world will increase exponentially as the population ages and becomes more obese. Indeed, the number of primary TKR procedures in the United States was projected to increase almost eight-fold, from 450,000 in 2005 to 3.48 million in 2030 (Kurtz 2007).

TKR is one of the most painful surgical procedures. Effective analgesia in the immediate postoperative phase is important to allow the patient to exercise and regain mobility, thereby facilitating recovery and decreasing the length of hospital stay (Capdevila 1999; Chelly 2001). Unrelieved severe postoperative pain can result in pathophysiological responses causing adverse postsurgical outcomes. These adverse outcomes have medical and economic implications, such as impaired early rehabilitation, delayed discharge, unscheduled rehospitalization, impaired health-related quality of life and increased risk of chronic pain (American Society of Anesthesiologists 2004; Carr 1999; Sinatra 2009; Twersky 1997).

Various analgesic techniques are used to relieve postoperative pain following TKR. Analgesic options include patient-controlled analgesia (PCA) using opioids, epidural analgesia, local infiltration analgesia and femoral nerve block (FNB) with local anaesthetic agents. PCA opioids have been associated with significant adverse effects such as respiratory depression, nausea, urinary retention and constipation (Chelly 2001; Sinatra 2009). Epidural analgesia, when given simultaneously with an anticoagulant, has been associated with spinal epidural haematoma as well as hypotension, urinary retention and pruritus (Capdevila 1999; Choi 2003; Sinatra 2009). Epidural analgesia also causes bilateral motor blockade to the same extent, which may interfere with early mobilization (Barrington 2005). Some randomized controlled trials (RCTs) have suggested that FNB provides better pain control and fewer opioid-related adverse effects compared with PCA opioid or epidural analgesia (Chelly 2001; Sundarathiti 2009; Szczukowski 2004; Wang 2002). Local infiltration analgesia is an emerging technique. The advantage of local infiltration analgesia is that pain conduction is blocked at its origin. A study evaluating the efficacy of a perioperative local infiltration analgesia consisting of local anaesthetic and morphine found a significant reduction in pain and opioid consumption during the first postoperative 48 hours compared with PCA morphine (Vendittoli 2006). To date, no systematic review has compared the effects of FNB with those of local infiltration analgesia.

Description of the intervention

Femoral, psoas (lumbar plexus) and fascia iliaca blocks have been used as postoperative analgesic techniques for TKR surgery. Among these nerve blocks, the FNB is used most commonly. An FNB may be given as a single shot or as a continuous block via a catheter and an infusion.

One method that is commonly used to deliver an FNB is the Winnie paravascular technique (Winnie 1973), in which a peripheral nerve stimulator or ultrasound guidance is often used to locate the nerve. When a peripheral nerve stimulator needle is used, the common femoral artery is palpated first with the patient in the supine position, and the stimulating needle is then inserted at the inguinal crease, approximately 1 cm lateral to the femoral artery pulse. Stimulation of the femoral nerve generates contraction of the quadriceps muscle. The needle position is optimized when contractions persist at an output of 0.3 mA to 0.5 mA. Approximately 20 ml to 30 ml of local anaesthetic is then injected. When using ultrasound, the operator places the transducer in the inguinal crease to locate the hyperechoic femoral nerve, which can be visualized lateral to the hypoechoic pulsative common femoral artery. Successful FNB is attained when the local anaesthetic is seen circumferentially around the nerve (Ballantyne 2010).

An FNB can also be performed as part of a more extensive nerve block, termed the *three-in-one block* (femoral, lateral femoral cutaneous and obturator nerves). In the three-in-one block, a greater volume of local anaesthetic is used, and pressure is applied just distal to the needle during administration of the local anaesthetic to help spread the local anaesthetic to the lateral femoral cutaneous and obturator nerves. However, the femoral nerve is the only nerve that is consistently blocked (Lang 1993). Consequently, the three-in-one block frequently is referred to simply as the FNB (Enneking 2009).

How the intervention might work

An FNB blocks sensation to the anteromedial aspect of the knee, thus reducing pain and muscle spasms. Compared with a single shot, continuous FNB provides a longer duration of postoperative analgesia (Sinatra 2009) with a lower concentration of local anaesthetic. These features may allow earlier rehabilitation with a continuous FNB, compared with a single-shot FNB, in that the degree of motor block is reduced. An FNB does not block sensation to the posterior aspect of the knee, as it is innervated by the sciatic nerve. To improve postoperative analgesia, a sciatic and/or obturator nerve blockade is sometimes added to the FNB (McNamee 2002; Morin 2005).

Why it is important to do this review

Traditionally, PCA opioid and epidural analgesia were the postoperative analgesic methods of choice following TKR. Recent years have seen a growing interest in the use of FNB to minimize the adverse effects associated with PCA opioid or epidural analgesic techniques. However, FNB after TKR surgery is not without risk; concerns about prolonged quadriceps weakness (Kandasami 2009) and complications such as femoral neuropathy or neuritis have been reported; an increased risk of falls has also been associated with continuous FNB, compared with single-shot block or no block (Feibel 2009; Johnson 2013; Sharma 2010). Additionally, continuous FNB catheters require specialized skills and additional time for insertion and management.

A sciatic nerve block is sometimes combined with an FNB to improve analgesia (Morin 2005). However, this combination could lead to complications such as increased risk of falls and heel ulceration. The combination could also mask peroneal nerve injury or an evolving sciatic nerve injury from compartment syndrome (Ben-David 2004; Kadic 2009; Todkar 2005). A selective obturator

nerve block is added at times to an FNB, but the additional analgesic benefit is conflicting (Macalou 2004; McNamee 2002).

To date, no strong evidence has been obtained from large RCTs evaluating the comparative efficacy and safety of FNB versus other forms of postoperative analgesia. A recent systematic review limited to English language publications up to 2009 concluded that single-shot or continuous FNB (with PCA opioid) was superior to PCA opioid alone for acute pain control in the first 72 hours after knee replacement (Paul 2010). Another systematic review compared nerve blockade with epidural analgesia and found no significant difference in pain in the first 24 hours post operation (Fowler 2008). Our work will extend that of these reviews by looking at all comparator analgesic regimens, including the emerging analgesic technique of local infiltration analgesia; by examining outcomes beyond the initial postoperative period and by not limiting our search to English language publications. The rapidly rising number of TKRs performed annually worldwide will considerably increase the burden on healthcare resources (March 2004). A systematic review is needed to evaluate current evidence for the short- and long-term safety and efficacy of FNB after TKR.

OBJECTIVES

To evaluate the benefits and risks of FNB used as a postoperative analgesic technique relative to other analgesic techniques among adults undergoing TKR.

METHODS

Criteria for considering studies for this review

Types of studies

We included RCTs comparing FNB (inserted preoperatively, operatively or postoperatively) versus postoperative analgesic techniques not requiring an FNB (intravenous PCA opioids, epidural analgesia, local infiltration analgesia and oral analgesia).

We excluded quasi-randomized trials (e.g. using alternation) and observational studies.

Types of participants

We included adults who have undergone TKR surgery.

Types of interventions

FNB involving groin injections of any type (performed in isolation or as part of a three-in-one block) used to provide postoperative analgesia after TKR surgery versus no FNB. We also included RCTs that compared continuous versus single-shot FNB.

Types of outcome measures

In this review, we intend to examine the following outcomes across a range of potential alternatives to FNB. We will present sequentially all outcomes for each comparison with FNB as follows: (1) PCA opioid; (2) epidural analgesia; (3) local infiltration analgesia and (4) oral analgesia; we will also compare two types of FNB: (5) continuous FNB versus (6) single-shot FNB.

Primary outcomes

1. Pain at rest and on movement

We considered pain at rest and on movement within the following postoperative time frames: first two hours, three to 12 hours, 24 hours, 48 hours, 72 hours and more than 72 hours. Pain outcomes were based on measures with reliable and validated psychometric properties such as the visual analogue scale (VAS). We converted data to a zero to 10 scale. Unless otherwise reported, pain was assumed to be experienced at rest.

2. Serious adverse events

We defined a serious adverse event as any untoward occurrence that results in death, is life threatening, requires rehospitalization or prolongation of existing hospitalisation, results in persistent or significant disability or is considered a medically important event or reaction. Examples include neurological injury, postoperative falls and thrombotic events. We used the time frame set by the study authors.

Secondary outcomes

3. Proportion of participants in significant pain postoperatively, as defined by the study authors

4. Time from end of surgery to first rescue analgesic request

5. Opioid consumption

We converted intravenous fentanyl and oral oxycodone to intravenous morphine dosing equivalents using the following computations: 1 mg oral oxycodone = 0.6 mg intravenous morphine; and 1 µg intravenous fentanyl = 0.067 mg intravenous morphine (Allman 2006; Berdine 2006). Data on opioid consumption were expected to be skewed in distribution, and it is recommended that opioid consumption be dichotomized (Moore 2011; PaPaS document 2011). However, opioid consumption was not reported as above or below a certain threshold. Instead, the included studies reported opioid consumption as mean (standard deviation (SD)) milligrams, and we had to follow suit. Some studies reported opioid consumption as cumulative consumption since its commencement, whereas others used consumption over 24 hours. When available, we analysed total cumulative consumption since commencement.

6. Adverse effects

a. Nausea or vomiting, or both, in the first 72 hours. If more than one time point was reported, we chose the data closest to the first 24 hours, when the adverse effects were most pronounced. When both nausea and vomiting were reported, we used the data for vomiting, with the assumption that all participants who vomited would have also experienced nausea. When nausea and vomiting were reported as mild, moderate or severe, the numbers experiencing severe nausea or vomiting were used.

b. Sedation in the first 72 hours using sedation score of at least one on a four-point scale where zero = alert; one = drowsy; two = sleeping, easy to arouse and three = sleeping, difficult to arouse; or using the author's definition of event/no event.

c. Urinary retention requiring catheterization or using the author's definition of event/no event in the first 72 hours.

d. Technical failure of the blocks.

7. Physical function

- a. Knee flexion range of motion during the first four postoperative days.
- b. Knee extension range of motion during the first four postoperative days.
- c. Time to first ambulation during the first four postoperative days.

When more than one time point was reported for range of motion, we chose the time point closest to the fourth day, when ambulation ability is more critical with impending hospital discharge.

8. Participant satisfaction with analgesia during the hospital stay

Participant satisfaction was reported on a continuous scale or as dichotomous data (i.e. number of participants in a study satisfied with treatment). We normalized continuous data to a zero to 10 scale.

Search methods for identification of studies

Electronic searches

We searched the Cochrane Central Register of Controlled Trials (CENTRAL) 2013, Issue 1 (see [Appendix 1](#)); MEDLINE (Ovid SP) (1948 to January 2013) (see [Appendix 2](#)); EMBASE (Ovid SP) (1980 to January 2013) (see [Appendix 3](#)); CINAHL (EBSCO host) (1982 to January 2013) (see [Appendix 4](#)); ISI Web of Science (1973 to January 2013) (see [Appendix 5](#)) and dissertation abstracts.

We combined the sensitive strategies suggested in Section 6.4 of the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)) to search for trials in MEDLINE. We adopted the search strategy for MEDLINE to search all other databases. The search strategies used are reported in Appendices 1 to 5. We applied no language restriction.

Searching other resources

One review author (EC) reviewed the references of relevant articles and textbooks for additional citations. We also searched trial registers (www.clinicaltrials.gov, www.controlled-trials.com and <http://www.anzctr.org.au>), Google Scholar and the Procedure Specific Postoperative Pain Management (PROSPECT) Website (www.postoppain.org) to ensure that significant papers were not missed. The date of the last search was 31 January 2013.

Data collection and analysis

Selection of studies

Two review authors (EC and MF) independently screened for eligibility the titles and abstracts of publications identified in the literature search. We obtained and assessed the full published manuscripts of clinical trials that appeared to be eligible to assess their relevance on the basis of prespecified inclusion criteria. Each of these review authors documented the reasons for study exclusion. The fifth review author (NC) resolved disagreements regarding study exclusion. All review authors have participated in trials that could potentially be eligible for this review (see [Declarations of interest](#)). To prevent conflict of interest, two independent reviewers (MH and FP; see [Acknowledgements](#)) determined the eligibility of these trials.

A copy of the 'Study selection form' is provided in [Appendix 6](#).

Data extraction and management

Two review authors (EC and MF) independently performed data extraction using a data extraction form (see [Appendix 7](#)). The fifth review author (NC) resolved disagreements. Two independent reviewers extracted the data of an included trial when the review authors were among the study investigators. We contacted trial authors to ask for additional details of their studies.

Assessment of risk of bias in included studies

Two review authors (EC and MF) independently assessed the methodological quality of eligible trials using the tool stated in Chapter 8 of the *Cochrane Handbook for Systematic Reviews of Interventions* ([Higgins 2011](#)) (see [Appendix 8](#)). We resolved disagreements by discussion with the fifth review author (NC). Two independent reviewers assessed the risk of bias of included trials when the review authors were among the study investigators.

In cases of insufficient reporting to enable judgement of 'low risk' or 'high risk,' we contacted the respective study authors to ask for more information. We rated the method as 'unclear' if we could not reach the study authors.

We conducted sensitivity analyses to determine the effect of excluding studies considered to be at high or unclear risk of bias. The results of studies with high and unclear risk of bias were not considered when the results of our sensitivity analyses were interpreted, although we presented them in the meta-analyses for completeness.

1. Random sequence generation

We considered random sequence generation as low risk if it was generated by using a computer, by using a random number table algorithm or by drawing lots or shuffling cards.

2. Concealment of allocation

We considered allocation concealment as low risk if participant recruiters, investigators and participants were unable to anticipate treatment assignment. Adequate methods included a central allocation system (telephone, Web-based or pharmacy controlled randomisation) or sequentially numbered sealed opaque envelopes.

3. Blinding of participants and personnel

We considered blinding as low risk of bias if:

- a. participants and personnel (healthcare providers) were blinded to the allocated intervention; and
- b. it was unlikely that the blinding could have been broken.

We considered blinding as high risk if:

- a. blinding of participants and personnel was attempted, but it was likely that the blinding could have been broken; and
- b. no blinding or incomplete blinding was applied.

4. Blinding of outcome assessment

We considered blinding as low risk if:

a. outcome assessors were blinded to the allocated intervention, and it was unlikely that the blinding could have been broken.

We considered blinding as high risk if:

a. outcome assessors were blinded, but it was likely that the blinding could have been broken; and

b. outcome assessors were not blinded.

5. Incomplete outcome data

We considered a study as having low risk of incomplete outcome data if:

a. no outcome data were missing;

b. the proportion of missing outcomes compared with observed event risks was not enough to have a clinically relevant impact on the intervention effect estimate (i.e. the percentage of missing data was small relative to the incidence proportion of the outcome);

c. reasons for missing outcome data were unlikely to be related to true outcomes and the percentage of missing data was small; and

d. missing outcome data were balanced in quantities across intervention groups with similar reasons for missing data across groups, and the percentage of missing data was small.

A study was considered as having high risk of incomplete outcome data if:

a. reasons for missing outcome data were likely to be related to true outcomes with imbalance in numbers or reasons for missing data across intervention groups;

b. the proportion of missing outcomes compared with the proportion of observed events was enough to induce clinically relevant bias in intervention effect estimates; and

c. 'as treated' analysis was performed with substantial differences between the numbers of participants contributing to the analysis and the numbers of participants randomly assigned.

6. Selective reporting

We considered low risk of selective reporting if all of the a priori outcomes of a study that are of interest in the review were reported in the prespecified way.

We considered high risk of selective reporting for any of the following.

a. Not all of the a priori outcomes of a study were reported.

b. One or more primary outcomes were reported using only measurements, analysis methods or subsets of data that were not prespecified.

c. One or more reported primary outcomes were not prespecified, unless clear justification of their reporting was provided.

d. One or more outcomes of interest in the review were reported incompletely and therefore could not be included in a meta-analysis.

7. Other sources of bias

We also assessed whether the study used intention-to-treat (ITT) analysis methods (Hollis 1999) for the primary outcomes.

Measures of treatment effect

We calculated risk ratios (RRs) with 95% confidence intervals (CIs) for dichotomous outcomes and mean differences (MDs) or standardized mean differences (SMDs) with 95% CIs for continuous outcomes. For comparisons based on continuous outcomes consisting of trials that reported only differences between groups, we converted the data for all relevant trials to estimate the intervention effect and its corresponding standard error (Borenstein 2009) and used generic inverse variance outcome type. A similar approach was used for comparisons including RCTs that reported a zero SD in one of the intervention arms.

For continuous outcomes, we interpreted intervention effects based on SMDs according to the rule of thumb where 0.2 or smaller represents a small effect, 0.5 a moderate effect and 0.8 or larger a large effect (Higgins 2011; see Section 12.6.2). To further assist in interpreting SMDs, we reexpressed them in their original units by multiplying the pooled SMD with the pooled SD of the control group differences (Higgins 2011; see Section 12.6.4).

SMD is a more sensible measure for synthesizing pain outcomes because one cannot verify whether 2-cm changes in VAS are equivalent across the scale. Thus the combination of VAS results that have sampled different parts of the scale might be misrepresented by combining MDs and might be better represented by using SMDs. We have used SMDs to report the outcome of participant satisfaction (measured using scale zero to 10) for the same reason..

We calculated number needed to treat for an additional beneficial outcome (NNTB) or an additional harmful outcome (NNTH) as and where appropriate. We obtained confidence intervals for NNTB/NNTH by inverting and exchanging the confidence limits for the absolute risk reduction (ARR). We used the Wilson score method to calculate the confidence limits of ARR and NNTB/NNTH (Bender 2001; Wilson 1927).

Unit of analysis issues

Some RCTs had three or four allocation groups. When the study contributed several independent comparisons (e.g. FNB vs PCA and FNB vs epidural), the interventions were analysed separately in the appropriate meta-analyses as if they were from different studies.

When the RCTs contributed several correlated comparisons, with similar type of FNB (e.g. continuous FNB with ropivacaine vs PCA and continuous FNB with bupivacaine vs PCA), to the same meta-analysis, we combined the groups to create a single pairwise comparison, as recommended in Chapter 16 of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011).

If the RCTs contributed several correlated comparisons, with different types of FNB (e.g. single-shot FNB vs PCA and continuous FNB vs PCA) to a meta-analysis, we split the control group into two groups (shared) and included two 'reasonably independent' comparisons. This was done to address the issues of double-counting and unit of analysis errors in the control group (Higgins 2011; see Chapter 16).

Dealing with missing data

We contacted the study authors if any required data were missing or unclear.

We calculated the missing estimates of SDs from other relevant statistics reported in the study (e.g. standard error, 95% CI, P value). If these statistics were reported, the missing SDs were imputed according to the recommendations of the *Cochrane Handbook for Systematic Reviews of Interventions* (Higgins 2011; see Chapter 16) using the RevMan calculator. Some studies presented results using a combination of median, interquartile range (25th and 75th quartiles) and/or range (minimum and maximum values). We calculated corresponding means and SDs from the presented results based on the recommendation of Hozo 2005. The required data were estimated as follows.

Mean = (Minimum + 2 * Median + Maximum)/4, if median and range were reported for each group; otherwise, Mean = Median.

SD = SQRT (1/12(((Min - 2 * Median + Max)²/4 + (Min - Max)²)), if group size was less than 15 and median and range were reported for each group.

SD = SQRT(((Min - Max)/4)), if the range was reported and the group size was greater than 15 but equal to or less than 70.

SD = SQRT((Min - Max)/6), if the range was reported and the group size was greater than 70.

SD = (75th Quartile - 25th Quartile)/1.35, if the interquartile range was reported.

Assessment of heterogeneity

We assessed heterogeneity of included RCTs on the basis of their clinical and methodological diversity (risk of bias assessment). Our a priori hypothesis for sources of clinical heterogeneity were as follows.

1. Different analgesic regimens (e.g. FNB as single-shot or continuous infusion, using different types of local anaesthetics, concentrations, infusion rates and timings of injections; and with or without sciatic or obturator nerve blocks).
2. Different standard co-analgesic regimens.

We presented primary analyses using the random-effects model to account for the impact of anticipated clinical and methodological heterogeneity. We assessed statistical heterogeneity using the Chi² test for heterogeneity and quantified heterogeneity using the I² statistic (Higgins 2011). We considered values of I² greater than 50% to represent significant between-study heterogeneity (Higgins 2003). If significant heterogeneity was demonstrated, we explored the data to test whether our planned subgroup analyses explained this heterogeneity.

Assessment of reporting biases

We used Orwin's fail-safe N test to evaluate the impact of potential publication bias on the robustness of the overall observed analgesic effect (Orwin 1983). We determined how many missing studies without an intervention effect (magnitude of SMD ≤ 0.01) would render a significant overall or pooled effect non-significant or trivial. The trivial effect was defined by an SMD value of 0.2 (Orwin 1983).

Data synthesis

We conducted a meta-analysis using Review Manager 5.2 (RevMan 5.2) when data from two or more RCTs were sufficient. We used the random-effects model of DerSimonian and Laird, as we anticipated that heterogeneity would be present in the interventions and in the outcomes (DerSimonian 1986).

VAS results that sampled different parts of the scale were combined using SMDs. As the dichotomous data for opioid consumption were not available, the means of opioid consumption were synthesized using SMDs. When it was not possible to conduct a meta-analysis, we discussed the results narratively.

Primary analyses include the following.

1. Single-shot or continuous FNB (± sciatic/obturator block ± PCA opioid) versus PCA opioid.
2. Single-shot or continuous FNB (± sciatic/obturator block) versus epidural analgesia.
3. Single-shot or continuous FNB versus local infiltration analgesia.
4. Single-shot or continuous FNB versus oral analgesia.
5. Continuous versus single-shot FNB.

We stratified the meta-analyses of included RCTs according to type of FNB (single-shot FNB, single-shot FNB + sciatic block, single-shot FNB + obturator block, continuous FNB, continuous FNB + sciatic block). For subgroup and sensitivity analyses, we combined the various types of FNB.

We did not adjust for multiplicity of the multiple primary analyses as, in general, this is not recommended (see Section 16.7.2 of the *Cochrane Handbook for Systematic Reviews of Interventions*) (Higgins 2011).

Subgroup analysis and investigation of heterogeneity

We limited subgroup and sensitivity analyses to the outcomes of pain at rest and on movement at 24 hours.

We performed the following subgroup analyses.

1. Type of FNB (i.e. single-shot FNB and continuous FNB, with or without an additional sciatic/obturator nerve block).
2. FNB (any type) with or without a concurrent parenteral opioid.
3. Type of local anaesthetic (i.e. FNB (any type) using ropivacaine and FNB (any type) using bupivacaine).

Sensitivity analysis

We performed sensitivity analyses to evaluate the effect on the overall primary result of removing trials with low and unclear methodological quality (i.e. allocation concealment, as well as blinding of participants, personnel and outcome assessors). We did not conduct sensitivity analyses to determine the effect of missing data imputation, as no substantial missing data issues were identified in the included RCTs. Also, sensitivity analysis based on multiple interventions within studies was not performed because we followed the recommendations provided in the *Cochrane Handbook for Systematic Reviews of Interventions* (see Chapter 16; Higgins 2011), as discussed in the Unit of analysis issues section.

Summary of findings tables

We used the principles of the GRADE system ([Guyatt 2008](#)) to assess the quality of the body of evidence associated with specific outcomes (pain at rest at 24 hours, pain on movement at 24 hours, neurological injury, opioid consumption at 24 hours, nausea and/or vomiting, knee flexion range of motion, participant satisfaction with analgesia) in our review, and we constructed a 'Summary of findings' (SoF) table using the GRADE software (see [Appendix 9](#)). The GRADE approach appraises the quality of a body of evidence based on the extent to which one can be confident that an estimate of effect or association reflects the item being assessed. The quality of a body of evidence considers within-study risk of bias (methodological quality), directness of the evidence, heterogeneity of the data, precision of effect estimates and risk of publication bias (see Chapter 12) ([Higgins 2011](#)).

RESULTS

Description of studies

See [Characteristics of included studies](#) and [Characteristics of excluded studies](#).

Results of the search

[Figure 1](#) shows the results of the literature search. We identified 87 publications for potential inclusion, of which we excluded 36 after reviewing the full-text reports and assigned as awaiting assessment to four conference abstracts. We identified 45 eligible RCTs from 47 publications (two sets of publications reported the same randomized trials ([Bergeron 2009](#) and [Kardash 2007](#); [Seet 2006](#) and [Shum 2009](#)). We managed to contact 21 authors of the included studies to clarify the study methodology or to obtain additional data (three of the authors no longer have the original data).

Figure 1. Study flow diagram.

Included studies

We included 45 RCTs with a total of 2710 participants. Details of the individual RCTs are provided in the [Characteristics of included studies](#) table. Three RCTs were reported in Chinese (Tang 2010; Wang 2010; Yu 2010), and one RCT was published in German (Fritze 2009). Two RCTs (Chan 2013; Widmer 2012), for which some of the review authors were named, were judged eligible for inclusion by two independent reviewers (MH and FP). In two RCTs (Martin 2008; Mistraletti 2006), the effects of FNB on inflammation and postoperative suppression of gluconeogenesis, respectively, were the main focus of the studies.

The sample size of the individual RCTs ranged from 18 (Ng 2012) to 200 (Chan 2013) participants, the age of study participants ranged from 29 to 86 years (Affas 2011) and the proportion of female participants ranged from 38% (Good 2007) to 100% (Park 2010).

Of the included RCTs, 20 (Adams 2002; Allen 1998; Chan 2012; Chan 2013; de Lima e Souza 2008; Fritze 2009; Ganapathy 1999; Hirst 1996; Hunt 2009; Kaloul 2004; Kardash 2007; Macalou 2004; McNamee 2001; Mistraletti 2006; Ng 2001; Park 2010; Seet 2006; Singelyn 1998; Tugay 2006; Xie 2012) evaluated more than two treatment groups (see [Unit of analysis issues](#) section).

In three of these RCTs, one of the treatment groups was excluded from the review because it was a non-randomized group (Hunt 2009) or because the treatment did not meet the review's inclusion criteria (psoas block group in Kaloul 2004; obturator block group in Kardash 2007).

The included RCTs made the following comparisons according to the aims of this review.

1. FNB (with or without PCA opioid) versus PCA opioid (29 RCTs with two RCTs having three allocation groups)

Of the included RCTs, 14 (Adams 2002; Allen 1998; Chan 2012; Chan 2013; Good 2007; Hirst 1996; Hunt 2009; Kardash 2007; Macalou 2004; Ng 2001; Ozen 2006; Szczukowski 2004; Tugay 2006; Wang 2002) compared single-shot FNB versus PCA.

Two RCTs (Allen 1998; McNamee 2001) compared single-shot FNB + sciatic block versus PCA, and one RCT (Macalou 2004) compared single-shot FNB + obturator versus PCA.

Of the included RCTs, 12 (Baranovic 2011; Chan 2013; Ganapathy 1999; Hirst 1996; Kadic 2009; Kaloul 2004; Seet 2006; Serpell 2001; Singelyn 1998; Tang 2010; Wang 2010; Yu 2010) compared continuous FNB versus PCA.

Two RCTs (Martin 2008; Mistraretti 2006) compared continuous FNB + sciatic versus PCA.

All except six RCTs (Adams 2002; Baranovic 2011; Chan 2013; Mistraretti 2006; Singelyn 1998; Yu 2010) had a concurrent PCA opioid in the FNB intervention.

2. FNB versus epidural analgesia (10 RCTs with one RCT contributing two comparisons)

One RCT (Adams 2002) compared single-shot FNB versus epidural, and another RCT (Davies 2004) compared single-shot FNB + sciatic versus epidural. One RCT (Lee 2011) compared single-shot FNB + epidural versus epidural; the results of this study were not pooled, as a concurrent epidural was included in the FNB intervention.

Five RCTs (Barrington 2005; Fritze 2009; Long 2006; Singelyn 1998; Sundarathiti 2009) compared continuous FNB versus epidural.

Three RCTs (Fritze 2009; Mistraretti 2006; Zaric 2006) compared continuous FNB + sciatic versus epidural.

3. FNB versus local infiltration analgesia (six RCTs)

One RCT (Parvataneni 2007) compared single-shot FNB versus local infiltration analgesia.

Three RCTs (Affas 2011; Carli 2010; Toftdahl 2007) compared continuous FNB versus local infiltration analgesia.

One RCT (Widmer 2012) compared single-shot FNB + local infiltration analgesia versus local infiltration analgesia alone; the results of this study were not pooled, as a concurrent local infiltration analgesia was included in the FNB intervention.

One cross-over RCT (Ng 2012) with knees as the unit of comparison compared continuous FNB versus local infiltration analgesia. The data from this RCT were not included in the meta-analysis, as we were uncertain about the cross-over effects and issues with analysis.

4. FNB versus oral analgesia (one RCT)

One RCT (Nader 2012) compared continuous FNB + oral analgesia versus oral analgesia.

5. Continuous FNB versus single-shot FNB (four RCTs)

Four RCTs (Chan 2013; Hirst 1996; Park 2010; Salinas 2006) made the comparison between continuous and single-shot FNB. Three of these studies (Hirst 1996; Park 2010; Salinas 2006) had a concurrent PCA opioid intervention in both the continuous and single-shot FNB groups, and one study (Chan 2013) had a concurrent PCA opioid intervention only in the single-shot FNB group.

Excluded studies

We excluded 36 trials for the reasons given in the [Characteristics of excluded studies](#). We assigned four trials reported as conference abstracts as awaiting assessment as they do not contain enough information to allow determination of eligibility, or they do not provide quantifiable outcome data ([Characteristics of studies awaiting classification](#)).

Risk of bias in included studies

The risk of bias assessment for the individual RCTs is shown in the 'Risk of bias' graph and the 'Risk of bias' summary in [Figure 2](#) and [Figure 3](#), respectively. Details are provided in the risk of bias tables in the [Characteristics of included studies](#).

Figure 2. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Adams 2002	+	?	-	-	+	+	+
Affas 2011	+	+	-	-	+	+	+
Allen 1998	?	?	+	+	+	+	+
Baranovic 2011	+	?	?	?	-	+	-
Barrington 2005	+	+	-	-	+	+	+
Carli 2010	+	+	+	+	+	+	+
Chan 2012	+	?	+	+	+	+	+
Chan 2013	+	+	-	?	+	+	+
Davies 2004	+	+	?	-	+	+	+
de Lima e Souza 2008	+	+	+	+	+	+	+
Fritze 2009	?	+	?	-	+	+	+
Ganapathy 1999	+	+	+	+	+	+	+
Good 2007	+	+	+	+	+	+	+
Hirst 1996	?	?	+	+	+	+	+
Hunt 2009	+	+	+	+	+	+	+
Kadic 2009	?	+	-	-	+	+	+
Kaloul 2004	+	?	?	-	+	+	+
Kardash 2007	+	+	+	+	+	+	+
Lee 2011	?	+	-	?	+	+	+
Long 2006	?	?	?	?	?	+	+

Figure 2. (Continued)

Long 2006	?	?	?	?	?	+	+
Macalou 2004	?	?	-	+	+	+	+
Martin 2008	+	+	-	-	+	+	+
McNamee 2001	?	+	+	+	+	+	+
Mistralletti 2006	+	?	?	-	+	+	+
Nader 2012	+	+	-	-	+	+	+
Ng 2001	?	?	+	+	+	+	+
Ng 2012	+	?	+	+	+	+	+
Ozen 2006	+	+	?	+	+	+	+
Park 2010	+	+	?	?	?	+	+
Parvataneni 2007	?	?	?	+	+	+	+
Salinas 2006	+	+	-	-	+	+	+
Seet 2006	+	+	-	-	?	+	+
Serpell 2001	?	?	-	-	+	+	+
Singelyn 1998	+	?	-	-	+	+	+
Sundarathiti 2009	+	?	-	-	+	+	+
Szczukowski 2004	+	+	+	+	+	+	+
Tang 2010	+	+	-	-	+	+	+
Toftdahl 2007	+	+	-	-	+	+	+
Tugay 2006	+	+	+	+	+	+	+
Wang 2002	+	+	+	+	+	+	+
Wang 2010	+	?	?	?	+	+	+
Widmer 2012	+	+	+	+	+	+	+
Xie 2012	?	+	+	+	?	+	+
Yu 2010	+	?	?	?	+	+	+
Zaric 2006	+	+	-	-	-	+	?

Figure 3. Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

Allocation

All studies stated that the participants were randomly allocated to treatment groups. The method of randomization was judged to have a low risk of bias for 33 (73%) RCTs and unclear risk for 12 (27%) RCTs, as the methods were not stated and we were unable to reach the study authors for details.

Concealment of allocation was adequate in 28 (62%) and unclear in 17 (38%) included RCTs. Concealment using sealed envelopes without mention of whether they were sequentially numbered and opaque was rated unclear.

Blinding

Blinding of participants and personnel was judged to be at low risk of bias for 17 (38%) RCTs, high risk of bias for 17 (38%) RCTs and unclear risk of bias for 11 (24%) RCTs. Twenty RCTs (44%) reported using blinded outcome assessors, 18 (40%) RCTs did not blind outcome assessors and seven (16%) RCTs were judged as having unclear risk, as they provided no information on assessor blinding.

Incomplete outcome data

The short follow-up period (first 72 hours post operation) in most of the RCTs reduced the risk of loss to follow-up. Hence, attrition bias was low in all except four (9%) RCTs (Long 2006; Park 2010; Seet 2006; Xie 2012), which were rated as unclear, as no information on post randomization exclusions was provided. Two (4%) RCTs (Baranovic 2011; Zaric 2006) were rated as high risk, as 18% to 20% post randomization exclusions were determined. See [Characteristics of included studies](#).

Selective reporting

We judged that selective reporting bias was avoided by the reporting of results for all outcomes listed in the methods section

and by the provision of additional data on request. The risk of reporting bias was low to moderate for the included RCTs.

Other potential sources of bias

All RCTs except two (4%) (Baranovic 2011; Zaric 2006) had a low risk of bias for intention-to-treat analysis, as the number of participants lost to follow-up for the outcomes of interest in this review was very small.

Effects of interventions

See: [Summary of findings for the main comparison FNB \(any type\) compared with PCA opioid for knee replacement surgery](#); [Summary of findings 2 FNB \(any type\) compared with epidural analgesia for knee replacement surgery](#); [Summary of findings 3 FNB \(any type\) compared with local infiltration analgesia for knee replacement surgery](#); [Summary of findings 4 Continuous FNB compared with single-shot FNB for knee replacement surgery](#)

See [Data and analyses](#); [Summary of findings for the main comparison \(FNB with or without PCA opioid vs PCA opioid\)](#); [Summary of findings 2 \(FNB vs epidural\)](#); [Summary of findings 3 \(FNB vs local infiltration analgesia\)](#) and [Summary of findings 4 \(continuous vs single-shot FNB\)](#).

For pain and participant satisfaction outcomes, we interpreted intervention effects based on SMDs according to the rule of thumb where 0.2 or smaller represents a small effect, 0.5 a moderate effect and 0.8 or larger a large effect (Higgins 2011; see Section 12.6.2).

I. FNB (with or without PCA opioid) versus PCA opioid

Primary outcomes

1. Pain at rest and on movement

(See [Analysis 1.1](#); [Analysis 1.2](#); [Analysis 1.3](#); [Analysis 1.4](#); [Analysis 1.5](#); [Analysis 1.6](#); [Analysis 1.7](#); [Figure 4](#); [Analysis 1.9](#); [Analysis 1.10](#).)

Figure 4. Forest plot of comparison: 1 FNB versus PCA opioid. Outcome: 1.8 Pain on movement at 24 hours.

Footnotes

(1) Study author provided data on standard deviations.

(2) Study author provided data on standard deviations.

Pooled results for FNB (any type, with or without a concurrent PCA opioid) vs PCA opioid demonstrated significantly lower pain at rest for FNB from zero to 72 hours: first two hours (11 RCTs, 706 participants, SMD -0.58, 95% CI -1.00 to -0.16, $I^2 = 84\%$), three to 12 hours (14 RCTs, 972 participants, SMD -0.97, 95% CI -1.42 to -0.52, $I^2 = 89\%$), 24 hours (19 RCTs, 1066 participants, SMD -0.72, 95% CI -0.93 to -0.51, $I^2 = 54\%$), 48 hours (17 RCTs, 957 participants, SMD -0.64, 95% CI -1.03 to -0.25, $I^2 = 86\%$), and 72 hours (eight RCTs, 560 participants, SMD -0.67, 95% CI -1.32 to -0.01, $I^2 = 92\%$).

Similarly, the pooled results for pain on movement also demonstrated significantly less pain for FNB (any type, with or without a concurrent PCA opioid) from zero to 48 hours: first two hours (four RCTs, 218 participants, SMD -1.29, 95% CI -2.12 to -0.46, $I^2 = 84\%$), three to 12 hours (eight RCTs, 462 participants, SMD -1.06, 95% CI -1.68 to -0.43, $I^2 = 88\%$), 24 hours (17 RCTs, 1017 participants, SMD -0.94, 95% CI -1.32 to -0.55, $I^2 = 86\%$) and 48 hours (13 RCTs, 742 participants, SMD -0.44, 95% CI -0.71 to -0.16, $I^2 = 65\%$). The results for 72 hours also favoured FNB, although not significantly (six RCTs, 438 participants, SMD -0.17, 95% CI -0.39 to 0.04, $I^2 = 11\%$).

Three RCTs (264 participants) (Chan 2013; Martin 2008; Mistraretti 2006) assessed pain intensity after 72 hours. Of these, one RCT (Chan 2013) compared continuous FNB (catheter duration less than 72 hours) and single-shot FNB with PCA, and two RCTs (Martin 2008; Mistraretti 2006) compared continuous FNB + sciatic versus PCA (catheter duration 48 hours). Chan 2013 found no significant difference in pain scores at rest and on movement at day five, week two and month three; Mistraretti 2006 reported no difference in pain at rest on discharge between the allocation groups. Martin 2008 reported significantly less pain at rest for FNB at day seven, but not at month one and month three, compared with PCA opioid.

For pain at rest and on movement at 48 hours, the result of test for differences between subgroups is statistically significant ($P < 0.05$), suggesting that analgesic effects at 48 hours are not consistent between the different types of FNB subgroups.

Pain at rest and on movement at 24 hours: subgroup analysis by type of FNB

(See Analysis 1.3; Analysis 1.8.)

We categorized RCTs according to type of FNB (i.e. single-shot FNB, single-shot FNB + sciatic/obturator block, continuous FNB, continuous FNB + sciatic block). For the single-shot FNB subgroup, pain at rest at 24 hours (nine RCTs, 416 participants, SMD -0.65, 95% CI -1.08 to -0.22, $I^2 = 75\%$) and pain on movement at 24 hours (six RCTs, 287 participants, SMD -0.50, 95% CI -0.81 to -0.19, $I^2 = 29\%$) were significantly less in the FNB subgroup compared with PCA. Pain scores were also significantly lower in the continuous FNB subgroup at rest at 24 hours (10 RCTs, 578 participants, SMD -0.74, 95% CI -0.97 to -0.51, $I^2 = 27\%$) and on movement at 24 hours (10 RCTs, 584 participants, SMD -1.09, 95% CI -1.74 to -0.43, $I^2 = 92\%$).

Only a few small RCTs evaluated the addition of sciatic or obturator block to the single-shot or continuous FNB. The single-shot FNB + sciatic block subgroup had significantly lower pain scores compared with the PCA opioid subgroup on movement at 24 hours (two RCTs, 92 participants, SMD -0.73, 95% CI -1.18 to -0.29, $I^2 = 0\%$) but not at rest at 24 hours (one RCT, 24 participants, SMD -0.73, 95% CI -1.74 to 0.28). The continuous FNB + sciatic block subgroup

favoured FNB for pain at rest at 24 hours (two RCTs, 54 participants, SMD -0.85, 95% CI -1.55 to -0.15, $I^2 = 28\%$) but not for pain on movement at 24 hours (two RCTs, 54 participants, SMD -2.08, 95% CI -4.74 to 0.58, $I^2 = 90\%$).

Generally, results of the subgroup analyses demonstrated that the type of FNB explained some of the heterogeneity of the overall results ($I^2 < 30\%$ for continuous FNB and continuous FNB + sciatic block for pain at rest at 24 hours; and $I^2 < 30\%$ for single-shot FNB and single-shot FNB + sciatic block for pain on movement at 24 hours).

Results of the various types of FNB at other timings are shown in these analyses (see Analysis 1.4; Analysis 1.5; Analysis 1.9; Analysis 1.10).

Pain at rest and on movement at 24 hours: subgroup analysis by FNB with and without concurrent PCA opioid

(See Analysis 1.11; Analysis 1.12.)

For pain at rest at 24 hours, FNB (any type) both with a concurrent PCA opioid (15 RCTs, 771 participants, SMD -0.67, 95% CI -0.94 to -0.41, $I^2 = 65\%$) and without a concurrent PCA opioid (five RCTs, 295 participants, SMD -0.93, 95% CI -1.31 to -0.55, $I^2 = 43\%$) demonstrated significantly less pain compared with PCA opioid alone. Similarly, for pain on movement at 24 hours, the groups given FNB with a concurrent PCA opioid (14 RCTs, 802 participants, SMD -0.67, 95% CI -0.99 to -0.35, $I^2 = 76\%$) and without a concurrent PCA opioid (four RCTs, 215 participants, SMD -2.26, 95% CI -3.95 to -0.57, $I^2 = 95\%$) had significantly less pain when compared with the group given PCA opioid alone.

Pain at rest and on movement at 24 hours: subgroup analysis by FNB with ropivacaine versus bupivacaine

(See Analysis 1.13; Analysis 1.14.)

For pain at rest at 24 hours, both FNB (any type) with ropivacaine (eight RCTs, 379 participants, SMD -1.30, 95% CI -2.03 to -0.58, $I^2 = 90\%$) and FNB (any type) with bupivacaine (11 RCTs, 649 participants, SMD -0.61, 95% CI -0.89 to -0.33, $I^2 = 61\%$) demonstrated significantly less pain compared with PCA opioid. For pain on movement at 24 hours, the group given FNB with ropivacaine (eight RCTs, 409 participants, SMD -0.98, 95% CI -1.67 to -0.29, $I^2 = 90\%$) and FNB with bupivacaine (10 RCTs, 632 participants, SMD -0.92, 95% CI -1.39 to -0.44, $I^2 = 85\%$) also had significantly less pain compared with the group given PCA opioid.

This subgroup analysis did not explain the heterogeneity in the main results.

Pain at rest and on movement at 24 hours: sensitivity analysis by adequacy of allocation concealment and blinding

(See Analysis 1.15; Analysis 1.16; Analysis 1.17; Analysis 1.18.)

Pooling only the RCTs with low risk of bias regarding allocation concealment showed a significant reduction in the FNB (any type) for pain at rest (nine RCTs, 523 participants, SMD -0.54, 95% CI -0.76 to -0.32, $I^2 = 28\%$) and on movement (nine RCTs, 583 participants, SMD -0.54, 95% CI -0.75 to -0.33, $I^2 = 30\%$) at 24 hours.

Pooling only the results of trials with low risk of bias for adequate blinding of participants, personnel and outcome assessors also demonstrated significantly less pain at rest at 24 hours (seven RCTs, 319 participants, SMD -0.62, 95% CI -1.08 to -0.17, $I^2 = 71\%$) and on movement at 24 hours (seven RCTs, 356 participants, SMD -0.57, 95% CI -0.79 to -0.35, $I^2 = 0$). Heterogeneity was explained by adequate blinding for pain on movement but not at rest at 24 hours.

2. Serious adverse events

Few RCTs reported on serious adverse events.

Five RCTs (474 participants) reported no neurological injury in the FNB group (Chan 2012; Chan 2013; Kadic 2009; McNamee 2001; Seet 2006). Persistent numbness of the anterior thigh at 48 hours was reported in one participant in the single-shot FNB group compared with none in the PCA opioid group (one RCT, 40 participants) (Kardash 2007). One RCT (42 participants) reported that in the single-shot FNB group, a participant had knee numbness, and in the PCA group, two participants had infection and one participant had dyspnoea (Good 2007). The only RCT (200 participants) that reported on thrombotic events had one event in the continuous FNB group (Chan 2013). Two RCTs (282 participants) reported no falls in the FNB or PCA group (during the hospital stay (Chan 2013) or during the first three postoperative days (Chan 2012)). Three RCTs (192 participants) reported that no local anaesthetic toxicity was noted in the FNB or PCA group (Kadic 2009; McNamee 2001; Seet 2006). Five RCTs reported on technical failure: 27% technical failure with continuous FNB 0.2% bupivacaine and 55% with continuous FNB 0.1% bupivacaine in one RCT (62 participants) (Ganapathy 1999), no technical failure with the continuous FNB in one RCT (58 participants) (Kadic 2009) and no technical failure with the single-shot FNB in three RCTs (171 participants) (Allen 1998; de Lima e Souza 2008; McNamee 2001).

Secondary outcomes

1. Proportion of participants with significant pain postoperatively

Six comparisons from five RCTs (511 participants) reported the proportion of participants with moderate/severe pain. FNB demonstrated significantly fewer participants with moderate/severe pain compared with PCA opioid (RR 0.73, 95% CI 0.65 to 0.82, $I^2 = 0$). The NNTB was five (95% CI four to 11).

2. Time from end of surgery to first rescue opioid request

Two RCTs (Chan 2012; McNamee 2001) reported that participants receiving single-shot FNB took significantly longer time to first opioid request compared with those receiving PCA. One RCT (McNamee 2001) (74 participants, MD 7.21 hour, 95% CI 6.88 to 7.54 hour) demonstrated a much larger FNB effect compared with the other (Chan 2012) (82 participants, MD 3.23 hour, 95% CI 1.88 to 4.58 hour). The former RCT (McNamee 2001) had the addition of a sciatic block. The findings of these two RCTs were not pooled because of the high heterogeneity ($I^2 = 99\%$).

3. Opioid consumption

(See Analysis 1.19; Analysis 1.20.)

Pooled results for opioid consumption (IV morphine equivalent) were significantly reduced for FNB at 24 hours (20 RCTs, 1152 participants, MD -14.74 mg, 95% CI -18.68 to -10.81 mg, $I^2 = 88\%$) and

at 48 hours (19 RCTs, 1001 participants, MD -14.53 mg, 95% CI -20.03 to -9.02 mg, $I^2 = 88\%$).

4. Adverse effects

4.1 Nausea and/or vomiting

(See Analysis 1.21.)

Pooled results showed that the risk of nausea and/or vomiting was significantly lower for FNB compared with PCA opioid (16 RCTs, 1100 participants, RR 0.47, 95% CI 0.33 to 0.68, $I^2 = 73\%$). The NNTH was four (95% CI three to six).

4.2 Sedation

(See Analysis 1.22.)

Pooled results showed that the risk of sedation was significantly lower in the FNB group as compared with the PCA opioid group (nine RCTs, 808 participants, RR 0.34, 95% CI 0.16 to 0.74, $I^2 = 85\%$; NNTH five, 95% CI five to eight).

4.3 Urinary retention

(See Analysis 1.23.)

Seven RCTs with 37 events from 490 participants evaluated urinary retention. These results showed no difference in risk of urinary retention between FNB and PCA opioid (RR 0.57, 95% CI 0.20 to 1.68, $I^2 = 40\%$).

5. Physical function

(See Analysis 1.24.)

During postoperative day two to four, the FNB group achieved greater knee flexion (10 RCTs, 541 participants, MD 6.48 degrees, 95% CI 4.27 to 8.69 degrees, $I^2 = 79\%$) and knee extension (three RCTs, 297 participants, MD -0.16 degrees, 95% CI -0.30 to -0.01 degrees, $I^2 = 0\%$) compared with the PCA opioid group. Two RCTs evaluated time to first ambulation and found no differences between groups (71 participants, MD 1.86 degrees, 95% CI -7.40 to 11.12 degrees, $I^2 = 0\%$).

6. Participant satisfaction with analgesia during hospital stay

(See Analysis 1.25.)

Four studies assessed participant satisfaction on a continuous scale, and their pooled results indicated greater participant satisfaction with FNB versus PCA opioid (180 participants, SMD 1.06, 95% CI 0.74 to 1.38, $I^2 = 0\%$). Two small RCTs reported participant satisfaction as a dichotomous variable, and their results were not pooled because of high heterogeneity ($I^2 = 96\%$). The findings of both studies were not statistically significant (42 participants, RR 1.00, 95% CI 0.91 to 1.09) (Adams 2002); (62 participants, RR 1.64, 95% CI 1.02 to 2.64) (de Lima e Souza 2008).

II. FNB versus epidural analgesia

Primary outcomes

1. Pain at rest and on movement

(See Analysis 2.3; Analysis 2.5.)

We found no differences in pooled pain scores at rest and on movement for all measured time points in the first 72 hours post operation between FNB and epidural analgesia. Pooled results for FNB (any type) versus epidural for pain at rest at 24 hours included the following: six RCTs, 328 participants, SMD -0.05, 95% CI -0.43 to 0.32, $I^2 = 61\%$; and for pain on movement at 24 hours: six RCTs, 317 participants, SMD 0.01, 95% CI -0.21 to 0.24, $I^2 = 0\%$. The data from one study with a concurrent epidural in the FNB intervention showed significantly lower pain scores for the FNB + epidural intervention for pain on movement at 24 hours compared with epidural alone (78 participants, SMD -2.94, 95% CI -3.59 to -2.29) but not for pain at rest at 24 hours (SMD -0.34, 95% CI -0.79 to 0.10) (Lee 2011). Details of the pooled results at rest at two hours,

three to 12 hours and 48 hours are shown in [Analysis 2.1](#); [Analysis 2.2](#); [Analysis 2.4](#) and [Analysis 2.6](#). The other non-significant findings were for pain at rest at 72 hours (two RCTs, 124 participants, SMD 0.18, 95% CI -0.35 to 0.71, $I^2 = 45\%$); pain on movement at three to 12 hours (one RCT, 59 participants, SMD 0.20, 95% CI -0.31 to 0.71) and pain on movement at 72 hours (one RCT, 54 participants, SMD 0.61, 95% CI -0.06 to 1.09).

The test results for subgroup differences were significant ($P < 0.05$) for pain at rest at two hours and at three to 12 hours,

Pain at rest and on movement at 24 hours: subgroup analysis by type of FNB

(See [Analysis 2.3](#); [Figure 5](#).)

Figure 5. Forest plot of comparison: 2 FNB versus epidural. Outcome: 2.5 Pain on movement at 24 hours.

No significant differences were detected between pain scores at rest and on movement at 24 hours for the subgroups of single-shot FNB + sciatic block, continuous FNB and continuous FNB + sciatic block. For pain at rest at 24 hours, statistical heterogeneity was reduced in the analyses by type of FNB for the continuous FNB + sciatic block subgroup (three RCTs, 93 participants, SMD 0.09, 95% CI -0.32 to 0.51, $I^2 = 0\%$). For pain on movement at 24 hours, no heterogeneity was found ($I^2 = 0\%$) in the subgroups of continuous FNB (three RCTs, 165 participants, SMD 0.12, 95% CI -0.19 to 0.43) and continuous FNB + sciatic block (three RCTs, 93 participants, SMD 0.06, 95% CI -0.36 to 0.47). Only one RCT was identified in the subgroup of single-shot FNB + sciatic block (59 participants, SMD -0.33, 95% CI -0.85 to 0.18) for this outcome.

Pain at rest and on movement at 24 hours: subgroup analysis by FNB with ropivacaine versus bupivacaine

(See [Analysis 2.7](#); [Analysis 2.8](#).)

For pain at rest at 24 hours, no significant differences were detected for FNB with ropivacaine (four RCTs, 190 participants, SMD -0.16, 95% CI -0.68 to 0.37, $I^2 = 65\%$) or FNB with bupivacaine (two RCTs, 216 participants, SMD 0.20, 95% CI -0.13 to 0.53, $I^2 = 0\%$) compared with epidural, respectively. Similarly, for pain on movement at 24 hours, no significant differences were found for FNB with ropivacaine (four RCTs, 120 participants, SMD 0.10, 95% CI -0.27 to 0.47, $I^2 = 0\%$) or FNB with bupivacaine (three RCTs, 197 participants, SMD -0.03, 95% CI -0.31 to 0.25, $I^2 = 0\%$) compared with epidural, respectively.

Pain at rest and on movement at 24 hours: sensitivity analysis by adequacy of allocation concealment and blinding

(See [Analysis 2.9](#); [Analysis 2.10](#).)

No significant difference between the groups was detected at 24 hours on the basis of pooled results from RCTs with low risk of bias for allocation concealment for pain at rest (three RCTs, 194 participants, SMD 0.20, 95% CI -0.07 to 0.47, $I^2 = 0\%$) and for pain on movement (four RCTs, 271 participants, SMD 0.03, 95% CI -0.21 to 0.27, $I^2 = 0\%$). Statistical heterogeneity was reduced in analyses by adequate allocation concealment for pain at rest and on movement at 24 hours. Sensitivity analysis on the adequacy of blinding was not conducted, as no RCTs with low risk of bias for blinding were identified.

2. Serious adverse events

Four RCTs reported on neurological injury: One RCT (59 participants) reported that a participant with epidural developed unilateral foot drop and sphincteric disturbance after operation, but the aetiology was not stated ([Davies 2004](#)), and three RCTs (247 participants) reported no incidence of neurological injury in either allocation group ([Barrington 2005](#); [Lee 2011](#); [Sundarathiti 2009](#)). One RCT (70 participant) with a follow-up of six months post operation reported a participant with a fall at home with wound dehiscence requiring reoperation in the continuous FNB group and none in the epidural group ([Long 2006](#)). This RCT also reported two participants in the continuous FNB group requiring close manipulation, one participant in the epidural group with bladder inflammation from reaction to latex of the Foley catheter and no incidence of haematoma formation of the knee or drainage in either group ([Long 2006](#)). Another RCT (50 participants) reported a participant with atrial fibrillation in the epidural group ([Zaric 2006](#)). There was no incidence of local anaesthetic toxicity from two RCTs (111 patients) ([Sundarathiti 2009](#); [Zaric 2006](#)), and no incidence of catheter-related infection from one RCT (61 patients) ([Sundarathiti 2009](#)). One RCT (108 patients) reported three cardiac events (one non-ST elevation myocardial infarction and one cardiac conduction defect requiring permanent pacemaker) in the continuous FNB group and one hypotension (60/40 mmHg and bradycardic, spinal-epidural block reached dermatome T3 upon 2 hours of the epidural infusion) in the epidural group ([Barrington 2005](#)). Another RCT (59 patients) reported frequent hypotension in both allocation groups (37% in the single-shot FNB+sciatic group and 27% in the epidural group) ([Davies 2004](#)). Conversely, another RCT (78 patients) reported no incidence of severe hypotension or cardiac complications ([Lee 2011](#)).

Secondary outcomes

1. Proportion of participants with significant pain postoperatively

The result from one small RCT (61 participants) favoured epidural between groups for the proportion of participants with moderate/severe pain (RR 2.07, 95% CI 1.28 to 3.33) ([Sundarathiti 2009](#)).

2. Time from end of surgery to first rescue analgesic request

We found no data on this outcome.

3. Opioid consumption

(See [Analysis 2.11](#); [Analysis 2.12](#).)

Results from five RCTs indicate that the FNB group consumed less opioid (IV morphine equivalent) compared with the epidural group

at 24 hours (341 participants, MD -4.35 mg, 95% CI -9.95 to 1.26 mg, $I^2 = 50\%$). Similarly, pooled results at 48 hours also favoured FNB (four RCTs, 233 participants, MD -1.28 mg, 95% CI -5.30 to 2.74 mg, $I^2 = 0\%$), although this finding was not statistically significant.

We did not pool the data from three RCTs because the data at 24 and 48 hours were not available ([Barrington 2005](#); [Sundarathiti 2009](#)) or because the analgesics were too varied ([Singelyn 1998](#)). In the first RCT (108 participants) ([Barrington 2005](#)), over 72 hours, the continuous FNB group required less morphine (mean (SD) 44 (30) vs 53 (28) mg, P value 0.45), but more oxycodone (mean (SD) 21 (15) vs 13 (12) mg, P value 0.005) compared with the epidural group. However the epidural group also received a mean dosage of fentanyl of 1.74 mg. In the second RCT (61 participants) ([Sundarathiti 2009](#)), the cumulative IV tramadol over 72 hours was higher in the continuous FNB group compared with the epidural group (median (range) = 150 (0 to 350) vs 50 (0 to 150) mg, P value 0.001). In the third RCT (30 participants) ([Singelyn 1998](#)), both the FNB and epidural groups received infusion bupivacaine with sufentanil and clonidine, and the supplemental analgesics required in the first 48 hours were comparable (propacetamol 1.7 (1.1) vs 1.1 (1.5) g, P value 0.37; IM piritramide 1.9 (4.1) vs 2.3 (6.2) mg, P value 0.29).

4. Adverse effects

4.1 Nausea and/or vomiting

(See [Analysis 2.13](#).)

Pooled results from four RCTs demonstrated that the risk of nausea and/or vomiting in the FNB group was less than in the epidural group (183 participants, RR 0.63, 95% CI 0.41 to 0.97, $I^2 = 0\%$). The NNTH was eight, 95% CI four to 53.

4.2 Sedation

The only RCT that evaluated sedation found no significant difference in risk of sedation between the allocation groups (50 participants, RR 0.43, 95% CI 0.04 to 4.40) ([Zaric 2006](#)).

4.3 Urinary retention

Pooled results from four RCTs revealed no significant difference between groups in risk of urinary retention (200 participants, RR 0.36, 95% CI 0.07 to 1.88, $I^2 = 49\%$).

4.4 Technical failure of the blocks

The meta-analysis of four RCTs (287 participants) found no difference in technical failure of the block (RR 0.87, 95% CI 0.49 to 1.55, $I^2 = 0\%$) or catheter disconnection/problem (RR 0.73, 95% CI 0.24 to 2.22, $I^2 = 15\%$) between the FNB and epidural groups.

5. Physical function

Pooled data of six RCTs revealed no significant difference in knee flexion during postoperative day two to four (328 participants, MD -1.65 degrees, 95% CI -5.14 to 1.84 degrees, $I^2 = 33\%$), and the results from one RCT ([Long 2006](#)) showed no significant difference in knee extension on postoperative day three between groups (115 participants, MD 5.00 degrees, 95% CI 1.62 to 8.38 degrees).

One small RCT evaluated the time to first ambulation and found no differences between groups (16 participants, MD 8.50 hours, 95% CI -5.50 to 22.50 hours) ([Mistrali 2006](#)).

6. Participant satisfaction with analgesia

Two RCTs reported on participant satisfaction as a continuous variable (scale zero to 10), and their pooled results indicate that participants with FNB were more satisfied (120 participants, SMD 0.60, 95% CI 0.23 to 0.97, $I^2 = 0\%$). The only RCT that evaluated participant satisfaction as a dichotomous outcome reported that all participants in both FNB and epidural groups were satisfied with their treatment (30 participants, RR 1, 95% CI 0.91 to 1.09) (Singelyn 1998).

III. FNB versus local infiltration analgesia

Primary outcomes

1. Pain at rest and on movement

(See Analysis 3.1; Analysis 3.2; Analysis 3.3; Analysis 3.4.)

Pooled results revealed no difference between FNB and local infiltration analgesia for pain at rest at 24 hours (four RCTs, 216 participants, SMD 0.06, 95% CI -0.61 to 0.72, $I^2 = 82\%$) and at 48 hours (two RCTs, 114 participants, SMD -0.26, 95% CI -0.94 to 0.43, $I^2 = 68\%$) or for pain on movement at 24 hours (three RCTs, 153 participants, SMD 0.38, 95% CI -0.10 to 0.86, $I^2 = 53\%$) and at 48 hours (two RCTs, 111 participants, SMD -0.14, 95% CI -0.71 to 0.43, $I^2 = 54\%$).

One RCT with FNB + local infiltration analgesia versus local infiltration analgesia also found no difference in pain at rest at 24 hours between allocation groups (55 participants, SMD -0.11, 95% CI -0.64 to 0.42) (Widmer 2012). Similarly, one cross-over RCT (16 participants) comparing collateral knees reported no significant differences between FNB and local infiltration analgesia for pain at rest and on movement at 24 hours, 48 hours and 72 hours (Ng 2012). One RCT assessed pain at rest at 12 hours and found that local infiltration analgesia yielded lower pain scores compared with FNB (76 participants, SMD 0.56, 95% CI 0.10 to 1.02) (Toftdahl 2007).

Pain at rest and on movement at 24 hours: sensitivity analysis by adequacy of blinding

Only one small RCT (40 participants) described adequate blinding of participants, personnel and outcome assessors for pain at rest and on movement at 24 hours (Carli 2010). The RCT favoured FNB for pain at rest at 24 hours (SMD -1.12, 95% CI -1.79 to -0.45) but not for pain on movement at 24 hours (SMD 0.26, 95% CI -0.37 to 0.88).

2. Serious adverse event

One RCT (76 participants) reported the following adverse events in the local infiltration group (40 participants) and none in the FNB group: two participants requiring open surgical intervention (one for deep surgical site infection and another for tissue necrosis), two participants with a cardiac event (one syncopal attack of cardiac origin and one episode of chest pain justifying investigation), two participants with gastric ulcer and two participants with urinary tract infection (Toftdahl 2007). Another RCT (40 participants) reported that one participant in the local infiltration group had a cardiac event (atrial fibrillation requiring treatment) (Carli 2010). Four RCTs (171 participants) reported no incidence of neurological injury, wound infection, surgical intervention, cardiac events or falls in both FNB and infiltration groups (Affas 2011; Ng 2012; Parvataneni 2007; Widmer 2012).

Secondary outcomes

1. Proportion of participants with significant pain postoperatively

No statistical significance was detected in the proportion of participants with moderate/severe pain from the pooling of two RCTs (114 participants, RR 1.86, 95% CI 0.50 to 6.92, $I^2 = 45\%$).

2. Time from end of surgery to first rescue analgesic request

We found no data on this outcome.

3. Opioid consumption

At 24 hours, a small RCT found that FNB had lower opioid consumption compared with local infiltration analgesia, but the results were not significant (40 participants, MD -11.50 mg, 95% CI -24.08 mg to 1.08) (Carli 2010). Another RCT found that FNB + local infiltration analgesia resulted in lower opioid consumption at 24 hours compared with local infiltration analgesia alone (55 participants, MD -35.44 mg, 95% CI -56.02 to -14.86 mg) (Widmer 2012). These two RCTs were not pooled, as one of them (Widmer 2012) had a concurrent local infiltration analgesia in the FNB group. At 48 hours, pooled results of two RCTs on opioid consumption revealed no significant differences (117 participants, MD 3.51 mg, 95% CI -23.43 to 30.46 mg). Similarly, one cross-over RCT (16 participants) reported no significant differences in opioid consumption on postoperative day one or day two between the allocation groups (Ng 2012).

4. Adverse effects

4.1 Nausea and/or vomiting

(See Analysis 3.5.)

Pooled results of three RCTs showed no statistical differences between groups in risk of nausea and vomiting (177 participants, RR 1.71, 95% CI 0.64 to 4.62, $I^2 = 47\%$).

5. Physical function

One RCT found no significant differences in knee flexion on postoperative day two between groups (40 participants, MD -0.50 degrees, -5.28 to 4.28 degrees) (Carli 2010). Another RCT failed to find any significant difference between groups in knee extension on postoperative day two (76 participants, MD 1.15 degrees, 95% CI -1.73 to 4.03 degrees) (Toftdahl 2007).

6. Participant satisfaction with analgesia

One RCT found that participants with FNB had lower satisfaction scores compared with participants with local infiltration analgesia (60 participants, mean 7.8 vs 9.0, SD not reported) (Parvataneni 2007).

IV. FNB versus oral analgesia

Primary outcomes

1. Pain at rest and on movement

Only one RCT (62 participants) compared FNB versus oral analgesia. In this RCT (Nader 2012), continuous FNB + oral analgesia was compared with oral analgesia alone on postoperative day one at discontinuation of the epidural analgesia. Investigators found that continuous FNB + oral analgesia led to significantly lower pain

scores at rest at 24 hours (MD -3.03, 95% CI -4.00 to -2.06) and 48 hours (MD -2.00, 95% CI -3.20 to -0.80), as well as on movement at 24 hours (MD -2.00, 95% CI -3.10 to -0.90), when compared with oral analgesia alone (Nader 2012). At 72 hours, no significant difference was detected for pain at rest (MD -1.00, 95% CI -2.18 to 0.18).

2. Serious adverse event

Following epidural, four of 31 participants in the oral opioid analgesia group had deep venous thrombosis or pulmonary embolus versus none in the continuous FNB + oral analgesia group (Nader 2012).

Secondary outcomes

Results from one RCT (62 participants) showed that following epidural, the FNB + oral analgesia group consumed less opioid at 24

hours (MD -15.00 mg, 95% CI -24.50 to -5.50 mg) and at 48 hours (MD -7.00 mg, 95% CI -21.25 to 7.25 mg), although the results were not statistically significant. FNB + oral analgesia also led to greater knee flexion compared with oral analgesia alone (MD 15.00 mg, 95% CI 5.96 to 24.04 mg). However, no significant difference was detected in participant satisfaction between FNB + oral analgesia and oral analgesia alone (MD 2.00 mg, 95% CI 1.06 to 2.94 mg) (Nader 2012). We found no data on the other secondary outcomes.

V. Continuous versus single-shot FNB

Primary outcomes

1. Pain at rest and on movement

(See Analysis 4.1; Analysis 4.2; Analysis 4.3; Analysis 4.4; Figure 6; Analysis 4.6.)

Figure 6. Forest plot of comparison: 4 Continuous FNB versus single-shot FNB. Outcome: 4.5 Pain on movement at 24 hours.

Continuous FNB showed significantly lower pain intensity when compared with single-shot FNB at rest at 24 hours (four RCTs, 272 participants, SMD -0.62, 95% CI -1.17 to -0.07, I² = 73%) and at 48 hours (four RCTs, 272 participants, SMD -0.96, 95% CI -1.69 to -0.22, I² = 84%), as well as on movement at 24 hours (four RCTs, 272 participants, SMD -0.42, 95% CI -0.67 to -0.17, I² = 0%) and at 48 hours (four RCTs, 272 participants, SMD -0.54, 95% CI -1.02 to -0.06, I² = 65%).

Pooled results of two RCTs (156 participants) at the following time points found no significant differences between allocation groups: pain at rest at two hours (SMD 0.15, 95% CI -0.17 to 0.46, I² = 0), at three to 12 hours (SMD -0.03, 95% CI -0.35 to 0.28, I² = 0) and at 72 hours (SMD 0.10, 95% CI -0.21 to 0.42, I² = 0); and pain on movement at 72 hours (SMD -0.09, 95% CI -0.41 to 0.22, I² = 0).

Pain at rest and on movement at 24 hours: subgroup analysis by continuous FNB with and without a PCA opioid intervention

(See Analysis 4.7; Analysis 4.8.)

For pain at rest at 24 hours, pooled RCTs with a concurrent PCA opioid intervention in both continuous and single-shot FNB groups (three RCTs, 138 participants, SMD -0.86, 95% CI -1.35 to -0.38, I² = 36%) and the one RCT with a concurrent PCA opioid intervention only in the single-shot FNB group (134 participants, SMD -0.16, 95% CI -0.50 to 0.18), respectively, showed that continuous FNB resulted in less pain compared with single-shot FNB. For pain on movement at 24 hours, results for continuous FNB with a concurrent PCA opioid (three RCTs, 138 participants, SMD -0.59, 95% CI -0.96 to -0.22, I² = 0) and without a concurrent PCA opioid (one RCT, 134 participants, SMD -0.27, 95% CI -0.61 to 0.07), respectively, also

showed continuous FNB to lead to less pain compared with single-shot FNB.

Pain at rest and on movement at 24 hours: sensitivity analysis by adequacy of allocation concealment and blinding

(See Analysis 4.9; Analysis 4.10.)

Pooling of only RCTs with low risk of bias for allocation concealment did not change the magnitude or the direction of the effect estimate for pain at 24 hours, at rest (three RCTs, 250 participants, SMD -0.73, 95% CI -1.41 to -0.04, I² = 82%) or on movement (three RCTs, 250 participants, SMD -0.47, 95% CI -0.79 to -0.15, I² = 25%).

Only one small RCT (Hirst 1996) (22 participants) had low risk of bias pertaining to blinding. Results indicated that differences between groups for pain at 24 hours were not statistically significant at rest (MD -0.50, 95% CI -2.25 to 1.25) and on movement (MD -0.70, 95% CI -3.09 to 1.69).

2. Serious adverse events

Two RCTs (170 participants) that compared continuous versus single-shot FNB reported no incidence of neurological injury (Chan 2013; Salinas 2006). One RCT (38 participants) reported no catheter-related infection (Salinas 2006).

Secondary outcomes

1. Proportion of participants in significant pain postoperatively

We found no data on this outcome.

2. Time from end of surgery to first rescue analgesic request

We found no data on this outcome.

3. Opioid consumption

(See [Analysis 4.11](#); [Analysis 4.12](#).)

Opioid consumption was significantly less in the continuous FNB group compared with the single-shot FNB group at 24 hours (three RCTs, 236 participants, MD -13.81 mg, 95% CI -23.27 to -4.35 mg, $I^2 = 88\%$) and at 48 hours (four RCTs, 269 participants, MD -14.59 mg, 95% CI -22.35 to -6.82 mg, $I^2 = 75\%$).

4. Adverse effects

4.1 Nausea and/or vomiting

Pooled results from two RCTs showed no differences in risk of nausea/vomiting between continuous and single-shot FNB groups (214 participants, RR 0.55, 95% CI 0.20 to 1.56, $I^2 = 78\%$).

4.2 Sedation

Two RCTs (214 participants) presented results on sedation ([Chan 2013](#); [Park 2010](#)). One of these RCTs (134 participants) reported zero events in both groups and did not contribute to the analysis ([Chan 2013](#)). Thus, only the results of one RCT were considered; these revealed no differences in risk of sedation between groups (80 participants, RR 4.33, 95% CI 0.60 to 31.07) ([Park 2010](#)).

4.3 Technical failure of the blocks

One RCT (36 participants) reported that no technical failure occurred ([Salinas 2006](#)).

5. Physical function

We found no data on this outcome.

6. Participant satisfaction with analgesia

We found no data on this outcome.

DISCUSSION

Summary of main results

This review has examined the published evidence comparing the use of FNB (any type) with or without concurrent treatments including PCA opioid versus a variety of alternative methods for reducing postoperative pain following knee replacement surgery. We found that pain at rest and pain on movement were less for FNB at all time intervals considered up to 72 hours post operation. At 24 hours, FNB (any type) had a moderate to large analgesic effect at rest and on movement compared with PCA. The moderate to large effects (SMD), when re-expressed as mean differences (zero to 10 scale), represented at 24 hours a 1.20-point decrease in mean pain at rest and a 1.66-point decrease in mean pain on movement for FNB compared with PCA ([Summary of findings for the main comparison](#)). FNB also resulted in less opioid consumption at 24 hours and 48 hours, and correspondingly lower risk of nausea/vomiting or sedation, compared with PCA opioid.

When compared with an epidural block, no differences in pain at rest or on movement were noted when FNB was used, although FNB demonstrated lower opioid consumption at 24 hours, lower risk of nausea and/or vomiting and greater participant satisfaction

compared with epidural. Similarly, no differences were detected between FNB and local infiltration analgesia for pain at rest and on movement at any evaluated time point.

The single study comparing FNB versus oral analgesia after TKR suggested that the former offered superior analgesia at rest from 24 to 48 hours, and on movement at 24 hours.

Using a continuous FNB was superior to using a single-shot technique, with moderate effects on pain intensity noted at 24 hours and 48 hours both at rest and on movement. Reexpressing SMD as MD on a zero to 10 scale suggests, at 24 hours, a 0.95-point reduction in mean pain at rest and a 0.74-point reduction in mean pain on movement ([Summary of findings 4](#)). Continuous FNB also led to lower opioid consumption at 24 hours and at 48 hours compared with single-shot FNB.

Serious adverse events are rare with FNB, and we have been unable to report definitive results concerning the risk of serious adverse events for any of the analgesic techniques examined. Nevertheless, the adverse events in the local infiltration analgesia group reported by [Toftdahl 2007](#) are a cause of concern. Larger trials examining two main issues are needed before this technique can be recommended. First, two reinterventions out of 40 participants for infection or tissue necrosis at the surgical site seems unusually high. Second, although blood concentrations have been reported to be low, the total amount of local anaesthetic administered with this technique is high. Cardiac toxicity in patients with underlying cardiac disease may occur with blood concentrations that would be well tolerated in normal patients. In the RCT, two participants given local infiltration analgesia had cardiac events ([Toftdahl 2007](#)). Therefore, before widespread use of this technique is considered, especially in centres with very short hospital stays (less than 24 hours), studies evaluating the effects of this technique on cardiac conduction may be warranted.

Overall completeness and applicability of evidence

Generally, application of the review's finding to clinical practice is possible, as adequate descriptions of inclusion and exclusion criteria were provided. Nevertheless, the included RCTs have variable reporting or inconsistent definitions for some of the secondary outcomes. Subgroups involving single-shot or continuous FNB with the addition of sciatic or obturator block should be interpreted cautiously because of the small numbers of trials and participants. For relatively uncommon events such as urinary retention, differences between groups would have been difficult to detect. Additionally, serious adverse events were rare, were reported in very few RCTs and most often were reported anecdotally.

In this review, the overall summary effects of FNB compared with PCA opioid, epidural or local infiltration analgesia, respectively, were derived from the combination of data from any type of FNB including single-shot FNB, if the data were derived from similar postoperative periods. Since the effects of the single-shot FNB possibly last less than 24 hours, the magnitude of the overall summary effect of the FNB (any type) after 24 hours likely underestimates the intervention effect of the continuous FNB.

Quality of the evidence

Generally, most RCTs were at low to moderate risk of bias for the aspects rated in our risk of assessment tool, except for the aspect of

blinding (see [Figure 3](#)). We rated 14 (31%) RCTs at high risk for both participant and assessor blinding, and eight (18%) RCTs were rated at high risk for one blinding aspect. The high risk of bias for blinding was expected, as the nature of the analgesic regimens in this review is such that it is challenging to implement blinding effectively. To do so would have required the insertion of sham catheters. Even with sham catheters, clinical signs of numbness of the affected areas would have broken the blinding to participants. The open-label nature of some of the RCTs may potentially bias the measurement of subjective outcomes such as pain and participant satisfaction.

For allocation concealment, sensitivity analyses pooling only RCTs at low risk of bias demonstrated that the primary results for comparisons of FNB (with or without PCA opioid) versus PCA opioid, FNB versus epidural and continuous FNB versus single-shot FNB were robust. Pooling only RCTs at low risk of bias for blinding of participants/personnel and outcome assessors also showed the findings to be robust for the comparison of FNB versus PCA opioid. As expected, pooling results of trials with low risk of bias resulted in a slight reduction of the magnitude of the mean intervention effect but no change in statistical significance. We were unable to assess the effect of blinding for the comparison of FNB versus epidural, as all included RCTs were at high risk of blinding bias. We also could not conduct sensitivity analyses for the other comparisons because of the limited number of available RCTs.

Potential biases in the review process

Publication bias

We may have missed trials that were not indexed in CENTRAL, MEDLINE, EMBASE, CINAHL or Web of Science or that remain unpublished in journals. The fail-safe N test indicated that our results were robust to publication bias (see [Appendix 10](#)). Nonetheless, we did approach trialists for information on existing RCTs. Although the possibility of publication bias cannot be excluded, a well-conducted non-significant trial would have had a reasonable chance of being published. The procedure we used to retrieve all RCTs was comprehensive and systematic, and two review authors conducted independent selection from results obtained by the searches. Thus the risk of publication bias is low.

Pooling and heterogeneity

We pooled data from RCTs testing the same comparisons (i.e. FNB vs PCA opioid, FNB vs epidural and FNB vs local infiltration analgesia) and presented the meta-analyses stratified by different types of FNB (single-shot or continuous FNB, with or without sciatic/obturator block). Notable differences were reported in the analgesic regimen across pooled FNB subgroups such as FNB with different local anaesthetics, drug concentrations and infusion rates; control groups with variations in drugs and dosages and differences in the supplemental analgesics used in these trials. These differences likely contributed to high heterogeneity in many of the meta-analyses in this review. The type of FNB and the methodological quality of the trials have explained some of the heterogeneity of pooled data.

We used the random-effects model to account for the anticipated heterogeneity of included RCTs. We also used risk ratios rather than odds ratios to report dichotomous outcomes. These decisions resulted in more conservative estimates of treatment effects with wider confidence intervals, compared with outcomes when fixed-

effect models and odds ratios were used. We did not reanalyse the data using a fixed-effect model because of the high heterogeneity of the included trials. Sensitivity analyses demonstrated that the primary findings were robust to potentially influential allocation concealment and blinding bias.

Small-study effects

We did not assess the influence of small trials on estimated treatment effects. Nevertheless, given that all the RCTs included in this review had fewer than 100 participants per arm, small-study effects could potentially have biased the results of meta-analyses towards an inflated treatment effect ([Nuesch 2010](#)).

Agreements and disagreements with other studies or reviews

The conclusions of the current review are similar to those provided by a previously published meta-analysis comparing FNB and opioids ([Paul 2010](#)). Both reviews found single-shot FNB to lead to lower pain scores during movement at 24 hours and at 48 hours; lower opioid requirement at 24 hours and at 48 hours and lower risk of nausea and vomiting. Unlike Paul's review, the current review also found that single-shot FNB and continuous FNB have lower pain intensity at rest at 24 hours and 48 hours, lower risk of sedation, greater knee range of motion and higher participant satisfaction. Both [Fischer 2008](#) and the authors of the current review found single-shot or continuous FNB to have superior effects on pain up to 48 hours. The current review also demonstrated that FNB reduced the risk of nausea and/or vomiting, although this outcome was not significant in the [Fischer 2008](#) review. Additionally, we demonstrated that continuous FNB was superior to single-shot FNB for postoperative analgesia and opioid consumption, unlike in [Paul 2010](#).

Any discordance with the two past reviews ([Fischer 2008](#); [Paul 2010](#)) may be due to the fact that the current review included a larger number of RCTs for meta-analysis and therefore had a greater possibility of detecting statistical significance. Moreover, for the continuous versus single-shot FNB comparisons, we conducted direct (head-to-head) comparisons of parallel trials, while [Paul 2010](#) used an indirect comparison.

Another review compared continuous peripheral nerve block and opioids ([Richman 2006](#)). That review concluded that continuous peripheral nerve block provided superior postoperative analgesia compared with opioids at 24 hours, 48 hours and 72 hours. It also reported that nerve block led to fewer minor complications, including nausea or vomiting, pruritus and sedation, and improved participant satisfaction. These conclusions were similar to ours, despite the broader inclusion criteria provided in the review by [Richman 2006](#), which extended inclusion to both upper and lower extremity operations, and catheter locations in places other than the femoral nerve.

Our results for FNB versus epidural were somewhat in agreement with those of [Fowler 2008](#), who compared FNB and epidural analgesia for major knee surgery. Both reviews found no differences in pain scores in the first 48 hours, or in physical activities, between FNB and epidural. Both reviews also found that participant satisfaction was greater with FNB compared with epidural analgesia. In contrast to [Fowler 2008](#), our review additionally demonstrated lower opioid consumption at 24 hours and lower risk

of nausea and/or vomiting in the FNB group. These differing results could have resulted from differences in the participant populations (TKR compared with major knee surgery).

AUTHORS' CONCLUSIONS

Implications for practice

This review found evidence that following TKR, FNB (with or without concurrent treatments including PCA opioid) provided superior analgesia compared with PCA opioid alone, similar analgesia compared with epidural and less nausea/vomiting compared with both PCA opioid alone and epidural analgesia. The review also found that continuous FNB provided a better analgesia and adverse effects profile compared with single-shot FNB. The analgesic benefits of FNB over PCA are moderate to large, and those of continuous FNB over single-shot FNB are small to moderate (Cohen 1988). These findings favour the use of FNB, in particular continuous FNB, over PCA opioid or epidural. The analgesic benefit offered by FNB, however, did not appear to extend beyond the first 72 hours.

We do not have enough evidence in this review to draw conclusions about the additional analgesic benefit of adding sciatic or obturator nerve blocks to the FNB, although results suggest that little analgesic benefit is gained by the addition of the sciatic block. Nevertheless, it was not the intent of this review to compare FNB + sciatic or FNB + obturator versus FNB. Any effects of FNB on serious adverse events were unclear because of the limited available evidence. Information from the identified RCTs was insufficient for definitive conclusions to be drawn on the comparisons between FNB and local infiltration analgesia or oral analgesia.

Implications for research

The findings of this review on local infiltration analgesia and oral analgesia were based on the results of a few small RCTs. More RCTs comparing FNB and local infiltration analgesia are warranted, given the increasing popularity of the latter and the increasing number of total knee replacement operations conducted annually worldwide. One problem of the existing evidence involves the varied components of the local infiltration analgesia used in the different studies, making it unclear which particular components

of the technique are the determinants of any observed effects. The consistent components have been the local anaesthetic (with ropivacaine or bupivacaine) and epinephrine, although in varying dosages. Conversely, the roles of the other components such as ketorolac, morphine and corticosteroid are less clear, and great variance in usage has been noted between studies. Moreover, safety concerns regarding surgical site infection or tissue necrosis have arisen, along with occurrences of cardiac toxicity in cardiac patients with the use of local infiltration analgesia. Hence, an important area for research is to determine the ideal components and safety of local infiltration analgesia. Indeed, large prospective observational studies are needed to quantify the risk of serious adverse events associated with the local infiltration analgesia.

ACKNOWLEDGEMENTS

We would like to thank Associate Prof Mike Bennett (content editor), Prof Nathan Pace (statistical editor), Prof Joanne Guay, Prof Andrew Moore (peer reviewers) and Dr Janet Wale (consumer editor) for help and editorial advice provided during the preparation of this systematic review.

We would like to thank Dr John Carlisle (content editor), Dr Marialena Trivella (statistical editor), Prof Nathan Pace (coordinating editor); Prof Joanne Guay, Prof Andrew Moore, Dr Gustavo Zanolli and Associate Prof Fiona Blyth (peer reviewers) and Dr Janet Wale and Suzanne Cunliffe (consumers) for help and editorial advice provided during the preparation of this systematic review protocol. We also thank the CARG Trials Search Co-ordinator, Dr Karen Hovhannisyan, for help with the search strategy, and the Managing Editor, Mrs Jane Cracknell, for efficient administrative support.

We would like to thank the 21 study authors who responded to our request for additional information, including three authors who regretted that the data were no longer available. We are grateful to Dr Markus Huebscher (MH) and Fereshteh Pourkazemi (FP) for determining the study inclusion and extracting data from studies for which the review authors are the named authors, Isabel Ng HL and Dr Zhang Jinbin for their help in extracting data from the Chinese studies (Tang 2010; Wang 2010; Yu 2010) and Dr Markus Huebscher for extracting data from the German study (Fritze 2009).

REFERENCES

References to studies included in this review

Adams 2002 {published data only}

Adams HA, Saatweber P, Schmitz CS, Hecker H. Postoperative pain management in orthopaedic patients: no differences in pain score, but improved stress control by epidural anaesthesia. *European Journal of Anaesthesiology* 2002;**19**(9):658-65. [PUBMED: 12243289]

Affas 2011 {published data only}

Affas F, Nygrds EB, Stiller CO, Wretenberg P, Olofsson C. Pain control after total knee arthroplasty: a randomized trial comparing local infiltration anesthesia and continuous femoral block. *Acta Orthopaedica* 2011;**82**(4):441-7. [PUBMED: 21561303]

Allen 1998 {published data only}

Allen HW, Liu SS, Ware PD, Nairn CS, Owens BD. Peripheral nerve blocks improve analgesia after total knee replacement surgery. *Anesthesia and Analgesia* 1998;**87**(1):93-7. [PUBMED: 9661553]

Baranovic 2011 {published data only}

Baranovic S, Maldini B, Milosevic M, Golubic R, Nikolic T. Peripheral regional analgesia with femoral catheter versus intravenous patient controlled analgesia after total knee arthroplasty: a prospective randomized study. *Collegium Antropologicum* 2011;**35**(4):1209-14. [PUBMED: 22397261]

Barrington 2005 {published data only}

Barrington MJ, Olive D, Low K, Scott DA, Brittain J, Choong P. Continuous femoral nerve blockade or epidural analgesia after total knee replacement: a prospective randomized controlled trial. *Anesthesia and Analgesia* 2005;**101**(6):1824-9. [PUBMED: 16301267]

Carli 2010 {published data only}

Carli F, Clemente A, Asenjo JF, Kim DJ, Mistracetti G, Gomarasca M, et al. Analgesia and functional outcome after total knee arthroplasty: periarticular infiltration vs continuous femoral nerve block. *British Journal of Anaesthesia* 2010;**105**(2):185-95. [PUBMED: 20551021]

Chan 2012 {published data only}

Chan M-H, Chen W-H, Tung Y-W, Liu K, Tan P-H, Chia Y-Y. Single-injection femoral nerve block lacks preemptive effect on postoperative pain and morphine consumption in total knee arthroplasty. *Acta Anaesthesiologica Taiwanica* 2012;**50**:54-8. [PUBMED: 22769858]

Chan 2013 {published and unpublished data}

Chan EY, Franssen M, Sathappan S, Chua NH, Chan YH, Chua N. Comparing the analgesia effects of single-injection and continuous femoral nerve blocks with patient controlled analgesia after total knee arthroplasty. *Journal of Arthroplasty* 2013;**28**(4):608-13. [PUBMED: 23142441]

Davies 2004 {published data only}

Davies AF, Segar EP, Murdoch J, Wright DE, Wilson IH. Epidural infusion or combined femoral and sciatic nerve blocks as perioperative analgesia for knee arthroplasty. *British Journal of Anaesthesia* 2004;**93**(3):368-74. [PUBMED: 15247111]

de Lima e Souza 2008 {published and unpublished data}

de Lima e Souza R, Correa CH, Henriques MD, de Oliveira CB, Nunes TA, Gomez RS. Single-injection femoral nerve block with 0.25% ropivacaine or 0.25% bupivacaine for postoperative analgesia after total knee replacement or anterior cruciate ligament reconstruction. *Journal of Clinical Anesthesia* 2008;**20**(7):521-7. [PUBMED: 19019663]

Fritze 2009 {published data only}

Fritze P, Anderl S, Marouf A, Cumlivski R, Muller C, Pernicka E, et al. Pain therapy using stimulating catheters after total knee arthroplasty. *Schmerz* 2009;**23**(3):292-8. [PUBMED: 19308464]

Ganapathy 1999 {published data only}

Ganapathy S, Wasserman RA, Watson JT, Bennett J, Armstrong KP, Stockall CA, et al. Modified continuous femoral three-in-one block for postoperative pain after total knee arthroplasty. *Anesthesia and Analgesia* 1999;**89**(5):1197-202. [PUBMED: 10553834]

Good 2007 {published data only}

Good RP, Snedden MH, Schieber FC, Polachek A. Effects of a preoperative femoral nerve block on pain management and rehabilitation after total knee arthroplasty. *American Journal of Orthopedics* 2007;**36**(10):554-7. [PUBMED: 18033568]

Hirst 1996 {published data only}

Hirst GC, Lang SA, Dust WN, Cassidy JD, Yip RW. Femoral nerve block. Single injection versus continuous infusion for total knee arthroplasty. *Regional Anesthesia* 1996;**21**(4):292-7. [PUBMED: 8837185]

Hunt 2009 {published and unpublished data}

Hunt KJ, Bourne MH, Mariani EM. Single-injection femoral and sciatic nerve blocks for pain control after total knee arthroplasty. *Journal of Arthroplasty* 2009;**24**(4):533-8. [PUBMED: 19026519]

Kadic 2009 {published data only}

Kadic L, Boonstra MC, De Waal Malefijt MC, Lako SJ, Van Egmond J, Driessen JJ. Continuous femoral nerve block after total knee arthroplasty. *Acta Anaesthesiologica Scandinavica* 2009;**53**(7):914-20. [PUBMED: 19388886]

Kaloul 2004 {published data only}

Kaloul I, Guay J, Cote C, Fallaha M. The posterior lumbar plexus (psoas compartment) block and the three-in-one femoral nerve block provide similar postoperative analgesia after total knee replacement. *Canadian Journal of Anaesthesia* 2004;**51**(1):45-51. [PUBMED: 14709460]

Kardash 2007 {published and unpublished data}

Bergeron SG, Kardash KJ, Huk OL, Zukor DJ, Antoniou J. Functional outcome of femoral versus obturator nerve block after total knee arthroplasty. *Clinical Orthopaedics & Related Research* 2009;**467**(6):1458-62. [PUBMED: 19224305]

* Kardash K, Hickey D, Tessler MJ, Payne S, Zukor D, Velly AM. Obturator versus femoral nerve block for analgesia after total knee arthroplasty. *Anesthesia and Analgesia* 2007;**105**(3):853-8. [PUBMED: 17717250]

Lee 2011 {published data only}

Lee AR, Choi DH, Ko JS, Choi SJ, Hahm TS, Kim GH, et al. Effect of combined single-injection femoral nerve block and patient-controlled epidural analgesia in patients undergoing total knee replacement. *Yonsei Medical Journal* 2011;**52**(1):145-50. [PUBMED: 21155047]

Long 2006 {published data only}

Long WT, Ward SR, Dorr LD, Raya J, Boutary M, Sirianni LE. Postoperative pain management following total knee arthroplasty: a randomized comparison of continuous epidural versus femoral nerve infusion. *The Journal of Knee Surgery* 2006;**19**(2):137-43. [PUBMED: 16642893]

Macalou 2004 {published data only}

Macalou D, Trueck S, Meuret P, Heck M, Vial F, Ouologuem S, et al. Postoperative analgesia after total knee replacement: the effect of an obturator nerve block added to the femoral 3-in-1 nerve block. *Anesthesia and Analgesia* 2004;**99**(1):251-4. [PUBMED: 15281539]

Martin 2008 {published data only}

Martin F, Martinez V, Mazoit JX, Bouhassira D, Cherif K, Gentili ME, et al. Antiinflammatory effect of peripheral nerve blocks after knee surgery: clinical and biologic evaluation. *Anesthesiology* 2008;**109**(3):484-90. [PUBMED: 18719447]

McNamee 2001 {published data only}

McNamee DA, Convery PN, Milligan KR. Total knee replacement: a comparison of ropivacaine and bupivacaine in combined femoral and sciatic block. *Acta Anaesthesiologica Scandinavica* 2001;**45**(4):477-81. [PUBMED: 11300387]

Mistraletti 2006 {published data only}

Mistraletti G, De La Cuadra-Fontaine JC, Asenjo FJ, Donatelli F, Wykes L, Schrickler T, et al. Comparison of analgesic methods for total knee arthroplasty: metabolic effect of exogenous glucose. *Regional Anesthesia & Pain Medicine* 2006;**31**(3):260-9. [PUBMED: 16701193]

Nader 2012 {published data only}

Nader A, Kendall MC, Wixson RL, Chung B, Polakow LM, McCarthy RJ. A randomized trial of epidural analgesia followed by continuous femoral analgesia compared with oral opioid analgesia on short- and long-term functional recovery after total knee replacement. *Pain Medicine* 2012;**13**(7):937-47. [PUBMED: 22680916]

Ng 2001 {published data only}

Ng HP, Cheong KF, Lim A, Lim J, Puhaindran ME. Intraoperative single-shot "3-in-1" femoral nerve block with ropivacaine 0.25%, ropivacaine 0.5% or bupivacaine 0.25% provides comparable 48-hr analgesia after unilateral total knee replacement. *Canadian Journal of Anaesthesia* 2001;**48**(11):1102-8. [PUBMED: 11744586]

Ng 2012 {published data only}

Ng FY, Ng JKF, Chiu KY, Yan CH, Chan CW. Multimodal periarticular injection vs continuous femoral nerve block after total knee arthroplasty. A prospective, crossover, randomized clinical trial. *Journal of Arthroplasty* 2012;**27**(6):1234-8. [PUBMED: 22325963]

Ozen 2006 {published data only}

Ozen M, Inan N, Tumer F, Uyar A, Baltaci B. The effect of 3-in-1 femoral nerve block with ropivacaine 0.375% on postoperative morphine consumption in elderly patients after total knee replacement surgery. *Agri Dergisi* 2006;**18**(4):44-50. [PUBMED: 17457713]

Park 2010 {published data only}

Park CK, Cho CK, Lee GG, Lee JH. Optimizing dose infusion of 0.125% bupivacaine for continuous femoral nerve block after total knee replacement. *Korean Journal of Anesthesiology* 2010;**58**(5):468-76. [PUBMED: 20532056]

Parvataneni 2007 {published data only}

Parvataneni HK, Shah VP, Howard H, Cole N, Ranawat AS, Ranawat CS. Controlling pain after total hip and knee arthroplasty using a multimodal protocol with local periarticular injections: a prospective randomized study. *Journal of Arthroplasty* 2007;**22**(6):33-8. [PUBMED: 17823012]

Salinas 2006 {published data only}

Salinas FV, Liu SS, Mulroy MF. The effect of single-injection femoral nerve block versus continuous femoral nerve block after total knee arthroplasty on hospital length of stay and long-term functional recovery within an established clinical pathway. *Anesthesia and Analgesia* 2006;**102**(4):1234-9. [PUBMED: 16551930]

Seet 2006 {published and unpublished data}

* Seet E, Leong WL, Yeo ASN, Fook-Chong S. Effectiveness of 3-in-1 continuous femoral block of differing concentrations compared to patient controlled intravenous morphine for post total knee arthroplasty analgesia and knee rehabilitation. *Anaesthesia and Intensive Care* 2006;**34**(1):25-30. [PUBMED: 16494145]

Shum CF, Lo NN, Yeo SJ, Yang KY, Chong HC, Yeo SN. Continuous femoral nerve block in total knee arthroplasty: immediate and two-year outcomes. *Journal of Arthroplasty* 2009;**24**(2):204-9. [PUBMED: 18534496]

Serpell 2001 {published data only}

Serpell MG, Millar FA, Thomson MF. Comparison of lumbar plexus block versus conventional opioid analgesia after total knee replacement. *Anaesthesia* 1991;**46**(4):275-7. [PUBMED: 2024744]

Singelyn 1998 {published data only}

Singelyn FJ, Deyaert M, Joris D, Pendevel E, Gouverneur JM. Effects of intravenous patient-controlled analgesia with morphine, continuous epidural analgesia, and continuous three-in-one block on postoperative pain and knee rehabilitation after unilateral total knee arthroplasty. *Anesthesia and Analgesia* 1998;**87**(1):88-92. [PUBMED: 9661552]

Sundarathiti 2009 {published data only}

Sundarathiti P, Ruananukul N, Channum T, Kitkunasathean C, Mantay A, Thammasakulsiri J, et al. A comparison of continuous femoral nerve block (CFNB) and continuous epidural infusion (CEI) in postoperative analgesia and knee rehabilitation after total knee arthroplasty (TKA). *Journal of the Medical Association of Thailand* 2009;**92**(3):328-34. [PUBMED: 19301724]

Szczukowski 2004 {published data only}

Szczukowski MJ Jr, Hines JA, Snell JA, Sisca TS. Femoral nerve block for total knee arthroplasty patients: a method to control postoperative pain. *Journal of Arthroplasty* 2004;**19**(6):720-5. [PUBMED: 15343531]

Tang 2010 {published and unpublished data}

Tang S, Xu ZH, Huang YG, He K, Ren LY, Qian WW, et al. Comparison of the influences of continuous femoral nerve block and patient controlled intravenous analgesia on total knee arthroplasty. [Chinese]. *Acta Academiae Medicinae Sinicae* 2010;**32**(5):574-8. [PUBMED: 21050565]

Toftdahl 2007 {published and unpublished data}

Toftdahl K, Nikolajsen L, Haraldsted V, Madsen F, Tonnesen EK, Soballe K. Comparison of peri- and intraarticular analgesia with femoral nerve block after total knee arthroplasty: a randomized clinical trial. *Acta Orthopaedica* 2007;**78**(2):172-9. [PUBMED: 17464603]

Tugay 2006 {published and unpublished data}

Tugay N, Saricaoglu F, Satilmis T, Alpar U, Akarcali I, Citaker S, et al. Single-injection femoral nerve block. Effects on the independence level in functional activities in the early postoperative period in patients with total knee arthroplasty. *Neurosciences* 2006;**11**(3):175-9. [PUBMED: 22266616]

Wang 2002 {published data only}

Wang H, Boctor B, Verner J. The effect of single-injection femoral nerve block on rehabilitation and length of hospital stay after total knee replacement. *Regional Anesthesia and Pain Medicine* 2002;**27**(2):139-44. [PUBMED: 11915059]

Wang 2010 {published data only}

Wang H-J, Zhang D-Z, Li S-Z. Comparing the analgesic efficacy of continuous femoral nerve blockade and continuous intravenous analgesia after total knee arthroplasty. [Chinese]. *Chung Hua I Hsueh Tsa Chih [Chinese Medical Journal]* 2010;**90**(33):2360-2. [PUBMED: 21092500]

Widmer 2012 {published data only}

Widmer BJ, Scholes CJ, Pattullo GG, Oussedik SI, Parker DA, Coolican MRJ. Is femoral nerve block necessary during total knee arthroplasty? A randomized controlled trial. *Journal of Arthroplasty* 2012;**27**:1800-5. [PUBMED: 22658231]

Xie 2012 {published data only}

Xie Z, Hussain W, Cutter TW, Apfelbaum JL, Drum ML, Manning DW. Three-in-one nerve block with different concentrations of bupivacaine in total knee arthroplasty. Randomized, placebo-controlled, double-blind trial. *Journal of Arthroplasty* 2012;**27**(5):673-8.e1. [PUBMED: 21945081]

Yu 2010 {published data only}

Yu H-P, Liu Z-H, Guo W-S, Jiang H-Y, Zhao J. Effect of continuous femoral nerve block in analgesia and the early rehabilitation after total knee replacement. [Chinese]. *Zhongguo Gushang [China Journal of Orthopaedics and Traumatology]* 2010;**23**(11):825-7. [PUBMED: 21254673]

Zaric 2006 {published and unpublished data}

Zaric D, Boysen K, Christiansen C, Christiansen J, Stephensen S, Christensen B. A comparison of epidural analgesia with combined continuous femoral-sciatic nerve blocks after total knee replacement. *Anesthesia and Analgesia* 2006;**102**(4):1240-6. [PUBMED: 16551931]

References to studies excluded from this review
Affas 2012 {published data only}

Affas F, Stiller CO, Nygard EB, Stephanson N, Wretenberg P, Olofsson C. A randomized study comparing plasma concentration of ropivacaine after local infiltration analgesia and femoral block in primary total knee arthroplasty. *Scandinavian Journal of Pain* 2012;**3**(1):46-51. [DOI: [10.1016/j.sjpain.2011.09.001](https://doi.org/10.1016/j.sjpain.2011.09.001)]

Bagry 2008 {published data only}

Bagry H, de la Cuadra Fontaine JC, Asenjo JF, Bracco D, Carli F. Effect of a continuous peripheral nerve block on the inflammatory response in knee arthroplasty. *Regional Anesthesia and Pain Medicine* 2008;**33**(1):17-23. [PUBMED: 18155052]

Blisard 2012 {published data only}

Blisard R, Haddad FS. Pre-emptive peripheral nerve block delays recovery after total knee arthroplasty. *Orthopedics Today* 2012; Vol. 32, issue 2:31-2.

Cappelleri 2011 {published data only}

Cappelleri G, Ghisi D, Fanelli A, Albertin A, Somalvico F, Aldegheri G. Does continuous sciatic nerve block improve postoperative analgesia and early rehabilitation after total knee arthroplasty? A prospective, randomized, double-blinded study. *Regional Anesthesia and Pain Medicine* 2011;**36**(5):489-92. [PUBMED: 21857276]

Carvalho 2012 {published data only}

Carvalho R, Braganca JP, Calixto L. Effect of a single shot sciatic nerve block combined with a continuous femoral block on pain scores after knee arthroplasty. A randomized controlled trial. *British Journal of Anaesthesia* 2012;**108**:23-4. [DOI: [10.4236/ojanes.2012.24025](https://doi.org/10.4236/ojanes.2012.24025)]

Chelly 2001 {published data only}

Chelly JE, Greger J, Gebhard R, Coupe K, Clyburn TA, Buckle R, et al. Continuous femoral blocks improve recovery and outcome of patients undergoing total knee arthroplasty. *Journal of Arthroplasty* 2001;**16**(4):436-45. [PUBMED: 11402405]

Combes 2000 {published data only}

Combes X, Cerf C, Bouleau D, Duvaldestin P, Dhonneur G. The effects of residual pain on oxygenation and breathing pattern during morphine analgesia. *Anesthesia and Analgesia* 2000;**90**(1):156-60. [PUBMED: 10624997]

Dang 2005 {published data only}

Pham Dang C, Gautheron E, Guilley J, Fernandez M, Waast D, Volteau C, et al. The value of adding sciatic block to continuous femoral block for analgesia after total knee replacement. *Regional Anesthesia and Pain Medicine* 2005;**30**(2):128-33. [PUBMED: 15765454]

Edwards 1992 {published data only}

Edwards ND, Wright EM. Continuous low-dose 3-in-1 nerve blockade for postoperative pain relief after total knee replacement. *Anesthesia and Analgesia* 1992;**75**(2):265-7. [PUBMED: 1632541]

Frassanito 2009 {published data only}

Frassanito L, Vergari A, Messina A, Pitoni S, Puglisi C, Chierichini A. Anaesthesia for total knee arthroplasty: efficacy of single-injection or continuous lumbar plexus associated with sciatic nerve blocks—a randomized controlled study. *European Review for Medical and Pharmacological Sciences* 2009;**13**(5):375-82. [MEDLINE: 19961044]

Frassanito 2010 {published data only}

Frassanito L, Vergari A, Zanghi F, Messina A, Bitondo M, Antonelli M. Post-operative analgesia following total knee arthroplasty: comparison of low-dose intrathecal morphine and single-shot ultrasound-guided femoral nerve block: a randomized, single blinded, controlled study. *European Review for Medical & Pharmacological Sciences* 2010;**14**(7):589-96. [PUBMED: 20707248]

Hakkalamani 2008 {published data only}

Hakkalamani S, Carroll FA, Ford C, Mereddy P, Jefferies G, Parkinson RW. Analgesia in total knee replacement: a comparison between femoral versus combined femoral and sciatic nerve block. *Journal of Bone and Joint Surgery (Proceedings)* 2008;**90b** (supp):328-E0A.

Ifeld 2008 {published data only}

Ifeld BM, Le LT, Meyer RS, Mariano ER, Vandenborne K, Duncan PW, et al. Ambulatory continuous femoral nerve blocks decrease time to discharge readiness after tricompartiment total knee arthroplasty: a randomized, triple-masked, placebo-controlled study. *Anesthesiology* 2008;**108**(4):703-13. [PUBMED: 18362603]

Ifeld 2009 {published data only}

Ifeld BM, Meyer RS, Le LT, Mariano ER, Williams BA, Vandenborne K, et al. Health-related quality of life after tricompartiment knee arthroplasty with and without an

extended-duration continuous femoral nerve block: a prospective, 1-year follow-up of a randomized, triple-masked, placebo-controlled study. *Anesthesia and Analgesia* 2009;**108**(4):1320-5. [PUBMED: 19299806]

Ifeld 2010 {published data only}

Ifeld BM, Mariano ER, Girard PJ, Loland VJ, Meyer RS, Donovan JF, et al. A multicenter, randomized, triple-masked, placebo-controlled trial of the effect of ambulatory continuous femoral nerve blocks on discharge-readiness following total knee arthroplasty in patients on general orthopaedic wards. *Pain* 2010;**150**(3):477-84. [PUBMED: 20573448]

Johnson 2011 {published data only}

Johnson CB, Steele-Moses SK. The use of continuous femoral nerve blocks versus extended release epidural morphine: a study comparing outcomes in total knee arthroplasty procedures. *Orthopedic Nursing* 2011;**30**(1):44-53. [MEDLINE: 21278555]

Koh 2011 {published data only}

Koh IJ, Kang YG, Chang CB, Do SH, Seong SC, Kim TK. Does periarticular injection have additional pain relieving effects during contemporary multimodal pain control protocols for TKA? A randomised, controlled study. *Knee* 2012;**19**(4):253-9. [PUBMED: 21507661]

McMeniman 2010 {published data only}

McMeniman TJ, McMeniman PJ, Myers PT, Hayes DA, Cavdarski A, Wong M, et al. Femoral nerve block vs fascia iliaca block for total knee arthroplasty postoperative pain control: a prospective, randomized controlled trial. *Journal of Arthroplasty* 2010;**25**(8):1246-9. [PUBMED: 20178889]

Meftah 2012 {published data only}

Meftah M, Wong AC, Nawabi DH, Yun RJ, Ranawat AS, Ranawat CS. Pain management after total knee arthroplasty using a multimodal approach. *Orthopedics* 2012;**35**(5):e660-4. [PUBMED: 22588407]

Morin 2005 {published data only}

Morin AM, Kratz CD, Eberhart LHJ, Dinges G, Heider E, Schwarz N, et al. Postoperative analgesia and functional recovery after total-knee replacement: comparison of a continuous posterior lumbar plexus (psoas compartment) block, a continuous femoral nerve block, and the combination of a continuous femoral and sciatic nerve block. *Regional Anesthesia and Pain Medicine* 2005;**30**(5):434-45. [PUBMED: 16135347]

Ng 2012b {published data only}

Ng F-Y, Chiu K-Y, Yan CH, Ng K-FJ. Continuous femoral nerve block versus patient-controlled analgesia following total knee arthroplasty. *Journal of Orthopaedic Surgery* 2012;**20**:23-6. [PUBMED: 22535806]

Niskanen 2005 {published data only}

Niskanen RO, Strandberg N. Bedside femoral block performed on the first postoperative day after unilateral total knee arthroplasty: a randomized study of 49 patients. *The Journal of Knee Surgery* 2005;**18**(3):192-6. [PUBMED: 16152867]

Rais 2009 {published data only}

Rais K, Ben Said A, Soussi M, Chahed S, Kaabachi O. Multimodal regional and systemic approach for postoperative analgesia in total knee arthroplasty. *European Journal of Anaesthesiology* 2009;**26**:123.

Rajeev 2007 {published data only}

Rajeev S, Batra YK, Panda NB, Kumar M, Nagi ON. Combined continuous "3-in-1" and sciatic nerve blocks provide improved postoperative analgesia with no correlation to catheter tip location after unilateral total knee arthroplasty. *Journal of Arthroplasty* 2007;**22**(8):1181-6. [PUBMED: 18078888]

Rasiah 2012 {published data only}

Rasiah R, Azhar AA, Thevanthiran MN. Single injection of femoral nerve block for bilateral total knee arthroplasty for post-operative pain management. *British Journal of Anaesthesia* 2012;**108**:ii411.

Safa 2011 {published data only}

Safa B, Haslam L, Gollish J, McCartney C. A prospective, randomized trial, comparing analgesic efficacy and postoperative functional recovery of either single shot sciatic nerve block or posterior capsule infiltration combined with femoral block for total knee arthroplasty. *Regional Anesthesia and Pain Medicine* 2011;**36**:508.

Serrano 2011 {published data only}

Serrano A, Santiveri X, Bisbe E, Ortiz P, Puig L, Castillo J. Analgesic efficacy of associating a sciatic block to a femoral block in the postoperative period of total knee arthroplasty. *European Journal of Anaesthesiology* 2011;**28**:119-20.

Sites 2004 {published data only}

Sites BD, Beach M, Gallagher JD, Jarrett RA, Sparks MB, Lundberg CJF. A single injection ultrasound-assisted femoral nerve block provides side effect-sparing analgesia when compared with intrathecal morphine in patients undergoing total knee arthroplasty. *Anesthesia and Analgesia* 2004;**99**(5):1539-43; table of contents. [PUBMED: 15502061]

Taninishi 2011 {published data only}

Taninishi H, Sato K, Morita K. Effects of anterior sciatic nerve block on intraoperative hemodynamics and pain relief in the postanesthesia care unit for patients undergoing total knee arthroplasty. *Regional Anesthesia and Pain Medicine* 2011;**36**(5):E263.

Tarkkila 1998 {published data only}

Tarkkila P, Tuominen M, Huhtala J, Lindgren L. Comparison of intrathecal morphine and continuous femoral 3-in-1 block for pain after major knee surgery under spinal anaesthesia. *European Journal of Anaesthesiology* 1998;**15**(1):6-9. [PUBMED: 9522133]

Tricarico 2009 {published data only}

Tricarico E, Tomasino S, D'Orlando L. Epidural analgesia compared with peripheral nerve blockade after major knee surgery. *Critical Care* 2009;**13**:S160.

Watson 2005 {published data only}

Watson MW, Mitra D, McLintock TC, Grant SA. Continuous versus single-injection lumbar plexus blocks: comparison of the effects on morphine use and early recovery after total knee arthroplasty. *Regional Anesthesia & Pain Medicine* 2005;**30**(6):541-7. [PUBMED: 16326339]

Wegener 2011 {published data only}

Wegener JT, van Ooij B, van Dijk CN, Hollmann MW, Preckel B, Stevens MF. Value of single-injection or continuous sciatic nerve block in addition to a continuous femoral nerve block in patients undergoing total knee arthroplasty: a prospective, randomized, controlled trial. *Regional Anesthesia & Pain Medicine* 2011;**36**(5):481-8. [PUBMED: 21857273]

Weston-Simons 2012 {published data only}

Weston-Simons JS, Pandit H, Haliker V, Dodd CA, Popat MT, Murray DW. Intra-articular local anaesthetic on the day after surgery improves pain and patient satisfaction after unicompartmental knee replacement: a randomised controlled trial. *Knee* 2012;**19**:352-5. [PUBMED: 21669534]

Zhang 2010 {published data only}

Zhang HH, Yan SH, Li X, Jin Y, Liu ZC. Analgesia following artificial joint replacement joint replacement: nerve block based on gait analysis. *Journal of Clinical Rehabilitative Tissue Engineering Research* 2010;**14**(22):4018-22.

Ziwenga 2010 {published data only}

Ziwenga O, Tsui B, Sriramatr D. Comparison of quadriceps weakness following total knee arthroplasty using analgesia by femoral nerve block: continuous vs patient controlled techniques. *Regional Anesthesia and Pain Medicine* 2010;**35**(5):E194.

References to studies awaiting assessment
Grider 2011 {published data only}

Grider JS, Harned ME, Paul S, Mauro G. Comparison of single shot versus continuous femoral local anesthetic nerve block in patients undergoing knee arthroplasty. *Regional Anesthesia and Pain Medicine* 2011;**36**(5):508-20.

Mullen 2008 {published data only}

Mullen M, Rooney BP, Kelly MP, Storey N. 24 hour infusion femoral and sciatic nerve block following total knee arthroplasty. *Scottish Medical Journal* 2008;**2**:59.

Tobin 2011 {published data only}

Tobin R, Singh MK, Sharma P, Girotra G, Arora D, Panigrahi B. Ultrasound guided in-plane continuous femoral nerve block for postoperative analgesia & early mobilisation in patients undergoing unilateral total knee arthroplasty—an Indian experience. *Regional Anesthesia and Pain Medicine* 2011;**36**(5):E263.

Yuksel 2011 {published data only}

Yuksel BE, Doger C, Erdogan N, Ornek D, Kadiogullari N. Continuous epidural analgesia versus combination of single dose sciatic nerve block and continuous femoral analgesia in

total knee arthroplasty. *Regional Anesthesia and Pain Medicine* 2011;**36**(5):E183-E4.

Additional references

Allman 2006

Allman KG, Wilson IH (editors). *Oxford Handbook of Anesthesia*. 2nd edition, Chapter 39. New York: Oxford University Press, 2006.

American Society of Anesthesiologists 2004

No authors listed. Practice guidelines for acute pain management in the perioperative setting: an updated report by the American Society of Anesthesiologists Task Force on acute pain management. *Anesthesiology* 2004;**100**(6):1573-81. [MEDLINE: 15166580]

Ballantyne 2010

Ballantyne JC, Rathmell JP, Fishman SM (editors). *Bonica's Management of Pain*. 4th Edition. Philadelphia: Lippincott, Williams & Wilkins, 2010.

Ben-David 2004

Ben-David B, Schmalenberger K, Chelly JE. Analgesia after total knee arthroplasty: is continuous sciatic blockade needed in addition to continuous femoral blockade?. *Anesthesia and Analgesia* 2004;**98**(3):747-9. [MEDLINE: 14980931]

Bender 2001

Bender R. Calculating confidence intervals for the number need to treat. *Controlled Clinical Trials* 2001;**22**:102-10.

Berdine 2006

Berdine HJ, Nesbit SA. Equianalgesic dosing of opioids. *Journal of Pain and Palliative Care Pharmacotherapy* 2006;**20**(4):79-84. [PUBMED: 17182514]

Bergeron 2009

Bergeron SG, Kardash KJ, Huk OL, Zukor DJ, Antoniou J. Functional outcome of femoral versus obturator nerve block after total knee arthroplasty. *Clinical Orthopaedics and Related Research* 2009;**467**(6):1458-62. [PUBMED: 19224305]

Borenstein 2009

Borenstein M. Effect sizes for continuous data. In: Cooper HM, Hedges LV, Valentine JC editor(s). *The Handbook of Research Synthesis and Meta-analysis*. Second Edition. New York: Russell Sage Foundation, 2009:225-7.

Capdevila 1999

Capdevila X, Barthelet Y, Biboulet P, Ryckwaert Y, Rubenovitch J, d'Athis F. Effects of perioperative analgesic technique on the surgical outcome and duration of rehabilitation after major knee surgery. *Anesthesiology* 1999;**91**(1):8-15. [MEDLINE: 10422923]

Carr 1999

Carr DB, Goudas LC. Acute pain. *Lancet* 1999;**353**(9169):2051-8. [MEDLINE: 10376632]

Choi 2003

Choi PT, Bhandari M, Scott J, Douketis J. Epidural analgesia for pain relief following hip or knee replacement. *Cochrane Database of Systematic Reviews* 2003, Issue 3. [DOI: 10.1002/14651858.CD003071]

Cohen 1988

Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. Second Edition. Hillsdale, NJ: Lawrence Erlbaum Associates, 1988.

DerSimonian 1986

DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials* 1986;**7**(3):177-88. [MEDLINE: 3802833]

Enneking 2009

Enneking FK, Wedel DJ, Horlocker TT. Chapter 14. The lower extremity: somatic blockade. In: Cousins MJ, Carr DB, Horlocker TT, Bridenbaugh PO editor(s). *Cousins and Bridenbaugh's Neural Blockade in Clinical Anesthesia and Pain Medicine*. Fourth. Philadelphia: Lippincott Williams & Wilkins, 2009:349-50.

Feibel 2009

Feibel RJ, Dervin GF, Kim PR, Beaulé PE. Major complications associated with femoral nerve catheters for knee arthroplasty: a word of caution. *The Journal of Arthroplasty* 2009;**24**:132-7. [PUBMED: 19553071]

Fischer 2008

Fischer HBJ, Simanski CJP, Sharp C, Bonnet F, Camu F, Neugebauer EAM, et al. A procedure-specific systematic review and consensus recommendations for postoperative analgesia following total knee arthroplasty. *Anaesthesia* 2008;**63**(10):1105-23. [PUBMED: 18627367]

Fowler 2008

Fowler SJ, Symons J, Sabato S, Myles PS. Epidural analgesia compared with peripheral nerve blockade after major knee surgery: a systematic review and meta-analysis of randomized trials. *British Journal of Anaesthesia* 2008;**100**(2):154-64. [PUBMED: 18211990]

Guyatt 2008

Guyatt GH, Oxman AD, Kunz R, Vist GE, Falck-Ytter Y, Schunemann HJ. What is "quality of evidence" and why is it important to clinicians. *BMJ* 2008;**336**:995-8. [MEDLINE: 18456631]

Higgins 2003

Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557-60. [MEDLINE: 12958120]

Higgins 2011

Higgins JPT, Green S (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. www.cochrane-handbook.org.

Hollis 1999

Hollis S, Campbell F. What is meant by intention to treat analysis? Survey of published randomised controlled trials. *BMJ* 1999;**319**:670-4. [MEDLINE: 10480822]

Hozo 2005

Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Medical Research Methodology* 2005; Vol. 5, issue 13. [PUBMED: 15840177]

Johnson 2013

Johnson RL, Kopp SL, Hebl JR, Erwin PJ, Mantilla CB. Falls and major orthopaedic surgery with peripheral nerve blockade: a systematic review and meta-analysis. *British Journal of Anaesthesia* 2013;**110**(4):518-28. [23440367]

Kandasami 2009

Kandasami M, Kinninmonth AW, Sarungi M, Baines J, Scott NB. Femoral nerve block for total knee replacement: a word of caution. *Knee* 2009;**16**:98-100. [MEDLINE: 19046884]

Kurtz 2007

Kurtz S, Ong K, Lau E, Mowat F, Halpern M. Projections of primary and revision hip and knee arthroplasty in the United States from 2005 to 2030. *Journal of Bone and Joint Surgery* 2007;**89**:780-5. [MEDLINE: 17403800]

Lang 1993

Lang SA, Yip RW, Chang PC, Gerard MA. The femoral 3-in-1 block revisited. *Journal of Clinical Anesthesia* 1993;**5**(4):292-6. [PUBMED: 8373606]

March 2004

March LM, Bagga H. Epidemiology of osteoarthritis in Australia. *Medical Journal of Australia* 2004;**180**:S6-10. [MEDLINE: 14984356]

McNamee 2002

McNamee DA, Parks L, Milligan KR. Post-operative analgesia following total knee replacement: an evaluation of the addition of an obturator nerve block to combined femoral and sciatic nerve block. *Acta Anaesthesiologica Scandinavica* 2002;**46**(1):95-9. [PUBMED: 11903080]

Moore 2011

Moore RA, Mhuirheartaigh RJ, Derry S, McQuay HJ. Mean analgesic consumption is inappropriate for testing analgesic efficacy in post-operative pain: analysis and alternative suggestion. *European Journal of Anaesthesiology* 2011;**28**(6):427-32. [PUBMED: 21445017]

Murphy 2008

Murphy L, Schwartz TA, Helmick CG, Renner JB, Tudor G, Koch G, et al. Lifetime risk of symptomatic knee osteoarthritis. *Arthritis and Rheumatism* 2008;**59**(9):1207-13. [MEDLINE: 18759314]

Nuesch 2010

Nuesch E, Trelle S, Reichenbach S, Rutjes A, Tschannen B, Altman DG, et al. Small study effects in meta-analyses of

osteoarthritis trials: meta-epidemiological study. *British Medical Journal* 2010;**341**:c3515. [PUBMED: 20639294]

Orwin 1983

Orwin RG. A fail-safe N for effect size in meta-analysis. *Journal of Educational Statistics* 1983;**8**(2):157-9.

PaPaS document 2011

Cochrane Pain, Palliative and Supportive Care Review Group. Authoring or Assessing a Cochrane Protocol, Review, or Review Update for the PaPaS Review Group. Version 1.0 [2011]. The Cochrane Collaboration, 2011. <http://papas.cochrane.org/sites/papas.cochrane.org/files/uploads/L%20-%20PaPaSAuthor%26RefereeGuidance.pdf>. Accessed 29 Mar 2013.

Paul 2010

Paul JE, Arya A, Hurlburt L, Cheng J, Thabane L, Tidy A, et al. Femoral nerve block improves analgesia outcomes after total knee arthroplasty: a meta-analysis of randomized controlled trials. *Anesthesiology* 2010;**113**(5):1144-62. [MEDLINE: 20966667]

RevMan 5.2 [Computer program]

The Nordic Cochrane Centre, The Cochrane Collaboration. Review Manager (RevMan). Version 5.2. Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2012.

Richman 2006

Richman JM, Liu SS, Courpas G, Wong R, Rowlingson AJ, McGready J, et al. Does continuous peripheral nerve block provide superior pain control to opioids? A meta-analysis. *Anesthesia and Analgesia* 2006;**102**:248-57. [MEDLINE: 16368838]

Sharma 2010

Sharma S, Iorio R, Specht LM, Davies-Lepie S, Healy WL. Complications of femoral nerve block for total knee arthroplasty. *Clinical Orthopaedics Related Research* 2010;**468**:135-40. [MEDLINE: 19680735]

Shum 2009

Shum CF, Lo NN, Yeo SJ, Yang KY, Chong HC, Yeo SN. Continuous femoral nerve block in total knee arthroplasty: immediate and two-year outcomes. *Journal of Arthroplasty* 2009;**24**(2):204-9. [PUBMED: 18534496]

Sinatra 2009

Sinatra RS, Leon-Casasola OA, Ginsberg B, Viscusi ER (editors). Acute Pain Management. Cambridge, MA: Cambridge University Press, 2009.

Todkar 2005

Todkar M. Sciatic nerve block causing heel ulcer after total knee replacement in 36 patients. *Acta Orthopaedica Belgica* 2005;**71**(6):724-5. [MEDLINE: 16459865]

Twersky 1997

Twersky R, Fishman D, Homel P. What happens after discharge? Return hospital visits after ambulatory surgery. *Anesthesia and Analgesia* 1997;**84**(2):319-24. [MEDLINE: 9024021]

Vendittoli 2006

Vendittoli P-A, Makinen P, Drolet P, Lavigne M, Fallaha M, Guertin M-C, et al. A multimodal analgesia protocol for total knee arthroplasty. A randomized, controlled study. *Journal of Bone and Joint Surgery* 2006;**88**(2):282-9. [PUBMED: 16452738]

Wilson 1927

Wilson EB. Probably inference, the law of succession, and statistical inference. *Journal of American Statistical Association* 1927;**22**:209-12.

Winnie 1973

Winnie AP, Ramamurthy S, Durrani Z. The inguinal paravascular technic of lumbar plexus anesthesia: the "3-in-1 block". *Anesthesia and Analgesia* 1973;**52**:989-96. [MEDLINE: 4796576]

References to other published versions of this review

Chan 2012

Chan E-Y, Fransen M, Parker DA, Assam PN, Chua N. Femoral nerve blocks for acute postoperative pain after knee replacement surgery. *Cochrane Database of Systematic Reviews* 2012, Issue 7. [DOI: [10.1002/14651858.CD009941](https://doi.org/10.1002/14651858.CD009941)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES

Characteristics of included studies [ordered by study ID]

Adams 2002

Methods	Randomized trial: drawing lots
Participants	<p>Single centre in Hannover, Germany</p> <p>63 participants undergoing unilateral TKR</p> <p>Inclusions: ASA I to III</p> <p>Exclusion: age younger than 18 years, endocrine disorders except diabetes mellitus, glucocorticoid therapy, use of etomidate up to 48 hours before the beginning of study and allergy towards one of the test substances. Patients requesting another method than the randomly intended postoperative pain management</p> <p>Mean age (years): 69 (range 56 to 86)</p> <p>Female: 70%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Three-in-one single-shot FNB with bupivacaine 40 ml 0.375% solution after detection of the nerve using a nerve stimulator</p> <p>2. Continuous epidural with perioperative bupivacaine test dose 3 ml 0.25% solution and postoperative bupivacaine 0.375% (1 ml per 10 cm body height, maximum 15 ml). Started at first expression of pain at PACU</p> <p>3. IV piritramide with initial small bolus injections of 0.1 to 0.2 mg/kg and diclofenac 100 mg rectally, and subsequently single doses of piritramide—about 0.025 mg/kg (2 mg at 80 kg) from a PCA pump with 'lockout' time of 10 minutes. Started at first expression of pain at PACU. Diclofenac 100 mg was administered rectally</p>
Outcomes	<p>Pain intensity at rest: visual analogue scale at one hour, two hours, three hours</p> <p>Nausea/vomiting: first postoperative day</p> <p>Participant satisfaction with postoperative management: first postoperative day</p>
Notes	Study author provided additional information on methods of randomization and blinding

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Adams 2002 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "random." Drawing lots (study author provided information)
Allocation concealment (selection bias)	Unclear risk	No mention (study author unable to provide additional information)
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	"None were blinded" (study author provided information)
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	"None were blinded" (study author provided information)
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Affas 2011

Methods	Randomized trial: mixing 40 tickets and drawing one envelope at a time
Participants	<p>Orthopaedics clinic at Karolinska University Hospital in Solna, Sweden</p> <p>40 participants with osteoarthritis or rheumatoid arthritis scheduled for primary unilateral elective TKR under spinal anaesthesia</p> <p>Inclusion criteria: ASA I to III and older than 18 years of age</p> <p>Exclusion criteria: allergy or intolerance to one of the study drugs, renal insufficiency, epilepsy, language difficulty, mental illness, dementia, QT interval on electrocardiogram > 450 ms before start</p> <p>Mean age (years): 68 (range 29 to 88)</p> <p>Female: 53%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB with 30 ml ropivacaine (2 mg/ml) was injected, followed by 15 ml of the same concentration every four-hourly for 24 hours (total dose 240 mg/24 h). Nerve identified via nerve stimulation</p> <p>2. Periarticular and intra-articular infiltration of a solution containing 150 ml ropivacaine (2 mg/ml), 1 ml ketorolac (30 mg/ml) and 5 ml epinephrine (0.1 mg/ml). Total dose of ropivacaine was 300 mg</p>

Affas 2011 (Continued)

Adjunct analgesics: IV ketorolac 30 mg given during the first 24 hours post operation, IV PCA morphine (2 mg/dose) given on demand with a lock-out time of six minutes and maximum dose of 35 mg over four hours and paracetamol 1 g × 4 over 24 hours

Outcomes	Pain intensity at rest and on movement: visual analogue scale at 24 hours Total opioid consumption (mg) at 24 hours Proportion of participants in pain at 24 hours Adverse events
Notes	One participant from the FNB group had a history of insensitivity to pain. He did not demand any PCA morphine, and pain severity was assessed as no pain at all time points

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "The randomization sequence was determined by mixing 40 tickets, 20 labelled 'F' and 20 labelled 'LIA' in sealed opaque envelopes, and drawing one envelope at a time. The anaesthesiologist performing the spinal anaesthesia and femoral block or supervising the local infiltration analgesia technique did not participate in the randomization procedure"
Allocation concealment (selection bias)	Low risk	Quote: "...sealed opaque envelopes, and drawing one envelope at a time"
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Quote: "open labelled"
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Quote: "open labelled"
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Allen 1998

Methods	Randomized trial: method not stated
Participants	Single centre from Washington, USA 36 participants undergoing unilateral TKR Exclusion criteria: age < 40 or > 80 years; ASA physical status greater than III; allergy to local anaesthetics; history of opioid dependence; contraindications to spinal anaesthesia, femoral nerve block or sciatic nerve block (coagulation defects, infection at puncture site, preexisting neurological deficits in the lower extremities); inability to use patient-controlled analgesia (IV PCA), contraindications for NSAID

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Allen 1998 (Continued)

use (NSAID or aspirin allergy, severe liver disease or serum creatinine 21.7 mg/dl), weight > 140 kg and recent steroid use

Mean age (years): 68

Female: 61%

Number excluded post randomization: none

Number lost to follow-up: none

Interventions	<p>1. Single-shot FNB (with sham sciatic) with 30 ml of 0.25% bupivacaine with 1:400,000 epinephrine was injected. Nerve identified via nerve stimulation. IV PCA morphine initial bolus of 1 mg and lockout interval of 10 minutes with no limit or background infusion; further adjustments at the discretion of the orthopaedic doctor</p> <p>2. Single-shot FNB + sciatic with 30 ml of 0.25% bupivacaine with 1:400,000 epinephrine was injected. Nerve identified via nerve stimulation. IV PCA morphine initial bolus of 1 mg and lockout interval of 10 minutes with no limit or background infusion; further adjustments at the discretion of the orthopaedic doctor</p> <p>3. IV PCA morphine initial bolus of 1 mg and lockout interval of 10 minutes with no limit or background infusion; further adjustments at the discretion of the orthopaedic doctor</p> <p>Adjunct analgesics: IV ketorolac either 15 (< 65 years or < 50 kg) or 30 mg every six hours for at least 24 hours. If oral medications were tolerated after 24 hours, then oral ibuprofen 600 mg every eight hours was substituted for ketorolac</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale at one hour upon arrival on ward and every four hours until 10 pm, and morning and evening of postoperative days one to five</p> <p>Total opioid consumption: postoperative days one and two</p> <p>Participant satisfaction: two weeks after discharge</p> <p>Nausea/vomiting</p> <p>Sedation</p> <p>Pruritus</p> <p>Other outcomes:</p> <p>Technical block to pinprick of saphenous nerve</p> <p>Technical block to pinprick of sural nerve</p>
Notes	<p>Additional information regarding the study method sought, but study authors could not be reached. For pain at rest, the data presented in the figures were not used for the meta-analysis, as quoted from the report: "Data were incomplete after the 8h measurement and are not presented"</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "random." Method not stated
Allocation concealment (selection bias)	Unclear risk	No mention

Allen 1998 (Continued)

Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Control group had sham injections with dressings. All participants told to expect numbness and motor weakness
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Blinded investigators collected data
Incomplete outcome data (attrition bias) All outcomes	Low risk	For pain severity at rest: report quoted: "Data were incomplete after the 8 h measurement and are not presented." Apart from this, no dropouts/withdrawals reported
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Baranovic 2011

Methods	Randomized trial: computer-generated list
Participants	<p>Single centre from Zagreb, Croatia</p> <p>80 participants undergoing TKR under spinal anaesthesia</p> <p>Inclusion criteria: ASA II to III</p> <p>Exclusion criteria: previous vascular surgery in the region of femoral veins or arteries; confirmed coagulopathy; local infection; hepatic and renal insufficiency; dementia; BMI > 30 kg/m²; allergy to local anaesthetic, morphine and NSAID drugs; a previously diagnosed neurological deficit</p> <p>Mean age (years): 70 (range 66 to 73)</p> <p>Female: 48%</p> <p>Number excluded post randomization: nine in continuous FNB because femoral catheter fell out</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB using nerve stimulation. Bolus dose 8 ml 0.35% levobupivacaine, then 5 to 6ml per hour of 0.25% levobupivacaine via catheter in addition to IV morphine 5 mg (wt < 60 kg) or 10 mg (wt > 60 kg), if posterior knee pain > three. Femoral catheter in situ for 24 hours</p> <p>2. PCA morphine concentration 1 mg/ml, basal rate 3 mg/h, bolus upon request 2 mg, lockout interval eight minutes</p> <p>Adjunct analgesics: IV diclofenac 75 mg 12-hourly</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale at two-hourly × 24 hours</p> <p>Opioid consumption: first 24 hours</p> <p>Nausea/vomiting: first 24 hours</p> <p>Sedation: first 24 hours</p> <p>Urinary retention</p>

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Baranovic 2011 (Continued)

Knee flexion: postoperative day two

Participant satisfaction

Other outcomes:

Haemodynamic instability

Length of stay

Notes	Additional information regarding study method sought, but study authors could not be reached
-------	--

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Quote: "...randomized using statistical software Medcalc for Windows"
Allocation concealment (selection bias)	Unclear risk	No mention
Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	No mention
Blinding of outcome assessment (detection bias) Subjective outcomes	Unclear risk	Quote: "The VAS scores were assessed by persons who were not involved in setting the femoral nerve catheter." It may be obvious that the FNB group did not receive an IV PCA
Incomplete outcome data (attrition bias) All outcomes	High risk	Nine participants (20%) in the continuous FNB group excluded post randomization because their catheter fell out
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	High risk	ITT an issue with the high risk of attrition bias

Barrington 2005

Methods	Randomized trial: computer-generated in permuted blocks of four
Participants	<p>Single centre from St Vincent's Hospital, Melbourne, Australia</p> <p>112 participants undergoing primary TKR</p> <p>Exclusion criteria: inability to give informed consent for language or cognitive reasons; contraindications to neuraxial blockade (including participant refusal, platelet count $100 \times 10^9/L$ or coagulopathy); contraindications to continuous FNB (e.g. infection overlying injection site or previous femoropopliteal bypass surgery); and contraindication to any study drugs</p> <p>Mean age (years): 70 ± 10</p> <p>Female: 53%</p> <p>Number excluded post randomization: none</p>

Barrington 2005 (Continued)

Number lost to follow-up: four (three protocol violations from continuous FNB group and one data lost from epidural group)

Interventions	<p>1. Continuous FNB with femoral infusion of bupivacaine 0.2% commenced at 0.1 ml/kg/h, with PCA bolus of 0.05 ml/kg and 60-minute lockout period. Nerve identified via nerve stimulation</p> <p>2. Continuous epidural with infusion of ropivacaine 0.2% plus fentanyl 4 g/ml commenced at 6 to 10 ml/h; subsequently the acute pain service varied the infusion rate to maintain sensory blockade of the surgical site. Both interventions continued until postoperative day three morning</p> <p>Adjunct analgesics: rofecoxib 50 mg daily and oxycodone 10 to 15 mg every four hours, as required. If analgesia was inadequate despite oral adjuncts, an IV morphine infusion was commenced</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale at postoperative days one and two at rest, and during CPM and active physiotherapy</p> <p>Nausea and vomiting (0 = no nausea, 1 = nausea only, 2 = nausea and vomiting)</p> <p>Knee flexion: postoperative days one and two</p> <p>Hypotension</p> <p>Neurological injury</p> <p>Technical failure of blocks</p> <p>Catheter dislodgement</p> <p>Sit out of bed on postoperative day one</p> <p>Other outcomes:</p> <p>Drainage blood</p> <p>Number of participants who required rescue analgesics</p> <p>Length of stay</p>
Notes	<p>Different local anaesthetics used in the two allocation groups (bupivacaine in continuous FNB group and ropivacaine in epidural group). Also epidural group had fentanyl in the infusion group but not in the continuous FNB group. Different modalities included continuous infusion in the epidural group and PCA in the continuous FNB group</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated in permuted blocks of four
Allocation concealment (selection bias)	Low risk	Sequentially numbered, opaque sealed envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Participants and treated clinicians not blinded to study group randomization
Blinding of outcome assessment (detection bias)	High risk	Quotes: "The motor block was assessed...by unblinded physiotherapists..."; "The limitations of this study include its non-blinded nature..."

Barrington 2005 (Continued)

Subjective outcomes

Incomplete outcome data (attrition bias) All outcomes	Low risk	Four participants not included in the analysis (three protocol violations from continuous FNB group and one data lost from epidural group). Apart from this, no other dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes prespecified
Other bias	Low risk	Appears to be free of other sources of bias

Carli 2010

Methods	Randomized trial: computer-generated tables
Participants	<p>Single centre from McGill University Health Centre (MUHC), Canada</p> <p>40 participants undergoing primary, unilateral, tricompartamental cemented TKR</p> <p>Inclusion criteria: older than 18 years of age and referred for TKR secondary to osteoarthritis</p> <p>Exclusion criteria: ASA IV and V, history of abnormal liver enzymes, hepatic failure, renal insufficiency, cardiac failure, organ transplant, morbid obesity (BMI 40 kg/m²), neuropathic pain, history of stroke or major neurological deficit, sensory and motor disorders in the operated limb, previous drug dependency, long-term use of opioids, allergy to local anaesthetics, inability to walk independently and inability to comprehend pain assessment</p> <p>Mean age (years): 71 (range 55 to 85)</p> <p>Female: 73%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB with loading dose of 8 ml of ropivacaine 0.2%, continuous infusion of ropivacaine 0.2% at a rate of 8 ml/h for 48 hours. Infiltration of the posterior capsule of the knee with a solution containing 100 mg of ropivacaine (50 ml of ropivacaine 0.2%), 0.5 ml of ketorolac (30 mg/ml) and 0.25 ml of epinephrine (1 mg/ml) (after all bony cuts and before cementing of implants). Sham periarticular and intra-articular infiltration with saline 0.9%</p> <p>2. Sham continuous FNB with saline 0.9%. Infiltration of posterior capsule of the knee with a solution containing 100 mg of ropivacaine (50 ml of ropivacaine 0.2%), 0.5 ml of ketorolac (30 mg/ml) and 0.25 ml of epinephrine (1 mg/ml) (after all bony cuts and before cementing of implants)</p> <p>Intra-articular injection with 100 ml of ropivacaine 0.2%, 1 ml of ketorolac (30 mg/ml) and 0.5 ml of epinephrine (1 mg/ml) infiltrated into remainder of capsule, cut quadriceps tendon, patellar ligament and soft tissues surrounding joint and subcutaneous tissues after closure of medial parapatellar arthrotomy; 50 ml of ropivacaine 0.5% with ketorolac (30 mg/ml) and 0.25 ml of epinephrine (1 mg/ml) infused over a two-hour period at 24 hours after the end of the operation via intra-articular catheter</p> <p>Both groups: solution containing 100 mg of ropivacaine (50 ml of ropivacaine 0.2%), 0.5 ml of ketorolac (30 mg/ml) and 0.25 ml of epinephrine (1 mg/ml) was injected into the posterior capsule of the knee</p> <p>PCA pump to deliver incremental doses of 1 mg of morphine, with lockout of seven minutes and no background infusion, discontinued 48 hours after surgery</p>

Carli 2010 (Continued)

Adjunct analgesics: slow-release oral oxycodone 10 mg every 12 hours. For breakthrough pain with NRS > 3, oral oxycodone 5 to 10 mg every three hours was prescribed for three days. Celecoxib 100 mg every 12 hours and acetaminophen 1 g every six hours, respectively, for five days

Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain severity at rest and on movement: numerical rating scale zero to 10 at first three days, upon discharge, six weeks after discharge</p> <p>Total opioid consumption: first 48 hours</p> <p>Nausea/vomiting: first 48 hours</p> <p>Knee flexion: postoperative days one and two</p> <p>Other outcomes:</p> <p>Length of stay</p> <p>Knee flexion and extension: week six</p> <p>SF-12</p> <p>WOMAC</p> <p>Knee Society Score</p> <p>Two-minute walking test</p> <p>Six-minute walking test</p> <p>Ambulated > 30 m</p>	
Notes		
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated tables
Allocation concealment (selection bias)	Low risk	Sealed brown envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Clinical staff and participants not aware of treatment allocations
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Research assistant unaware of the study hypothesis. Different treatment groups may not be obvious, as both groups received IV PCA opioid
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Chan 2012

Methods	Randomized trial: computer random number generator
Participants	<p>Single centre from Taipei, Taiwan</p> <p>88 participants undergoing unilateral primary TKR</p> <p>Exclusion criteria: < 40 and > 80 years old, ASA > III, known hypersensitivities to test substances used in study, history of substance abuse, contraindications to spinal anaesthesia, having femoral neuropathy or communication difficulties</p> <p>Mean age (years): single-shot FNB 67.7 ± 8.9 versus PCA 71.4 ± 7.9</p> <p>Female: single-shot FNB 75.6% versus PCA 70.7%</p> <p>Number excluded post randomization: six excluded because of severe allergy to morphine, communication difficulties, multiple operations and loss to follow-up</p> <p>Number lost to follow-up: unclear (from the number of withdrawals, could range from one to three participants)</p>
Interventions	<p>1. Single-shot FNB 0.4 ml/kg 0.375% bupivacaine and 1:200,000 epinephrine after spinal anaesthesia but before surgery. Used ultrasound imaging and nerve stimulators. IV PCA morphine bolus 1 mg with five-minute lockout (four-hour maximum dose of 30 mg) started at PACU. Dosage adjusted by 0.4 mg every one to two hours, if pain control was unsatisfactory</p> <p>2. Single-shot FNB 0.4 ml/kg 0.375% bupivacaine and 1:200,000 epinephrine immediately after surgery. Used ultrasound imaging and nerve stimulators. IV PCA morphine bolus 1 mg with five-minute lockout (four-hour maximum dose of 30 mg) started at PACU. Dosage adjusted by 0.4 mg every one to two hours, if pain control was unsatisfactory</p> <p>3. Sham FNB 0.4 ml/kg saline and 1:200,000 epinephrine after spinal anaesthesia but before surgery. IV PCA morphine bolus 1 mg with five-minute lockout (four-hour maximum dose of 30 mg) started at PACU. Dosage adjusted by 0.4 mg every one to two hours, if pain control was unsatisfactory. Sham FNB with saline given</p> <p>4. Sham FNB 0.4 ml/kg saline and 1:200,000 epinephrine immediately after surgery. IV PCA morphine bolus 1 mg with five-minute lockout (four-hour maximum dose of 30 mg) started at PACU. Dosage adjusted by 0.4 mg every one to two hours, if pain control was unsatisfactory. Sham FNB with saline given</p> <p>Adjunct analgesics: not mentioned</p>
Outcomes	<p>Pain intensity at rest and on movement: visual analogue scale at two hours, four hours, six hours, 24 hours, 48 hours, 72 hours</p> <p>Opioid consumption: 24 hours, 48 hours</p> <p>Time to first morphine request</p> <p>Nausea/vomiting</p> <p>Pruritus</p> <p>Dizziness</p> <p>Knee flexion</p> <p>Neuropathy of FNB</p> <p>Postoperative falls</p>

Chan 2012 (Continued)

Notes For the meta-analysis, we combined single-shot FNB groups (before and after surgery); similarly we combined sham FNB groups (before and after surgery) (i.e. interventions one and two vs interventions three and four)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated random numbers
Allocation concealment (selection bias)	Unclear risk	Sealed envelopes. Not mentioned whether sequentially labelled and opaque
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Quote: "Double blind"; "The patients, the pain management team, and the chart analysts were not informed of the patients' group assignment"
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Quote: "Double blind"; "The patients, the pain management team, and the chart analysts were not informed of the patients' group assignment"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Six participants excluded. Missing outcome data due to not meeting inclusion criteria
Selective reporting (reporting bias)	Low risk	Analysis predefined
Other bias	Low risk	Appears to be free of other sources of bias

Chan 2013

Methods	Randomized trial: computer random number generator
Participants	<p>Single centre from Singapore</p> <p>200 participants undergoing unilateral primary TKR</p> <p>Inclusion criteria: 40 to 85 years of age; ASA I to III; BMI < 35 kg/m²; competent in using VAS pain score</p> <p>Exclusion criteria: contraindication to femoral block; abnormal coagulation studies; thrombocytopenia < 100,000/cc; renal insufficiency; lower extremities thromboembolic diseases; allergy to study medications; taking daily opioid past two weeks before surgery</p> <p>Mean age (years): continuous FNB: 66.4 ± 8.3 versus single-shot FNB: 66.1 ± 7.6 versus PCA: 64.7 ± 8.4</p> <p>Female: continuous FNB 81.5% versus single-shot FNB 82.6% versus PCA 80.3%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	1. Continuous FNB 20 ml of 0.25% bupivacaine with 1:400,000 adrenaline (2.5 mcg/ml) followed by continuous infusion bupivacaine 0.125% 4 ml/h started at PACU for 72 hours. Oral oxycodone 5 mg four-hourly when necessary

Chan 2013 (Continued)

2. Single-shot FNB 20 ml of 0.25% bupivacaine with 1:400,000 adrenaline (2.5 mcg/ml); IV PCA morphine bolus 1 mg with five-minute lockout (maximum 10 mg/h) until day three; for participants at risk of developing renal impairment, fentanyl 20 mcg dose with five-minute lockout (maximum dose of 240 mcg/h)

3. IV PCA morphine bolus 1 mg with five-minute lockout (maximum 10 mg/h) until day three; for participants at risk of developing renal impairment, fentanyl 20 mcg dose with five-minute lockout (maximum dose of 240 mcg/h)

Adjunct analgesics: days one to six: oral paracetamol 1 g six-hourly; and equianalgesic dose of ibuprofen or COX-2 inhibitor. From 72 hours: oral oxycodone 5 mg six-hourly when necessary

Outcomes

Outcomes of interest for the review:

Pain intensity at rest and on movement: visual analogue scale at one hour, six hours, 12 hours, 24 hours, 48 hours, 72 hours, 120 hours, week two, month three

Opioid consumption: postoperative days one and two

Proportion of participants in significant pain: 24 hours, 48 hours

Nausea/vomiting: 24 hours, 48 hours

Pruritus

Urinary retention

Adverse events

Participant satisfaction

Knee flexion: daily until discharge

Knee extension: daily until discharge

Other outcomes:

Day achieved 90 degrees knee flexion

Day achieved rehabilitation indices

Straight leg raised

Length of stay

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer random number generator
Allocation concealment (selection bias)	Low risk	Opaque sealed envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Unblinded

Chan 2013 (Continued)

Blinding of outcome assessment (detection bias) Subjective outcomes	Unclear risk	Data collector unaware of study objectives. May be obvious that continuous FNB group did not receive an IV PCA
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Davies 2004

Methods	Randomized trial: computer random number generator
Participants	<p>Single centre from Exeter, UK</p> <p>60 participants undergoing unilateral primary TKR</p> <p>Exclusion criteria: ASA > III or had a contraindication to use of non-steroidal anti-inflammatory drugs, local anaesthetic agent, neuraxial blockade or tourniquet usage; painful polyarthralgia</p> <p>Mean age (years): 72 ± 9</p> <p>Female: 70%</p> <p>Number excluded post randomization: one from epidural group because of insertion failure</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Single-shot FNB + sciatic with 0.375% bupivacaine 30 ml for the femoral component and 25 ml for the sciatic component</p> <p>2. Continuous lumbar epidural analgesia 0.5% bupivacaine, test dose 3 ml, a further 7 ml and an infusion of bupivacaine 0.25% until postoperative day two</p> <p>Adjunct analgesics: 1-mg bolus of morphine sulphate with lockout of five minutes; diclofenac 50 mg every eight hours</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity on movement: visual analogue scale at six hours, 24 hours, 48 hours</p> <p>Opioid consumption: 24 hours, 48 hours</p> <p>Urinary retention</p> <p>Hypotension</p> <p>Participant satisfaction with analgesics: 24 hours, 48 hours</p> <p>Straight leg raised</p> <p>Knee flexion</p> <p>Other outcomes:</p> <p>Motor block</p>

Davies 2004 (Continued)

Complete analgesia at recovery unit

Total perioperative blood loss

Total drain loss

Length of stay

Notes Additional information regarding the study method sought, but study authors could not be reached

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer random number generator
Allocation concealment (selection bias)	Low risk	Codes stored by third party in opaque sealed envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	Surgeon blinded. Not stated for participants
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Clinical staff uninvolved with the study
Incomplete outcome data (attrition bias) All outcomes	Low risk	One participant excluded. No other dropouts/exclusions
Selective reporting (reporting bias)	Low risk	Analysis predefined
Other bias	Low risk	Appears to be free of other sources of bias

de Lima e Souza 2008

Methods	Randomized trial: computer-assisted randomization
Participants	<p>Single centre from Minas Gerais, Brazil</p> <p>61 participants undergoing unilateral TKR</p> <p>Inclusion criteria: ASA II to III</p> <p>Exclusion criteria: patient refusal of spinal anaesthesia or FNB; age < 18 or > 90 years; weight less than 40 kg; BMI > 45 kg/m²; allergy to local anaesthetics; preexisting neurological deficit; or inability to understand VAS scoring</p> <p>Mean age (years): 65 ± 9,2 (author provided information)</p> <p>Female: 72% (author provided information)</p> <p>Number excluded post randomization: none</p>

de Lima e Souza 2008 (Continued)

Number lost to follow-up: none

Interventions	<p>1. Single-shot FNB with 40 ml of 0.25% ropivacaine 100 mg, nerve stimulation; IV PCA morphine 2 mg on participant demand</p> <p>2. Single-shot FNB with 40 ml of 0.25% bupivacaine with epinephrine 1:200,000 (100 mg); nerve stimulation; IV PCA morphine 2 mg on participant demand</p> <p>3. IV PCA morphine 2 mg on participant demand; no sham FNB</p> <p>Adjunct analgesics: 30 mg oral codeine plus acetaminophen 0.5 g every six hours starting at the end of surgery, then 20 mg of tenoxicam 12/12 hours</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Proportion of pain at rest and on movement: visual analogue scale, zero to six hours, six to 10 hours, 10 to 24 hours</p> <p>Adverse effects: urinary retention requiring catheterization, nausea, vomiting and sedation</p> <p>Other outcomes:</p> <p>Pruritus</p> <p>Use of morphine (yes/no)</p> <p>Participant satisfaction (yes/no)</p>
Notes	Study author provided additional information on proportion of participants in pain, use of morphine and pain satisfaction

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-assisted, randomized treatment assignments
Allocation concealment (selection bias)	Low risk	Sequentially ordered sealed envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Quote (from report): "double blind." Participants and anaesthesiologist blinded (data from study author)
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Blinded research assistant
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Fritze 2009

Methods	Randomized trial: method not stated
Participants	<p>Single hospital (Orthopaedic Department) in Vienna, Austria</p> <p>54 participants undergoing TKR</p> <p>Inclusion criteria: ASA I to III, no contraindications against the procedure (allergies, skin alterations, blood-clotting disorder, previous hip or spine surgery, no contraindications against used medications), over 50 years</p> <p>Exclusion criteria: insufficient cooperation, major surgery (e.g. osteotomy, replacement TKR implant), contraindications against used medications, scheduled spinal anaesthesia</p> <p>Mean age (years): 70 (range 50 to 85)</p> <p>Female: 78%</p> <p>Number excluded post randomization: two (catheter dislocation)</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB with initial 10 ml 1% prilocaine and 20 ml 0.75% ropivacaine; postoperative infusion 0.2% ropivacaine 5 ml/h until postoperative day three</p> <p>2. Continuous FNB + sciatic with initial 10 ml 1% prilocaine and 20 ml 0.75% ropivacaine; postoperative infusion 0.2% ropivacaine 5 ml/h until postoperative day three</p> <p>3. Continuous epidural catheter with test dose 4 ml 1% xylocaine, bolus 10 ml 0.75% ropivacaine; post-operative infusion 0.2% ropivacaine 7 ml/h</p> <p>Adjunct analgesics: oral lornoxicam twice 8 mg, tramadol thrice 100 mg, in cases of severe pain post operation: IV piritramid 3 mg every 10 minutes (if in intensive care unit during first post operation night) until pain relief or SC 15 mg every four hours until pain relief in general ward</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale at one hour, 24 hours, 48 hours, 72 hours</p> <p>Knee flexion: 48 hours</p> <p>Pulmonary embolism</p> <p>Other outcomes:</p> <p>Time in bed (hours)</p> <p>Haematoma at catheter site</p>
Notes	Additional information regarding the study method sought, but study authors could not be reached

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not stated
Allocation concealment (selection bias)	Low risk	Sealed opaque envelopes

Fritze 2009 (Continued)

Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	Not stated
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Performed by physician on duty
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Ganapathy 1999

Methods	Randomized trial: closed envelope system by the pharmacy
Participants	<p>Single centre from London, Ontario, Canada</p> <p>62 participants undergoing TKR</p> <p>Inclusion criteria: > 18 and < 80 years, ASA I to III</p> <p>Exclusion criteria: significant medical or psychiatric problems, unable to co-operate with study protocol, contraindications to spinal anaesthesia or had allergy to bupivacaine, indomethacin, morphine or codeine</p> <p>Mean age (years): 70 ± 9</p> <p>Female: 50%</p> <p>Number excluded post randomization:none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB with 0.2% bupivacaine via the femoral catheter 10 ml/h and continued for 48 hours. Nerve identified via nerve stimulation. IV PCA morphine 1.5 mg with lockout time of six minutes, increased to 2 mg if pain control poor and VAS pain score > 50/100-mm scale</p> <p>2. Continuous FNB with 0.1% bupivacaine via the femoral catheter 10 ml/h and continued for 48 hours. Nerve identified via nerve stimulation. IV PCA morphine 1.5 mg with lockout time of six minutes, increased to 2 mg if pain control poor and VAS pain score > 50/100-mm scale; sham block</p> <p>3. IV PCA 1.5 mg with lockout time of six minutes, increased to 2 mg if pain control poor and VAS pain score > 50/100-mm scale; sham block</p> <p>Adjunct analgesics: 100 mg of indomethacin rectally every 12 hours for 48 hours</p>
Outcomes	<p>Pain intensity: visual analogue scale at day of surgery, at postoperative days one and two, on discharge</p> <p>Total opioid consumption: day of surgery to postoperative day one</p> <p>Range of flexion</p>

Ganapathy 1999 (Continued)

Nausea and vomiting

Notes	Study author provided additional information on methods of randomization, allocation concealment and blinding
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Closed envelope system by the pharmacy" (study author provided information)
Allocation concealment (selection bias)	Low risk	Drug labelled as study drug by the pharmacy. Randomization not broken unless medical emergencies (study author provided information)
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	"Patients and personnel were blinded" (study author provided information)
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	A blinded independent observer evaluated VAS scores, a blinded physiotherapist recorded range of flexion and a blinded surgeon assessed range of flexion at six weeks
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Good 2007

Methods	Randomized trial: coded and supplied by hospital pharmacy
Participants	<p>Single centre from Bryn Mawr Hospital, Pennsylvania, USA</p> <p>42 participants undergoing TKR</p> <p>Inclusion criteria: not stated</p> <p>Exclusion criteria: not stated</p> <p>Mean age (years): 65</p> <p>Female: 38%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Single-shot FNB with 40 ml 0.5% bupivacaine with epinephrine 1:200,000 before surgery; nerve identified via nerve stimulator. IV morphine 1 mg/h, max 10 mg/h, continued until postoperative day one</p> <p>2. IV morphine 1 mg/h, max 10 mg/h, continued until postoperative day one with 40 ml 0.5% saline given as placebo block</p>

Good 2007 (Continued)

Adjunct analgesics: not stated

Outcomes	Outcomes of interest for the review: Pain intensity on movement: visual analogue scale at every eight hours on day of surgery, daily from postoperative day one to three Total opioid consumption Nausea/vomiting Urinary retention Infection Dyspnoea Knee flexion and extension: postoperative days two and three Other outcomes: Difficulty in ambulation Fever Ambulation distance based on graded scale
Notes	Study author provided additional information on method of randomization

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Coded and supplied by hospital pharmacy (study author provided information)
Allocation concealment (selection bias)	Low risk	Coded and supplied by hospital pharmacy. Unblinded at end of study
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Blinded ortho surgeon, anaesthesiologist, physical therapist, clinical nursing staff, participants
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Blinded outcome assessor.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Hirst 1996

Methods	Randomized trial: method not stated
Participants	<p>Single centre from Saskatchewan, Canada.</p> <p>33 participants undergoing TKR</p> <p>Inclusion criteria: ASA I to III</p> <p>Exclusion criteria: not stated</p> <p>Mean age (years): 69</p> <p>Female: 79%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Single-shot FNB with 20 ml 0.05% bupivacaine with 1:200,000 adrenaline. Nerve identified via nerve stimulation. IV PCA morphine 1 mg/ml, dose 1.5 mg, lockout seven minutes, four-hour dose limit 30 mg. If inadequate, increase to 2 mg/ml, lockout five minutes</p> <p>2. Continuous FNB with 20 ml 0.05% bupivacaine with 1:200,000 adrenaline and Infusion 0.125% bupivacaine 6 ml/h for 48 hours. IV PCA morphine 1 mg/ml, dose 1.5 mg, lockout seven minutes, four-hour dose limit 30 mg. If inadequate, increase to 2 mg/ml, lockout five minutes</p> <p>3. Sham FNB. IV PCA morphine 1 mg/ml, dose 1.5 mg, lockout seven minutes, four-hour dose limit 30 mg. If inadequate, increase to 2 mg/ml, lockout five minutes</p> <p>Adjunct analgesics: not stated</p>
Outcomes	<p>Pain intensity at rest and on movement: visual analogue scale two hours, six hours, 12 hours, 18 hours, 24 hours, 48 hours, 72 hours</p> <p>Total opioid consumption: 24 hours, 48 hours</p> <p>Nausea: reported as incidence</p>
Notes	Study author contacted for additional information but no longer has original data

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not stated. Study author unable to provide additional information
Allocation concealment (selection bias)	Unclear risk	Method not stated. Study author unable to provide additional information
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Quote: "Double blind." Study author unable to provide additional information
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Not stated but presumably low because sham FNB in control group and IV PCA in all groups. Study author unable to provide additional information
Incomplete outcome data (attrition bias)	Low risk	No dropouts/withdrawals

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Hirst 1996 (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Hunt 2009

Methods	Randomized trial: Web-based random number generator
Participants	<p>Single centre from Salt Lake City, Utah, USA</p> <p>57 participants TKR for degenerative joint disease</p> <p>Exclusion criteria: allergy to local anaesthetics, inability to use a PCA pump, history of opioid dependence, severe liver disease, renal insufficiency, bilateral procedure and contraindication to non-steroidal anti-inflammatory drugs</p> <p>Mean age (years): 69 ± 7</p> <p>Female: 67%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Single-shot FNB with 10 to 15 ml 0.5% bupivacaine. Nerve identified via nerve stimulation. IV PCA morphine 1 mg to 2 mg every 10 minutes (lockout 10 minutes) with maximum of 30 mg in four hours</p> <p>2. IV PCA morphine 1 mg to 2 mg every 10 minutes (lockout 10 minutes) with maximum of 30 mg in four hours until postoperative day two; sham FNB with 10 to 15 ml saline</p> <p>Adjunct analgesics: standard dose of vicodin. In cases of allergy, an alternate analgesic was used</p>
Outcomes	<p>Pain severity at rest: visual analogue scale at day of surgery, postoperative days one and two</p> <p>Total opioid consumption: postoperative days one and two</p>
Notes	Single-shot FNB + sciatic group excluded from data extraction, as it was a non-random arm. Study author provided additional information on method of randomization, allocation concealment, persons blinded and co-analgesics used

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Web-based random number generator (study author provided information)
Allocation concealment (selection bias)	Low risk	Sealed envelopes (study author provided information)
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Participants and surgeon blinded (study author provided information)

Hunt 2009 (Continued)

Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Ward nurses blinded to treatment groups
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Kadic 2009

Methods	Randomized trial: method not stated
Participants	<p>Single centre from Radboud University Nijmegen Medical Centre, The Netherlands</p> <p>58 participants undergoing primary unilateral TKR</p> <p>Exclusion criteria: preexisting neurological deficit, anticoagulant therapy, long-term use of opioids, bilateral TKR, revision knee surgery, infection at the puncture site or known intolerance to any of the drugs to be used according to the study protocol</p> <p>Mean age (years): 67</p> <p>Female: 67%</p> <p>Number excluded post randomization: five (one from continuous FNB and two from PCA due to protocol violations, one from PCA due to severe adverse events and one from continuous FNB due to not meeting inclusion criteria)</p> <p>Number lost to follow-up: for month three: 41% in continuous FNB and 38% in PCA</p>
Interventions	<p>1. Continuous FNB with 20 to 25 ml ropivacaine 0.75%, postoperatively, a continuous infusion of 0.2% ropivacaine at 5 ml/h to a max of 10 ml/h for the next 48 hours. Nerve identified via nerve stimulation. IV PCA morphine 1 mg on demand, lockout period of six minutes</p> <p>2. IV PCA morphine 1 mg on demand, lockout period of six minutes</p> <p>Adjunct analgesics: paracetamol 1000 mg four times daily and diclofenac 50 mg three times daily</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain severity on movement: numerical rating scale of zero to 10. Day of surgery, days one and two</p> <p>Total opioid consumption: first 48 hours</p> <p>Nausea/vomiting</p> <p>Sedation</p> <p>Constipation</p> <p>Urinary retention</p> <p>Neurological injury (not applicable for PCA group)</p> <p>Local anaesthetic toxicity</p>

Kadic 2009 (Continued)

Infection

Participant satisfaction with analgesia: on discharge

Other outcomes:

Knee flexion

Knee Society Score: knee; function

WOMAC: pain; stiffness

90 degrees knee flexion

Blood loss during surgery

Technical failure of block (not applicable to PCA group)

Notes Additional information regarding the study method sought, but study authors could not be reached

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "random." Method not stated
Allocation concealment (selection bias)	Low risk	Sealed envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Unblinded trial
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Unblinded independent pain team: Physician and physical therapist conducted assessments
Incomplete outcome data (attrition bias) All outcomes	Low risk	Loss to follow-up for month three; knee flexion and functional outcome measures were 41% in continuous FNB and 38% in PCA. However, the review authors judged that this will not bias the review because these outcomes are not of interest for the review
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Kaloul 2004

Methods	Randomized trial: blocks of six with allocation drawn from envelopes
Participants	Single centre in Montreal, Canada
	40 participants with osteoarthritis for primary TKR
	Inclusion: 20 to 80 years of age, ASA I to III

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Kaloul 2004 (Continued)

Exclusion: allergic to amide local anaesthetics, fentanyl or midazolam; had a history of hepatic or renal failure (serum creatinine > 150 µmol/L); had a contraindication to regional anaesthesia (acquired or congenital coagulopathy, systemic or local infection, neurological disease affecting lower limbs) or use of PCA (drug dependence, inability to understand the use of the PCA device), or weight greater than 110 kg

Mean age (years): 68

Female: 63%

Number excluded post randomization: nine in continuous FNB as catheter fell out

Number lost to follow-up: none

Interventions	<p>1. Continuous FNB with 30 ml of ropivacaine 0.5% with epinephrine 1:200,000 injected, followed by ropivacaine 0.2% infusion at 12 ml/h for 48 hours. IV morphine 1 mg infused over two minutes with five-minute lockout period</p> <p>2. IV morphine 1 mg infused over two minutes with five-minute lockout period</p> <p>Adjunct analgesics: rectal indomethacin 100 mg twice daily for 48 hours after surgery. Subsequently, IV PCA or oral analgesia (oxycodone or a combination of acetaminophen and codeine)</p>
Outcomes	<p>Pain intensity at rest and on movement: visual analogue scale at six hours, 24 hours, 48 hours and daily during physiotherapy for postoperative days one and two</p> <p>Total opioid consumption: 24 hours, 48 hours</p> <p>Participant satisfaction</p>
Notes	Allocation group with psoas block was excluded from data extraction. Data on morphine consumption at 48 hours in abstract differed from graph: data from abstract used

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomly assigned in blocks of six with allocation drawn from an envelope to one of three allocation groups
Allocation concealment (selection bias)	Unclear risk	Allocation drawn from an envelope. Not stated whether envelopes were opaque and sealed
Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	Not stated
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Unblinded research assistant
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Kardash 2007

Methods	Randomized trial: computer-generated in blocks of four
Participants	<p>Single centre from Montreal, Canada</p> <p>40 participants undergoing elective unilateral TKR</p> <p>Exclusion criteria: < 18 years, inability to communicate lucidly in English or French, revision knee replacement, morbid obesity (body mass index > 35), contraindication to spinal anaesthesia (back fusion, coagulopathy, local infection), renal failure or hypersensitivity to bupivacaine, fentanyl or NSAIDs</p> <p>Mean age (years): 65 ± 2 (single-shot FNB); 67 ± 1 (IV PCA)</p> <p>Female: 75%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: 29 for outcomes measured at week six and year one</p>
Interventions	<p>1. Single-shot FNB with 20 ml bupivacaine 0.5% with 1/200,000 epinephrine, nerve stimulator; IV PCA fentanyl 50 g/ml set to deliver 25 g every five minutes as needed</p> <p>2. IV PCA fentanyl 50 g/ml set to deliver 25 g every five minutes as needed; sham FNB (area prepared and dressing applied, no injection)</p> <p>3. Obturator nerve block (not considered for this review)</p> <p>Adjunct analgesics: celecoxib 100 mg and acetaminophen 650 mg PO were given in the recovery room, then every 12 and six hours. If pain at rest was rated > six/10, breakthrough medication with IM ketorolac 10 mg given every four hours prn</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain severity at rest and on movement: visual analogue scale at 24 hours, four hours, discharge</p> <p>Total opioid consumption (fentanyl): every four hours for 48 hours</p> <p>Time to first dose of PCA</p> <p>Time to first breakthrough analgesics</p> <p>Nausea/vomiting at 48 hours</p> <p>Pruritus at 48 hours</p> <p>Urinary retention at 48 hours</p> <p>Persistent numbness at 48 hours</p> <p>Knee flexion: postoperative day two and at discharge</p> <p>Other outcomes:</p> <p>Knee scores at six weeks and one year</p> <p>Length of stay</p>
Notes	<p>Data for the obturator nerve block group were not extracted, as they are not of interest for the review. Study author provided additional information on methods of randomization and allocation concealment</p>

Risk of bias

Kardash 2007 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomization computer-generated in blocks of four according to a list prepared by the study epidemiologist (study author provided information)
Allocation concealment (selection bias)	Low risk	"Allocation was concealed in sealed envelopes" (study author provided information)
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Quotes (from report): "double-blind"; "Both patients and nurse observers collecting data were blinded to the intervention group"; "For all groups, a bandage was placed over the inguinal area, to maintain double-blinding in the recovery period"
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Outcome assessors blinded to the intervention groups
Incomplete outcome data (attrition bias) All outcomes	Low risk	29 participants lost to follow-up for the outcome of knee scores at six weeks and one year. However the review authors judged that this will not bias the review because these outcomes are not of interest for the review
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Lee 2011

Methods	Randomized trial: method not stated
Participants	<p>Single centre from Seoul, Korea</p> <p>80 participants undergoing unilateral TKR</p> <p>Exclusion criteria: study drug allergy, opioid dependence, history of previous postoperative nausea, vomiting (PONV) and motion sickness, contraindication for epidural block (previous back surgery, bleeding diathesis and neurological dysfunction), contraindication to FNB (infection at injection site), inability to use the patient-controlled epidural analgesia device and to comprehend the VAS for pain assessment</p> <p>Mean age (years): 68 (single-shot FNB group) versus 70 (epidural group)</p> <p>Female: 91%</p> <p>Number excluded post randomization: two participants from FNB group excluded from analysis because of catheter dislodgement</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Single-shot FNB with 0.25% levobupivacaine (30 ml) with 5 µg/ml epinephrine; patient-controlled epidural analgesia (PCEA) via epidural catheter for 48 hours</p> <p>2. PCEA via epidural catheter; infusion comprised 0.2% ropivacaine plus fentanyl 3 µg/ml, and programmed to deliver at basal rate of 4 ml/h, 3-ml bolus dose, with lockout duration of 10 minutes, for 48 hours</p> <p>Adjunct analgesics: meperidine (50 mg) IV if participants reported pain ≥ 50 mm on VAS at rest</p>

Lee 2011 (Continued)

Outcomes

Outcomes of interest for the review:

Pain intensity at rest and on movement: visual analogue scale at zero to six hours, six to 24 hours, 24 to 48 hours

Proportion of participants in pain: from zero to 48 hours

Nausea/vomiting

Participant satisfaction: 48 hours

Neurological injury

Other outcomes:

Total patient-controlled epidural analgesia consumption at 24 hours and 48 hours

Supplementary meperidine

Notes

Additional information regarding study methods sought, but study authors could not be reached

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not stated
Allocation concealment (selection bias)	Low risk	Sealed envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Quotes: "Patients and treating clinicians were not blinded as to study group randomization"; "Anesthesiologist who visited patients during the postoperative period was not aware of patients' assignments"
Blinding of outcome assessment (detection bias) Subjective outcomes	Unclear risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Long 2006

Methods

Randomized trial: method not stated

Participants

Single centre from California, USA

80 participants undergoing unilateral TKR

Long 2006 (Continued)

Exclusion criteria: systemic arthritis, history of substance abuse, chronic pain syndromes, diabetes mellitus peripheral neuropathy, allergy to study medication

Mean age (years): 69 (range 41 to 85)

Female: 50%

Number excluded post randomization: 10 because of catheter failure

Number lost to follow-up: none

Interventions	<p>1. Continuous FNB with bolus 0.2% ropivacaine 100 mg, infusion 5 ml/h × 36 hours</p> <p>2. Epidural catheter inserted with 20.2% ropivacaine infused at 5 ml/h × 36 hours. FNB and epidural catheters discontinued at 6 am postoperative day two</p> <p>Adjunct analgesics: hydrocodone or hydrocodone bitartrate and acetaminophen given four hourly for postoperative day, then on request for the rest of the hospitalization. Oral oxycodone 6 pm day of surgery unless comfortable. IV ketorolac 30 mg six-hourly × first 24 hours, as needed, for second 24 hours for breakthrough pain. COX-2 morning of hospitalization</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain severity at rest: visual analogue scale, four-hourly</p> <p>Total opioid consumption: daily until postoperative day three</p> <p>Knee flexion at days two and three, on discharge</p> <p>Knee extension at days two and three, on discharge</p> <p>Postoperative falls</p> <p>Deep vein thrombosis</p> <p>Other outcomes:</p> <p>Bladder infection</p> <p>Straight leg raise</p> <p>Walking distance</p>
Notes	Additional information regarding the study method sought, but study authors could not be reached

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not stated
Allocation concealment (selection bias)	Unclear risk	Method not stated
Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	Not stated
Blinding of outcome assessment (detection bias)	Unclear risk	Not stated

Long 2006 (Continued)

Subjective outcomes

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Quote: "10 were eliminated from the study because of catheter failures"; unknown if numbers were equal across groups
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Macalou 2004

Methods	Randomized trial: "randomly assigned by envelope by clinician in charge"
Participants	<p>Single centre from Nancy, France</p> <p>90 participants undergoing unilateral TKR</p> <p>Exclusion criteria: age < 18 or >85 years, ASA I to III, morbid obesity, allergy to local anaesthetics or other medications used in this study, contraindications to regional anaesthesia, preexisting neurological deficits in lower extremities, pregnancy, breast-feeding and inability to use a PCA or to comprehend pain scales</p> <p>Mean age (years): 68 ± 9 (single-shot FNB) versus 71 ± 9 (single-shot FNB + obturator) versus 70 ± 7 (IV PCA)</p> <p>Female: 79%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Single-shot FNB with 0.5% bupivacaine and 2% lidocaine with 1:200 000 epinephrine. Nerve identified via nerve stimulator. IV PCA morphine delivering 1-mg doses with seven-minute lockout period and maximum dose of 25 mg in four hours</p> <p>2. Single-shot FNB with selective obturator nerve block. Nerve identified via nerve stimulator. IV PCA morphine delivering 1-mg doses with seven-minute lockout period and maximum dose of 25 mg in four hours</p> <p>3. IV PCA morphine delivering 1-mg doses with seven-minute lockout period and maximum dose of 25 mg in four hours; sham FNB</p> <p>Adjunct analgesics: During the first 24 hours after surgery, all participants received 2 g propacetamol and 50 mg ketoprofen infused IV over 15 minutes at six-hour intervals</p>
Outcomes	<p>Pain intensity at rest: visual analogue scale, hourly × six</p> <p>Total morphine consumption: first six hours</p> <p>Nausea/vomiting</p> <p>Sedation</p> <p>Hypotension</p> <p>Bradycardia</p> <p>Respiratory depression</p>

Macalou 2004 (Continued)

Notes Additional information regarding the study method sought, but study authors could not be reached

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Randomly assigned by envelope (unable to reach study authors for additional information)
Allocation concealment (selection bias)	Unclear risk	Quote: "...assigned by envelope .." (unable to reach study authors for additional information). Unclear whether envelopes were sequentially numbered, sealed and opaque
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Quotes: "single-blind"; "Pain was evaluated at rest by another blinded investigator"
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Quotes: "single-blind"; "Pain was evaluated at rest by another blinded investigator"
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Martin 2008

Methods	Randomized trial: randomization list
Participants	<p>Two institutions in Paris, France</p> <p>40 participants undergoing TKR, regardless of age</p> <p>Inclusion criteria: indication for TKR due to knee arthritis and surgery performed during general anaesthesia</p> <p>Exclusion criteria: previous surgery or trauma of the knee or preoperative use of corticosteroids</p> <p>Mean age (years): 67 ± 2 (continuous FNB); 70 ± 2 (IV PCA)</p> <p>Female: 70%</p> <p>Number excluded post randomization: two participants in PCA group excluded for absence of preoperative blood measurement</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB + sciatic (single-shot) with bolus 20 ml of 0.75% ropivacaine each for femoral and sciatic blocks, followed by continuous infusion of 0.2% ropivacaine, 0.15 ml/kg/h via femoral catheter for 48 hours, performed preoperatively; IV PCA morphine 1-mg bolus with five-minute lockout time; if insufficient, increased to 2 mg. Discontinued 72 hours after surgery</p>

Martin 2008 (Continued)

2. IV PCA morphine 1-mg bolus with five-minute lockout time; if insufficient, increased to 2 mg. Discontinued 72 hours after surgery

Adjunct analgesics: IV acetaminophen (1 g every six hours)

Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale, days one, four, seven, months one, three</p> <p>Total opioid consumption: 24 hours, 48 hours, 72 hours</p> <p>Knee flexion: days one, four, seven, months one and three</p> <p>Other outcomes:</p> <p>Stand with and without assistance</p> <p>Uses toilet with and without assistance</p>
Notes	<p>Main aim of the study was to evaluate the effects of FNB on inflammation (clinical, cytokine levels). Study author provided additional information on method of allocation concealment</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomization list
Allocation concealment (selection bias)	Low risk	Opaque sealed envelopes (study author provided information)
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Quote: "We could not perform a blinded study because of the sensory effects of the block and the visibility of the catheter during postoperative evaluation"; review authors judge that these outcomes are not likely to be influenced by lack of blinding
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Quote: "We could not perform a blinded study because of the sensory effects of the block and the visibility of the catheter during postoperative evaluation"
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

McNamee 2001

Methods	Randomized trial: method not stated
Participants	<p>Single centre from Belfast, UK</p> <p>75 participants undergoing TKR, regardless of age</p>

McNamee 2001 (Continued)

Inclusion criteria: ASA I to III, 40 to 85 years of age, weighed 40 to 95 kg and had no contraindication to regional anaesthesia

Mean age (years): 69 (range 54 to 84)

Female: 64%

Number excluded post randomization: one participant excluded because of vasovagal collapse

Number lost to follow-up: none

Interventions	<p>1. Single-shot FNB + sciatic with 2 mg/kg of bupivacaine 7.5 mg/ml divided equally between femoral and sciatic nerves. Nerve identified via nerve stimulator. IV PCA morphine 1-mg bolus with five-minute lockout time</p> <p>2. Single-shot FNB + sciatic with 2 mg/kg ropivacaine 7.5 mg/ml divided equally between femoral and sciatic nerves. Nerve identified via nerve stimulator. IV PCA morphine 1-mg bolus with five-minute lock-out time</p> <p>3. IV PCA morphine 1-mg bolus with five-minute lockout time. No femoral nerve blockade, but the area was prepared and a dressing applied</p> <p>Co-analgesics: not stated</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity on movement: visual analogue scale, four-hourly × 48 hours</p> <p>Total opioid consumption at 24 hours and 48 hours</p> <p>Time to first morphine request</p> <p>Motor blockade</p> <p>Proportion of participants in pain > 30/100 mm</p> <p>Other outcomes:</p> <p>Emesis scores</p>

Notes	Additional information regarding study methods sought, but study authors could not be reached
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not stated
Allocation concealment (selection bias)	Low risk	Sealed envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Quotes: "...blinding of the patient"; "All blocks performed by two authors blinded to which local anaesthetic was being administered"
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Quote: "...observations performed by nursing staff, unaware as to which group the patient had been allocated"

McNamee 2001 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Mistraletti 2006

Methods	Randomized trial: computer-generated randomization
Participants	<p>Single centre from McGill, Montreal, Quebec, Canada</p> <p>27 participants undergoing primary unilateral TKR</p> <p>Exclusion criteria: age < 18 or > 85 years; ASA > III; morbid obesity (BMI > 40); diabetes or any metabolic disease; loss of body weight > 10% over the preceding three months; serum albumin < 35 g/l; cardiac, renal or hepatic failure; anaemia (haematocrit < 30%); organ transplant; rheumatoid arthritis; allergy to local anaesthetics or other medications used in the study; contraindication to use of PCA, epidural and CPNB; preexisting neurological deficits in lower extremities; and inability to use PCA or to comprehend pain scales</p> <p>Mean age (years): continuous FNB + sciatic: 67 ± 8; epidural: 64 ± 11; IV PCA: 71 ± 11</p> <p>Female: 59%</p> <p>Number excluded post randomization/lost to follow-up: three participants excluded from analysis (one participant in epidural group for removal of epidural catheter, one participant in continuous FNB group for cutting of femoral catheter and one participant in PCA group for refusal to continue the study)</p>
Interventions	<p>1. Continuous FNB + sciatic with bolus, lignocaine 2% with epinephrine 2.5 µg/ml, 0.25 ml/kg. Infusion ropivacaine 0.2% initially at 8 ml/h in femoral catheter and 4 ml/h in sciatic. Subsequent adjustments performed to obtain VAS score < four and least possible motor block. Administered for 48 hours. Nerve identified via nerve stimulation</p> <p>2. Epidural with bupivacaine 0.1% with fentanyl 3 µg/ml starting at 10 ml/h and adjusted if VAS at rest > three or VAS at knee flexion > four</p> <p>3. IV PCA morphine with bolus doses from 2.5 to 5 mg (max 30 mg) until VAS score at rest < four, infusion at incremental doses of 1 mg, with lockout of seven minutes</p> <p>Adjunct analgesics: oral naproxen 500 mg every eight hours and oral acetaminophen 650 mg every six hours</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale, 24 hours, 48 hours, at discharge</p> <p>Knee flexion: 48 hours</p> <p>Time to walking</p>
Notes	<p>Main aim of study was to assess whether postoperative suppression of glyconeogenesis by dextrose infusion would be influenced by continuous nerve blocks compared with epidural or PCA. Additional information regarding the study method sought, but study authors could not be reached</p>

Mistraletti 2006 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization
Allocation concealment (selection bias)	Unclear risk	Method not stated
Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	Not stated
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Investigators not involved in perioperative management collected VAS and other data
Incomplete outcome data (attrition bias) All outcomes	Low risk	Three participants excluded from analysis (one participant in epidural group for removal of epidural catheter, one participant in continuous FNB group for cutting of femoral catheter and one participant in PCA group for refusal to continue the study)
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Nader 2012

Methods	Randomized trial: computer-generated randomization
Participants	<p>Single centre from Chicago, USA</p> <p>62 participants undergoing elective unilateral TKR by a single surgeon</p> <p>Exclusion criteria: contraindication to the use of neuraxial analgesia, pregnancy, infection, prior surgery in inguinal/lumbar region, haemostatic abnormality, preexisting neuropathy, currently taking antidepressant or antiepileptic drugs, opioid-tolerant patients or neuromuscular disease that could interfere with data collection</p> <p>Median (IQR) age (years): continuous FNB: 65 (60 to 76) versus IV PCA: 64 (60-71)</p> <p>Female: 67.7%</p> <p>Number excluded post randomization/loss to follow-up: two participants excluded from analysis for data > one month (one participant in each group)</p>
Interventions	<p>1. Continuous FNB 10-ml bolus of ropivacaine 0.25%, 5 ml/h infusion of ropivacaine 0.1%, commenced on postoperative day one morning at discontinuation of epidural analgesia (infusion bupivacaine 1 mg/ml and hydromorphone 10 mg/ml, bolus 3 ml of the same solution, lockout of 15 minutes, maximum 15 ml/h). Oral opioid hydrocodone 10 mg plus acetaminophen 325 mg every four to six hours</p> <p>2. Oral opioid hydrocodone 10 mg plus acetaminophen 325 mg every four to six hours, commenced on postoperative day one morning after discontinuation of epidural analgesia</p>

Nader 2012 (Continued)

Adjunct analgesics: If unable to achieve verbal rating score for pain < four, analgesic regimen was switched to sustained-release oxycodone 10 mg every 12 hours with oral hydromorphone 2 mg every four hours for breakthrough pain

Outcomes	Pain intensity at rest and on movement: visual analogue scale, hospitalisation, months one, six and 12 Opioid consumption: 24 hours, 48 hours, 72 hours Knee flexion: postoperative days one and two, months one, six and 12 Participant satisfaction: at discharge Postoperative falls Venous thrombotic events
----------	---

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization
Allocation concealment (selection bias)	Low risk	Sealed opaque envelope, not opened until after informed consent was obtained
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Not blinded
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Not blinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals for short-term data. One dropout from each group after one month
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Ng 2001

Methods	Randomized trial: method not stated
Participants	National University Hospital, Singapore 48 participants undergoing primary unilateral TKR Inclusion criteria: ASA I to II Exclusion criteria: age < 40 or > 80 years, weight < 50 or > 100 kg, allergy to local anaesthetics or acetaminophen, preexisting neurological deficit or inability to understand pain scale or PCA device usage

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Copyright © 2016 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

Ng 2001 (Continued)

Mean age (years): 64

Female: 85%

Number excluded post randomization: none

Number lost to follow-up: none

Interventions	<p>1. Single-shot FNB with 30 ml of ropivacaine 0.25% (75 mg), nerve identified via stimulation; IV PCA morphine 1 mg with lockout time of five minutes, continuous background infusion at 0.01 mg/kg/h set for first 24 hours</p> <p>2. Single-shot FNB with 30 ml of ropivacaine 0.5% (150 mg), nerve identified via nerve stimulation; IV PCA morphine 1 mg with lockout time of five minutes, continuous background infusion at 0.01 mg/kg/h set for first 24 hours</p> <p>3. Single-shot FNB with 30 ml of bupivacaine 0.25% (75 mg), nerve identified via nerve stimulation. IV PCA morphine 1 mg with lockout time of five minutes, continuous background infusion at 0.01 mg/kg/h set for first 24 hours</p> <p>4. Sham block. IV PCA morphine 1 mg with lockout time of five minutes, continuous background infusion at 0.01 mg/kg/h set for first 24 hours</p> <p>Adjunct analgesics: oral acetaminophen 1 g six-hourly started 24 hours after surgery</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Proportion of pain at rest and on movement: four-point verbal pain score, one hour, four hours, eight hours, 24 hours, 48 hours</p> <p>Total opioid consumption: 24 hours</p> <p>Respiratory depression</p> <p>Sedation</p> <p>Other outcomes:</p> <p>Length of stay</p>
Notes	Study author contacted for additional information but could not provide requested information

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not stated
Allocation concealment (selection bias)	Unclear risk	Sealed envelope. Unclear whether envelopes were sequentially numbered and opaque
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Presumably low risk as quote: "double blind." Also, sterile syringes containing test solutions were prepared by one of the authors not involved in the nerve block, and FNBs were performed by anaesthesiologists unaware of the test solution used
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Assessment done by an investigator blinded to participant grouping

Ng 2001 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Ng 2012

Methods	Randomized trial: randomization table
Participants	<p>Hong Kong</p> <p>16 participants undergoing bilateral osteoarthritis of the knee scheduled for staged total knee (staged operations performed at a minimum interval of three months)</p> <p>Exclusion criteria: known allergies to any of the test drugs, those with major systemic illnesses (heart failure, renal impairment and coagulopathy), long-term users of opioids and participants who had failed femoral catheter insertions</p> <p>Mean age (years): 70 ± 7.6</p> <p>Female: 83%</p> <p>Number excluded post randomization: two</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB 20-ml bolus 0.2% ropivacaine, followed by continuous infusion at 10 ml/h, starting in the recovery room and continuing for 72 hours after the operation. Sham multimodal periarticular injection</p> <p>2. Multimodal periarticular soft tissue injection with ropivacaine (300 mg in 30 ml), adrenaline (1 mg in 0.5 ml), saline (70 ml), triamcinolone acetonide (40 mg in 1 ml), injected into posterior joint capsules, periarticular soft tissue. Triamcinolone infiltrated into the skin to avoid fat atrophy. Sham continuous FNB</p> <p>Adjunct analgesics: patient-controlled intravenous morphine (1-mg bolus, lockout time of five minutes, maximum 6 mg/h)</p>
Outcomes	<p>Pain intensity: visual analogue scale, postoperative days one, two and three</p> <p>Morphine consumption: postoperative days one, two and three</p> <p>Knee flexion: daily × three</p> <p>Participant satisfaction with pain management at 72 hours</p>
Notes	<p>Cross-over trial comparing collateral knees. Pair analysis not reported. Means (SDs) are available only for the two groups separately. Unit of analysis error may exist if we consider the data as from a parallel-group trial, and carry-over effect may exist for outcomes such as functionality and participant satisfaction. The population in this trial may be quite different (i.e. fitter) from that of parallel-arm RCTs. Hence, the review authors decided not to extract the data for meta-analysis</p>

Risk of bias

Ng 2012 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Randomization table to allow 50% of participants treated to receive true FNB first and 50% of participants to receive true MPI first, using SPSS software (version 14.0; SPSS Inc, Chicago, Illinois)
Allocation concealment (selection bias)	Unclear risk	Not mentioned
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Participants and nurses responsible for recording pain scores blinded to treatment allocation
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Nurses recording pain scores blinded. Sham FNB/local infiltration analgesia used
Incomplete outcome data (attrition bias) All outcomes	Low risk	Two participants withdrew but unlikely to affect study conclusions as the study recruited two extra participants compared with required sample size
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Ozen 2006

Methods	Randomized trial: computer assisted
Participants	<p>Single centre from Ankara, Turkey</p> <p>34 participants undergoing unilateral TKR</p> <p>Inclusion criteria: 50 to 75 years of age, ASA I to III</p> <p>Exclusion criteria: weight < 50 or > 100 kg, undergoing tumour surgery, immunosuppressive therapy, preexisting neuropathy or neurological deficits, allergy to local anaesthetics, inability to understand pain scale or PCA device usage</p> <p>Mean age (years): 64</p> <p>Female: 62%</p> <p>Number excluded post randomization: four</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Three-in-one single-shot FNB with 40 ml 0.375% ropivacaine. Nerve identified via nerve stimulation. IV PCA morphine 0.3 mg/ml, 1-mg bolus, 15-minute lockout</p> <p>2. IV PCA morphine 0.3 mg/ml, 1-mg bolus, 15-minute lockout</p> <p>Co-analgesics: not stated</p>

Ozen 2006 (Continued)

Outcomes	Pain intensity at rest. Visual analogue scale, hourly × four, then two-hourly × three, 18 hours, 24 hours, 36 hours, 48 hours
	Total opioid consumption: 24 hours, 48 hours
	Nausea/vomiting
	Hypotension

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer assisted
Allocation concealment (selection bias)	Low risk	Sealed envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	Not stated
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Team monitoring postop pain blind to treatment group
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Park 2010

Methods	Randomized trial: drawing of lots
Participants	Single centre from Daejeon, Korea
	80 participants undergoing elective unilateral TKR
	Inclusion criteria: female, ASA I to II under spinal anaesthesia
	Exclusion criteria: contraindications to regional anaesthetic technique (e.g. local infection, sepsis, coagulation abnormality), allergy to local anaesthetic or morphine, contraindications for NSAID use, pre-existing neurological deficit in lower extremities and inability to comprehend pain scales or to use IV PCA device
	Mean age (years): 67 ± 6
	Female: 100%

Park 2010 (Continued)

Number excluded post randomization: eight

Number lost to follow-up: none

Interventions	<p>Three arms of different concentrations of continuous FNB versus single-shot FNB</p> <ol style="list-style-type: none"> 1. Continuous FNB with bolus 20 ml of 0.25% bupivacaine with epinephrine 1:200,000, followed by continuous infusion 0.125% bupivacaine 2 ml/h until postoperative day three or four 2. Continuous FNB with bolus 20 ml of 0.25% bupivacaine with epinephrine 1:200,000, followed by continuous infusion 0.125% bupivacaine 4 ml/h until postoperative day three or four 3. Continuous FNB with bolus 20 ml of 0.25% bupivacaine with epinephrine 1:200,000, followed by continuous infusion 0.125% bupivacaine 6 ml/h until postoperative day three or four 4. Single-shot FNB with bolus 20 ml of 0.25% bupivacaine with epinephrine 1:200,000, followed by continuous infusion 0.125% bupivacaine 0 ml/h until postoperative day three or four <p>Adjunct analgesics: IV PCA morphine-ketorolac at 1-ml bolus of 0.5 mg/ml morphine, 1.5 mg/ml ketorolac solution, eight-minute lockout interval and no continuous background infusion, bolus dose on demand when VAS > three out of 10</p> <p>Rescue doses of 90 mg IM diclofenac</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale, PACU, 6 pm on day of surgery, postoperative days one and two</p> <p>Total opioid consumption: postoperative days one and two</p> <p>Nausea/vomiting</p> <p>Sedation</p> <p>Pruritus</p> <p>Rescue analgesics required</p>
Notes	<p>Study author provided additional information on methods of randomization and allocation concealment</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Drawing of lots (study author provided information)
Allocation concealment (selection bias)	Low risk	Sealed opaque envelopes (study author provided information)
Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	No mention
Blinding of outcome assessment (detection bias) Subjective outcomes	Unclear risk	No mention

Park 2010 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Eight participants excluded post randomization; not mentioned whether they were equally distributed across groups
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Parvataneni 2007

Methods	Randomized trial: method not stated
Participants	<p>Centre in USA</p> <p>60 participants undergoing TKR</p> <p>Inclusion criteria: diagnosis of non-inflammatory osteoarthritis. Participant factors such as BMI and co-morbidities were not considered during the enrolment process</p> <p>Mean age (years): 69</p> <p>Female: 48%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Single-shot FNB with IV PCA morphine for 24 hours</p> <p>2. Intra-articular injection with 0.5% bupivacaine 200 to 400 mg, morphine sulfate (0.4 to 1.0 ml) 4 to 10 mg, epinephrine 1:1000 (0.3 ml) 300 µg, methylprednisolone acetate 40 mg, cefuroxime (10 ml) 750 mg, normal saline 22 ml. Clonidine transdermal patch applied in operating room: 100 µg/24 h</p> <p>Adjunct analgesics: IM ketorolac every six hours as needed (15 mg if > 65 years, 30 mg if < 65 years, hold if renal impairment), if ketorolac is ineffective, IM morphine 2 to 4 mg every two to four hours, celecoxib 200 mg orally daily for 10 days, oxycodone SR 10/20 mg orally every 12 hours for 48 hours, oxycodone 5 mg orally every six hours as needed, acetaminophen 1000 mg orally every six hours</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest: visual analogue scale, postoperative day one</p> <p>Knee ROM: postoperative day two</p> <p>Nausea/vomiting</p> <p>Participant satisfaction: postoperative day one</p> <p>Other outcomes:</p> <p>Straight leg raised</p> <p>Pain less than expected</p> <p>Confusion</p> <p>Discharged home</p>
Notes	Additional information regarding the study method sought, but study authors could not be reached

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Parvataneni 2007 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not stated
Allocation concealment (selection bias)	Unclear risk	Method not stated
Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	Not stated
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Research personnel blinded to randomization
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Salinas 2006

Methods	Randomized trial: computer random number generator
Participants	<p>Single centre in Seattle, USA</p> <p>36 participants undergoing unilateral TKR under spinal anaesthesia</p> <p>Inclusion criteria: ASA I to III participants</p> <p>Exclusion criteria: age < 18 or > 85 years; BMI > 45; renal insufficiency; contraindications (localized infection, sepsis, preexisting lower extremity neurological abnormality) or refused spinal anaesthesia or femoral nerve block; allergy to local anaesthetics, morphine or oxycodone; long-term opioid use; and difficulties in comprehending VAS pain scores or in using IV PCA device</p> <p>Mean age (years): 67 (9) versus 68 (6)</p> <p>Female: 50%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB with 30 ml of ropivacaine 0.375% with epinephrine 2.5 g/ml, followed by continuous infusion of ropivacaine 0.2% at 10 ml/h via catheter six hours later until 8 am on the morning of postoperative day two; nerve stimulation</p> <p>2. Single-shot FNB with 30 ml of ropivacaine 0.375% with epinephrine 2.5 g/ml; nerve stimulation</p>

Salinas 2006 (Continued)

Adjunct analgesics: IV PCA morphine 1-mg bolus with five-minute lockout, to titrate VAS scores to $\leq$ three until morning postoperative day one. Oral ibuprofen 600 mg three times a day (starting date of surgery). Breakthrough pain ($>$ four/10) treated on demand with oral oxycodone 5 mg

Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale, twice daily</p> <p>Total opioid consumption: 48 hours</p> <p>Nerve injury</p> <p>Catheter-related infection</p>
----------	--

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer random number generator
Allocation concealment (selection bias)	Low risk	Sealed envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Unblinded
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Unblinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	Presumably low, as no mention of withdrawals/dropouts
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Seet 2006

Methods	Randomized trial: on-line randomization software
Participants	<p>Single centre at Singapore General Hospital, Singapore</p> <p>60 participants undergoing elective unilateral TKR under subarachnoid block</p> <p>Inclusion criteria: ASA I to III</p> <p>Exclusion criteria: contraindication to a lumbar subarachnoid block (e.g. coagulopathy, local infection, sepsis, previous lumbar spine operation), patients who preferred general anaesthesia, body weight $<$ 40 kg or $>$ 120 kg, allergies to study drugs, rheumatoid arthritis, preexisting lower limb neurological deficits</p>

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Seet 2006 (Continued)

Mean age (years): 66 ± 7

Female: 73%

Number excluded post randomization: three from continuous FNB 0.15% ropivacaine and two from continuous FNB 0.2% ropivacaine (one unable to locate the subarachnoid space, one inadequate subarachnoid, three slipped femoral catheters postoperatively)

Number lost to follow-up: none

Interventions

1. Three-in-one continuous FNB with 0.15% ropivacaine at 10 ml/h first 24 hours and 5 ml/h next 24 hours. Nerve identified via nerve stimulation. IV PCA morphine 1-mg/ml bolus doses with lockout period of five minutes and maximum dose of 8 mg/h

2. 3-in-1 continuous FNB with 0.2% ropivacaine at 10 ml/h first 24 hours and 5 ml/h next 24 hours. Nerve identified via nerve stimulation. IV PCA morphine 1-mg/ml bolus doses with lockout period of five minutes and maximum dose of 8 mg/h

3. IV PCA morphine 1-mg/ml bolus doses with lockout period of five minutes and maximum dose of 8 mg/h

Adjunct analgesics: oral paracetamol 1 g six-hourly. Oral rofecoxib 50 mg each morning

Outcomes
Outcomes of interest for the review:

Pain severity at rest and on movement: visual analogue scale, six hours, 12 hours, 24 hours, 48 hours, 72 hours

Total opioid consumption: 24 hours, 48 hours, 72 hours

Nausea/vomiting

Urinary retention

Pruritus

Sedation

Hypotension

Giddiness

Participant satisfaction: 72 hours

Knee flexion

Time to ambulation

Time to discharge

Length of stay

Other outcomes:

Knee Society Clinical Rating System: knee scores, function scores

Notes

Study author provided additional information on methods of randomization and allocation concealment

Risk of bias
Bias
Authors' judgement
Support for judgement

Seet 2006 (Continued)

Random sequence generation (selection bias)	Low risk	"On-line randomization software" (study author provided information)
Allocation concealment (selection bias)	Low risk	Sealed opaque envelopes (study author provided information)
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Unblinded
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Unblinded
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Five participants excluded post randomization
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Serpell 2001

Methods	Randomized trial: method not stated
Participants	<p>Single centre in Ninewells Hospital, Dundee, UK</p> <p>30 participants undergoing elective TKR under spinal anaesthesia</p> <p>Inclusion criteria: ASA I to II, 18 to 85 years</p> <p>Exclusion criteria: history of neurological disease, hypersensitivity to bupivacaine or unsuitable for spinal anaesthesia</p> <p>Mean age (years): 69 ± 3</p> <p>Female: 77%</p> <p>Number excluded post randomization: one participant from continuous FNB excluded because of post-operative urinary retention</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB with 0.3 ml/kg 0.5% bupivacaine and adrenaline 1:200,000, administered in 5-ml increments. Nerve identified via nerve stimulation, block inserted post surgery, cannula left in situ to enable top-ups of same dose of bupivacaine at six- to eight-hour intervals during the next 48 hours; IV PCA morphine 2-mg boluses to a maximum of 12 mg/h with lockout interval of nine minutes</p> <p>2. IV PCA morphine 2-mg boluses to a maximum of 12 mg/h with lockout interval of nine minutes</p> <p>Adjunct analgesics: IM morphine 10 mg or paracetamol 1 g orally</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest: visual analogue scale, 24 hours, 48 hours</p>

Serpell 2001 (Continued)

Total morphine consumption: 48 hours

Nausea/vomiting

Other outcomes:

Total number of IM supplements 10 mg

Total number of paracetamol supplements 1 g

Total number of doses of prochlorperazine

Notes	Additional information regarding study method sought, but study authors could not be reached
-------	--

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not stated
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Unblinded
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Unblinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	One participant from continuous FNB excluded because of postoperative urinary retention. No other dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Singelyn 1998

Methods	Randomized trial: computer-generated list of random permutations
Participants	<p>Single centre in Brussels, Belgium</p> <p>45 participants undergoing elective TKR under general anaesthesia</p> <p>Inclusion criteria: ASA I to III</p> <p>Exclusion criteria: contraindications to regional anaesthetic technique (e.g. local infection, sepsis, coagulation abnormality), age < 18 or > 80 years, weight < 50 or > 100 kg, allergy to local anaesthetic and/or opioid, preexisting neurological deficit, diabetes or inability to comprehend pain scales or to use a PCA device</p> <p>Mean age: not stated</p>

Singelyn 1998 (Continued)

Female: percentage not stated

Number excluded post randomization: none

Number lost to follow-up: none

Interventions	<p>1. Continuous FNB with bolus 37 ml of 0.25% bupivacaine with epinephrine 1:200,000, followed by continuous infusion 0.125% bupivacaine with sufentanil 0.1 µg/ml and clonidine 1 µg/ml, 10 ml/h, for 48 hours. Nerve identified via nerve stimulator</p> <p>2. IV PCA with morphine 2 mg/ml, dose 1.5 mg, lockout time of eight minutes, for 48 hours</p> <p>3. Epidural with test dose 3 ml 0.25% bupivacaine with epinephrine 1:200,000, 10 ml of same solution and 10 µg/ml sufentanil, followed by infusion 0.125% bupivacaine with sufentanil 0.1 µg/ml and clonidine 1 µg/ml 10 ml/h, for 48 hours</p> <p>Adjunct analgesics: If postoperative pain score $\geq$ one (ranged from zero to three) in FNB and epidural groups: 1g IV propacetamol (Prodafalgan), followed by 10 to 20 mg of IM piritramide, a synthetic p-agonist opioid (Dipidolor) (15 mg piritramide = 10 mg morphine—information provided by study author)</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale, four hours, 24 hours, 48 hours</p> <p>Knee flexion: daily until discharge</p> <p>Nausea/vomiting</p> <p>Urinary retention</p> <p>Hypotension</p> <p>Technical problem: kinked cath</p> <p>Time to achieve 90 degrees knee flexion</p> <p>Day able to walk</p> <p>Other outcomes:</p> <p>Hospital stay (including rehabilitation)</p> <p>Supplemental analgesia (propacetamol)</p> <p>IM piritramide</p>
Notes	Study author contacted for additional information but no longer has original data

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated list of random permutations
Allocation concealment (selection bias)	Unclear risk	Method not stated
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Not stated but likely unblinded

Singelyn 1998 (Continued)

Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Not stated but likely unblinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Sundarathiti 2009

Methods	Randomized trial: random number of tables
Participants	<p>Single centre in Bangkok, Thailand</p> <p>61 participants undergoing elective unilateral TKR under spinal anaesthesia</p> <p>Inclusion criteria: ASA physical status I to III</p> <p>Exclusion criteria: < 35 or > 80 years of age, BMI > 45, renal insufficiency (creatinine > 1.5 mg/dl), preexisting neurological deficit, inability to comprehend pain scales, long-term opioid use and contraindications to neuraxial block or FNB</p> <p>Mean age (years): 68 ± 9</p> <p>Female: 85%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB with bolus 20 ml of 0.25% levobupivacaine, continuous infusion of 0.125% levobupivacaine at 8 ml/h for 24 hours postoperatively, then reduced to 6 ml/h if VAS > three, nerve identified via nerve stimulator</p> <p>2. Continuous epidural with bolus 0.125% levobupivacaine 10 ml plus 2 mg morphine, continuous infusion of 0.125% levobupivacaine with morphine 0.0125 mg/ml at 4 ml/h for 24 hours postop, then reduced to 3 ml/h if VAS > three</p> <p>Adjunct analgesics: oral ultracet one tab three times a day, oral acetaminophen 500 mg four times a day. Breakthrough pain (defined as VAS > four): IV tramadol 50 mg every four hours until discharge</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Proportion of participants in pain. At PACU, six hours, 12 hours, 24 hours, 36 hours, 48 hours and 72 hours</p> <p>Nausea/vomiting</p> <p>Pruritus</p> <p>Urinary retention</p> <p>Dizziness</p> <p>Numbness</p>

Sundarathiti 2009 (Continued)

Catheter-related infection

Nerve injury

Local anaesthesia toxicity

Complete motor blockade

Participant satisfaction (one = poor, two = fair, three = good, four = excellent)

Other outcomes:

Stand with help

Walker mobilization and transfer to chair with and without help

Hospital length of stay

Notes Study author contacted but could not provide additional information

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number of tables
Allocation concealment (selection bias)	Unclear risk	Method not mentioned
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Unblinded
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Unblinded
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Szczukowski 2004

Methods	Randomized trial: research randomizer application
Participants	Single centre in Maryland, USA
	40 participants undergoing unilateral TKR
	Inclusion criteria: age 40 to 80 years, ASA I to III

Szczukowski 2004 (Continued)

Exclusion criteria: weighed > 300 lb or < 110 lb, or had an allergy to medications such as bupivacaine, epinephrine, morphine or meperidine. Other exclusions included history of narcotic abuse, preexisting neurological deficit in the lower extremity, inability to use the PCA, a bleeding disorder, a local or systemic infection, a bilateral procedure, uncontrolled hypertension or history of severe arrhythmia

Mean age (years): 67

Female: 63%

Number excluded post randomization: none

Number lost to follow-up: none

Interventions	<p>1. Single-shot FNB with 30 ml of 0.5% bupivacaine with epinephrine 1:200,000. Nerve identified via stimulation. IV PCA morphine loading dose of 3 mg and initial bolus dose of 1.5 mg, with lockout interval of 15 minutes and four-hour maximum of 30 mg; morphine infusion discontinued after 24 hours</p> <p>2. Sham block with 30 ml of preservative-free normal saline. IV PCA morphine loading dose of 3 mg and initial bolus dose of 1.5 mg, with lockout interval of 15 minutes and four-hour maximum of 30 mg; morphine infusion discontinued after 24 hours</p> <p>Adjunct analgesics: Oxycodone (5 mg)/acetaminophen (325 mg) oral tablets (1 tablet) every four hours as needed for breakthrough pain while on the PCA. When PCA morphine was discontinued, started on oxycodone (5 mg)/acetaminophen (325 mg) (one to two tablets) every four hours as needed</p>
Outcomes	<p>Pain severity at rest: visual analogue scale, day of surgery, daily post operation until discharge</p> <p>Total opioid consumption: 24 hours after surgery</p> <p>Sedation</p> <p>Nausea/vomiting</p> <p>Urinary retention</p> <p>Disorientation</p> <p>Pruritus and any other complications</p> <p>Sensory block</p> <p>Participant satisfaction</p>
Notes	Study author contacted for additional information but no longer has original data

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Research randomizer application (www.randomizer.org/form.htm)
Allocation concealment (selection bias)	Low risk	Quote: "Each vial was labelled in a similar manner, and the surgeons, anaesthesiologists, nurses, and all operating room staff were blinded to the contents of the vials. Each vial was labelled with a patient protocol number. The records were maintained and locked in the pharmacy-services research closet"
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Quote: "...surgeons, anaesthesiologists, nurses, and all operating room staff were blinded to the contents of the vials"

Szczukowski 2004 (Continued)

Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Quote: "double-blind." Presumably low risk as it will not be obvious which was the intervention group because both groups were given IV PCA
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Tang 2010

Methods	Randomized trial: computer-generated randomization
Participants	<p>Single centre in Beijing, China</p> <p>34 participants undergoing elective TKR under general anaesthesia</p> <p>Inclusion criteria: ASA I to II, age 18 to 80 years, weight 40 to 70 kg</p> <p>Exclusion criteria: impaired blood clotting, generalized sensitive skin, hypersensitivity to study drugs, uncooperative patients, impaired immune system</p> <p>Mean age (years): 64 ± 7</p> <p>Female: 79%</p> <p>Number excluded post randomization: one from continuous FNB group because catheter fell out</p> <p>Number lost to follow-up: one dropout from PCA morphine group</p>
Interventions	<p>1. Continuous FNB with bolus 25 ml 0.4% ropivacaine, post operation infusion 0.2% ropivacaine loading dose 10 ml, background infusion 10 ml/h; IV PCA morphine bolus 1 mg/ml; lockout period of five minutes, no background infusion</p> <p>2. IV PCA morphine bolus 1 mg/ml; lockout period of five minutes, no background infusion</p> <p>Adjunct analgesics: not stated</p>
Outcomes	<p>Outcomes of interest to the study:</p> <p>Pain intensity at rest and on movement: visual analogue scale, PACU, four hours, eight hours, 12 hours, 24 hours, 48 hours</p> <p>Total opioid consumption: PACU, four hours, eight hours, 12 hours, 24 hours, 48 hours</p> <p>Nausea/vomiting</p> <p>Sedation</p> <p>Pruritus</p> <p>Knee flexion: 24 hours, 48 hours</p> <p>Participant satisfaction</p> <p>Other outcomes:</p>

Tang 2010 (Continued)

Haemodynamic status

Blood results

Notes	Study author provided additional information on methods of allocation concealment and blinding
-------	--

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Excel to generate the sequence
Allocation concealment (selection bias)	Low risk	Sequentially numbered opaque, sealed envelopes (study author provided information)
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Unblinded (study author provided information)
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Unblinded (study author provided information)
Incomplete outcome data (attrition bias) All outcomes	Low risk	One participant from continuous FNB group withdrew from the study because the catheter slipped out, and one participant from the PCA group dropped out after randomization. Apart from these, no other dropouts/withdrawals reported
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Toftdahl 2007

Methods	Randomized trial: method: thorough shuffling of sealed envelopes
Participants	<p>Department of Orthopaedic Surgery, Aarhus Hospital, Denmark</p> <p>80 participants undergoing primary TKR under spinal anaesthesia</p> <p>Exclusion criteria: lack of mental ability to provide informed consent, neuropathic pain or sensory disorders in the leg to be operated, previous major bone surgery in the knee joint, intolerance to study drugs, failure of spinal anaesthesia and reoperation or trauma to the knee within the study period</p> <p>Mean age (years): continuous FNB: 72 ± 9; periarticular injections: 70 ± 9</p> <p>Female: 61%</p> <p>Number excluded post randomization: three from continuous FNB group because of conversion of failed spinal anaesthesia to general anaesthesia.</p> <p>Number lost to follow-up: none</p>

Toftdahl 2007 (Continued)

Interventions	<p>1. Continuous FNB with bolus 20 ml ropivacaine 10 mg/ml, infusion 10 ml/h ropivacaine (2 mg/ml) for 48 hours. If needed, a bolus of 20 ml could be given every eight hours. After skin closure, 4 mg morphine (0.4 mg/ml) and 50 mg bupivacaine (5 mg/ml) given intra-articularly through the drain</p> <p>2. Periarticular injections: 150 ml ropivacaine (2 mg/ml), 1 ml ketorolac (30 mg/ml) and 1 ml epinephrine (0.5 mg/ml) by infiltration of the knee at the end of surgery and two postoperative injections of these substances through an intra-articular catheter</p> <p>Adjunct analgesics: Paracetamol (1 g × 4), ibuprofen (400 mg × 3) and controlled-release oxycodone (20 mg × 2) daily. Immediate-release oxycodone (5 to 10 mg) used for pain treatment if pain at rest exceeded three on the zero to 10 numeric rating scale. IV morphine (0.05 to 0.1 mg/kg) was used for the treatment of severe pain (> seven)</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain severity at rest and on movement: visual analogue scale, one hour, two hours, thrice daily until postoperative day three and discharge</p> <p>Number of participants in pain (scores ≥ three from surgery to the evening of postoperative day one)</p> <p>Total opioid consumption: postoperative day one</p> <p>Nausea/vomiting</p> <p>Dizziness</p> <p>Pruritus</p> <p>Urinary track infection</p> <p>Constipation (on the evening of day three)</p> <p>Wound infection</p> <p>Flexion > 90 degrees</p> <p>Other outcomes:</p> <p>Bullae around wounds</p> <p>Knee ROM: median extension defect postoperative days one and two</p> <p>Walking distance</p> <p>Quadriceps function</p> <p>Ability to hold quadriceps tension > five seconds postoperative day one</p> <p>Ability to lift leg for > five seconds postoperative day two</p> <p>Length of stay</p>
Notes	<p>Study author provided additional information/data on method of randomization and individual participant data on pain intensity scores, nausea and vomiting, opioid consumption and knee range of motion</p>
Risk of bias	
Bias	Authors' judgement Support for judgement
Random sequence generation (selection bias)	<p>Low risk</p> <p>Thorough shuffling of sealed envelopes in batches of 20 envelopes per shuffle (study author provided information)</p>

Toftdahl 2007 (Continued)

Allocation concealment (selection bias)	Low risk	Sequentially numbered opaque, sealed envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Quote: "Blinding of patients and caregivers was not attempted. It was considered impossible to do properly, as the partial motor block present in patients in group F would be obvious to patients and staff"
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Quote: "Data collection and analysis was carried out by assessor who was not blinded since catheter placement was obvious"
Incomplete outcome data (attrition bias) All outcomes	Low risk	Four participants who discontinued their allocated interventions were included in the data analysis
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Tugay 2006

Methods	Randomized trial: drawing envelopes
Participants	<p>Single centre: Department of Orthopaedics and Traumatology, Hacettepe Hospital, Turkey</p> <p>23 participants undergoing TKR for osteoarthritis</p> <p>Inclusion criteria: ASA I to III</p> <p>Exclusion criteria: significant medical/ psychiatric problems, unable to co-operate with study protocol, contraindication to RA</p> <p>Mean age (years): 66 (range 51 to 78)</p> <p>Female: 87%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Single-shot FNB (preemptive) with 40 ml 0.25% bupivacaine administered before muscle relaxant. Nerve identified via nerve stimulator. IV PCA morphine infusion 1 mg, lockout time of five minutes</p> <p>2. Single-shot FNB (post operation) with 40 ml 0.25% bupivacaine administered after reversal of muscle relaxant. Nerve identified via nerve stimulator. IV PCA morphine infusion 1 mg, lockout time of five minutes</p> <p>3. IV PCA morphine infusion 1 mg, lockout time of five minutes</p> <p>Adjunct analgesics: none (study author provided information)</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity: changes in group means. Visual analogue scale, one hour, four hours, six hours, 12 hours, 24 hours, 48 hours</p> <p>Total opioid consumption: one hour, four hours, six hours, 12 hours, 24 hours, 48 hours</p>

Tugay 2006 (Continued)

Other outcomes:

Ambulation of velocity

Iowa University Level of Assistance scale

Length of stay

Notes

No difference in pain reduction between preemptive and postop FNB. Study author provided additional information on co-analgesics and methods of randomization, allocation concealment and blinding

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The femoral nerve block protocols and control were written and enveloped. Prior to each operation an envelope was taken by the anaesthesiologist ..." (study author provided information)
Allocation concealment (selection bias)	Low risk	"Only the anaesthesiologists knew the femoral nerve block method used on each patient" (study author provided information)
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	"Both patients and the rehabilitation staff were all blinded to the femoral nerve block protocol" (study author provided information)
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	"The physiotherapists were not informed about which protocol was applied to which patients until the end of the study. The functional evaluations and pain evaluations were done by these physiotherapist authors" (study author provided information)
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Wang 2002

Methods	Randomized trial: drawing marked cards from a deck
Participants	<p>Single centre in Henry Ford Hospital, Detroit, USA</p> <p>30 participants undergoing TKR</p> <p>Inclusion criteria: age 40 to 70 years</p> <p>Exclusion criteria: hypersensitivity to the test substance or co-morbid condition that might compromise safety or ability to comply with study procedures</p> <p>Mean age (years): single-shot FNB: 67 ± 8; IV PCA: 66 ± 10</p> <p>Female: 63%</p> <p>Number excluded post randomization: none</p>

Femoral nerve blocks for acute postoperative pain after knee replacement surgery (Review)

Wang 2002 (Continued)

Number lost to follow-up: none

Interventions	<p>1. Single-shot FNB with 0.25% bupivacaine with epinephrine 1:100,000. Nerve identified via nerve stimulator. IV PCA morphine 1-mg doses with five-minute lockout</p> <p>2. IV PCA morphine 1-mg doses with five-minute lockout; placebo FNB with 40 ml N/saline</p> <p>Co-analgesics: not stated</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale, PACU, postoperative days one to three</p> <p>Total opioid consumption: postoperative days one and two</p> <p>Morphine related adverse effects</p> <p>Knee flexion postoperative day one, discharge</p> <p>Other outcomes:</p> <p>Distance of ambulation</p> <p>Distance goal met</p> <p>Length of stay</p>
Notes	Study author contacted for additional data on morphine consumption but could not provide the information

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Drawing marked cards from mixed pack with equal numbers of each designation
Allocation concealment (selection bias)	Low risk	Blindly drawn marked cards from mixed pack with equal numbers of each designation
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Quote: "placebo-controlled, double-blind study"
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	All study personnel including outcome assessors blinded to treatment group assignment .
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Wang 2010

Methods	Randomized trial: computer-generated list
Participants	<p>Single centre in Beijing, China</p> <p>126 participants undergoing unilateral TKR</p> <p>Inclusion criteria: ASA I to II, 40 to 84 years, 45 to 95 kg</p> <p>Exclusion criteria: hypersensitivity to anaesthetics used for the study, congenital myopathy and neuropathies, severe obesity (BMI > 35), electrolyte derangement, severe diabetes mellitus, redo surgery, liver/renal failure, bleeding diathesis</p> <p>Mean age (years): 67 ± 13</p> <p>Female: 64%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB with bolus 20 ml 0.5% ropivacaine, infusion: 0.2% ropivacaine 5 ml/h, PCA bolus 5 ml, lockout of 30 minutes for 48 hours</p> <p>2. IV PCA with fentanyl 30 mcg/kg + tropisetron 5 mg in normal saline 100 ml, infusing 2 ml/h, bolus 2 ml, lockout time of 10 minutes</p> <p>Adjunct analgesics: IM pethidine 50 mg or oral Celebrex 200 mg</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale, four hours, eight hours, 12 hours, 16 hours for pain at rest, 24 hours, 36 hours, 48 hours for pain at rest and on movement</p> <p>Proportion of participants in pain</p> <p>Nausea/vomiting</p> <p>Participant satisfaction</p> <p>Other outcomes:</p> <p>Supplemental analgesia (pethidine)</p> <p>Supplemental analgesia (Celebrex)</p>
Notes	Study author provided additional information on method of randomization

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated list (study author provided information)
Allocation concealment (selection bias)	Unclear risk	Method not stated
Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	Not stated. It would have been difficult to blind participants without a placebo block

Wang 2010 (Continued)

Blinding of outcome assessment (detection bias) Subjective outcomes	Unclear risk	Outcome assessor unaware of participant allocation. However it may be obvious that the FNB group has a catheter and the other group does not
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Widmer 2012

Methods	Randomized trial: method not stated
Participants	<p>Single centre in Sydney, Australia</p> <p>55 participants undergoing unilateral TKR</p> <p>Inclusion criteria: 18 to 85 years</p> <p>Exclusion criteria: known allergies to study medication, anatomical aberrations in inguinal area preventing FNB, history of drug or alcohol abuse, significant cognitive impairment or psychiatric disorder, postoperative endotracheal intubation, preoperative use > 40 mg oral morphine or severe cardiac, hepatic or renal disease</p> <p>Median (IQR) age (years): 72.1 (64.4 to 76.5) versus 69.4 (63.4 to 75.5)</p> <p>Female: 44%</p> <p>Number excluded post randomization: none</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. FNB with bolus 30 ml 0.375% ropivacaine and periarticular anaesthetic infiltration 100 ml 0.2% ropivacaine with 0.5 mg adrenaline, to posterior, medial, lateral and anterior capsule, as well as arthrotomy margins, subcutaneous tissue and skin. IV PCA fentanyl 20 µg at five-minute intervals on demand until postoperative day two. Additional boluses allowed</p> <p>2. Sham setup for FNB with skin preparation and dressing, and periarticular anaesthetic infiltration 100 ml 0.2% ropivacaine with 0.5 mg adrenaline, to posterior, medial, lateral and anterior capsule, as well as arthrotomy margins, subcutaneous tissue and skin. IV PCA fentanyl 20 µg at five-minute intervals on demand until postoperative day two. Additional boluses allowed</p> <p>Adjunct analgesics: COX-2 inhibitor and paracetamol 1 g six-hourly. Postoperative day two: oral oxycodone SR 10 mg 12-hourly. Breakthrough pain with oxycodone immediate release 5 to 10 mg three-hourly as needed</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain at rest (most severe pain during first 24 hours on scale of one to four). Visual analogue scale, first 24 hours</p> <p>Total opioid consumption (fentanyl) 24 hours</p> <p>Adverse events</p> <p>Other outcomes:</p>

Widmer 2012 (Continued)

Knee function scores at one year

SF-36

Notes

Study author provided additional information on methods of randomization and allocation concealment

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization (study author provided information)
Allocation concealment (selection bias)	Low risk	Coded and opaque envelopes (study author provided information)
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Quote: "Both the investigators and patients were blinded to treatment modality, and the procedural anaesthetists did not reveal group allocation to any party associated with the study"
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Quote: "blinded study personnel"
Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Xie 2012

Methods	Randomized trial: method not stated
Participants	<p>Single centre in Chicago, USA</p> <p>105 participants undergoing unilateral TKR</p> <p>Exclusion criteria: age < 18 or > 90 years, ASA > III, coagulopathy, cellulitis at the injection site, severe morbid obesity defined as BMI > 40, seizure disorder, severe liver or renal disease, opioid dependency, mental or psychiatric disease affecting ability to comprehend pain scale, preexisting neurological deficits or neuropathy in the lower extremities</p> <p>Mean age (years): 69</p> <p>Female: 72%</p> <p>Number excluded post randomization: six for surgery cancelled, PCA not ordered or missing PCA data sheet</p> <p>Number lost to follow-up: none</p>

Xie 2012 (Continued)

Interventions	<p>1. Single-shot FNB with 30 ml 0.25% bupivacaine with 1:200,000 epinephrine. Nerve identified via nerve stimulation. IV PCA morphine converted to oral pain medication on postoperative day one or two</p> <p>2. Single-shot FNB with 30 ml 0.50% bupivacaine with 1:200,000 epinephrine. Nerve identified via nerve stimulation. IV PCA morphine converted to oral pain medication on postoperative day one or two</p> <p>3. Sham FNB with 30 ml normal saline. IV PCA morphine converted to oral pain medication on postoperative day one or two</p> <p>Co-analgesics: not stated</p>
Outcomes	<p>Pain severity: six-point Likert scale from zero to five “none or minimal” (scores zero and one), “moderate” (scores two and three) and “considerable” (scores four and five)</p> <p>Total opioid consumption: 24 hours</p> <p>Nausea</p> <p>Sedation</p> <p>Participant satisfaction</p>
Notes	Additional information regarding study method sought, but study authors could not be reached

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Method not stated
Allocation concealment (selection bias)	Low risk	Formulated by pharmacists and labelled with numbers only
Blinding of participants and personnel (performance bias) Subjective outcomes	Low risk	Anaesthesiologists, surgeons and participants blinded to treatment assignment
Blinding of outcome assessment (detection bias) Subjective outcomes	Low risk	Not stated
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Six participants withdrawn for surgery cancelled, PCA not ordered or missing PCA data sheet
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Yu 2010

Methods	Randomized trial: random number table
Participants	Single centre in Beijing, China

Yu 2010 (Continued)

80 participants undergoing TKR

Exclusion criteria: lumbar spondylosis, diabetes mellitus, mentally subnormal, insomnia, unable to understand VAS

Mean age (years): 65.0 ± 4

Female: 88%

Number excluded post randomization: none

Number lost to follow-up: none

Interventions	<p>1. Continuous FNB with initial dose of 5 ml 2% lignocaine, infusion 0.2% ropivacaine 200 ml, 4 ml/h for 72 hours</p> <p>2. PCA with sufentanil 200 µg, ondansetron 12 mg in saline 200 ml, infusing 2 ml/h</p> <p>Adjunct analgesics: COX-2 100 mg BD, NSAIDs or opioids</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest: visual analogue scale, two hours, six hours, 24 hours, 36 hours, 48 hours, 56 hours, 72 hours</p> <p>Supplemental analgesics: number of times opioid used</p> <p>Supplemental analgesics: number of times NSAIDs used</p> <p>Nausea/vomiting</p> <p>Giddiness</p> <p>Postoperative active motion of knee joint</p> <p>Other outcomes:</p> <p>Abdominal discomfort</p> <p>Hospital for Special Surgery Knee Score: motor function</p> <p>Restful sleep at 72 hours</p>
Notes	Additional information regarding study method sought, but study authors could not be reached

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number table
Allocation concealment (selection bias)	Unclear risk	Not stated
Blinding of participants and personnel (performance bias) Subjective outcomes	Unclear risk	Not stated
Blinding of outcome assessment (detection bias) Subjective outcomes	Unclear risk	Outcome assessor not involved in surgery. Group with catheter would have been obvious

Yu 2010 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	No dropouts/withdrawals
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Low risk	Appears to be free of other sources of bias

Zaric 2006

Methods	Randomized trial: computerized random number tables
Participants	<p>Single centre in Copenhagen, Denmark</p> <p>60 participants undergoing TKR with general anaesthesia</p> <p>Inclusion criteria: ASA I to III</p> <p>Exclusion criteria: morphine intolerance, neurological disease, coagulation disturbance and chronic pain and rheumatoid arthritis</p> <p>Mean age (years): 66 ± 7</p> <p>Female: 53%</p> <p>Number excluded post randomization: 11 participants (four in continuous FNB and seven in epidural group)</p> <p>Number lost to follow-up: none</p>
Interventions	<p>1. Continuous FNB + sciatic with bolus 30 ml of ropivacaine 7.5 mg/ml and Infusion ropivacaine 2 mg/ml and sufentanil 1 µg/ml at 5 ml/h through femoral nerve catheter; and bolus 30 ml of ropivacaine 7.5 mg/ml and Infusion ropivacaine 0.5 mg/ml at 5 ml/h through sciatic nerve catheter</p> <p>2. Epidural with ropivacaine 7.5 mg/ml given in 5-ml aliquots to attain a level of analgesia at Th 10, and infusion ropivacaine 2 mg/ml and sufentanil 1 µg/ml at 5 ml/h</p> <p>All infusions were continued for 55 hours</p> <p>Adjunct analgesics: IV PCA morphine bolus 2 mg with lockout period of six minutes and maximum dose of 20 mg/h, paracetamol 1 g four times daily. If VAS > three at rest, 5 ml ropivacaine 7.5-mg/ml bolus given through catheters. If pain score > five, 5 ml lidocaine 20-mg/ml bolus given. After cessation of infusions, rofecoxib 25 mg, sustained-release morphine (Contalgin) 20 mg (10 mg for participants > 70 years) twice daily and morphine 10 mg</p>
Outcomes	<p>Outcomes of interest for the review:</p> <p>Pain intensity at rest and on movement: visual analogue scale, four hours post operation, 11 am and 14.30 pm on first, second and morning of third postoperative days</p> <p>Total opioid consumption: postoperative days one and two</p> <p>Nausea/vomiting</p> <p>Dizziness (moderate to severe)</p> <p>Sedation</p> <p>Pruritus</p>

Zaric 2006 (Continued)

Urinary retention

Adverse events

Knee flexion: postoperative day one, discharge

Other outcomes:

Rehab indices

Length of stay

Notes Study author provided additional data on mean (SD) for pain intensity on movement and at rest with knee ROM flexion

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computerized random number tables
Allocation concealment (selection bias)	Low risk	Sealed opaque envelopes
Blinding of participants and personnel (performance bias) Subjective outcomes	High risk	Blinding of groups not attempted
Blinding of outcome assessment (detection bias) Subjective outcomes	High risk	Blinding of groups not attempted
Incomplete outcome data (attrition bias) All outcomes	High risk	11 participants (18%) excluded post randomization (four in continuous FNB and six in epidural group because of insufficient analgesia; one in epidural group had an acute myocardial infarction followed by coronary artery bypass graft)
Selective reporting (reporting bias)	Low risk	Outcomes predefined
Other bias	Unclear risk	Potential impact of high attrition on ITT bias may be an issue, as the study did not use ITT analysis

Abbreviations:

ASA: American Society of Anaesthesiologists; BD: twice a day; BMI: body mass index; COX: cyclo-oxygenase; CPM: continuous passive motion; CPNB: continuous peripheral nerve block; FNB: femoral nerve block; IQR: interquartile range; ITT: intention-to-treat; NSAID: non-steroidal anti-inflammatory drug; PACU: postanesthesia care unit; PCA: patient-controlled analgesia; PCEA: patient-controlled epidural analgesia; ROM: range of motion; SD: standard deviation; SF: Short Form; TKR: total knee replacement; VAS: visual analogue scale; WOMAC: Western Ontario and McMaster Osteoarthritis Index.

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Affas 2012	Secondary analysis of Affas 2011 (an included study). Wrong outcomes.

Study	Reason for exclusion
Bagry 2008	Wrong intervention: lumbar plexus versus PCA opioid
Blisard 2012	Not an RCT
Cappelleri 2011	Population included arthrolysis participants with no separate data for TKR participants
Carvalho 2012	No non-FNB group (continuous FNB vs continuous FNB + sciatic)
Chelly 2001	Not an RCT
Combes 2000	Population included knee surgery participants with no separate data for TKR participants
Dang 2005	No non-FNB group (continuous FNB vs continuous FNB + sciatic)
Edwards 1992	Wrong comparator (intramuscular papaveretum)
Frassanito 2009	Wrong intervention (single-shot lumbar plexus block vs continuous lumbar plexus/sciatic blocks)
Frassanito 2010	Wrong comparator (intrathecal morphine)
Hakkalamani 2008	Not an RCT. No non-FNB comparator (FNB vs FNB + sciatic)
Ilfeld 2008	Wrong intervention
Ilfeld 2009	Wrong intervention
Ilfeld 2010	Wrong intervention
Johnson 2011	Wrong comparator (DepoDur extended-release epidural morphine)
Koh 2011	Wrong intervention (periarticular multimodal drug injection + multimodal pain control vs multimodal pain control)
McMeniman 2010	No non-FNB group (FNB vs fascia iliaca block)
Meftah 2012	Wrong intervention (FNB/epidural)
Morin 2005	No non-FNB group (continuous posterior lumbar plexus block vs continuous FNB vs continuous FNB + sciatic nerve block)
Ng 2012b	Not an RCT
Niskanen 2005	Wrong intervention (bedside femoral blocks on first postoperative day vs IM oxycodone/tramadol/epidural and additional femoral blocks)
Rais 2009	No non-FNB comparator
Rajeev 2007	No non-FNB comparator
Rasiah 2012	Compared collateral knees in participants undergoing simultaneous bilateral total knee replacement
Safa 2011	Interim analysis
Serrano 2011	No non-FNB comparator

Study	Reason for exclusion
Sites 2004	Wrong control group (intrathecal morphine)
Taninishi 2011	No non-FNB comparator
Tarkkila 1998	Population included major knee surgery participants with no separate data for TKR participants. Wrong comparator (intrathecal morphine)
Tricarico 2009	Population included major knee surgery participants with no separate data for TKR participants
Watson 2005	Wrong intervention (continuous vs single-shot lumbar plexus blocks)
Wegener 2011	No non-FNB comparator (FNB vs FNB + single-shot sciatic nerve block vs FNB/continuous sciatic nerve block)
Weston-Simons 2012	No non-FNB comparator
Zhang 2010	Not an RCT. Population included artificial joint replacement participants with no separate data for TKR participants
Ziwenga 2010	No non-FNB comparator (continuous infusion vs patient-controlled FNB)

Abbreviations:

FNB: femoral nerve block; PCA: patient-controlled analgesia; RCT: randomized controlled trial; TKR: total knee replacement.

Characteristics of studies awaiting assessment *[ordered by study ID]*

[Grider 2011](#)

Methods	Randomized trial: method not stated
Participants	Number of centre: not stated. Study done in USA 80 participants undergoing knee replacements
Interventions	1. Continuous FNB with bolus 30 ml 0.5% ropivacaine followed by postoperative infusion 8 ml/h 0.2% ropivacaine 2. Single-shot FNB with bolus 30 ml 0.5% ropivacaine Postoperation: Both groups received rescue IV PCA morphine first 48 hours followed by oral oxy-codone
Outcomes	VAS pain scores at rest Morphine consumption Knee range of motion Local anaesthetic or opioid analgesic complications Hospital length of stay
Notes	Conference abstract: possibly eligible but not in format that is easy to appraise

Mullen 2008

Methods	Unable to obtain abstract or to contact study author for details
Participants	Ditto
Interventions	Ditto
Outcomes	Ditto
Notes	Conference abstract: awaiting information to determine eligibility

Tobin 2011

Methods	Randomized trial: method not stated
Participants	Number of centre: not stated. Study done in India 63 participants undergoing unilateral TKR Inclusions: ASA I to II
Interventions	1. Continuous FNB with bolus 30 ml 0.25% bupivacaine, followed by postoperative infusion 8 ml/h 0.125% bupivacaine 2. Epidural with bolus 8 ml 0.25% bupivacaine, infusion 8 ml/h 0.125% bupivacaine
Outcomes	VAS pain scores Adverse effects Motor blockade Fentanyl and bupivacaine consumption Rehabilitation indices Oxford knee scores
Notes	Conference abstract with no quantifiable outcome data provided

Yuksel 2011

Methods	Randomized trial: method not stated.
Participants	Number of centre: not stated. Study done in Turkey 100 participants undergoing unilateral TKR Inclusions: ASA I to III
Interventions	1. Continuous FNB with single-shot sciatic, and PCA 0.125% bupivacaine, bolus 0.05 ml/kg, lockout time of 30 minutes, infusion 0.1 ml/kg/h for 48 hours 2. Continuous epidural with PCA 0.125% bupivacaine, bolus 0.05 ml/kg, lockout time of 30 minutes, infusion 0.1 ml/kg/h Both groups: if VAS > four: subcutaneous morphine 0.1 mg/kg

Yuksel 2011 (Continued)

Outcomes	VAS pain scores
	Total bupivacaine consumption
	Degree of knee flexion
	Participant satisfaction
Notes	Conference abstract with no quantifiable outcome data provided

Abbreviations:

ASA: American Society of Anaesthesiologists; FNB: femoral nerve block; PCA: patient-controlled analgesia; TKR: total knee replacement; VAS: visual analogue scale.

DATA AND ANALYSES

Comparison 1. FNB versus PCA opioid

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain at rest first 2 hours	11		Std. Mean Difference (Random, 95% CI)	-0.58 [1.00, -0.16]
1.1 Single-shot FNB vs PCA opioid	7		Std. Mean Difference (Random, 95% CI)	-0.44 [-1.16, 0.28]
1.2 Single-shot FNB + obturator vs PCA opioid	1		Std. Mean Difference (Random, 95% CI)	-1.25 [-1.92, -0.57]
1.3 Continuous FNB vs PCA opioid	5		Std. Mean Difference (Random, 95% CI)	-0.62 [-1.25, 0.00]
1.4 Continuous FNB + sciatic vs PCA opioid	1		Std. Mean Difference (Random, 95% CI)	-0.63 [-1.63, 0.38]
2 Pain at rest 3 to 12 hours	14	972	Std. Mean Difference (IV, Random, 95% CI)	-0.97 [-1.42, -0.52]
2.1 Single-shot FNB vs PCA opioid	8	388	Std. Mean Difference (IV, Random, 95% CI)	-0.66 [-1.20, -0.11]
2.2 Single-shot FNB + sciatic vs PCA opioid	1	18	Std. Mean Difference (IV, Random, 95% CI)	-1.38 [-2.48, -0.27]
2.3 Single-shot FNB + obturator vs PCA opioid	1	47	Std. Mean Difference (IV, Random, 95% CI)	-0.37 [-1.01, 0.26]
2.4 Continuous FNB vs PCA opioid	8	519	Std. Mean Difference (IV, Random, 95% CI)	-1.29 [-2.06, -0.52]
3 Pain at rest 24 hours	19		Std. Mean Difference (Random, 95% CI)	-0.72 [-0.93, -0.51]
3.1 Single-shot FNB vs PCA opioid	9		Std. Mean Difference (Random, 95% CI)	-0.65 [-1.08, -0.22]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
3.2 Single-shot FNB + sci-atic vs PCA opioid	1		Std. Mean Difference (Random, 95% CI)	-0.73 [-1.74, 0.28]
3.3 Continuous FNB vs PCA opioid	10		Std. Mean Difference (Random, 95% CI)	-0.74 [-0.97, -0.51]
3.4 Continuous FNB + sci-atic vs PCA opioid	2		Std. Mean Difference (Random, 95% CI)	-0.85 [-1.55, -0.15]
4 Pain at rest 48 hours	17	957	Std. Mean Difference (IV, Random, 95% CI)	-0.64 [-1.03, -0.25]
4.1 Single-shot FNB vs PCA opioid	9	416	Std. Mean Difference (IV, Random, 95% CI)	-0.38 [-0.83, 0.07]
4.2 Single-shot FNB + sci-atic vs PCA opioid	1	18	Std. Mean Difference (IV, Random, 95% CI)	0.21 [-0.77, 1.20]
4.3 Continuous FNB vs PCA opioid	9	507	Std. Mean Difference (IV, Random, 95% CI)	-0.89 [-1.58, -0.20]
4.4 Continuous FNB + sci-atic vs PCA opioid	1	16	Std. Mean Difference (IV, Random, 95% CI)	-1.96 [-3.21, -0.71]
5 Pain at rest 72 hours	8	560	Std. Mean Difference (IV, Random, 95% CI)	-0.67 [-1.32, -0.01]
5.1 Single-shot FNB vs PCA opioid	5	272	Std. Mean Difference (IV, Random, 95% CI)	-0.26 [-0.61, 0.08]
5.2 Continuous FNB vs PCA opioid	4	250	Std. Mean Difference (IV, Random, 95% CI)	-1.24 [-3.07, 0.60]
5.3 Continuous FNB + sci-atic vs PCA opioid	1	38	Std. Mean Difference (IV, Random, 95% CI)	-0.56 [-1.21, 0.09]
6 Pain on movement first 2 hours	4	218	Std. Mean Difference (IV, Random, 95% CI)	-1.29 [-2.12, -0.46]
6.1 Single-shot FNB vs PCA opioid	2	98	Std. Mean Difference (IV, Random, 95% CI)	-0.85 [-1.27, -0.43]
6.2 Continuous FNB vs PCA opioid	3	120	Std. Mean Difference (IV, Random, 95% CI)	-1.60 [-3.39, 0.19]
7 Pain on movement 3 to 12 hours	8		Std. Mean Difference (Random, 95% CI)	-1.06 [-1.68, -0.43]
7.1 Single-shot FNB vs PCA opioid	2		Std. Mean Difference (Random, 95% CI)	-1.22 [-1.66, -0.79]
7.2 Single-shot FNB + sci-atic vs PCA opioid	1		Std. Mean Difference (Random, 95% CI)	-0.34 [-0.83, 0.15]
7.3 Continuous FNB vs PCA opioid	6		Std. Mean Difference (Random, 95% CI)	-1.18 [-2.13, -0.23]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
8 Pain on movement 24 hours	17		Std. Mean Difference (Random, 95% CI)	-0.94 [-1.32, -0.55]
8.1 Single-shot FNB vs PCA opioid	6		Std. Mean Difference (Random, 95% CI)	-0.50 [-0.81, -0.19]
8.2 Single-shot FNB + sci-atic vs PCA opioid	2		Std. Mean Difference (Random, 95% CI)	-0.73 [-1.18, -0.29]
8.3 Continuous FNB vs PCA opioid	10		Std. Mean Difference (Random, 95% CI)	-1.09 [-1.74, -0.43]
8.4 Continuous FNB + sci-atic vs PCA opioid	2		Std. Mean Difference (Random, 95% CI)	-2.08 [-4.74, 0.58]
9 Pain on movement 48 hours	13	742	Std. Mean Difference (IV, Random, 95% CI)	-0.44 [-0.71, -0.16]
9.1 Single-shot FNB vs PCA opioid	6	287	Std. Mean Difference (IV, Random, 95% CI)	-0.24 [-0.48, -0.00]
9.2 Single-shot FNB + sci-atic vs PCA opioid	2	92	Std. Mean Difference (IV, Random, 95% CI)	0.34 [-0.10, 0.78]
9.3 Continuous FNB vs PCA opioid	7	347	Std. Mean Difference (IV, Random, 95% CI)	-0.60 [-0.95, -0.24]
9.4 Continuous FNB + sci-atic vs PCA opioid	1	16	Std. Mean Difference (IV, Random, 95% CI)	-2.81 [-4.30, -1.32]
10 Pain on movement 72 hours	6	438	Std. Mean Difference (IV, Random, 95% CI)	-0.17 [-0.39, 0.04]
10.1 Single-shot FNB vs PCA opioid	4	230	Std. Mean Difference (IV, Random, 95% CI)	-0.11 [-0.38, 0.16]
10.2 Continuous FNB vs PCA opioid	3	170	Std. Mean Difference (IV, Random, 95% CI)	-0.09 [-0.40, 0.23]
10.3 Continuous FNB + sci-atic vs PCA opioid	1	38	Std. Mean Difference (IV, Random, 95% CI)	-0.78 [-1.44, -0.12]
11 Pain at rest 24 hours—subgrouped by FNB with and without concurrent PCA	19		Std. Mean Difference (Random, 95% CI)	Subtotals only
11.1 FNB with concurrent PCA	15		Std. Mean Difference (Random, 95% CI)	-0.67 [-0.94, -0.41]
11.2 FNB without concurrent PCA	5		Std. Mean Difference (Random, 95% CI)	-0.93 [-1.31, -0.55]
12 Pain on movement 24 hours—subgrouped by	17		Std. Mean Difference (Random, 95% CI)	Subtotals only

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
FNB with and without concurrent PCA				
12.1 Concurrent PCA in FNB	14		Std. Mean Difference (Random, 95% CI)	-0.67 [-0.99, -0.35]
12.2 No concurrent PCA in FNB	4		Std. Mean Difference (Random, 95% CI)	-2.26 [-3.95, -0.57]
13 Pain at rest 24 hours—subgrouped by FNB ropivacaine vs bupivacaine	19		Std. Mean Difference (Random, 95% CI)	Subtotals only
13.1 FNB with ropivacaine	8		Std. Mean Difference (Random, 95% CI)	-1.30 [-2.03, -0.58]
13.2 FNB with bupivacaine	11		Std. Mean Difference (Random, 95% CI)	-0.61 [-0.89, -0.33]
14 Pain on movement 24 hours—subgrouped by FNB ropivacaine vs bupivacaine	17		Std. Mean Difference (Random, 95% CI)	Subtotals only
14.1 FNB with ropivacaine	8		Std. Mean Difference (Random, 95% CI)	-0.98 [-1.67, -0.29]
14.2 FNB with bupivacaine	10		Std. Mean Difference (Random, 95% CI)	-0.92 [-1.39, -0.44]
15 Pain at rest 24 hours—sensitivity analysis by low bias for allocation concealment	9		Std. Mean Difference (Random, 95% CI)	-0.54 [-0.76, -0.32]
16 Pain on movement 24 hours—sensitivity analysis by low bias for allocation concealment	9		Std. Mean Difference (Random, 95% CI)	-0.54 [-0.75, -0.33]
17 Pain on rest 24 hours—sensitivity analysis by blinding of participants, personnel and outcome assessor	7		Std. Mean Difference (Random, 95% CI)	-0.62 [-1.08, -0.17]
18 Pain on movement 24 hours—sensitivity analysis by blinding of participants, personnel and outcome assessor	7		Std. Mean Difference (Random, 95% CI)	-0.57 [-0.79, -0.35]
19 Opioid consumption 24 hours (mg)	20		Mean Difference (Random, 95% CI)	-14.74 [-18.68, -10.81]
19.1 SFNB vs PCA opioid	12		Mean Difference (Random, 95% CI)	-12.86 [-18.65, -7.08]
19.2 SFNB/Sciatic vs PCA opioid	2		Mean Difference (Random, 95% CI)	-17.93 [-21.43, -14.43]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
19.3 SFNB/Obturator vs PCA opioid	1		Mean Difference (Random, 95% CI)	-13.7 [-18.28, -9.12]
19.4 CFNB vs PCA opioid	8		Mean Difference (Random, 95% CI)	-16.89 [-24.01, -9.77]
19.5 CFNB/Sciatic vs PCA opioid	1		Mean Difference (Random, 95% CI)	-13.0 [-20.17, -5.83]
20 Opioid consumption 48 hours (mg)	19	1001	Mean Difference (IV, Random, 95% CI)	-14.53 [-20.03, -9.02]
20.1 Single-shot FNB vs PCA opioid	11	485	Mean Difference (IV, Random, 95% CI)	-13.21 [-21.99, -4.44]
20.2 Single-shot FNB + sciatic vs PCA opioid	2	92	Mean Difference (IV, Random, 95% CI)	-10.10 [-18.26, -1.94]
20.3 Continuous FNB vs PCA opioid	8	386	Mean Difference (IV, Random, 95% CI)	-19.14 [-27.53, -10.76]
20.4 Continuous FNB + sciatic vs PCA opioid	1	38	Mean Difference (IV, Random, 95% CI)	-6.0 [-17.39, 5.39]
21 Nausea and/or vomiting	16	1100	Risk Ratio (M-H, Random, 95% CI)	0.47 [0.33, 0.68]
21.1 Single-shot FNB vs PCA opioid	8	479	Risk Ratio (M-H, Random, 95% CI)	0.67 [0.44, 1.02]
21.2 Single-shot FNB + obturator vs PCA opioid	1	47	Risk Ratio (M-H, Random, 95% CI)	0.25 [0.11, 0.56]
21.3 Continuous FNB vs PCA opioid	9	574	Risk Ratio (M-H, Random, 95% CI)	0.35 [0.18, 0.68]
22 Sedation	9	808	Risk Ratio (M-H, Random, 95% CI)	0.34 [0.16, 0.74]
22.1 Single-shot FNB vs PCA opioid	4	326	Risk Ratio (M-H, Random, 95% CI)	0.52 [0.16, 1.75]
22.2 Single-shot FNB + obturator vs PCA opioid	1	47	Risk Ratio (M-H, Random, 95% CI)	0.42 [0.15, 1.24]
22.3 Continuous FNB vs PCA opioid	6	435	Risk Ratio (M-H, Random, 95% CI)	0.20 [0.05, 0.77]
23 Urinary retention	7	490	Risk Ratio (M-H, Random, 95% CI)	0.57 [0.20, 1.68]
23.1 Single-shot FNB vs PCA opioid	3	183	Risk Ratio (M-H, Random, 95% CI)	1.66 [0.43, 6.45]
23.2 Continuous FNB vs PCA opioid	5	307	Risk Ratio (M-H, Random, 95% CI)	0.29 [0.06, 1.36]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
24 Knee flexion range of motion (postoperative day 2 to 4)	10	541	Mean Difference (IV, Random, 95% CI)	6.48 [4.27, 8.69]
24.1 Single-shot FNB vs PCA opioid	5	295	Mean Difference (IV, Random, 95% CI)	3.74 [1.45, 6.04]
24.2 Continuous FNB vs PCA opioid	4	192	Mean Difference (IV, Random, 95% CI)	10.32 [4.70, 15.93]
24.3 Continuous FNB + sciatic vs PCA opioid	2	54	Mean Difference (IV, Random, 95% CI)	12.78 [5.96, 19.60]
25 Participant satisfaction with analgesia during hospital stay	4	180	Mean Difference (IV, Random, 95% CI)	1.68 [0.79, 2.58]
25.1 Continuous FNB vs PCA opioid	4	180	Mean Difference (IV, Random, 95% CI)	1.68 [0.79, 2.58]

Analysis 1.1. Comparison 1 FNB versus PCA opioid, Outcome 1 Pain at rest first 2 hours.

Analysis 1.2. Comparison 1 FNB versus PCA opioid, Outcome 2 Pain at rest 3 to 12 hours.

Analysis 1.3. Comparison 1 FNB versus PCA opioid, Outcome 3 Pain at rest 24 hours.

Analysis 1.4. Comparison 1 FNB versus PCA opioid, Outcome 4 Pain at rest 48 hours.

Analysis 1.5. Comparison 1 FNB versus PCA opioid, Outcome 5 Pain at rest 72 hours.

Analysis 1.6. Comparison 1 FNB versus PCA opioid, Outcome 6 Pain on movement first 2 hours.

Analysis 1.7. Comparison 1 FNB versus PCA opioid, Outcome 7 Pain on movement 3 to 12 hours.

Analysis 1.8. Comparison 1 FNB versus PCA opioid, Outcome 8 Pain on movement 24 hours.

Analysis 1.9. Comparison 1 FNB versus PCA opioid, Outcome 9 Pain on movement 48 hours.

Analysis 1.10. Comparison 1 FNB versus PCA opioid, Outcome 10 Pain on movement 72 hours.

Study or subgroup	FNB		PCA Opioid		Std. Mean Difference Random, 95% CI	Weight	Std. Mean Difference Random, 95% CI			
	N	Mean(SD)	N	Mean(SD)						
Heterogeneity: Tau²=0.01; Chi²=7.89, df=7(P=0.34); I²=11.29%										
Test for overall effect: Z=1.61(P=0.11)										
Test for subgroup differences: Chi²=3.72, df=1 (P=0.16), I²=46.17%										
Favours FNB					-4	-2	0	2	4	Favours PCA Opioid

Analysis 1.11. Comparison 1 FNB versus PCA opioid, Outcome 11 Pain at rest 24 hours—subgrouped by FNB with and without concurrent PCA.

Study or subgroup	Experi- mental	Control	Std. Mean Difference	Std. Mean Difference	Weight	Std. Mean Difference
	N	N	(SE)	IV, Random, 95% CI		IV, Random, 95% CI
1.11.1 FNB with concurrent PCA						
Allen 1998	12	6	-0.7 (0.514)		4.05%	-0.73[-1.74,0.28]
Allen 1998	12	6	0 (0.5)		4.19%	0.03[-0.95,1.01]
Chan 2012	0	0	-1.7 (0.26)		7.17%	-1.75[-2.26,-1.24]
Chan 2013	69	33	-0.3 (0.213)		7.86%	-0.29[-0.71,0.12]
Good 2007	22	20	-0.3 (0.311)		6.44%	-0.34[-0.95,0.27]
Hirst 1996	11	5	-0.4 (0.511)		4.09%	-0.35[-1.36,0.65]
Hirst 1996	11	6	-0.2 (0.541)		3.82%	-0.24[-1.3,0.82]
Hunt 2009	33	24	-0.7 (0.276)		6.93%	-0.71[-1.25,-0.16]
Kaloul 2004	20	20	-0.7 (0.326)		6.23%	-0.72[-1.36,-0.08]
Kardash 2007	19	20	0 (0.32)		6.31%	0.02[-0.61,0.65]
Martin 2008	20	18	-0.6 (0.332)		6.15%	-0.6[-1.25,0.05]
Ozen 2006	15	15	-1.1 (0.393)		5.35%	-1.13[-1.9,-0.36]
Seet 2006	35	20	-0.3 (0.282)		6.85%	-0.31[-0.86,0.24]
Serpell 2001	13	16	-0.3 (0.375)		5.58%	-0.29[-1.03,0.44]
Tang 2010	16	16	-1 (0.376)		5.57%	-1.03[-1.77,-0.29]
Wang 2002	15	15	-1.2 (0.396)		5.32%	-1.18[-1.96,-0.41]
Wang 2010	63	63	-1.3 (0.196)		8.09%	-1.29[-1.68,-0.91]
Subtotal (95% CI)					100%	-0.67[-0.94,-0.41]
Heterogeneity: Tau ² =0.19; Chi ² =45.37, df=16(P=0); I ² =64.73%						
Test for overall effect: Z=4.97(P<0.0001)						
1.11.2 FNB without concurrent PCA						
Baranovic 2011	35	36	-0.9 (0.25)		26.59%	-0.94[-1.43,-0.45]
Chan 2013	65	33	-0.5 (0.217)		29.91%	-0.5[-0.92,-0.07]
Mistraletti 2006	8	8	-1.5 (0.375)		17.08%	-1.55[-2.28,-0.81]
Singelyn 1998	15	15	-1.4 (0.555)		9.66%	-1.36[-2.45,-0.27]
Yu 2010	40	40	-0.8 (0.38)		16.76%	-0.82[-1.57,-0.08]
Subtotal (95% CI)					100%	-0.93[-1.31,-0.55]
Heterogeneity: Tau ² =0.08; Chi ² =7.07, df=4(P=0.13); I ² =43.4%						
Test for overall effect: Z=4.83(P<0.0001)						
Test for subgroup differences: Chi ² =1.18, df=1 (P=0.28), I ² =15.34%						
<div style="display: flex; justify-content: space-between;">Favours FNB-4 -2 0 2 4Favours PCA Opioid</div>						

Analysis 1.12. Comparison 1 FNB versus PCA opioid, Outcome 12 Pain on movement 24 hours—subgrouped by FNB with and without concurrent PCA.

Analysis 1.13. Comparison 1 FNB versus PCA opioid, Outcome 13 Pain at rest 24 hours—subgrouped by FNB ropivacaine vs bupivacaine.

Analysis 1.14. Comparison 1 FNB versus PCA opioid, Outcome 14 Pain on movement 24 hours—subgrouped by FNB ropivacaine vs bupivacaine.

Analysis 1.15. Comparison 1 FNB versus PCA opioid, Outcome 15 Pain at rest 24 hours—sensitivity analysis by low bias for allocation concealment.

Analysis 1.16. Comparison 1 FNB versus PCA opioid, Outcome 16 Pain on movement 24 hours—sensitivity analysis by low bias for allocation concealment.

Analysis 1.17. Comparison 1 FNB versus PCA opioid, Outcome 17 Pain on rest 24 hours—sensitivity analysis by blinding of participants, personnel and outcome assessor.

Analysis 1.18. Comparison 1 FNB versus PCA opioid, Outcome 18 Pain on movement 24 hours—sensitivity analysis by blinding of participants, personnel and outcome assessor.

Analysis 1.19. Comparison 1 FNB versus PCA opioid, Outcome 19 Opioid consumption 24 hours (mg).

Analysis 1.20. Comparison 1 FNB versus PCA opioid, Outcome 20 Opioid consumption 48 hours (mg).

Analysis 1.21. Comparison 1 FNB versus PCA opioid, Outcome 21 Nausea and/or vomiting.

Analysis 1.22. Comparison 1 FNB versus PCA opioid, Outcome 22 Sedation.

Analysis 1.23. Comparison 1 FNB versus PCA opioid, Outcome 23 Urinary retention.

**Analysis 1.24. Comparison 1 FNB versus PCA opioid, Outcome
24 Knee flexion range of motion (postoperative day 2 to 4).** **Analysis 1.25. Comparison 1 FNB versus PCA opioid, Outcome
25 Participant satisfaction with analgesia during hospital stay.**

Comparison 2. FNB versus epidural

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain at rest first 2 hours	3		Std. Mean Difference (Random, 95% CI)	0.53 [-0.56, 1.62]
1.1 Single-shot FNB vs epidural	1		Std. Mean Difference (Random, 95% CI)	2.08 [1.33, 2.83]
1.2 Continuous FNB vs epidural	1		Std. Mean Difference (Random, 95% CI)	0.08 [-0.72, 0.88]
1.3 Continuous FNB + sciatic vs epidural	2		Std. Mean Difference (Random, 95% CI)	-0.05 [-0.67, 0.57]
2 Pain at rest 3 to 12 hours	2	120	Std. Mean Difference (IV, Random, 95% CI)	-0.20 [-1.08, 0.67]
2.1 Continuous FNB vs epidural	1	70	Std. Mean Difference (IV, Random, 95% CI)	-0.64 [-1.12, -0.16]
2.2 Continuous FNB + sciatic vs epidural	1	50	Std. Mean Difference (IV, Random, 95% CI)	0.25 [-0.30, 0.81]
3 Pain at rest 24 hours	6		Std. Mean Difference (Random, 95% CI)	-0.05 [-0.43, 0.32]
3.1 Continuous FNB vs epidural	4		Std. Mean Difference (Random, 95% CI)	-0.11 [-0.72, 0.50]
3.2 Continuous FNB + sciatic vs epidural	3		Std. Mean Difference (Random, 95% CI)	0.09 [-0.32, 0.51]
4 Pain at rest 48 hours	6	328	Std. Mean Difference (IV, Random, 95% CI)	-0.14 [-0.51, 0.22]
4.1 Continuous FNB vs epidural	4	235	Std. Mean Difference (IV, Random, 95% CI)	-0.28 [-0.77, 0.21]
4.2 Continuous FNB + sciatic vs epidural	3	93	Std. Mean Difference (IV, Random, 95% CI)	0.07 [-0.34, 0.49]
5 Pain on movement 24 hours	6	317	Std. Mean Difference (IV, Random, 95% CI)	0.01 [-0.21, 0.24]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
5.1 Single-shot FNB + sciatic vs epidural	1	59	Std. Mean Difference (IV, Random, 95% CI)	-0.33 [-0.85, 0.18]
5.2 Continuous FNB vs epidural	3	165	Std. Mean Difference (IV, Random, 95% CI)	0.12 [-0.19, 0.43]
5.3 Continuous FNB + sciatic vs epidural	3	93	Std. Mean Difference (IV, Random, 95% CI)	0.06 [-0.36, 0.47]
6 Pain on movement 48 hours	6	317	Std. Mean Difference (IV, Random, 95% CI)	-0.10 [-0.32, 0.13]
6.1 Single-shot FNB + sciatic vs epidural	1	59	Std. Mean Difference (IV, Random, 95% CI)	-0.41 [-0.92, 0.11]
6.2 Continuous FNB vs epidural	3	165	Std. Mean Difference (IV, Random, 95% CI)	-0.06 [-0.37, 0.25]
6.3 Continuous FNB + sciatic vs epidural	3	93	Std. Mean Difference (IV, Random, 95% CI)	0.07 [-0.41, 0.54]
7 Pain at rest 24 hours—subgrouped by FNB ropivacaine vs bupivacaine	6		Std. Mean Difference (Random, 95% CI)	Subtotals only
7.1 FNB with ropivacaine	4		Std. Mean Difference (Random, 95% CI)	-0.16 [-0.68, 0.37]
7.2 FNB with bupivacaine	2		Std. Mean Difference (Random, 95% CI)	0.20 [-0.13, 0.53]
8 Pain on movement 24 hours—subgrouped by FNB ropivacaine vs bupivacaine	6		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
8.1 FNB with ropivacaine	3	120	Std. Mean Difference (IV, Random, 95% CI)	0.10 [-0.27, 0.47]
8.2 FNB with bupivacaine	3	197	Std. Mean Difference (IV, Random, 95% CI)	-0.03 [-0.31, 0.25]
9 Pain at rest 24 hours—sensitivity analysis by low bias for allocation concealment	3		Std. Mean Difference (Random, 95% CI)	0.20 [-0.07, 0.47]
10 Pain on movement 24 hours—sensitivity analysis by low bias for allocation concealment	4	271	Std. Mean Difference (IV, Random, 95% CI)	0.03 [-0.21, 0.27]
11 Opioid consumption 24 hours (mg)	5	341	Mean Difference (IV, Random, 95% CI)	-4.35 [-9.95, 1.26]
11.1 Single-shot FNB + sciatic vs epidural	1	59	Mean Difference (IV, Random, 95% CI)	0.0 [-6.28, 6.28]
11.2 Continuous FNB vs epidural	3	205	Mean Difference (IV, Random, 95% CI)	-9.06 [-19.46, 1.34]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
11.3 Continuous FNB + sciatic vs epidural	2	77	Mean Difference (IV, Random, 95% CI)	-0.56 [-8.24, 7.12]
12 Opioid consumption 48 hours (mg)	4	233	Mean Difference (IV, Random, 95% CI)	-1.28 [-5.30, 2.74]
12.1 Single-shot FNB + sciatic vs epidural	1	59	Mean Difference (IV, Random, 95% CI)	-4.0 [-18.88, 10.88]
12.2 Continuous FNB vs epidural	2	97	Mean Difference (IV, Random, 95% CI)	-3.54 [-12.42, 5.34]
12.3 Continuous FNB + sciatic vs epidural	2	77	Mean Difference (IV, Random, 95% CI)	0.62 [-5.51, 6.75]
13 Nausea and/or vomiting	4	183	Risk Ratio (M-H, Random, 95% CI)	0.63 [0.41, 0.97]
13.1 Single-shot FNB vs epidural	1	42	Risk Ratio (M-H, Random, 95% CI)	0.64 [0.36, 1.15]
13.2 Continuous FNB vs epidural	2	91	Risk Ratio (M-H, Random, 95% CI)	0.67 [0.22, 2.07]
13.3 Continuous FNB + sciatic vs epidural	1	50	Risk Ratio (M-H, Random, 95% CI)	0.57 [0.10, 3.11]

Analysis 2.1. Comparison 2 FNB versus epidural, Outcome 1 Pain at rest first 2 hours.

Analysis 2.2. Comparison 2 FNB versus epidural, Outcome 2 Pain at rest 3 to 12 hours.

Analysis 2.3. Comparison 2 FNB versus epidural, Outcome 3 Pain at rest 24 hours.

Analysis 2.4. Comparison 2 FNB versus epidural, Outcome 4 Pain at rest 48 hours.

Analysis 2.5. Comparison 2 FNB versus epidural, Outcome 5 Pain on movement 24 hours.

Analysis 2.6. Comparison 2 FNB versus epidural, Outcome 6 Pain on movement 48 hours.

Analysis 2.7. Comparison 2 FNB versus epidural, Outcome 7 Pain at rest 24 hours—subgrouped by FNB ropivacaine vs bupivacaine.

Analysis 2.8. Comparison 2 FNB versus epidural, Outcome 8 Pain on movement 24 hours—subgrouped by FNB ropivacaine vs bupivacaine.

Analysis 2.9. Comparison 2 FNB versus epidural, Outcome 9 Pain at rest 24 hours—sensitivity analysis by low bias for allocation concealment.

Analysis 2.10. Comparison 2 FNB versus epidural, Outcome 10 Pain on movement 24 hours—sensitivity analysis by low bias for allocation concealment.

Analysis 2.11. Comparison 2 FNB versus epidural, Outcome 11 Opioid consumption 24 hours (mg). **Analysis 2.12. Comparison 2 FNB versus epidural, Outcome 12 Opioid consumption 48 hours (mg).**

Analysis 2.13. Comparison 2 FNB versus epidural, Outcome 13 Nausea and/or vomiting.

Comparison 3. FNB versus local infiltration analgesia

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain at rest 24 hours	4	216	Std. Mean Difference (IV, Random, 95% CI)	0.06 [-0.61, 0.72]
1.1 Single-shot FNB vs local infiltration	1	60	Std. Mean Difference (IV, Random, 95% CI)	0.50 [-0.01, 1.02]
1.2 Continuous FNB vs local infiltration	3	156	Std. Mean Difference (IV, Random, 95% CI)	-0.11 [-1.01, 0.80]
2 Pain at rest 48 hours	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
2.1 Continuous FNB vs local infiltration	2	114	Std. Mean Difference (IV, Random, 95% CI)	-0.26 [-0.94, 0.43]
3 Pain on movement 24 hours	3		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
3.1 Continuous FNB vs local infiltration	3	153	Std. Mean Difference (IV, Random, 95% CI)	0.38 [-0.10, 0.86]
4 Pain on movement 48 hours	2		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
4.1 Continuous FNB vs local infiltration	2	111	Std. Mean Difference (IV, Random, 95% CI)	-0.14 [-0.71, 0.43]
5 Nausea and/or vomiting	3	177	Risk Ratio (M-H, Random, 95% CI)	1.71 [0.64, 4.62]
5.1 SFNB vs local infiltration	1	60	Risk Ratio (M-H, Random, 95% CI)	2.14 [0.42, 10.80]
5.2 CFNB vs local infiltration	2	117	Risk Ratio (M-H, Random, 95% CI)	1.97 [0.35, 11.04]

Analysis 3.1. Comparison 3 FNB versus local infiltration analgesia, Outcome 1 Pain at rest 24 hours.

Analysis 3.2. Comparison 3 FNB versus local infiltration analgesia, Outcome 2 Pain at rest 48 hours.

Analysis 3.3. Comparison 3 FNB versus local infiltration analgesia, Outcome 3 Pain on movement 24 hours.

Analysis 3.4. Comparison 3 FNB versus local infiltration analgesia, Outcome 4 Pain on movement 48 hours.

Analysis 3.5. Comparison 3 FNB versus local infiltration analgesia, Outcome 5 Nausea and/or vomiting.

Comparison 4. Continuous FNB versus single-shot FNB

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Pain at rest first 2 hours	2	156	Std. Mean Difference (IV, Random, 95% CI)	0.15 [-0.17, 0.46]
2 Pain at rest 3 to 12 hours	2	156	Std. Mean Difference (IV, Random, 95% CI)	-0.03 [-0.35, 0.28]
3 Pain at rest 24 hours	4	272	Std. Mean Difference (IV, Random, 95% CI)	-0.62 [-1.17, -0.07]
4 Pain at rest 48 hours	4	272	Std. Mean Difference (IV, Random, 95% CI)	-0.96 [-1.69, -0.22]
5 Pain on movement 24 hours	4	272	Std. Mean Difference (IV, Random, 95% CI)	-0.42 [-0.67, -0.17]
6 Pain on movement 48 hours	4	272	Std. Mean Difference (IV, Random, 95% CI)	-0.54 [-1.02, -0.06]
7 Pain on rest 24 hours—subgrouped by continuous FNB with or without concurrent PCA	4		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
7.1 Continuous FNB with concurrent PCA	3	138	Std. Mean Difference (IV, Random, 95% CI)	-0.86 [-1.35, -0.38]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
7.2 Continuous FNB without concurrent PCA	1	134	Std. Mean Difference (IV, Random, 95% CI)	-0.16 [-0.50, 0.18]
8 Pain on movement 24 hours—subgrouped by continuous FNB with or without concurrent PCA	4		Std. Mean Difference (IV, Random, 95% CI)	Subtotals only
8.1 Continuous FNB with concurrent PCA	3	138	Std. Mean Difference (IV, Random, 95% CI)	-0.59 [-0.96, -0.22]
8.2 Continuous FNB without concurrent PCA	1	134	Std. Mean Difference (IV, Random, 95% CI)	-0.27 [-0.61, 0.07]
9 Pain at rest 24 hours—sensitivity analysis by low bias for allocation concealment	3	250	Std. Mean Difference (IV, Random, 95% CI)	-0.73 [-1.41, -0.04]
10 Pain on movement 24 hours—sensitivity analysis by low bias for allocation concealment	3	250	Std. Mean Difference (IV, Random, 95% CI)	-0.47 [-0.79, -0.15]
11 Opioid consumption 24 hours (mg)	3	236	Mean Difference (IV, Random, 95% CI)	-13.81 [-23.27, -4.35]
12 Opioid consumption 48 hours (mg)	4	269	Mean Difference (IV, Random, 95% CI)	-14.59 [-22.35, -6.82]

Analysis 4.1. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 1 Pain at rest first 2 hours.

Analysis 4.2. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 2 Pain at rest 3 to 12 hours.

Study or subgroup	Continuous FNB		Single-shot FNB		Std. Mean Difference Random, 95% CI	Weight	Std. Mean Difference Random, 95% CI
	N	Mean(SD)	N	Mean(SD)			
Heterogeneity: Tau²=0; Chi²=0.2, df=1(P=0.65); I²=0%							
Test for overall effect: Z=0.2(P=0.84)							
Favours Continuous					-4	-2	0
					2	4	Favours Single-shot

Analysis 4.3. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 3 Pain at rest 24 hours.

Study or subgroup	Continuous FNB		Single-shot FNB		Std. Mean Difference Random, 95% CI	Weight	Std. Mean Difference Random, 95% CI
	N	Mean(SD)	N	Mean(SD)			
Chan 2013	65	0.6 (1.2)	69	0.8 (1.5)		31.32%	-0.16[-0.5,0.18]
Hirst 1996	11	1.9 (2.7)	11	2.4 (1.3)		19.49%	-0.23[-1.07,0.61]
Park 2010	60	2.3 (1.4)	20	3.9 (1.6)		26.63%	-1.09[-1.63,-0.56]
Salinas 2006	18	1.7 (1.4)	18	3.3 (1.6)		22.56%	-1.04[-1.74,-0.34]
Total ***	154		118			100%	-0.62[-1.17,-0.07]
Heterogeneity: $\tau^2=0.22$; $\chi^2=11.24$, $df=3$ ($P=0.01$); $I^2=73.31\%$ Test for overall effect: $Z=2.21$ ($P=0.03$)							
Favours Continuous							Favours Single-shot

Analysis 4.4. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 4 Pain at rest 48 hours.

Study or subgroup	Continuous FNB		Single-shot FNB		Std. Mean Difference Random, 95% CI	Weight	Std. Mean Difference Random, 95% CI
	N	Mean(SD)	N	Mean(SD)			
Chan 2013	65	0.4 (0.9)	69	0.7 (1.1)		29.08%	-0.27[-0.61,0.07]
Hirst 1996	11	1.4 (2.3)	11	2.8 (2.3)		21.75%	-0.58[-1.44,0.28]
Park 2010	60	2.4 (1.3)	20	4.2 (1.7)		26.5%	-1.24[-1.78,-0.7]
Salinas 2006	18	0.9 (1.3)	18	3.2 (1.1)		22.66%	-1.87[-2.66,-1.07]
Total ***	154		118			100%	-0.96[-1.69,-0.22]
Heterogeneity: $\tau^2=0.45$; $\chi^2=18.34$, $df=3$ ($P=0$); $I^2=83.64\%$ Test for overall effect: $Z=2.56$ ($P=0.01$)							
Favours Continuous							Favours Single-shot

Analysis 4.5. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 5 Pain on movement 24 hours.

Study or subgroup	Continuous FNB		Single-shot FNB		Std. Mean Difference Random, 95% CI	Weight	Std. Mean Difference Random, 95% CI
	N	Mean(SD)	N	Mean(SD)			
Chan 2013	65	4 (1.7)	69	4.5 (1.8)		54.1%	-0.27[-0.61,0.07]
Hirst 1996	11	6.1 (3.3)	11	6.8 (2.3)		8.9%	-0.24[-1.07,0.6]
Park 2010	60	6 (2.3)	20	7.2 (1.5)		23.72%	-0.57[-1.08,-0.05]
Salinas 2006	18	4.7 (1.7)	18	6.3 (1.9)		13.28%	-0.87[-1.55,-0.18]
Total ***	154		118			100%	-0.42[-0.67,-0.17]
Heterogeneity: $\tau^2=0$; $\chi^2=2.86$, $df=3$ ($P=0.41$); $I^2=0\%$ Test for overall effect: $Z=3.27$ ($P=0$)							
Favours Continuous							Favours Single-shot

Analysis 4.6. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 6 Pain on movement 48 hours.

Analysis 4.7. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 7 Pain on rest 24 hours—subgrouped by continuous FNB with or without concurrent PCA.

Analysis 4.8. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 8 Pain on movement 24 hours—subgrouped by continuous FNB with or without concurrent PCA.

Analysis 4.9. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 9 Pain at rest 24 hours—sensitivity analysis by low bias for allocation concealment.

Analysis 4.10. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 10 Pain on movement 24 hours—sensitivity analysis by low bias for allocation concealment.

Analysis 4.11. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 11 Opioid consumption 24 hours (mg).

Analysis 4.12. Comparison 4 Continuous FNB versus single-shot FNB, Outcome 12 Opioid consumption 48 hours (mg).

APPENDICES

Appendix 1. Search strategy for CENTRAL, part of *The Cochrane Library*

- #1 MeSH descriptor Arthroplasty, Replacement, Knee explode all trees
- #2 MeSH descriptor Arthroplasty, Replacement explode all trees
- #3 MeSH descriptor Arthroplasty explode all trees
- #4 arthroplast* or (knee near (replacement or surg*))
- #5 (#1 OR #2 OR #3 OR #4)
- #6 MeSH descriptor Nerve Block explode all trees
- #7 nerve block* or FNB or ((femoral or psoas or (lumbar plexus) or (fascia iliaca)) near block*)
- #8 (#6 OR #7)
- #9 (#5 AND #8)

Appendix 2. Search strategy for MEDLINE (Ovid SP)

1. exp Arthroplasty, Replacement, Knee/ or exp Arthroplasty, Replacement/ or exp Arthroplasty/ or arthroplast*.af. or (knee adj3 (replacement or surg*)).af.
2. nerve block*.af. or ((femoral or psoas or lumbar plexus or fascia iliaca) adj3 block*).mp. or FNB.mp. or exp Nerve block/
3. 1 and 2
4. ((randomized controlled trial or controlled clinical trial).pt. or randomized.ab. or placebo.ab. or drug therapy.fs. or randomly.ab. or trial.ab. or groups.ab.) not (animals not (humans and animals)).sh.
5. 3 and 4

Appendix 3. Search strategy for EMBASE (Ovid SP)

1. exp knee arthroplasty/ or exp arthroplasty/ or arthroplast*.af. or (knee adj3 (replacement or surg*)).mp.
2. nerve block*.af. or ((femoral or psoas or lumbar plexus or fascia iliaca) adj3 block*).mp. or FNB.mp. or exp nerve block/
3. 1 and 2

4. (randomized-controlled-trial/ or randomization/ or controlled-study/ or multicenter-study/ or phase-3-clinical-trial/ or phase-4-clinical-trial/ or double-blind-procedure/ or single-blind-procedure/ or (random* or cross?over* or multicenter* or factorial* or placebo* or volunteer*).mp. or ((singl* or doubl* or trebl* or tripl*) adj3 (blind* or mask*)).ti,ab. or (latin adj square).mp.) not (animals not (humans and animals)).sh.

5. 3 and 4

Appendix 4. Search strategy for CINAHL (EBSCO host)

((MH "Arthroplasty, Replacement, Knee+") OR (MH "Arthroplasty, Replacement") OR (MH "Arthroplasty")) or arthroplast* or (knee and (replacement or surg*)) AND ((MM "Nerve Block") or (nerve block* or FNB) or ((femoral or psoas or (lumbar plexus) or (fascia iliaca)) and block*))

Appendix 5. Search strategy for Web of Science

#1 TS= arthroplast* or TS=(knee SAME (replacement or surg*))

#2 TS= (nerve block* or FNB) or TS=((femoral or psoas or (lumbar plexus) or (fascia iliaca)) SAMEblock*)

#3 (#1 and #2)

#4 TS=random* or TS=((clinical or controlled) SAME trial*)

#5 (#4 and #3)

Appendix 6. Study selection form

Title of article				
Study ID (surname of first author and year published):				
Report ID (if different from Study ID) (e.g. duplicate publications, follow-up studies)				
Publication type (e.g. full report, abstract)				
Study author contact details				
Date form completed (dd/mm/yy)				
Name of person extracting data				
Eligibility criteria				
Type of study	Participants	Interventions	Comparisons	Outcomes
RCT	Total knee replacement surgery	FNB (single-shot or continuous) +/- sciatic/obturator block	<ul style="list-style-type: none"> PCA opioids Epidural analgesia Infiltration +/- intra-articular injection Oral analgesia Placebo 	Circle: Pain intensity, neurological injury, proportion of participants in pain postoperatively, time to first rescue analgesia, opioid consumption, supplemental postoperative analgesic requirements, knee range of motion, early mobilization, adverse effects, adverse events (postoperative falls, technical failures of block, reintervention), time to achieve discharge criteria and participant satisfaction with analgesia
Criteria met?	Criteria met?	Criteria met?	Criteria met?	Other outcome/s?
	Yes/No/Unclear	Yes/No/Unclear		

(Continued)

Yes/No/Un-
clear

Yes/No/Un-
clear

Include: Yes/No/Unclear

If unclear, comment:

Reason for exclusion:
Notes:

Appendix 7. Data extraction form

Study ID (surname of first author and year published)

Study characteristics
Descriptions (include comparative information for each intervention or comparison group if available)

Further details
Source (pg and/ fig/table/other)
Single centre/Multi-centre
Start and end dates of data collection
Duration of participation (from recruitment to last follow-up)

Power (e.g. power and sample size calculation, level of power achieved)

Notes:
Participant characteristics (record all reported measures)

Population description (from which study participants were drawn)

Setting (include location)

Inclusion criteria
Exclusion criteria
Method of recruitment (e.g. phone, mail, clinic patients)

Total no. randomly assigned
Baseline imbalances
Withdrawals and exclusions
Age

(Continued)

Sex
BMI
No. of participants in each group

Intervention 1: _____;

Intervention 2: _____

Control 1: _____; Con-

trol 2: _____

No. of participants who received intended treatment

Intervention 1: _____;

Intervention 2: _____

Control 1: _____; Con-

trol 2: _____

No. of participants who were analysed

Intervention 1: _____;

Intervention 2: _____

Control 1: _____; Con-

trol 2: _____

Notes:
Interventions
Intervention group 1
FNB technique and type (i.e. continuous or single-shot FNB)

Additional blocks (i.e. sciatic or obturator)

Description of FNB (include type of drug, dosage and regimen used)

Duration of treatment period
PCA opioid in FNB group?

Yes/No. If yes, describe

Type of co-analgesia/Supplemental analgesia
Loss to follow-up
Others
Intervention group 2
FNB technique and type (i.e. continuous or single-shot FNB)

Additional blocks (i.e. sciatic or obturator)

Description of FNB (include type of drug, dosage and regimen used)

Duration of treatment period

(Continued)

PCA opioid in FNB group?	Yes/No. If yes, describe
Type of co-analgesia/Supplemental analgesia	
Loss to follow-up	
Others	
Control group 1	
Technique and type (i.e. PCA, epidural, infiltration +/- intra-articular injection, oral or placebo)	
Drugs, dosage and regimen used	
Placebo block in PCA group?	Yes/No. If yes, describe
Type of co-analgesia/Supplemental analgesia	
Loss to follow-up	
Others	
Control group 2	
Technique and type (i.e. PCA, epidural, infiltration +/- intra-articular injection, oral or placebo)	
Drugs, dosage and regimen used	
Placebo block in PCA group?	Yes/No. If yes, describe
Type of co-analgesia/Supplemental analgesia	
Loss to follow-up	
Others	

*NA: not appropriate; NR: not required.

Outcomes	Description (include outcome definition)	Time points reported	Person measuring/re-reporting	Imputation of missing data	Further details
Pain intensity					

(Continued)

(describe scale used, scale upper and lower limits and if high or low score is good, and whether scale is validated)

Proportion of participants in pain

Time to first rescue analgesia

Total opioid consumption

Adjunct oral analgesia

Knee range of motion

Early mobilization

Time to achieve discharge criteria

Adverse effects (nausea, vomiting, urinary retention, etc)

Technical failures of the block

Neurological injury

Mortality

Postoperative falls

Reintervention

Participant satisfaction with analgesia

(include scale used, scale upper and lower limits and if high or low score is good, and whether scale is validated)

Others:

Others:

Any outcomes collected but not reported? Yes/No

If yes, what outcome/s:

For continuous data (with a separate copy for each relevant intervention/control)									
Outcome	Unit of measurement/Time point	Intervention group		Control group		No. of missing participants in each group. Reasons missing?	No. of participants moved from other group. Reasons moved?	Details if outcome only described in text, or other summary statistics (SE, 95% CI)/If required statistics are not reported	Source (pg and/fig/table/other)
		n	Mean (SD) (specify if other variance)	n	Mean (SD) (specify if other variance)				
Pain intensity on movement (first 2 hours)									
Pain intensity on movement (> 2 to 12 hours)									
Pain intensity on movement (> 12 to 24 hours)									
Pain intensity on movement (> 24 to 72 hours)									
Pain intensity on movement (> 72 hours)									
Pain intensity at rest (first 2 hours)									
Pain intensity at rest (> 2 to 12 hours)									
Pain intensity at rest (> 12 to 24 hours)									
Pain intensity at rest (> 24 to 72 hours)									
Pain intensity at rest (> 72 hours)									
Time to first rescue analgesia									
Total opioid consumption

(Continued)

Supplemental postoperative analgesic requirements

Knee range of motion

Early mobilization

Time to achieve discharge criteria

Participant satisfaction with analgesia

Others

For dichotomous data (with a separate copy for each relevant intervention/control)

Outcomes	Intervention group (n/N) where n = no. of participants with outcome, N = no. of participants randomly assigned	Control group (n/N) where n = no. of participants with outcome, N = no. of participants randomly assigned	No. of missing participants in each group. Reasons missing?	No. of participants moved from other group. Reasons moved?	Location in text or source (pg and/fig/table/other)
Proportion of participants in pain					
Nausea and/or vomiting					
Sedation					
Urinary retention					
Technical failure of blocks					
Neurological injury					
Mortality					
Postoperative falls					
Reintervention					
Others:					

Other information relevant to the results

Indicate whether any data were obtained from the primary author and whether results were estimated from graphs etc or were calculated by you using a formula (the formula should be given). Provide information here if results not reported in the article are obtained.

Key conclusions of study authors:
Study funding sources:

Correspondence required for further study information (from whom, what and when)

References to other RCTs:

References to **published reports** of potentially eligible RCTs not already identified for this review? If yes, give details.

First author	Journal/Conference	Year of publication
--------------	--------------------	---------------------

References to **unpublished data** from potentially eligible RCTs not already identified for this review? If yes, give details.

Appendix 8. Quality assessment of eligible trials form

Domain	Risk of bias (Low/High/ Unclear)	Support for judgement (Quotes/ Comments)	Location in text or source (pg and/fig/ta- ble/ other)
Random sequence generation			
Allocation concealment			
Blinding of participants and personnel			
Blinding of outcome assessment			
Incomplete outcome data addressed (short-term outcomes—during hospital admission)			
Incomplete outcome data addressed (long-term outcomes—after discharge)			
Selective outcome reporting			
Intention-to-treat			
Other bias			
Notes:			

Were withdrawals described? Yes/No/Unclear

Comment if any:

Appendix 9. Summary of findings

Outcomes	Illustration comparative risks (95% CI)		Relative effect (95% CI)	No. of participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
Pain at rest at 24 hours						
Pain on movement at 24 hours						
Neurological injury						
Total opioid consumption						
Nausea and/or vomiting						
Knee range of motion						
Participant satisfaction with analgesia						

Appendix 10. Fail-safe-N analysis

Comparison	Outcome	Subgroup	Effect size (SMD)	No. of studies (N)	Fail-safe-n
1. FNB vs PCA opioid	Analysis 1.1 —Pain at rest at 24 hours	Overall	-0.72	19	60
1. FNB vs PCA opioid	Analysis 1.2 —Pain at rest at 24 hours by allocation concealment	Low risk of bias RCTs	-0.54	9	18
1. FNB vs PCA Opioid	Analysis 1.5 —Pain on movement at 24 hours	Overall	-0.94	17	78
1. FNB vs PCA opioid	Analysis 1.6 —Pain on movement at 24 hours by allocation concealment	Low risk of bias RCTs	-0.54	9	18
2. FNB vs epidural	Analysis 2.1 —Pain at Rrest at 24 hours	< not significant, not applicable >			
2. FNB vs epidural	Analysis 2.5 —Pain on movement at 24 hours	< not significant, not applicable >			
3. FNB vs local infiltration	Analysis 2.1 —Pain at rest at 24 hours	< not significant, not applicable >			
3. FNB vs local infiltration	Analysis 2.5 —Pain on movement at 24 hours	< not significant, not applicable >			

(Continued)

4. Continuous vs single-shot FNB	Analysis 4.1 —Pain at rest at 24 hours	Overall	-0.62	4	9
4. Continuous vs single-shot FNB	Analysis 4.3 —Pain on movement at 24 hours	Overall	-0.42	4	5

WHAT'S NEW

Date	Event	Description
6 January 2016	Amended	Typo corrected in main objectives section (an additional full stop was removed).

CONTRIBUTIONS OF AUTHORS

Ee-Yuee Chan (EC), Marlene Fransen (MF), David A Parker (DA), Pryseley N Assam (PA), Nelson Chua (NC)

Conceiving of the review: EC

Co-ordinating the review: EC

Undertaking manual searches: EC

Screening search results: EC, MF

Organizing retrieval of papers: EC

Screening retrieved papers against inclusion criteria: EC, MF, NC

Appraising quality of papers: EC, MF

Abstracting data from papers: EC, MF, NC

Writing to authors of papers to ask for additional information: EC

Providing additional data about papers: EC

Obtaining and screening data on unpublished studies: EC, MF

Managing data for the review: EC

Entering data into Review Manager ([RevMan 5.2](#)): EC

Handling RevMan statistical data: EC, PA

Performing other statistical analysis not using RevMan: EC, PA

Interpreting data: EC, MF, PA, NC, DP

Making statistical inferences: EC, MF, PA, NC, DP

Writing the review: EC with support from the other review authors

Performing previous work that was the foundation of the present study: EC

Serving as guarantor for the review (one review author): EC

Taking responsibility for reading and checking the review before submission: EC

DECLARATIONS OF INTEREST

Ee-Yuee Chan, Marlene Fransen and Nelson Chua were the study authors of an RCT ([Chan 2013](#)) that was eligible for inclusion in this Cochrane review.

Pryseley N Assam is a co-author of a possible publication arising from the RCT authored by Chan et al ([Chan 2013](#)).

David A Parker co-authored an RCT ([Widmer 2012](#)) that was also included in this review.

SOURCES OF SUPPORT

Internal sources

- Ee-Yuee Chan was supported by an International Postgraduate Research Scholarship, University of Sydney, Australia.

External sources

- No sources of support supplied

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

We made the following changes to the published protocol ([Chan 2012](#)).

1. Types of interventions.

We added a new comparison called continuous versus single-shot FNB for a more comprehensive review on FNB.

2. Primary outcomes: pain at rest and on movement.

We have reclassified pain at rest and on movement for the postoperative time frame '> 24 to 72' to two categories: '48 hours' and '72 hours.'

3. Data synthesis.

For better clarity, our text in the 'Data synthesis' section was changed as follows.

The primary analyses include:

- single-shot or continuous FNB ± sciatic/obturator block versus PCA opioid;
- single-shot or continuous FNB ± sciatic/obturator block versus epidural analgesia;
- single-shot or continuous FNB versus local infiltration analgesia;
- single-shot or continuous FNB versus oral analgesia; and
- continuous versus single-shot FNB.

4. Unit of analysis issues.

A total of 20 RCTs were included with three intervention arms: multiple independent comparisons (e.g. FNB vs PCA and FNB vs epidural); multiple correlated comparisons with similar types of FNB (e.g. continuous FNB with ropivacaine vs PCA and continuous FNB with bupivacaine vs PCA) or multiple correlated comparisons with different types of FNB (e.g. single-shot FNB vs PCA and continuous FNB vs PCA). We have reworded our text in [Unit of analysis issues](#) to clarify.

5. Dealing with missing data.

Some of the included RCTs presented results using summary statistics other than mean (SD). In the review, we added details on how we handled these missing data.

6. Subgroup analysis.

Because of limited data, subgroup analysis was restricted to:

- type of FNB (single-shot or continuous FNB, with or without an additional sciatic/ obturator nerve block);
- FNB with and without a concurrent parenteral opioid; and
- type of FNB local anaesthetics (i.e. ropivacaine or bupivacaine).

7. Sensitivity analysis.

Sensitivity analysis based on multiple interventions within studies was not performed because we followed the recommendations in the *Cochrane Handbook for Systematic Reviews of Interventions* (see Chapter 16; Higgins 2011), as discussed in the [Unit of analysis issues](#) section.

8. SoF tables.

For the SoF tables, we have reduced the number of outcomes to seven: pain at rest at 24 hours, pain on movement at 24 hours, neurological injury, opioid consumption, nausea and vomiting, knee range of motion and participant satisfaction with analgesia, by excluding 'time to first rescue analgesia' and 'early mobilisation'.

9. We changed the term 'surgical infiltration with/without intra-articular injection' to 'local infiltration analgesia,' as the latter appears to be commonly used in recent literature.

10. We defined 'early ambulation' as 'time to first ambulation.'

INDEX TERMS

Medical Subject Headings (MeSH)

*Arthroplasty, Replacement, Knee; *Femoral Nerve; Analgesia [methods]; Analgesia, Epidural; Analgesia, Patient-Controlled; Analgesics, Opioid [therapeutic use]; Nerve Block [*methods]; Pain, Postoperative [*drug therapy]; Randomized Controlled Trials as Topic

MeSH check words

Humans