Special Article

Cite this article: Nevarez-Flores AG, Sanderson K, Breslin M, Carr VJ, Morgan VA, Neil AL (2019). Systematic review of global functioning and quality of life in people with psychotic disorders. *Epidemiology and Psychiatric Sciences* **28**, 31–44. <https://doi.org/10.1017/S2045796018000549>

Received: 29 May 2018

Revised: 24 August 2018

Accepted: 31 August 2018

First published online: 1 October 2018

Key words:

Functioning; psychosis; quality of life; Schizophrenia

Author for correspondence:

Dr Amanda L. Neil, E-mail: Amanda.Neil@utas.edu.au

Systematic review of global functioning and quality of life in people with psychotic disorders

A. G. Nevarez-Flores¹, K. Sanderson^{2,1}, M. Breslin¹, V. J. Carr^{3,4}, V. A. Morgan^{5,1} and A. L. Neil¹

¹Menzies Institute for Medical Research, The University of Tasmania, 17 Liverpool St, Hobart Tasmania 7000, Australia; ²School of Health Sciences, University of East Anglia, Norwich Research Park Norwich, NR4 7TJ, UK; ³Research Unit for Schizophrenia Epidemiology, School of Psychiatry, University of New South Wales, St Vincent's Hospital, Sydney, New South Wales, Australia; ⁴Department of Psychiatry, School of Clinical Sciences, Monash Medical Centre, Monash University, Clayton Melbourne, Victoria, Australia and ⁵Neuropsychiatric Epidemiology Research Unit, Division of Psychiatry, Medical School, The University of Western Australia, Crawley, Western Australia, Australia

Abstract

Aims. People with psychotic disorders face impairments in their global functioning and their quality of life (QoL). The relationship between the two outcomes has not been systematically investigated. Through a systematic review, we aim to explore the presence and extent of associations between global functioning and QoL and establish whether associations depend on the instruments employed.

Methods. In May 2016, ten electronic databases were searched using a two-phase process to identify articles in which associations between global functioning and QoL were assessed. Basic descriptive data and correlation coefficients between global functioning and QoL instruments were extracted, with the strength of the correlation assessed according to the specifications of Cohen 1988. Results were reported with reference to the Meta-analysis of Observational Studies in Epidemiology guidelines and PRISMA standards. A narrative synthesis was performed due to heterogeneity in methodological approaches.

Results. Of an initial 15 183 non-duplicate articles identified, 756 were deemed potentially relevant, with 40 studies encompassing 42 articles included. Fourteen instruments for measuring global functioning and 22 instruments for measuring QoL were used. Twenty-nine articles reported linear associations while 19 assessed QoL predictors. Correlations between overall scores varied in strength, primarily dependent on the QoL instrument employed, and whether QoL was objectively or subjectively assessed. Correlations observed for objective QoL measures were consistently larger than those observed for subjective measures, as were correlations for an interviewer than self-assessed QoL. When correlations were assessed by domains of QoL, the highest correlations were found for social domains of QoL, for which most correlations were moderate or higher. Global functioning consistently predicted overall QoL as did depressive and negative symptoms.

Conclusions. This review is the first to explore the extent of associations between global functioning and QoL in people with psychotic disorders. We consistently found a positive association between global functioning and QoL. The strength of the association was dependent on the QoL instrument employed. QoL domains strongly associated with global functioning were highlighted. The review illustrates the extensive array of instruments used for the assessment of QoL and to a lesser extent global functioning in people with psychotic disorders and provides a framework to understand the different findings reported in the literature. The findings can also inform the future choice of instruments by researchers and/or clinicians. The observed associations reassure that interventions for improving global functioning will have a positive impact on the QoL of people living with a psychotic disorder.

Introduction

Psychotic disorders are of special interest due to the severity of their symptoms, the surrounding stigma and the consequences of dysfunction, discrimination and costs. The importance of functioning to psychotic disorders was reaffirmed when psychosis was described as an imprecise group of symptoms, of sufficient severity to disrupt everyday functioning (Petho and Ban, 1988).

According to the International Classification of Functioning, Disability and Health (World Health Organization, 2001), functioning denotes the positive features of the relationship between a health condition and the environmental and personal context of the individual,

Fig. 1. Timeline of functioning and quality of life instruments development.

while disability indicates negative features of that relationship. Thus, functioning is concerned with the ability of an individual to perform their roles and participate in life (Bowling, 2005). Global functioning should encompass the measurement of several types of functioning (Aas, 2010). Since 1962, a number of instruments have been created to measure global functioning, as well as specific dimensions (or types) of functioning e.g., social functioning, executive functioning, etc. (see Fig. 1).

Almost parallel to the development of instruments to assess functioning, and shortly after quality of life (QoL) in health care was raised by Elkinton, when he asked: 'What is the harmony within a man, and between a man and his world –the quality of life– to which the patient, the physician, and society aspires?' (Elkinton, 1966), interest in QoL as an outcome of people with psychosis began to emerge (Fig. 1). This interest occurred alongside the implementation of community support programmes after deinstitutionalisation (Baker and Intagliata, 1982; Lehman, 1988).

QoL has been defined as 'an individual's perception of their position in life in the context of the culture and value systems in which they live and in relation to their goals, expectations, standards and concerns' (World Health Organization, 1997). Despite this, there is no agreement on what aspects and how QoL should be assessed. More than 50 QoL instruments have been used in patients with mental disorders (Prigent *et al.*, 2014).

Functioning and more recently QoL have thus been identified as important outcomes in people with psychosis. Reviews focused on the assessment of global functioning and QoL in people with psychotic disorders are scarce (Awad *et al.*, 1997; Pinikahana *et al.*, 2002) and the relationship between them has not been systematically assessed. As impairment in the global functioning of people with psychosis is expected, it is important to understand the impact of this impairment on QoL. Establishing the relationship between global functioning and QoL measures would support the future choice of instruments for the assessment of

these outcomes and, in turn, identify strategies to diminish the societal burden of psychotic disorders.

The aim of the present study is to explore the presence and extent of associations between global functioning and QoL in people with psychotic disorders and establish whether this relationship is dependent on the instruments employed.

Methods

Selection criteria

This systematic review was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) (Moher *et al.*, 2009). Studies included for data extraction were full-text articles with a cross-sectional design or a follow-up design that provided required information at baseline. Baseline information only was sought, as changes in the functioning of people with psychosis over time is well documented (Harvey and Davidson, 2002; Ascher-Svanum *et al.*, 2013; Harvey, 2014). Articles could be written in English or Spanish (given native English and Spanish speakers within the authorship team). The sample needed to comprise people with schizophrenia and schizophrenia spectrum disorders with or without people with mood disorders with psychotic symptoms (bipolar disorder type I, major or severe depressive disorder with psychotic symptoms) assessed according to the Ninth/Tenth Revision of the International Classification of Diseases (ICD 9/10) or the Diagnostic and Statistical Manual of Mental Disorders Fourth/Fifth Edition (DSM-IV/5), and aged between 18 and 64 years, in which associations between global functioning and QoL were assessed. There were no time-period restrictions.

Due to the lack of a universal definition of global functioning, instruments that assessed several dimensions of functioning as an inclusive outcome were deemed a global functioning instrument. Likewise, given the absence of a universally accepted definition of QoL, instruments reporting on a group of outcomes that contribute to an individual's satisfaction with life and/or overall health (Fayers and Machin, 2016) were accepted for inclusion in the review.

Intervention studies, reviews and meta-analyses were excluded as well as studies based on populations with organic or induced psychosis, psychosis due to other mental disorders or other medical conditions and populations at high risk of psychosis but not yet diagnosed. Intervention studies were excluded as functioning levels differ in artificial environments such as clinical trials (Patterson *et al.*, 2001; Bellack *et al.*, 2007; Ascher-Svanum *et al.*, 2013).

Search strategy

In May 2016 Annual Reviews, Cochrane Library, CINAHL, EconLit, Embase, Medline, PsycARTICLES, PsycINFO, PubMed and ScienceDirect were searched using a two-phase identification process. Search terms were introduced with corresponding MeSH Terms, synonyms and stem words, as well as appropriate filters and use of Boolean operators. The detailed search strategy is available as Supplementary material.

Searches were divided into phases in order to identify potential publication bias. This approach was adopted because of concern that only strong associations would be reported in title and abstract. In Phase A, all keyword terms were searched in title and abstract. In Phase B, all keyword terms except functioning

were searched in title and abstract, with stem 'function' then searched for in the main text.

Study selection

Two independent reviewers (ANF and AN) assessed studies for inclusion, with inconsistencies and disagreements resolved by consensus. After deletion of duplicates, Phase A records were screened for inclusion criteria in their titles and abstracts. Full-text articles of included studies were obtained and assessed in full for eligibility. A similar process was used for Phase B records, except that screening for inclusion included a search for functioning in the main text. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement (von Elm *et al.*, 2007) was used to assess whether analytical observational studies reported STROBE elements determined *a priori* as essential. These items were 3, 7, 12, 16 and 18 (i.e., objectives, variables, statistical methods, main and key results).

Data extraction

Data extracted from each article encompassed basic descriptive data and correlation coefficients between global functioning and QoL instruments. Extraction was undertaken by ANF under the guidance of AN. After extraction, socio-demographic variables were coded for summarisation in frequency tables.

The correlation coefficients between global functioning and QoL extracted were: correlations between global functioning and other variables; and correlations between QoL and other variables comprising the domains and items covered for each instrument. This analysis reports on the strength of the correlation between global functioning and QoL and between global functioning and individual domains of QoL. Strength of correlation was assessed according to the specifications of Cohen (1988) established as $0.10 \leq r < 0.3$ small effect, $0.3 \leq r < 0.5$ medium effect and $0.50 \leq r$ large effect.

A narrative synthesis was performed given the heterogeneity in methodological approaches, including instruments employed in the assessment of global functioning and QoL, and statistical analyses employed. QoL instruments were defined as an objective if comprised objective items only (usually intended for interviewer-assessment), subjective if comprised subjective items only (usually intended for self-assessment), or subjective and objective. Results were reported with reference to the Meta-analysis of Observational Studies in Epidemiology guidelines (Stroup *et al.*, 2000) and PRISMA standards.

Results

Search results

Across Phases A and B 15 183 records were initially identified, of which 8673 were duplicates and excluded from further review. In Phase A, upon screening of title and abstract, 637 articles were then assessed as requiring full-text review for eligibility, with 34 articles assessed as eligible. In Phase B upon screening of title and abstract, 2601 articles were assessed as requiring full-text screening for functioning, with 119 then identified as requiring full-text review for eligibility. Eight additional articles were identified as eligible for inclusion (see Fig. 2). Thus 42 articles were included in this review as listed in Table 1.

Fig. 2. PRISMA flow diagram of Phase A and Phase B search.

The articles

Articles were published between 1997 and 2016, all met the required STROBE criteria for inclusion. Three-fifths (25 articles) were from Europe. Four articles related to the European Psychiatric Services: Inputs Linked to Outcome Domains and Needs (EPSILON) study: Gaite *et al.* (2002) and Becker *et al.* (2005) reported on the entire sample while Meijer *et al.* (2002) reported on the Amsterdam participants at baseline, and at 18 months follow up (2009), as part of a broader Netherlands' study. Each of these articles undertook different analyses of the data and as there was no pooling of results, the results for each are reported to maximise the comprehensiveness of the narrative synthesis. Four more articles (Holloway and Carson, 1999; Kusel *et al.*, 2007; Brissos *et al.*, 2011; Hunter and Barry, 2012) were published as part of larger studies. The systematic review thus comprised articles from 40 independent studies.

Data extracted overview

The median number of participants across all studies was 135, with the range 36–971. Schizophrenia was the single diagnosis in 26 articles (62%), only four articles (Greenley *et al.*, 1997; Holloway and Carson, 1999; Lasebikan and Owoaje, 2015; Stubbs *et al.*, 2015) (10%) included participants with bipolar disorder or depressive disorder with psychotic symptoms. Over half of the articles (57%) utilised one instrument for measuring functioning and one for measuring QoL. Six articles, employed two functioning instruments, 11 articles two QoL instruments and one article (Reine *et al.*, 2005) three QoL instruments. Fujino

et al. (2016) utilised two instruments, which together assessed global functioning.

Within the included studies, we identified 14 instruments for measuring global functioning and 22 for QoL. The Global Assessment of Functioning (GAF) was the most utilised functioning instrument (29 articles) and the WHO Quality of Life Instruments (WHOQOL-BREF) together with its Portuguese version the most utilised QoL instrument (12 articles) (see Supplementary material).

Of the 42 articles, 29 (69%) reported linear associations between global functioning and QoL overall scores and/or domain scores and presented correlation coefficients, four articles (10%) assessed associations between global functioning and QoL using alternate statistical methods; and 19 articles (45%) assessed predictors of QoL (Table 2). Outcomes for each are examined below.

Linear associations between global functioning and QoL (overall scores)

Twenty articles (48%) provided correlations between overall scores of global functioning and QoL instruments. Two of these articles (Mas-Exposito *et al.*, 2011; Bai *et al.*, 2014) also provided correlations between global functioning with QoL domains and among global functioning and QoL domains (Table 2).

Correlations between overall scores ranged in strength from strong to weak with nearly half of the correlations reported as moderate ($r = 0.34$ to 0.49). Most moderate correlations involved the GAF which is consistent with the frequent use of the instrument. The two largest correlations assessed QoL using the QoL Scale (QLS), the largest assessed functioning with the Personal and Social Performance (PSP) ($r = 0.84$), the second largest with the GAF ($r = 0.83$). The smallest correlation reported as significant were for the GAF and the QoL Questionnaire (QLQ) and the GAF and Manchester Short Assessment of QoL (MANSA) ($r = 0.16$ for each) (Fig. 3).

Amongst the QoL instruments, nearly one-third of the correlations were assessed in relation to the QLS primarily the complete 21-item version, with the 7-item and 5-item versions also assessed in Ritsner *et al.* (2005). Half the correlations were large, including that for the 7-item version. One-quarter of correlations were assessed in relation to the self-assessed WHOQOL-BREF. Correlations were reported for four domain scores and two separate scored items that assessed the individual's overall perception of QoL and health (World Health Organization, 1998; The WHOQOL Group, 1998) in Galuppi *et al.* (2010), and just for overall QoL in three articles (Chino *et al.*, 2009; Bai *et al.*, 2014; Ito *et al.*, 2015). Some articles reported correlations for the four domains and a total score (Miclutia *et al.*, 2008; Mas-Exposito *et al.*, 2011; Bai *et al.*, 2014). Ito *et al.* (2015) reported non-significant associations for the overall score in insidious onset and acute onset ($r = -0.24$, -0.21), respectively. Correlations observed for objective QoL measures such as QLS were consistently larger ($r = 0.20$ to 0.84) than those observed for subjective measures such as WHOQOL-BREF ($r = -0.21$ to 0.58). Interviewer-assessment was also associated with larger correlations than self-assessment as reflected in Riedel *et al.* (2011).

Linear association between overall scores of global functioning and QoL domains

Thirteen articles (31%) provided correlations between global functioning and QoL domains, as listed in Table 3. Global

Table 1. Main characteristics of included studies

Study	Conducted	Country of publication	Participants			Instruments	
			N	Age range	Dx ^a (%)	Functioning	Quality of Life
Greenley <i>et al.</i> (1997)	NA	USA	971	NA	NS	GAF	QLQ
ELCCT: Holloway and Carson (1999)	NA	England	70	NA	NS	WHO/DAS	LQOLP; LEC
Norman <i>et al.</i> (2000)	1989–1993	Canada	128	17–57	100	LSP	QLS; GWB
EPSILON: Gaite <i>et al.</i> (2002) Becker <i>et al.</i> (2005)	1997–1998	Denmark; England; Italy; Spain; The Netherlands	404	18–65	NS	GAF	LQoLP-EU (European version)
Meijer <i>et al.</i> (2009, 2002)	1997–1998	The Netherlands ^b	143	18–65	100	GAF	SF-36; LQoLP Dutch version
Mubarak (2005)	NA	Malaysia	258	NA	100	WHO/DAS	QoLI
Reine <i>et al.</i> (2005)	2000	France	205	18–70	100	GAF	SF-36; QoLI brief version; S-QoL
Ritsner <i>et al.</i> (2005)	NA	Israel	133	18–60	100	GAF	QLS; Q-LES-Q
König <i>et al.</i> (2007)	2003–2004	Germany	166	21–80	72	GAF; SOFAS	EQ-5D; WHOQOL-BREF
UK-SCAP: Kusel <i>et al.</i> (2007)	1999–2000	UK	442	NA	NS	GAF	QLS; MANSA
Prince (2007)	1994–1996	USA	264	17–65	NS	GAS	QoLI
Miclutia <i>et al.</i> (2008)	NA	Romania	50	18–55	100	GAF	WHOQOL-BREF
Adewuya and Makanjuola (2009)	2006	Nigeria	99	NA	100	GAF	WHOQOL-BREF
Chino <i>et al.</i> (2009)	NA	Japan	36	NA	100	GAF	WHOQOL-BREF
Kuo <i>et al.</i> (2009)	NA	Taiwan	100	18–65	100	GAF	SQSL-R4; LQOLP Taiwanese
Galuppi <i>et al.</i> (2010)	2008	Italy	104	NA	100	FPS	WHOQOL-BREF
Woon <i>et al.</i> (2010)	NA	Singapore	83	NA	100	GAF	WHOQOL-BREF
PSP Portuguese validation study: Brissos <i>et al.</i> (2011)	2009–2010	Portugal	76	18–65	100	PSP Portuguese version	WHOQOL-BREF Portuguese version
Hosseini and Yousefi (2011)	1999–2000	Iran	100	21–60 ^c	100	GAF	QLS
Karadayi <i>et al.</i> (2011)	NA	Turkey	102	18–65	100	PSP	QLS
Mas-Exposito <i>et al.</i> (2011)	2006–2008	Spain	241	NA	100	GAF; WHO DAS-S	WHOQOL-BREF
Riedel <i>et al.</i> (2011)	2007	Germany	136	18–65	78	GAF	QLS; RSM-Scale
Roe <i>et al.</i> (2011)	2007–2008	Israel	159	19–66	NS	GAF	MANSA
Guilera <i>et al.</i> (2012)	2007–2009	Spain	352	18–55	88	WHODAS II; SOFAS	EQ-5D
EGOFORS: Hunter and Barry (2012)	NA	Belgium; France; Germany; Israel; Italy; Spain; Sweden; Turkey; UK	295	NA	100	GAF; PSP	QLS
Nafees <i>et al.</i> (2012)	NA	UK	73	18–65	100	PSP; GAF	QLS
Pitkanen <i>et al.</i> (2012)	2005–2006	Finland	311	18–65	41	GAF	EQ-5D; Q-LES-Q short form
Razali and Wahid (2012)	NA	Malaysia	206	18–60	100	PSP	QoLI brief version
Medeiros-Ferreira <i>et al.</i> (2013)	2008–2009	Spain	76	NA	NS	GAF	EQ-5D
Akinsulore <i>et al.</i> (2014)	2010	Nigeria	100	NA	100	GAF	WHOQOL-BREF

(Continued)

Table 1. (Continued.)

Study	Conducted	Country of publication	Participants			Instruments	
			N	Age range	Dx ^a (%)	Functioning	Quality of Life
Bai <i>et al.</i> (2014)	NA	Taiwan	108	20–60	100	PSP; SRG-PSP	WHOQOL-BREF
Rocca <i>et al.</i> (2014)	2009–2011	Italy	92	18–65	100	PSP	QLS
Zendjidian <i>et al.</i> (2014)	NA	France	91	NA	100	GAF	SF-36
Dima <i>et al.</i> (2015)	2009–2010	Romania	131	18–65	NS	GAF	SF-36; Q-LES-Q short form
Ito <i>et al.</i> (2015)	2008–2011	Japan	168	16–55	NS	GAF	WHOQOL-BREF
Lasebikan and Owoaje (2015)	2008	Nigeria	652	NA	56	GAF	WHOQOL-BREF
Stubbs <i>et al.</i> (2015)	2010–2012	UK	438	NA	NS	GAF	EQ-5D 3 levels
Alessandrini <i>et al.</i> (2016)	2010–2014	France	271	NA	100	FROGS	S-QoL 18
Fujino <i>et al.</i> (2016)	NA	Japan	93	NA	100	UPSA-B; SFS	JSQLS
Rocca <i>et al.</i> (2016)	2008–2011	Italy	323	18–65	100	GAF	QLS

EGOFORS, European Group on Functional Outcomes and Remission in Schizophrenia study; ELCCT, East Lambeth Continuing Care Team study; EPSILON, European Psychiatric Services: Inputs Linked to Outcome Domains and Needs study; EQ-5D, EuroQol five dimensions questionnaire; FPS, Personal and Social Functioning Scale; FROGS, Functional Remission Of General Schizophrenia; GAF, Global Assessment of Functioning; GAS, Global Assessment Scale; GWB, General Well-Being Scale; JSQLS, Schizophrenia Quality of Life Scale Japanese version; LEC, Life Experiences Checklist; LQOLP, Lancashire Quality of Life Profile; LSP, Life Skills Profile; MANSA, Manchester Short Assessment of Quality of Life; NS, not specified; PSP, Personal and Social Performance scale; Q-LES-Q, Quality of Life Enjoyment and Satisfaction Questionnaire; QLQ, Quality of Life Questionnaire; QLS, Quality of Life Scale; QoLI, Lehman's Quality of Life Interview; RSM-Scale, Riedel-Spellmann-Musil-Scale; SF-36, Medical Outcomes Study (MOS) 36-Item Short-Form Health Survey; SFS, Social Functioning Scale; SOFAS, Social and Occupational Functioning Assessment Scale; SQLS-R4, Schizophrenia Quality of Life Scale Revision 4 Chinese version; S-QoL, Schizophrenia Quality of Life; SRG-PSP, Self-reported version of the graphic PSP; UK-SCAP, United Kingdom Schizophrenia Care and Assessment Programme; UPSA-B, University of California, San Diego (UCSD) Performance-based Skills Assessment-Brief version; USA, United States of America; WHO DAS-S, World Health Organization Short Disability Assessment Schedule; WHO/DAS, World Health Organization Disability Assessment Schedule; WHOQOL-BREF, World Health Organization Quality of Life abbreviated version; NA, not available.

^aPercentage of schizophrenia diagnosis.

^bIncluding EPSILON's participants from Amsterdam and other participants from The Netherlands.

^cLower – Upper age of participants included in the study.

functioning was measured with GAF in over half of the articles (62%), and with the exception of Bai *et al.* (2014) which employed the self-reported version of the graphic PSP (SRG-PSP), the interviewer undertook all assessments of functioning. The most utilised QoL instrument was the WHOQOL-BREF, followed by the Medical Outcomes Study 36-Item Short-Form Health Survey (SF-36). Assessments of QoL were self-assessed, with four exceptions (Meijer *et al.*, 2002; Reine *et al.*, 2005; Riedel *et al.*, 2011; Rocca *et al.*, 2014).

In total, we recorded 100 correlation coefficients across 39 domains within included QoL instruments. The strength of correlations varied widely and were primarily weak (48%). Just over one-quarter of correlations were presented in a single article (Reine *et al.*, 2005), with the GAF compared with three different QoL instruments, the SF-36, Schizophrenia QoL (S-QoL) and Lehman's QoL Interview brief version (QoLI brief) (subjective items). All associations reported in this study were weak and in some instances lower than for other comparable assessments.

The largest correlation coefficient ($r = 0.72$) was between functioning assessed with GAF and 'subjective wellbeing' assessed by the interviewer with the Riedel-Spellmann-Musil-Scale (RSM-Scale) (Riedel *et al.*, 2011). The next four highest correlations were also assessed between these instruments ($r = 0.61$ to 0.65) (Riedel *et al.*, 2011). The smallest significant correlation ($r = 0.10$) was between functioning assessed with GAF and interviewer-assessed 'mental and physical health' of the QoLI brief, subjective items only (Reine *et al.*, 2005).

Most correlations between global functioning and QoL domains were positive. Exceptions were for Mas-Exposito *et al.*

(2011), which employed the WHO Short Disability Assessment Schedule (WHO DAS-S) for evaluating functioning, and Fujino *et al.* (2016) and Kuo *et al.* (2009), which employed the Schizophrenia QLS Japanese version (JSQLS) and Revision 4 Chinese version (SQLS-R4) to assess QoL respectively. For each of these three instruments lower scores signify better outcomes.

Correlations for the WHOQOL-BREF tended to be larger than those for the SF-36. The WHOQOL-BREF gave rise to correlations that were primarily moderate in strength for each of the four domains of the instrument, although large correlations were also reported for the physical (Galuppi *et al.*, 2010) and psychological domains (Miclutia *et al.*, 2008). In regard to the SF-36, of the 24 correlations all were small except for four correlations from two of three articles: one correlation for the physical domain (Dima *et al.*, 2015), one for the mental health domain (Meijer *et al.*, 2002) and two for the social domain (Meijer *et al.*, 2002; Dima *et al.*, 2015).

The three most frequently assessed domains of QoL were the physical, psychological and social components. The vast majority of correlations for the physical component were small (70%), half were small for the psychological component and 45% small for the social component. Around a third of these small correlations were assessed in Reine *et al.* (2005) which reported just over a quarter of all correlations. Over half (56%) of the correlations reported for occupational and environmental issues, as well as for components related to treatment and satisfaction with life, in general, were small.

When correlations were limited to broad domains (i.e., included several items), over half of the correlations were

Table 2. Summary of associations provided by article

Articles (total 42)	N	%	References
Correlations			
Overall global functioning– Overall quality of life (QoL)	20	48	Karadayi <i>et al.</i> (2011), Mas-Exposito <i>et al.</i> (2011), Riedel <i>et al.</i> (2011), Hosseini and Yousefi (2011), Roe <i>et al.</i> (2011), Hunter and Barry (2012), Nafees <i>et al.</i> (2012), Gaite <i>et al.</i> (2002), Reine <i>et al.</i> (2005), König <i>et al.</i> (2007), Kusel <i>et al.</i> (2007), Miclutia <i>et al.</i> (2008), Galuppi <i>et al.</i> (2010), Dima <i>et al.</i> (2015), Chino <i>et al.</i> (2009), Greenley <i>et al.</i> (1997), Ritsner <i>et al.</i> (2005), Norman <i>et al.</i> (2000), Bai <i>et al.</i> (2014), Ito <i>et al.</i> (2015).
Overall global functioning – Domains of QoL	13	31	Mas-Exposito <i>et al.</i> (2011), Riedel <i>et al.</i> (2011), Meijer <i>et al.</i> (2002), Reine <i>et al.</i> (2005), Miclutia <i>et al.</i> (2008), Galuppi <i>et al.</i> (2010), Rocca <i>et al.</i> (2014), Dima <i>et al.</i> (2015), Kuo <i>et al.</i> (2009), Woon <i>et al.</i> (2010), Akinsulore <i>et al.</i> (2014), Bai <i>et al.</i> (2014), Fujino <i>et al.</i> (2016).
Domains of global functioning – Overall QoL	5	12	Mas-Exposito <i>et al.</i> (2011), Guilera <i>et al.</i> (2012), Holloway and Carson (1999), Chino <i>et al.</i> (2009), Bai <i>et al.</i> (2014).
Domains of global functioning– Domains of QoL	5	12	Mas-Exposito <i>et al.</i> (2011), Rocca <i>et al.</i> (2014), Alessandrini <i>et al.</i> (2016), Bai <i>et al.</i> (2014), Fujino <i>et al.</i> (2016).
Alternate statistical methods	4	10	Mubarak (2005), Pitkanen <i>et al.</i> (2012), Becker <i>et al.</i> (2005), Medeiros-Ferreira <i>et al.</i> (2013).
Associations estimated by multivariate analysis			
Global functioning – Overall QoL	10	24	Roe <i>et al.</i> (2011), Meijer <i>et al.</i> (2009), Kusel <i>et al.</i> (2007), Stubbs <i>et al.</i> (2015), Alessandrini <i>et al.</i> (2016), Rocca <i>et al.</i> (2016), Norman <i>et al.</i> (2000), Woon <i>et al.</i> (2010), Prince (2007), Lasebikan and Owoaje (2015).
Global functioning – Domains of QoL	4	10	Brissos <i>et al.</i> (2011), Zendjidian <i>et al.</i> (2014), Woon <i>et al.</i> (2010), Akinsulore <i>et al.</i> (2014).

moderate for the environmental, psychological and social components and small for over half the correlations for physical components.

Over two-thirds of the total correlations between global functioning and QoL domains were self-assessed and nearly half were small ($r = -0.03$ to 0.29). The remaining 29 correlations were interviewer-assessed and over half were small ($r = 0.10$ to 0.21).

Associations resulting from alternate statistical approaches

In the four articles assessing associations between global functioning and QoL using alternate statistical methods, four separate approaches were used. Mubarak (2005) which compared means of dysfunction with a dichotomisation of QoL found that people with high dysfunction (low functioning) had low QoL. Pitkanen *et al.* (2012) which compared medians of QoL with a dichotomisation of functioning reported that lower functioning indicated poorer QoL. Becker *et al.* (2005) which compared the means of QoL across three levels of functioning showed an increase in the mean of QoL across levels of functioning from low to high. Medeiros-Ferreira *et al.* (2013) which compared the means of HRQoL and functioning in subgroups of people with or without metabolic syndrome reported no association.

Functioning as a predictor of QoL

Nineteen articles (45%) assessed predictors of QoL. Of these, Fujino *et al.* (2016) and Rocca *et al.* (2014) did not include global functioning as an initial predictor, 13 modelled global functioning as a predictor in their final models and four (Gaite *et al.*, 2002; Adewuya and Makanjuola, 2009; Kuo *et al.*, 2009; Razali and Wahid, 2012) did not include it in the final models (Table 2).

Of the 13 articles that modelled global functioning as a predictor of QoL, nine considered QoL overall and three specific domains of QoL. Woon *et al.* (2010) tested both. Global

functioning predicted QoL overall in most analyses (Norman *et al.*, 2000; Kusel *et al.*, 2007; Meijer *et al.*, 2009; Woon *et al.*, 2010; Roe *et al.*, 2011; Lasebikan and Owoaje, 2015; Alessandrini *et al.*, 2016; Rocca *et al.*, 2016). Exceptions were Stubbs *et al.* (2015) and Prince (2007).

Discussion

This systematic review is the first to explore the presence and extent of associations between global functioning and QoL in people with psychotic disorders. The appraisal proved difficult given a lack of similarities between studies, and differences in methodological approaches including instruments employed, and inconsistencies in results for given instruments. Despite these difficulties, we found that most of the included articles reported positive associations between higher global functioning and better QoL, and through a narrative review, we were able to clarify the extent of these associations alongside important explanatory factors.

Our results showed that the strength of the association was primarily dependent on the QoL instrument used and whether QoL was being objectively or subjectively assessed. The largest correlations were given by objective QoL instruments completed by an interviewer, the RSM-Scale in particular. The RSM-Scale covers social, occupational and psychological functioning and includes physical functioning and subjective well-being (Riedel *et al.*, 2011). The other objective QoL instrument, the QLS, covers social (interpersonal relations), occupational (instrumental role) and psychological domains (intrapsychic foundations) as well as common objects and activities (Heinrichs *et al.*, 1984). Thus, domains covered by both objective QoL instruments, overlap with domains encompassed by measures of global functioning, which account for the strong associations observed. Furthermore, both, objective QoL instruments and measures of global functioning were assessed by the one interviewer leading to further consistency in

Fig. 3. Correlations between global functioning and quality of life.

assessment. In contrast, subjective QoL instruments are intended to be completed through self-assessment and given that the patient's perspective can differ from an evaluator's (Sainfort *et al.*, 1996; Atkinson *et al.*, 1997; Bengtsson-Tops *et al.*, 2005), the resulting differences will affect the strength of the correlation.

Even with the application of the same instruments, correlation coefficients will vary as a result of sampling variation. This was evident for studies that used the GAF and MANSA (Kusel *et al.*, 2007; Roe *et al.*, 2011). A smaller correlation was found when participants were all living in a psychiatric rehabilitation residential facility in Israel (Roe *et al.*, 2011) than recruited from inpatient and outpatient settings in the UK (Kusel *et al.*, 2007). Likewise, the importance of country and in turn differences in cultural and possibly health system structure and functioning is arguably reflected in Hosseini and Yousefi (2011). Two-thirds of participants in this study lived in an Iranian institution, and while QoL was measured with an objective QoL instrument (QLS), a small correlation was assessed. In comparison strong to moderate associations were assessed in other studies using the QLS from Europe (Kusel *et al.*, 2007; Karadayi *et al.*, 2011; Hunter and Barry, 2012; Nafees *et al.*, 2012), Canada (Norman *et al.*, 2000) and Israel (Ritsner *et al.*, 2005).

Variation in QoL of people living with psychosis based on sociodemographic characteristics is well documented (Browne *et al.*, 1996; Chan *et al.*, 2003; Caron *et al.*, 2005). Ethnicity has also been acknowledged as a contributor to the QoL of people with psychosis within a given cultural setting (Lehman, 1995; Ben-Zur *et al.*, 2014), which may in part be due to the impact of racism and discrimination on an individual's expectations (Lehman, 1995; Prince, 2007). Religious beliefs and spirituality

may also contribute directly to a better QoL (Cohen *et al.*, 2010; Grover *et al.*, 2014; Caqueo-Urizar *et al.*, 2016). It is thus evident that social and demographic issues will affect associations between global functioning and QoL of people living with psychosis.

Our results support respondent burden as a potential confounder in the assessment of associations between functioning and QoL (Ulrich *et al.*, 2005; Fricker *et al.*, 2014). We observed small correlations only when three QoL instruments were employed (Reine *et al.*, 2005), and predominantly small correlations when two QoL instruments were employed (Meijer *et al.*, 2002; Dima *et al.*, 2015).

We also observed that some authors (Miclutia *et al.*, 2008; Mas-Exposito *et al.*, 2011; Bai *et al.*, 2014) reported a total score for the WHOQOL-BREF when there is no such approved score for this instrument. The WHOQOL-BREF generates four domain scores and two separately scored items (overall perception of QoL and health) (World Health Organization, 1998; University of Washington, 2011). As the appropriate use of an instrument is essential for a valid outcome, all results pertaining to the WHOQOL-BREF total score are not considered reliable. In another study (Medeiros-Ferreira *et al.*, 2013), standard scoring techniques for the EQ-5D were not applied. Reine *et al.* (2005) reported the physical and mental composite scores of the SF-36, and while assessed (Ware *et al.*, 1995), it has been suggested they provide an imprecise summary of profile scores (Taft *et al.*, 2001). For these reasons, results of these studies were not considered reliable, reinforcing the importance of the proper use of an instrument, and the necessity of adhering to standardised scoring protocols.

Table 3. Correlations between overall scores of global functioning and domains scores of quality of life

Components	QoL instrument construct	Subjective									Both		Objective	
	Articles	Woon <i>et al.</i> (2010)	Akinsulore <i>et al.</i> (2014)	Miclutia <i>et al.</i> (2008)	Mas–Exposito <i>et al.</i> (2011)	Galuppi <i>et al.</i> (2010)	Bai <i>et al.</i> (2014)	Fujino <i>et al.</i> (2016)	Kuo <i>et al.</i> (2009)	Dima <i>et al.</i> (2015)	Reine <i>et al.</i> (2005)	Meijer <i>et al.</i> (2002)	Riedel <i>et al.</i> (2011)	Rocca <i>et al.</i> (2014)
	Instruments	GAF	GAF	GAF	WHODAS-S	FPS	SRG-PSP	UPSA-B	GAF	GAF	GAF	GAF	GAF	PSP
	QoL domains (number of items)	WHOQOL-BREF						JSQLS	SQLS -R4	SF-36 Q-LES-Q SF	SF-36 QoLI S-QoL	SF-36 LQoLP Dutch	RSM-Scale ^d	QLS
Environmental	Environment (8)	0.34 [♦]	0.19	0.42 ^{**}	−0.36 [♦]	0.43 [♦]	0.39 ^{**}							
	Living conditions (11) (RSM Sub_3)												0.26 [♦] 0.47 ^{♦e}	
	Living situation (4)											0.04		
	Residence (3)										0.07 ^b			
	Finances (4)											0.11		
	Disposable income (3)										−0.05 ^b			
	Personal security (3)										0.10 ^b			
	Safety (2)											0.08		
Life in general	Subjective well-being (16) (RSM Sub_1)												0.36 [♦] 0.72 ^{♦e}	
	Life satisfaction (1)									0.23 [*]				
	Goals (Fulfilment) (13)											0.21 ^{**}		
	Framework (10)											0.14		
	Symptoms/Side Effects (8)							−0.34 ^{**}						
	Satisfaction with treatment (1)									0.23 [*]				
Occupational	Occupational functioning (8)												0.34 [♦] 0.59 ^{♦e}	
	Instrumental role (4)													0.44 [♦]
	Job satisfaction (3)										−0.10 ^b			
	Common Objects and activities (2)													0.17
	Leisure activities (4)										0.12 ^b			

(Continued)

Table 3. (Continued.)

Components	QoL instrument construct	Subjective									Both		Objective	
	Articles	Woon et al. (2010)	Akinsulore et al. (2014)	Miclutia et al. (2008)	Mas–Exposito et al. (2011)	Galuppi et al. (2010)	Bai et al. (2014)	Fujino et al. (2016)	Kuo et al. (2009)	Dima et al. (2015)	Reine et al. (2005)	Meijer et al. (2002)	Riedel et al. (2011)	Rocca et al. (2014)
	Instruments	GAF	GAF	GAF	WHODAS-S	FPS	SRG-PSP	UPSA-B	GAF	GAF	GAF	GAF	GAF	PSP
	QoL domains (number of items)	WHOQOL-BREF									SF-36 QoLI S-QoL	SF-36 LQoLP Dutch	RSM-Scale ^d	QLS
Psychological	Psychological (6)	0.31 [♦]	0.42 ^{**}	0.58 ^{**}	−0.31 [♦]	0.34 [*]	0.39 ^{**}							
	Psychological wellbeing (10)										0.26 ^{c♦}			
	Mental health (5)									0.28 ^{a**}	0.24 ^{a♦}	0.38 ^{a**}		
	Composite Mental score										0.21 ^{a**}			
	Resilience (5)										0.26 ^{c♦}			
	Emotional functioning (10)												0.35 [♦] 0.65 ^{♦e}	
	Role Emotional (3)									0.12 ^{a*}	0.17 ^{a*}	0.25 ^{a**}		
	Self-esteem (6)										0.24 ^{c♦}			
	Negative esteem (5)											0.35 ^{**}		
	Positive esteem (5)											0.21 [*]		
	Psychosocial (15)							−0.17	−0.03					
Physical	Health (7)											0.33 ^{**}		
	Health (1)					0.30 [*]								
	General Health Perceptions (5)										0.15 ^{a*}	0.20 ^{a*}		
	Physical Health (7)	0.29 ^{**}	0.42 ^{**}	0.29 ^{**}	−0.30 [♦]	0.57 [†]	0.44 ^{**}							
	Physical functioning (10)									0.40 ^{a**}	0.14 ^a	0.28 ^{a**}		
	Physical functioning (7)												0.22 [♦] 0.41 ^{♦e}	
	Physical well-being (4)										0.21 ^{c♦}			
	Role physical (4)									0.20 ^{a**}	0.18 ^{a*}	0.26 ^{a**}		
	Composite physical score										0.14 ^a			

	Cognition (4)							0.28 [♦] 0.61 ^{♦e}
	Autonomy (4)						0.28 ^{c♦}	
	Energy/Fatigue (4)				0.21 ^{a**}	0.24 ^{a♦}	0.17 ^{a*}	
	Motivation energy (7)			−0.16	−0.11			
	Bodily Pain (2)						0.15 ^{a*}	0.19 ^{a**}
	Mental and physical health (3)						0.10 ^{b**}	
Social	Interpersonal relationships (8)							0.56 [♦]
	Social relationships (3)	NP	0.21 [*]	0.47 ^{**}	−0.31 [♦]	0.37 [*]	0.37 ^{**}	
	Social relations (3)						0.20 ^{b**}	
	Functioning in social roles (9) (RSM Sub_2)							0.35 [♦] 0.65 ^{♦e}
	Social functioning (7)							0.38 [♦] 0.65 ^{♦e}
	Social functioning (2)					0.32 ^{a**}	0.18 ^{a*}	0.37 ^{a**}
	Sentimental life (2)						0.23 ^{c♦}	
	Leisure and social (6)							0.18 [*]
	Friends relationships (5)						0.16 ^{c*}	
	Family relationships (5)						0.09 ^c	
	Family relations (2)							0.12
	Family relations (2)							−0.05 ^b

* $p < 0.05$, ** $p < 0.01$, [†] $p < 0.005$, [♦] $p < 0.001$.

Note: Quality of life domains were extracted from each of the QoL instruments. The number of items for the assessment of every domain is within parenthesis.

NP, not provided.

^aSF-36.

^bQoLI brief. Only subjective items were used.

^cS-QoL.

^dRSM-scale allocates their 36 items into the five-dimensions model, it also assigns the items to three subscores: Items 1–16 (Sub_1), items 17–20, 23, 32–35 (Sub_2), items 21–22, 24–31, 36 (Sub_3).

^eInterviewer-assessment.

Another important finding was that the domains covered by an instrument are key to the associations obtained. Further, the items included within an instrument will lead to differences in correlations as reflected in the results for the three variations of the QLS (the complete 21 items and abbreviated 7 and 5 items). We consider that in the assessment of QoL of people with psychosis it is important to include items that broadly encompass the mental or psychological domain, otherwise, the outcome will not fully reflect the potential experiences of this population. Thus, the assessment of mental health in the EQ-5D is considered problematic given that the dimension comprises a single item regarding the presence of anxiety and/or depression. In turn, while moderate correlations were obtained between the EQ-5D and global functioning, correlations themselves should not be considered sufficient in determining a reliable and valid QoL instrument.

Findings regarding the strength of the associations between global functioning and QoL domains highlighted social components. These had the highest correlations, with more than half moderate or higher even when subjectively assessed. However, the breadth of domains also impacted correlations and was smaller for narrow domains. As with comparisons between overall scores, the QoL instrument was the primary determinant of the strength of the correlations between global functioning and QoL domains.

Our systematic review has highlighted the extensive array of instruments for the assessment of QoL, and to a lesser extent global functioning in people living with a psychotic disorder. Further, given that both outcomes are commonly used, our systematic review provides the framework to understand the different findings reported in the literature, and inform the future choice of instruments by researchers and/or clinicians.

We found no patterns in the use of instruments in regard to either year, a country where studies were conducted, diagnostic criteria or other characteristics. This variability could be in part due to the absence of a universal definition of global functioning and QoL. However, the diversity is likely at least in part driven by cultural issues leading to modification of available instruments. Also limiting study findings was the need to undertake a narrative review given the heterogeneity of study findings. We did not include articles published in other than English or Spanish. All articles identified employed English, although over three-quarters were from non-English speaking countries. Therefore, it is considered that this limitation will have minimal impact if any on our findings. We did not register our protocol with PROSPERO: International prospective registered systematic reviews (University of York, 2011) as we considered our review fell under stated exclusion criteria: 'looking at the reporting of and/or use of outcomes in research would not be included'.

We believe that clearer and precise definitions of global functioning and QoL are required so these outcomes can be concisely and uniformly measured, and we can identify the domains of life that need to be targeted for improving these outcomes. Further, only by having standard/homogeneous instruments, can we consistently assess the impact of interventions aimed at improving these outcomes, and thereby contribute to the ongoing development and implementation of strategies for improving global functioning and QoL in people living with psychotic disorders.

Overall, most articles identified reported moderate and positive associations between global functioning and QoL. The strength of correlation was dependent upon the instruments employed and the respondent (e.g., a clinician or the individual

living with psychosis). However, the moderate associations between global functioning and QoL reassure that interventions that improve functioning in people with a psychotic disorder will have a positive impact on their QoL. Policy makers and clinicians should make improvement of QoL of people with psychosis a priority alongside symptom remission. Happiness and satisfaction, fulfilment of goals and expectations, are essential to people living with psychotic illness.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S2045796018000549>

Acknowledgements. None.

Financial Support. This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Conflict of Interest. None.

Availability of Data and Materials. Data supporting our findings can be available under request.

References

- Aas IM (2010) Global Assessment of Functioning (GAF): properties and frontier of current knowledge. *Annals of General Psychiatry* 9, 20.
- Adewuya AO and Makanjuola ROA (2009) Subjective quality of life of Nigerian schizophrenia patients: sociodemographic and clinical correlates. *Acta Psychiatrica Scandinavica* 120, 160–164.
- Akinsulore A, Aloba OO, Mapayi BM, Oloniniyi IO, Fatoye FO and Makanjuola RO (2014) Relationship between depressive symptoms and quality of life in Nigerian patients with schizophrenia. *Social Psychiatry and Psychiatric Epidemiology* 49, 1191–1198.
- Alessandrini M, Lançon C, Fond G, Faget-Agius C, Richieri R, Faugere M, Metairie E, Boucekine M, Llorca PM, Auquier P and Boyer L (2016) A structural equation modelling approach to explore the determinants of quality of life in schizophrenia. *Schizophrenia Research* 171, 27–34.
- Ascher-Svanum H, Novick D, Haro JM, Aguado J and Cui Z (2013). Empirically driven definitions of “good,” “moderate,” and “poor” levels of functioning in the treatment of schizophrenia. *Quality of Life Research* 22, 2085–2094.
- Atkinson M, Zibin S and Chuang H (1997) Characterizing quality of life among patients with chronic mental illness: a critical examination of the self-report methodology. *American Journal of Psychiatry* 154, 99–105.
- Awad AG, Voruganti LN and Heslegrave RJ (1997) Measuring quality of life in patients with schizophrenia. *PharmacoEconomics* 11, 32–47.
- Bai YM, Hsiao CY, Chen KC, Huang KL, Lee IH, Hsu JW, Chen PS and Yang YK (2014) The development of a self-reported scale for measuring functionality in patients with schizophrenia-Self-reported version of the graphic Personal and Social Performance (SRG-PSP) scale. *Schizophrenia Research* 159, 546–551.
- Baker F and Intagliata J (1982) Quality of life in the evaluation of community support systems. *Evaluation and Program Planning* 5, 69–79.
- Becker T, Leese M, Krumm S, Ruggeri M and Vazquez-Barquero JL (2005) Needs and quality of life among patients with schizophrenia in five European centres: what is the impact of global functioning scores? *Social Psychiatry and Psychiatric Epidemiology* 40, 628–634.
- Bellack AS, Green MF, Cook JA, Fenton W, Harvey PD, Heaton RK, Laughren T, Leon AC, Mayo DJ, Patrick DL, Patterson TL, Rose A, Stover E and Wykes T (2007) Assessment of community functioning in people with schizophrenia and other severe mental illnesses: a white paper based on an NIMH-sponsored workshop. *Schizophrenia Bulletin* 33, 805–822.
- Ben-Zur H, Duvdevany I and Saffoury Issa D (2014) Ethnicity moderates the effects of resources on quality of life for persons with mental illness living in community settings. *Psychiatric Rehabilitation Journal* 37, 309–315.
- Bengtsson-Tops A, Hansson L, Sandlund M, Bjarnason O, Korkeila J, Merinder L, Nilsson L, Sörgaard KW, Vinding HR and Middelboe T (2005) Subjective versus interviewer assessment of global quality of life

- among persons with schizophrenia living in the community: a Nordic multicentre study. *Quality of Life Research* **14**, 221–229.
- Bowling A** (2005) *Measuring Health: a Review of Quality of Life Measurement Scales*. Berkshire, England: Open University Press.
- Brissos S, Balanza-Martinez V, Dias VV, Carita AI and Figueira ML** (2011) Is personal and social functioning associated with subjective quality of life in schizophrenia patients living in the community? *European Archives of Psychiatry & Clinical Neuroscience* **261**, 509–517.
- Browne S, Roe M, Lane A, Gervin M, Morris M, Kinsella A, Larkin C and O'Callaghan E** (1996) Quality of life in schizophrenia: relationship to sociodemographic factors, symptomatology and tardive dyskinesia. *Acta Psychiatrica Scandinavica* **94**, 118–124.
- Caqueo-Urizar A, Urzúa A, Boyer L, Williams D, Caqueo-Urizar A, Urzúa A and Williams DR** (2016) Religion involvement and quality of life in patients with schizophrenia in Latin America. *Social Psychiatry and Psychiatric Epidemiology* **51**, 521–528.
- Caron J, Lecomte Y, Stip E and Renaud S** (2005) Predictors of quality of life in schizophrenia. *Community Mental Health Journal* **41**, 399–417, 19p.
- Chan GWL, Ungvari GS, Shek DTL and Leung JJP** (2003) Hospital and community-based care for patients with chronic schizophrenia in Hong Kong: quality of life and its correlates. *Social Psychiatry & Psychiatric Epidemiology* **38**, 196–203.
- Chino B, Nemoto T, Fujii C and Mizuno M** (2009) Subjective assessments of the quality of life, well-being and self-efficacy in patients with schizophrenia: regular article. *Psychiatry and Clinical Neurosciences* **63**, 521–528.
- Cohen J** (1988) *Statistical Power Analysis for the Behavioral Sciences*. L. Hillsdale, New Jersey: Erlbaum Associates.
- Cohen CI, Jimenez C and Mittal S** (2010) The role of religion in the well-being of older adults with schizophrenia. *Psychiatric Services* **61**, 917–922.
- Dima L, Vasile D, Rogozea L, Zia-Ul-Haq M, Bukhari SA and Moga M** (2015) Self-perception of quality of life in patients treated with antipsychotics. *Turkish Journal of Medical Sciences* **45**, 782–788.
- Elkinton JR** (1966) Medicine and the quality of life. *Annals of Internal Medicine* **64**, 711–714.
- Fayers PM and Machin D** (2016) *Quality of Life: The Assessment, Analysis and Reporting of Patient-Reported Outcomes*. Chichester, Sussex UK: Wiley Blackwell.
- Fricker S, Yan T and Tsai S** (2014) Response burden: what predicts it and who is burdened out? In AAPOR Proceedings, American Association for Public Opinion Research, Anaheim, California, May 15–18, 2014, pp. 4568–4577.
- Fujino H, Sumiyoshi C, Sumiyoshi T, Yasuda Y, Yamamori H, Ohi K, Fujimoto M, Hashimoto R, Takeda M and Imura O** (2016) Predicting employment status and subjective quality of life in patients with schizophrenia. *Schizophrenia Research: Cognition* **3**, 20–25.
- Gaite L, Vazquez-Barquero JL, Borra C, Ballesteros J, Schene A, Welcher B, Thornicroft G, Becker T, Ruggeri M and Herran A** (2002) Quality of life in patients with schizophrenia in five European countries: the EPSILON study. *Acta Psychiatrica Scandinavica* **105**, 283–292.
- Galuppi A, Turola MC, Nanni MG, Mazzoni P and Grassi L** (2010) Schizophrenia and quality of life: how important are symptoms and functioning? *International Journal of Mental Health Systems* **4**, 31.
- Greenley JR, Greenberg JS and Brown R** (1997) Measuring quality of life: a new and practical survey instrument. *Social Work* **42**, 244–254.
- Grover S, Davuluri T and Chakrabarti S** (2014) Religion, spirituality, and schizophrenia: a review. *Indian Journal of Psychological Medicine* **36**, 119.
- Guilera G, Gomez-Benito J, Pino O, Rojo JE, Cuesta MJ, Martinez-Aran A, Safont G, Tabares-Seisdedos R, Vieta E, Bernardo M, Crespo-Facorro B, Franco M and Rejas J** (2012) Utility of the World Health Organization Disability Assessment Schedule II in schizophrenia. *Schizophrenia Research* **138**, 240–247.
- Harvey PD** (2014). What is the evidence for changes in cognition and functioning over the lifespan in patients with schizophrenia? *The Journal of Clinical Psychiatry* **75**(Suppl. 2), 34–38.
- Harvey PD and Davidson M** (2002) Schizophrenia: Course Over the Lifetime. In *Neuropsychopharmacology: The Fifth Generation of Progress* (ed. KL Davis, D Charney, JT Coyle and C Nemeroff), pp. 641–655. Philadelphia: Lippincott Williams & Wilkins.
- Heinrichs DW, Hanlon TE and Carpenter WT** (1984) The Quality of Life Scale: an instrument for rating the schizophrenic deficit syndrome. *Schizophrenia Bulletin* **10**, 388–398.
- Holloway F and Carson J** (1999) Subjective quality of life, psychopathology, satisfaction with care and insight: an exploratory study. *International Journal of Social Psychiatry* **45**, 259–267.
- Hosseini SH and Yousefi MK** (2011) Quality of life and GAF in schizophrenia: correlation between quality of life and global functioning in schizophrenia. *Iranian Journal of Psychiatry and Behavioral Sciences* **5**, 120–125.
- Hunter R and Barry S** (2012) Negative symptoms and psychosocial functioning in schizophrenia: neglected but important targets for treatment. *European Psychiatry* **27**, 432–436.
- Ito S, Nemoto T, Tsujino N, Ohmuro N, Matsumoto K, Matsuoaka H, Tanaka K, Nishiyama S, Suzuki M, Kinoshita H, Ozawa H, Fujita H, Shimodera S, Kishimoto T, Matsumoto K, Hasegawa T and Mizuno M** (2015) Differential impacts of duration of untreated psychosis (DUP) on cognitive function in first-episode schizophrenia according to mode of onset. *European Psychiatry* **30**, 995–1001.
- Karadayi G, Emiroglu B and Uçok A** (2011) Relationship of symptomatic remission with quality of life and functionality in patients with schizophrenia. *Comprehensive Psychiatry* **52**, 701–707.
- König H-H, Roick C and Angermeyer MC** (2007) Validity of the EQ-5D in assessing and valuing health status in patients with schizophrenic, schizotypal or delusional disorders. *European Psychiatry* **22**, 177–187.
- Kuo PJ, Ma HI, Kuo CC, Huang WC and Chung MS** (2009) Factor analysis of the schizophrenia quality of life scale revision 4 (SQLS-R4) Chinese version and related factors. *International Journal of Psychiatry in Clinical Practice* **13**, 278–284.
- Kusel Y, Laugharne R, Perrington S, McKendrick J, Stephenson D, Stockton-Henderson J, Barley M, McCaul R and Burns T** (2007) Measurement of quality of life in schizophrenia: a comparison of two scales. *Social Psychiatry and Psychiatric Epidemiology* **42**, 819–823.
- Lasebikan VO and Owoaje ET** (2015) Quality of life in psychosis: prevalence and associated factors in a Nigerian clinical population. *Community Mental Health Journal* **51**, 491–496.
- Lehman AF** (1988) A quality of life interview for the chronically mentally ill. *Evaluation and Program Planning* **11**, 51–62.
- Lehman AF** (1995) Demographic influences on quality of life among persons with chronic mental illnesses. *Evaluation and Program Planning* **18**, 155–164.
- Mas-Exposito L, Amador-Campos JA, Gomez-Benito J and Lalucat-Jo L** (2011) The World Health Organization Quality of Life Scale Brief Version: a validation study in patients with schizophrenia. *Quality of Life Research* **20**, 1079–1089.
- Medeiros-Ferreira L, Obiols JE, Navarro-Pastor JB and Zuniga-Lagares A** (2013) Metabolic syndrome and health-related quality of life in patients with schizophrenia. *Actas Espanolas de Psiquiatria* **41**, 17–26.
- Meijer CJ, Schene AH and Koeter MW** (2002) Quality of life in schizophrenia measured by the MOS SF-36 and the Lancashire Quality of Life Profile: a comparison. *Acta Psychiatrica Scandinavica* **105**, 293–300.
- Meijer CJ, Koeter MW, Sprangers MA and Schene AH** (2009) Predictors of general quality of life and the mediating role of health related quality of life in patients with schizophrenia. *Social Psychiatry & Psychiatric Epidemiology* **44**, 361–368.
- Miclutia IV, Popescu CA and Macrea RS** (2008) Sexual dysfunctions of chronic schizophrenic female patients. *Sexual & Relationship Therapy* **23**, 119–129.
- Moher D, Liberati A, Tetzlaff J and Altman DG, and the PRISMA Group** (2009). Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Annals of Internal Medicine* **151**, 264–269.
- Mubarak AR** (2005) Social functioning and quality of life of people with schizophrenia in the northern region of Malaysia. *Australian e-Journal for the Advancement of Mental Health* **4**, 1–10.
- Nafees B, van Hanswijck de Jonge P, Stull D, Pascoe K, Price M, Clarke A and Turkington D** (2012) Reliability and validity of the Personal and Social Performance scale in patients with schizophrenia. *Schizophrenia Research* **140**, 71–76.
- Norman RMG, Malla AK, McLean T, Voruganti LPN, Cortese L, McIntosh E, Cheng S and Rickwood A** (2000) The relationship of

- symptoms and level of functioning in schizophrenia to general wellbeing and the Quality of Life Scale. *Acta Psychiatrica Scandinavica* **102**, 303–309.
- Patterson TL, Goldman S, McKibbin CL, Hughs T and Jeste DV** (2001) UCSD performance-based skills assessment: development of a new measure of everyday functioning for severely mentally ill adults. *Schizophrenia Bulletin* **27**, 235–245.
- Petho B and Ban TA** (1988) DCR Budapest-Nashville in the diagnosis and classification of functional psychoses. *Psychopathology* **21**, 149–240.
- Pinikahana J, Happell B, Hope J and Keks NA** (2002) Quality of life in schizophrenia: a review of the literature from 1995 to 2000. *International Journal of Mental Health Nursing* **11**, 103–111.
- Pitkanen A, Valimäki M, Endicott J, Katajisto J, Luukkaala T, Koivunen M, Kuosmanen L and Hatonen H** (2012) Assessing quality of life in patients with schizophrenia in an acute psychiatric setting: reliability, validity and feasibility of the EQ-5D and the Q-LES-Q. *Nordic Journal of Psychiatry* **66**, 19–25.
- Prigent A, Simon S, Durand-Zaleski I, Leboyer M and Chevreur K** (2014) Quality of life instruments used in mental health research: properties and utilization. *Psychiatry Research* **215**, 1–8.
- Prince JD** (2007) Education, ethnicity, and subjective life quality of recently discharged inpatients with schizophrenia. *Journal of Nervous & Mental Disease* **195**, 560–565.
- PROSPERO: International prospective register of systematic reviews.** Centre for Reviews and Dissemination, University of York. (<http://www.crd.york.ac.uk>). Accessed 13 August 2018.
- Razali SM and Wahid MA** (2012) Quality of life and depressive symptoms in patients with schizophrenia. *International Medical Journal* **19**, 130–134.
- Reine G, Simeoni MC, Auquier P, Loundou A, Aghababian V and Lançon C** (2005) Assessing health-related quality of life in patients suffering from schizophrenia: a comparison of instruments. *European Psychiatry* **20**, 510–519.
- Riedel M, Spellmann I, Schennach-Wolff R, Obermeier M and Musil R** (2011) The RSM-scale: a pilot study on a new specific scale for self- and observer-rated quality of life in patients with schizophrenia. *Quality of Life Research* **20**, 263–272.
- Ritsner M, Kurs R, Ratner Y and Gibel A** (2005) Condensed version of the Quality of Life Scale for schizophrenia for use in outcome studies. *Psychiatry Research* **135**, 65–75.
- Rocca P, Montemagni C, Zappia S, Pitera R, Sigaucho M and Bogetto F** (2014) Negative symptoms and everyday functioning in schizophrenia: a cross-sectional study in a real world-setting. *Psychiatry Research* **218**, 284–289.
- Rocca P, Montemagni C, Mingrone C, Crivelli B, Sigaucho M and Bogetto F** (2016) A cluster-analytical approach toward real-world outcome in outpatients with stable schizophrenia. *European Psychiatry* **32**, 48–54.
- Roe D, Mashiach-Eizenberg M and Lysaker PH** (2011) The relation between objective and subjective domains of recovery among persons with schizophrenia-related disorders. *Schizophrenia Research* **131**, 133–138.
- Sainfort F, Becker M and Diamond R** (1996) Judgments of quality of life of individuals with severe mental disorders: patient self-report versus provider perspectives. *The American Journal of Psychiatry* **153**, 497–502.
- Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, Moher D, Becker BJ, Sipe TA and Thacker SB** (2000) Meta-analysis of observational studies in epidemiology: a proposal for reporting. *Journal of the American Medical Association* **283**, 2008–2012.
- Stubbs B, Gardner-Sood P, Smith S, Ismail K, Greenwood K, Patel A, Farmer R and Gaughran F** (2015) Pain is independently associated with reduced health related quality of life in people with psychosis. *Psychiatry Research* **230**, 585–591.
- Taft C, Karlsson J and Sullivan M** (2001) Do SF-36 summary component scores accurately summarize subscale scores? *Quality of Life Research* **10**, 395–404.
- The WHOQOL Group** (1998) Development of the World Health Organization WHOQOL-BREF quality of life assessment. *Psychological Medicine* **28**, 551–558.
- Ulrich CM, Wallen GR, Feister A and Grady C** (2005) Respondent burden in clinical research: when are we asking too much of subjects? *IRB: Ethics & Human Research* **27**, 17–20.
- von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC and Vandenbroucke JP** (2007) The strengthening of reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *Bulletin of the World Health Organization* **85**, 867–872.
- Ware Jr JE, Kosinski M, Bayliss MS, McHorney CA, Rogers WH and Raczek A** (1995). Comparison of methods for the scoring and statistical analysis of SF-36 health profile and summary measures: summary of results from the Medical Outcomes Study. *Medical Care* **33**, AS264–AS279.
- Woon PS, Chia MY, Chan WY and Sim K** (2010) Neurocognitive, clinical and functional correlates of subjective quality of life in Asian outpatients with schizophrenia. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* **34**, 463–468.
- World Health Organization. Division of Mental Health and Prevention of Substance Abuse** (1997) *WHOQOL : measuring quality of life*. Geneva: World Health Organization. <http://www.who.int/iris/handle/10665/63482>
- World Health Organization. Division of Mental Health and Prevention of Substance Abuse** (1998) *Programme on mental health: WHOQOL user manual*. Geneva: WHO.
- World Health Organization** (2001) *International Classification of Functioning, Disability and Health: ICF*. Geneva: WHO.
- Zendjidian XY, Baumstarck K, Auquier P, Loundou A, Lançon C and Boyer L** (2014) Satisfaction of hospitalized psychiatry patients: why should clinicians care? *Patient Preference and Adherence* **8**, 575–583.