Efficient Transmission of Mixed *Plasmodium falciparum*/*vivax* Infections From Humans to Mosquitoes

Sujata Balasubramanian,¹ Rifat S. Rahman,¹ Chanthap Lon,² Christian Parobek,¹ Ratawan Ubalee,³ Nicholas Hathaway,⁵ Worachet Kuntawunginn,⁴ Mok My,⁶ Dav Vy,⁶ Jeremy Saxe,⁷ Charlotte Lanteri,⁴ Feng-Chang Lin,⁷ Michele Spring,⁴ Steven R. Meshnick,⁸ Jonathan J. Juliano,¹ David L. Saunders,^{4,9} and Jessica T. Lin¹

¹Institute of Global Health and Infectious Diseases, University of North Carolina, Chapel Hill; ²Armed Forces Research Institute of Medical Sciences, Phnom Penh, Cambodia; Departments of ³Entomology and ⁴Bacterial and Parasitic Diseases, Armed Forces Research Institute of Medical Sciences, Bangkok, Thailand; ⁵Department of Bioinformatics and Integrated Biology, University of Massachusetts, Worcester; ⁶Royal Cambodian Armed Forces, Phnom Penh, Cambodia; Departments of ⁷Biostatistics and ⁸Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill; ⁹US Army Medical Materiel Development Activity, Fort Detrick, Maryland

Background. In Southeast Asia, people are often coinfectd with different species of malaria (*Plasmodium falciparum* [Pf] and *Plasmodium vivax* [Pv]) as well as with multiple clones of the same species. Whether particular species or clones within mixed infections are more readily transmitted to mosquitoes remains unknown.

Methods. Laboratory-reared *Anopheles dirus* were fed on blood from 119 Pf-infected Cambodian adults, with 5950 dissected to evaluate for transmitted infection. Among 12 persons who infected mosquitoes, polymerase chain reaction and amplicon deep sequencing were used to track species and clone-specific transmission to mosquitoes.

Results. Seven of 12 persons that infected mosquitoes harbored mixed Pf/Pv infection. Among these 7 persons, all transmitted Pv with 2 transmitting both Pf and Pv, leading to Pf/Pv coinfection in 21% of infected mosquitoes. Up to 4 clones of each species were detected within persons. Shifts in clone frequency were detected during transmission. However, in general, all parasite clones in humans were transmitted to mosquitoes, with individual mosquitoes frequently carrying multiple transmitted clones.

Conclusions. Malaria diversity in human hosts was maintained in the parasite populations recovered from mosquitoes fed on their blood. However, in persons with mixed Pf/Pv malaria, Pv appears to be transmitted more readily, in association with more prevalent patent gametocytemia.

Keywords. *Anopheles dirus*; deep sequencing; gametocyte; malaria transmission; *Plasmodium vivax*.

In Southeast Asia, it is not uncommon for persons with malaria to be simultaneously coinfectd with *Plasmodium falciparum* and *Plasmodium vivax* (Pf/Pv). In addition, those with Pv infection frequently harbor more than 1 genetically distinct strain or clone [1, 2]. However, finding wild mosquitoes coinfectd with Pf/Pv is rare [3, 4], and limited study of genetic diversity in mosquitoes suggests a propensity for clonal transmission and inbreeding [5]. These findings raise the question of whether certain species or clones are favored during transmission.

Humans transmit malaria to *Anopheles* mosquitoes when the latter ingest a blood meal containing gametocytes. Gametocytes are sexual stage parasites that once activated, form oocysts in the mosquito midgut, which later rupture to release thousands of sporozoites that are delivered to another human host in the

mosquito's next blood meal. After replicating to great numbers in the human host, parasite numbers reach a nadir at the mosquito oocyst stage [6, 7]. Thus, human-to-mosquito transmission represents a bottleneck in the *Plasmodium* life cycle that may influence how parasite genetic diversity is transmitted through a population.

We hypothesized that a selective bottleneck exists in the mosquito that affects the transmission of species and clones within mixed species infections. To explore this question, we studied 7 persons with Pf/Pv coinfection and 5 persons with Pf monoinfection who infected *Anopheles dirus* mosquitoes as part of membrane feeding studies in Cambodia. Polymerase chain reaction (PCR) of individual mosquitoes was used to find the proportion of mosquitoes infected with each species after blood feeding. Amplicon deep sequencing was used to track the relative composition of individual parasite clones in human blood versus mosquitoes. We found that when transmission occurred, it was remarkably efficient, with the majority of clones from each species maintained through human-to-mosquito transmission. However, Pv appeared to be transmitted more readily from Pf/Pv-infected hosts, owing to frequent patent gametocytemia.

Received 16 April 2019; editorial decision 15 July 2019; accepted 23 July 2019; published online September 24, 2019.

Correspondence: Jessica T. Lin, MD MSCR, 130 Mason Farm Rd, CB 7030, Chapel Hill, NC 27599 (jessica_lin@med.unc.edu).

The Journal of Infectious Diseases® 2020;221:428–37

© The Author(s) 2019. Published by Oxford University Press for the Infectious Diseases Society of America. All rights reserved. For permissions, e-mail: journals.permissions@oup.com. DOI: 10.1093/infdis/jiz388

METHODS

Human Clinical Samples

Samples were collected in those presenting with uncomplicated malaria as part of clinical studies carried out in north-western Cambodia [8–10]. Participants were diagnosed with mono-*Pf* or mixed *Pf/Pv* infection by microscopy, with confirmation by PCR. Blood drawn before treatment was passed through a plasmodipur filter to collect leukocyte depleted blood or passed through a magnetic MACs (Miltenyi Biotec) system for gametocyte enrichment based on hemozoin content [11]. Deoxyribonucleic acid (DNA) was extracted using the QIAGEN Mini Blood DNA extraction kit. Molecular studies were approved by the Institutional Review Board at the University of North Carolina, the Walter Reed Army Institute of Research Institutional Review Board, and the Cambodian National Ethical Committee for Health Research.

Membrane Feeding Assays

Mosquito infectivity was measured by membrane feeding assay. Insectary-reared 5- to 8-day-old female *A. dirus* mosquitoes were membrane-fed patient whole blood, maintained on 10% sucrose, and harvested 9 and 16 days after feeding, to isolate parasite oocyst and sporozoite stages, respectively. At day 9 after feeding, 50 mosquitoes per patient were dissected for oocyst counting and another 30–50 were preserved in 95% ethanol, individually for mixed infections and in groups of 10 for monoinfections (Supplementary Figure S1). Mosquitoes harvested at day 9 were preserved whole, whereas only heads and thoraces of mosquitoes harvested at day 16 were preserved in 95% ethanol. The DNA extraction from mosquitoes was performed using a simplified chelex protocol [12]. Species-specific *Pf* and *Pv* real-time PCR (qPCR), targeting the 18S ribosomal ribonucleic acid gene, was performed as previously described [13, 14]. In a given subject, successful human-to-mosquito transmission was defined as a positive qPCR result in at least 1 of 30 blood-fed mosquitoes (or 1 of 5 pools of 10 mosquitoes) at both day 9 and day 16 postfeeding, corresponding to the presence of oocysts and sporozoites, respectively.

Amplicon Deep Sequencing

Polymerase chain reaction amplification of *Pf* apical membrane antigen 1 (*PfAMA1*) and/or *Pv* merozoite surface protein 1 (*PvMSP1*) was carried out in duplicate from each DNA sample using Roche Fast Start Taq DNA Polymerase. Amplified targets were as follows: for *PfAMA1*, a 170-base pair (bp) fragment corresponding to domain 2 [15]; for *PvMSP1*, a 117-bp variable portion of the 33-kDa subunit of the 42-kDa region of *MSP1* [16]. Deoxyribonucleic acid from leukocyte depleted or MACs-enriched blood from participants and DNA from mosquitoes identified as positive by real-time PCR were pooled or used individually as template for nested PCR using *PfAMA1* or *PvMSP1* primers (Supplementary Table S1). Most amplicons

required nested (in the case of *PvMSP1*) or heminested (in the case of *PfAMA1*) PCR amplification to visualize PCR product on ethidium bromide-stained 1% agarose gel. Polymerase chain reaction products were column purified (PureLink Pro 96-well PCR purification kit; Thermo Fisher Scientific), quantified, pooled, then processed using the Bioo Scientific NextFlex Rapid DNA sequencing kit or NEBNext Ultra DNA Library Prep Kit. Sequencing was performed on an Illumina MiSeq using 150-bp paired-end chemistry.

Haplotype Determination From Deep Sequencing

Analysis of sequences was done using SeekDeep version 2.3.4 (<http://seekdeep.brown.edu/>) [17]. Sequences were initially demultiplexed and filtered by fragment size and base call quality. To avoid false-positive haplotype calls, replicates were clustered, setting thresholds to eliminate variants with lower than 1% frequency for averaged cluster fraction in both replicates and at lower than 0.5% frequency within each replicate. Samples with fewer than 1000 reads in 1 replicate were also eliminated. A leniency of 1 high-quality base mismatch was allowed during clustering between 2 replicates so as to err in favor of stringency in calling haplotypes rather than overreport the multiplicity of infection (MOI). We observed good concordance in haplotype calling between PCR replicates for each sample (Supplementary Table S2).

Data Analysis

Multiplicity of infection was defined as the number of unique *PfAMA* or *PvMSP1* haplotypes detected per isolate. The DNA alignments were generated using Clustal Omega, and figures were made with GenVison software (DNASar, Madison, WI). Minimum spanning trees, showing the number of single-nucleotide polymorphisms (SNPs) and insertion/deletion events between haplotypes, were generated in SeekDeep and labeled using Inkscape 0.92. For multiclonal *Pv* infections, the proportion of sequence reads aligned to individual haplotypes were compared between blood and mosquito samples using χ^2 testing. A log base 2 transformation of the number of reads was used to reduce variance before performing a χ^2 test of size ($\alpha = 0.05$). Other graphs and statistical analyses were made using Graph-Pad Prism 6.

RESULTS

Human-to-Mosquito Transmission of *Plasmodium falciparum*/*Plasmodium vivax* Malaria

Mixed species *Pf/Pv* human blood-stage infection was readily transmitted to individual mosquitoes when gametocytes of both species were present. As previously reported, laboratory-reared *A. dirus* fed on blood from 107 *Pf* and 12 *Pf/Pv*-infected Cambodian adults yielded positive mosquito infections from 12 persons, including 5 with *Pf* monoinfection (Supplementary Figure S2) [8]. Among 12 subjects with *Pf/Pv*,

5 of 12 successfully transmitted *Pv* infection to mosquitoes, and 2 of 12 transmitted both *Pf* and *Pv* (Figure 1). More frequent *Pv* transmission reflected more prevalent *Pv* gametocytemia, because all transmitting subjects harbored microscopically patent gametocytemia corresponding to the species transmitted. Of the 5 *Pf/Pv* subjects who were noninfectious, 2 also harbored patent gametocytemia, but at low densities (<50 gametocytes/ μ L) (Supplementary Table S3). In general, gametocytemia seemed to correlate with the burden of mosquito infection, in terms of both oocyst prevalence and density (Figure 1 and Supplementary Table S3).

In 1 of the 2 subjects who transmitted both *Pf* and *Pv* (SN119), approximately one quarter of mosquitoes became infected (14 of 60, 23%), with most mosquitoes infected by either *Pf* or *Pv* (12 of 14), although 2 of 14 were infected with both species. In the other subject (SN063), almost all mosquitoes showed PCR evidence of infection (57 of 60), and the majority (45 of 57) were doubly infected (Figure 1). In total, the 12 subjects with mixed *Pf/Pv* malaria infected one third of mosquitoes fed on the cohort (223 of 709). *Plasmodium vivax* was more efficiently transmitted by these patients (211 of 709 or 30% *Pv*-infected mosquitoes vs 59 of 709 or 8.3% *Pf*-infected

mosquitoes), corresponding to the more frequent presence of microscopic *Pv* gametocytemia (8 of 12 patients with patent *Pv* gametocytes (Pvg) vs 3 of 12 patients with patent *Pf* gametocytes). When both *Pf* and *Pv* gametocytes were present, doubly infected mosquitoes were not uncommon, making up 21% (47 of 223) of all infected mosquitoes in the cohort.

Amplicon Deep Sequencing of Blood-Stage Infection

To compare MOI (number of coinfecting clones) in those with mixed *Pf/Pv* infection versus single-species infection, we applied amplicon deep sequencing to leukodepleted blood samples from 14 subjects with *Pf* or *Pv* monoinfection, in addition to the 12 subjects with mixed *Pf/Pv*. A median of 23 411 $\times$ coverage and 33 541 $\times$ coverage was achieved for *Pf*AMA1 and *Pv*MSP1 amplicons, respectively (Supplementary Table S2). Only those haplotypes that were present in duplicate PCRs, existed at $\geq 1.0\%$ frequency within each sample, and differed by >1 SNP across replicates were counted as individual clones. This led to detection of 5 unique *Pf*AMA1 haplotypes among 21 *Pf*-infected subjects and 12 *Pv*MSP1 haplotypes among 17 *Pv*-infected subjects (Supplementary Figures S3 and S4). All haplotypes matched sequences in GenBank. All differed by at least 2 SNPs in their sequences, with a median of 4.5 SNPs different from the next most similar haplotype (Supplementary Figure S5). One *Pf*AMA1 haplotype and 3 *Pv*MSP1 haplotypes were detected only as minor variants (present at in-host frequency <20%). At least 2 of these were unlikely to be false haplotypes because they were found in multiple samples as well as in the blood-fed mosquitoes.

The presence of coinfecting species did not appear to restrict the genetic complexity of mixed infections (Figure 2). The majority of *Pf/Pv* subjects simultaneously harbored multiple *Pf*AMA1 haplotypes and *Pv*MSP1 haplotypes. Overall, the mean *Pf* MOI was 2.5 in the mixed *Pf/Pv* infections vs 2.4 in the mono-*Pf* infections, whereas the mean *Pv* MOI was 2.4 in the mixed *Pf/Pv* infections vs 2.4 in the mono-*Pv* infections. Haplotypes occurring within individual subjects were usually differentiated by several SNPs, making it unlikely that PCR error, sequencing error, or single mutations falsely elevated the measured complexity of infection (Figure 3).

Although MOI was similar between species, multiclonal *Pv* infections were composed of haplotypes with a wider range of in-host frequencies (Figure 2). The dominant *Pv*MSP1 haplotype variant in each *Pv* infection accounted for a median of 76% (interquartile range [IQR], 57%–93%) of sequencing reads, whereas the dominant *Pf*AMA1 variant in each *Pf* infection accounted for a median of 97% (IQR, 94%–98%) of reads.

By increasing the in-host frequency threshold at which SeekDeep called unique haplotypes from 1% up to 30%, we predicted how MOI might be measured by genotyping methods that are less sensitive at detecting minority clones (Figure 4). An increasing threshold cutoff resulted in a steady decrease in

Figure 1. Transmission of mixed *Plasmodium falciparum* (*Pf*)/*Plasmodium vivax* (*Pv*) malaria from 7 Cambodian adults to membrane-fed *Anopheles dirus* mosquitoes: schematic of species-specific parasite burden. (a) Parasitemia scale. The number of + signs indicates the number of species-specific parasites/ μ L: 1–999 (+); 1000–9999 (++); 10 000+ (+++). (b) Gametocytemia scale. The number of gametocyte images indicates the number of species-specific gametocytes/ μ L: 1–49 (1); 50–149 (2); 150+ (3). (c) Oocyst-positive mosquitoes out of 30, as determined by species-specific quantitative polymerase chain reaction (qPCR) at day 9 postfeeding. (d) Sporozoite-positive mosquitoes out of 30, as determined by species-specific qPCR at day 16 postfeeding. *Pf*, orange; *Pv*, blue; *Pf/Pv* coinfection, green.

Figure 2. Percentage distribution of *PfAMA1* and *PvMSP1* haplotypes in the blood among mono-*Plasmodium falciparum* (*Pf*) and *Plasmodium vivax* (*Pv*) infections (A) and mixed *Pf/Pv* infections (B). Individual haplotypes (CAM_AV numbers) are represented by different colors, with width of bars corresponding to their within-host frequency (%) on the x-axis. Blood from 9 *Pf*, 5 *Pv*, and 12 *Pf/Pv* infections are shown. CAM_AV numbers refer to Cambodia_AnlongVeng, with number indicating the population prevalence of the haplotype (CAM_AV0 is the most common haplotype in the cohort, CAM_AV1 the next most common, etc).

diversity and MOI due to elimination of lower frequency haplotypes. At a threshold of 10% in-host frequency, the mean *Pf* MOI dropped to 1.1, suggesting that most of the minor haplotypes detected would be missed by other genotyping methods.

Human-to-Mosquito Transmission of *Plasmodium falciparum* Clones

Seven *Pf*-infected subjects in the cohort infected mosquitoes, including 5 with *Pf* monoinfection and 2 with *Pf/Pv* infection. In both groups, *PfAMA1* haplotypes detected in the blood were generally also detected in mosquitoes. Haplotypes that were dominant in blood remained dominant in mosquitoes at both oocyst and sporozoite stages, and minority haplotypes maintained a similar frequency through transmission (Figure 5). Instances of low-level minority haplotypes (<2% frequency) disappearing in the mosquito were observed in 2 subjects but were likely due to sampling limitations. For SN119-M, low-level mosquito infection (1–2 oocysts were found in each of the 3 of 20 *Pf*-infected mosquitoes) resulted in only 1 infected

mosquito being successfully deep-sequenced (Figure 1 and Supplementary Table S3). In contrast, for SN108, there was heavy mosquito infection (49 of 50 mosquitoes infected, each with median 149 oocysts), and all exhibited the same dominant *AMA1* haplotype found in the blood (SN108).

In addition, minor variant gametocyte haplotypes not readily detected in the blood were found to be transmitted to mosquitoes. Deep sequencing of DNA extracted from gametocytes isolated in 3 subjects (SN002, SN060, SN063-M) revealed the same dominant *PfAMA1* haplotype as that detected in the blood (Figure 5A). However, 1 gametocyte sample (SN063-M) revealed 3 minor variant haplotypes not found in the corresponding blood sample, 2 of which were also detected at low frequency in the mosquitoes (Figure 5B). Thus, haplotypes that might have appeared to be novel in the mosquito due to their absence in the blood were detected upon sequencing of gametocytes. Overall, the composition of variants in the mosquito reflected those found in the human host.

Figure 3. Minimum spanning trees depicting the relationship of individual *PfAMA1* (A) and *PvMSP1* (B) haplotypes within persons with mixed *Plasmodium falciparum/Plasmodium vivax* malaria. For each subject, the minimum spanning tree provides a graphical representation of the in-host frequency of each haplotype (denoted by size of colored orbs) and the number of single-nucleotide polymorphisms and insertion/deletions that separate each haplotype from the next most similar one (denoted by number of pink dots along mutational path between haplotypes).

Human-to-Mosquito Transmission of *Plasmodium vivax* Clones

Compared with the *Pf* infections, the 7 *Pv*-transmitting subjects showcased more diversity of haplotypes in both blood and mosquito stages (Figure 6). Again, haplotypes in the blood were also detected in mosquitoes, with those that were missed in mosquito pools being detected in individual mosquitoes (SN055-M and SN063-M in Figure 6B). However, haplotype frequencies dynamically changed through transmission, especially upon examination of individual mosquitoes. Nondominant haplotypes in the blood sometimes became the dominant haplotype in individual mosquitoes (for example, Ms13 in SN019-M, Mo24

in SN055-M, and Mo24 in SN073-M). In 2 of 4 subjects with multiclonal blood-stage *Pv* infection, the composition of haplotype frequencies observed at the oocyst and sporozoite stages differed from that in the blood, with the dominant haplotype rising in frequency in one (SN055-M) and decreasing in another (SN063-M) (Supplementary Table S4).

These shifts in haplotype frequency from blood to mosquitoes seemed to be associated with the efficiency of transmission. In 1 subject with abundant gametocytes (SN063, 204 Pvg/ μ L), 2 minority haplotypes in the blood (light and dark blue in Figure 6B) increased in frequency through transmission, together

Figure 4. Measured genetic diversity in the cohort according to sensitivity of detection of minority haplotypes for *Plasmodium falciparum* (*Pf*) (A) and *Plasmodium vivax* (*Pv*) (B). For infections of each species ($n = 21$ *Pf* and 17 *Pv*), the change in the mean multiplicity of infection ([MOI] left axis) and total number of haplotypes (right axis) in the cohort is shown to decrease as the threshold frequency below which haplotypes are not called in SeekDeep is increased. Although the mean MOI of each species is the same when minority haplotypes at 1% frequency are detected, the number of *Pv* haplotypes is greater and the measured *Pv* MOI is more robust to less sensitive genotype detection.

Figure 5. *PfAMA1* haplotypes in patient blood through transmission to mosquitoes. For each subject, the composition of haplotypes and their in-host frequencies are depicted in each stage of the parasite life cycle—blood-stage in human, gametocytes isolated from blood (when available), oocysts in mosquito, and sporozoites in head and thorax of mosquito (A). Haplotypes were detected in pooled mosquito samples, except for subject SN063-M, where individual mosquitoes (Mo2 through Mo20) at oocyst and sporozoite stages were also analyzed (B). On the left, the number of gametocytes/μL of blood detected by microscopy is shown. On the right, the percentage of mosquitoes with oocyst-positive midguts is shown along with the median (range) oocyst count per midgut. The suffix -M depicts a subject with mixed *Plasmodium falciparum*/*Plasmodium vivax* infection.

making up 21% of sequencing reads in the blood, but rising to 52% and 46% in pooled oocyst and sporozoite-stage mosquitoes, respectively; however, the majority haplotype in the blood (light green) remained the majority haplotype in fewer than half of individual mosquitoes. In contrast, in a subject who harbored fewer gametocytes (SN055, 42 Pvg/μL) that led to single-oocyst mosquito infections, the majority haplotype, (CAM_AV3, dark red) prevailed: although all 4 haplotypes in blood were detected in individual oocyst-stage mosquitoes, by the time parasites progressed to the sporozoite stage, only CAM_AV3 was detected in both pooled and individual mosquitoes (Figure 6). Thus, in association with a lower gametocyte burden, MOI fell from 4 in the human host to 1–2 in all but 1 of the individually sequenced mosquitoes.

Parasite Uptake in Individual Mosquitoes

Mosquitoes sampling multiclonal infections readily acquired multiple parasite haplotypes and sometimes displayed new haplotypes not detected in the blood. Even in the setting of mixed-species infection, only 3 of 69 individual *A. dirus* sequenced at

either oocyst or sporozoite stages after blood feeding displayed a single *PvMSP1* haplotype by deep sequencing, with the rest acquiring multiple parasite haplotypes in a single blood meal (Figure 6B). In addition, in 16 of 70 (23%) *Pv*-infected mosquitoes, sequencing revealed new haplotypes not detected in the blood, usually at very low frequencies (<5%). In all but 2 mosquitoes, these new haplotypes also appeared in multiple other mosquitoes. For one *Pf*-infected subject (SN063), a seemingly new minority haplotype appearing in oocyst-stage mosquitoes was in fact detectable in gametocyte-enriched blood spots from the subject, as previously mentioned. For the other *Pf* subject in which new haplotypes appeared in the mosquito (SN086), mosquito infection arose from submicroscopic gametocytes, and gametocyte enrichment was not attempted.

DISCUSSION

In this study, we directly examine the transmission of mixed species and mixed genotype infections from humans to mosquitoes using sensitive deep-sequencing methods. Our first

Figure 6. *PvMSP1* haplotypes in patient blood through transmission to mosquitoes. *Plasmodium vivax* haplotypes within mixed infections are depicted through human-to-mosquito transmission (A). For 4 subjects, data are shown for both pooled and individual mosquitoes, with individual mosquitoes sometimes displaying haplotypes not apparent in pooled samples (B).

observation, that *Pv* appears to be more readily transmitted than *Pf* from persons coinfecting with *Pf/Pv* is largely attributable to more prevalent *Pv* gametocytemia in the cohort. Our sample size is small, but this finding is consistent with previous observations from Southeast Asia that patent *Pf* gametocytemia is infrequent but usually necessary for *Pf* transmission [8, 18], whereas the *Pv* parasite densities that cause symptomatic infection are frequently sufficient to infect mosquitoes [19–21]. Thus, *Pv* enjoys a transmission advantage via early gametocytogenesis, whereby *Pv* gametocytes appear in tandem with the first generation of blood-stage asexual parasites, as opposed to the longer maturation period and delayed appearance of *Pf* gametocytes in the peripheral blood [22–24]. We show that this advantage persists in the setting of mixed infections, without evidence of interspecies competition hindering *Pv* transmission to mosquitoes once gametocytes are formed [25].

Second, we show that from a single blood meal from a *Pf/Pv*-infected person, *A. dirus* mosquitoes frequently carry both species to life cycle completion. Impressively, wild-caught *A. dirus* have been found to harbor as many as 3–4 different *Plasmodium* spp at once, including human and primate malaria species, although whether these arose from single or multiple blood meals is unknown [26]. In a recent study, a wild-caught *Anopheles stephensi* mosquito in Djibouti, Horn of Africa was also reported to be coinfecting with *Pf/Pv* [27].

Finally, we observed that the malaria genetic diversity present in human blood infections is efficiently maintained through transmission to mosquitoes, with *Plasmodium* clones existing at very low frequencies in the blood also detected in mosquitoes. Previous studies have suggested that the capacity of mosquitoes to sample minority alleles often exceeds our capacity to detect them in the blood [28–30]. We observed new haplotypes in mosquitoes as well, but to a much lesser degree than that seen in other studies, we speculate for two reasons. First, deep sequencing was applied to leukodepleted blood samples from symptomatic subjects. A higher parasite density in the blood found with symptomatic infections combined with enrichment of parasite versus human DNA maximized the chance of capturing all circulating clones at sequencing. Second, the polyclonal infections we tracked into the mosquito were mostly due to *Pv*. It may be that for *Pv*, a greater temporal and quantitative association between asexual and sexual-stage parasitemia [20, 23] results in more proportionate representation of all gametocyte genotypes in the blood.

The maintenance of plasmodial diversity from the human to mosquito reservoir has important implications for malaria control and elimination efforts. The capacity to maintain genetic diversity through transmission enhances the parasite's capacity to evolve in response to control interventions, including drugs, bednets, vaccines, and diagnostic tests. Previous studies have variously found substantial or negligible impact of drug resistance mutations and vaccine-induced escape on parasite

survival in the mosquito [31–35]. Our small sample size and especially the limited *Pf* genetic diversity found at our study site in Cambodia, where intensive drug pressure has selected for specific artemisinin-resistant subpopulations of parasites [36], may not reflect the range of selective factors that may impact upon parasite fitness in the mosquito.

Despite the efficiency of mosquito sampling, changing clone frequencies through transmission was commonplace. There was some indication that a lower gametocyte density translated into preferential transmission of the dominant haplotype and decreased MOI in individual mosquitoes. Although our sample size is too small to draw conclusions, it may be that transmission of genetic diversity is maximized in the majority of gametocyte carriers, until intensive malaria interventions result in a mostly asymptomatic reservoir of very low-density infections that allows intermittent transmission of a few dominant highly transmissible parasite strains [37–39].

There are methodological limitations that will be addressed in future iterations of this work. Amplicon sequencing of a single polymorphic marker may have underestimated the level of genetic diversity present in our samples. However, MOIs calculated by these methods previously aligned well with diversity estimates from genome-wide sequencing [1, 16, 24]. The use of different genetic markers between *Pf* and *Pv* requires caution when making any comparisons between species. We did not isolate and genotype single oocysts [5, 30, 40]. Future application of multilocus deep sequencing to single clonal oocysts could help determine which clones sampled from the blood are actually meiotic siblings related by descent and thus are more likely to have arisen through single inoculation rather than superinfection from multiple mosquito bites [41–43]. Findings may well differ with different vectors in Asia versus Africa. Finally, membrane feeding may not faithfully replicate mosquito sampling via skin feeding, which is likely to be even more efficient [44], although still ultimately limited by gametocyte numbers in the blood.

CONCLUSIONS

Overall, our detailed portrait of the transmission of multiple-clone *Pf/Pv* malaria infections from naturally infected Cambodian adults to *A. dirus* mosquitoes highlights the efficiency with which *Plasmodium* gametocytes maintain population-level genetic diversity through a vulnerable part of the malaria life cycle. Although we did observe changes in haplotype frequencies through transmission, overall, our data failed to show a selective bottleneck, although one may exist at lower gametocyte densities, such as those often found in asymptomatic carriers [45–47]. The methods shown here may yet reveal at what densities such a bottleneck appears and whether it leads to stochastic transmission of clones or a selective process that shapes population genetic diversity and response to malaria control interventions [34, 40]. Such tools and insights will

be central to understanding how changes in transmission intensity [48], genetic complexity [30], and parasite densities over time will affect the evolution of drug and vaccine resistance [31, 34, 40] and the composition of the human infectious reservoir [21, 47]. The limited data presented here support the notion that more readily transmitted *Pv* infections with preserved genetic diversity will ultimately prove the greater challenge for malaria elimination.

Supplementary Data

Supplementary materials are available at *The Journal of Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copyedited and are the sole responsibility of the authors, so questions or comments should be addressed to the corresponding author.

Notes

Acknowledgments. We thank the study participants as well as the Cambodian and Thai field staff who made the study possible.

Disclaimer. The views expressed in this manuscript are those of the authors and do not reflect official policy of the Department of the Army, Department of Defense, or the United States Government.

Financial support. This work was funded by National Institutes of Health Grant K08 AI110651 (to J. T. L.) and the American Society of Tropical Medicine and Hygiene and Burroughs Wellcome Fund (postdoctoral fellowship to J. T. L.).

Potential conflicts of interest. All authors: No reported conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest.

References

- Friedrich LR, Popovici J, Kim S, et al. Complexity of infection and genetic diversity in Cambodian *Plasmodium vivax*. *PLoS Negl Trop Dis* **2016**; 10:e0004526.
- Hong NV, Delgado-Ratto C, Thanh PV, et al. Population genetics of *Plasmodium vivax* in four rural communities in Central Vietnam. *PLoS Negl Trop Dis* **2016**; 10:e0004434.
- Imwong M, Nakeesathit S, Day NP, White NJ. A review of mixed malaria species infections in anopheline mosquitoes. *Malar J* **2011**; 10:253.
- Marchand RP, Culleton R, Maeno Y, Quang NT, Nakazawa S. Co-infections of *Plasmodium knowlesi*, *P. falciparum*, and *P. vivax* among humans and *Anopheles dirus* mosquitoes, Southern Vietnam. *Emerg Infect Dis* **2011**; 17:1232–9.
- Soontarawirat I, Andolina C, Paul R, et al. *Plasmodium vivax* genetic diversity and heterozygosity in blood samples and resulting oocysts at the Thai-Myanmar border. *Malar J* **2017**; 16:355.
- Smith RC, Vega-Rodríguez J, Jacobs-Lorena M. The *Plasmodium* bottleneck: malaria parasite losses in the mosquito vector. *Mem Inst Oswaldo Cruz* **2014**; 109:644–61.
- Zollner GE, Ponsa N, Garman GW, et al. Population dynamics of sporogony for *Plasmodium vivax* parasites from western Thailand developing within three species of colonized *Anopheles* mosquitoes. *Malar J* **2006**; 5:68.
- Lin JT, Ubalee R, Lon C, et al. Microscopic *Plasmodium falciparum* gametocytemia and infectivity to mosquitoes in Cambodia. *J Infect Dis* **2016**; 213:1491–4.
- Lin JT, Lon C, Spring MD, et al. Single dose primaquine to reduce gametocyte carriage and *Plasmodium falciparum* transmission in Cambodia: an open-label randomized trial. *PLoS One* **2017**; 12:e0168702.
- Chaorattanakawee S, Saunders DL, Sea D, et al. Ex vivo drug susceptibility and molecular profiling of clinical *Plasmodium falciparum* isolates from Cambodia in 2008–2013 suggest emerging piperazine resistance. *Antimicrob Agents Chemother* **2015**; 59:4631–43.
- Ribaut C, Berry A, Chevalley S, et al. Concentration and purification by magnetic separation of the erythrocytic stages of all human *Plasmodium* species. *Malar J* **2008**; 7:45.
- Musapa M, Mulenga M, Taida K, et al. A Simple chelex protocol for DNA extraction from *Anopheles* spp. *J Vis Exp* **2013**. Available at: <http://dx.doi.org/10.3791/3281>. Accessed 1 April 2019.
- Taylor SM, Juliano JJ, Trottman PA, et al. High-throughput pooling and real-time PCR-based strategy for malaria detection. *J Clin Microbiol* **2010**; 48:512–9.
- Veron V, Simon S, Carme B. Multiplex real-time PCR detection of *P. falciparum*, *P. vivax* and *P. malariae* in human blood samples. *Exp Parasitol* **2009**; 121:346–51.
- Chaorattanakawee S, Lanteri CA, Sundrakes S, et al. Attenuation of *Plasmodium falciparum* in vitro drug resistance phenotype following culture adaptation compared to fresh clinical isolates in Cambodia. *Malar J* **2015**; 14:486.
- Lin JT, Hathaway NJ, Saunders DL, et al. Using amplicon deep sequencing to detect genetic signatures of *Plasmodium vivax* relapse. *J Infect Dis* **2015**; 212:999–1008.
- Hathaway NJ, Parobek CM, Juliano JJ, Bailey JA. SeekDeep: single-base resolution de novo clustering for amplicon deep sequencing. *Nucleic Acids Res* **2018**; 46:e21.
- Vantaux A, Samreth R, Piv E, et al. Contribution to malaria transmission of symptomatic and asymptomatic parasite carriers in Cambodia. *J Infect Dis* **2018**; 217:1561–8.
- Kiattibutr K, Roobsoong W, Sriwichai P, et al. Infectivity of symptomatic and asymptomatic *Plasmodium vivax* infections to a Southeast Asian vector, *Anopheles dirus*. *Int J Parasitol* **2017**; 47:163–70.
- Koepfli C, Robinson LJ, Rarau P, et al. Blood-stage parasitaemia and age determine *Plasmodium falciparum* and *P. vivax* gametocytaemia in Papua New Guinea. *PLoS One* **2015**; 10:e0126747.
- Tadesse FG, Slater HC, Chali W, et al. The relative contribution of symptomatic and asymptomatic *Plasmodium vivax*

- and *Plasmodium falciparum* infections to the infectious reservoir in a low-endemic setting in Ethiopia. Clin Infect Dis **2018**; 66:1883–91.
22. Adapa SR, Taylor RA, Wang C, Thomson-Luque R, Johnson LR, Jiang RH. *Plasmodium vivax* readiness to transmit: implication for malaria eradication. BMC Syst Biol **2019**; 13:5.
 23. Bousema T, Drakeley C. Epidemiology and infectivity of *Plasmodium falciparum* and *Plasmodium vivax* gametocytes in relation to malaria control and elimination. Clin Microbiol Rev **2011**; 24:377–410.
 24. Parobek CM, Lin JT, Saunders DL, et al. Selective sweep suggests transcriptional regulation may underlie *Plasmodium vivax* resilience to malaria control measures in Cambodia. Proc Natl Acad Sci U S A **2016**; 113:E8096–105.
 25. Paul RE, Nu VA, Krettli AU, Brey PT. Interspecific competition during transmission of two sympatric malaria parasite species to the mosquito vector. Proc Biol Sci **2002**; 269:2551–7.
 26. Maeno Y, Quang NT, Culleton R, et al. Humans frequently exposed to a range of non-human primate malaria parasite species through the bites of *Anopheles dirus* mosquitoes in South-central Vietnam. Parasit Vectors **2015**; 8:376.
 27. Seyfarth M, Khairah BA, Abdi AA, Bouh SM, Faulde MK. Five years following first detection of *Anopheles stephensi* (Diptera: Culicidae) in Djibouti, Horn of Africa: populations established—malaria emerging. Parasitol Res **2019**. Available at: <https://doi.org/10.1007/s00436-019-06213-0>. Accessed 1 April 2019.
 28. Nwakanma D, Kheir A, Sowa M, et al. High gametocyte complexity and mosquito infectivity of *Plasmodium falciparum* in the Gambia. Int J Parasitol **2008**; 38:219–27.
 29. Grignard L, Gonçalves BP, Early AM, et al. Transmission of molecularly undetectable circulating parasite clones leads to high infection complexity in mosquitoes post feeding. Int J Parasitol **2018**; 48:671–7.
 30. Morlais I, Nsango SE, Toussile W, et al. *Plasmodium falciparum* mating patterns and mosquito infectivity of natural isolates of gametocytes. PLoS One **2015**; 10:e0123777.
 31. Mharakurwa S, Kumwenda T, Mkulama MA, et al. Malaria antifolate resistance with contrasting *Plasmodium falciparum* dihydrofolate reductase (DHFR) polymorphisms in humans and *Anopheles* mosquitoes. Proc Natl Acad Sci U S A **2011**; 108:18796–801.
 32. Goodman CD, Siregar JE, Mollard V, et al. Parasites resistant to the antimalarial atovaquone fail to transmit by mosquitoes. Science **2016**; 352:349–53.
 33. Conrad MD, Mota D, Musiime A, et al. Comparative prevalence of *Plasmodium falciparum* resistance-associated genetic polymorphisms in parasites infecting humans and mosquitoes in Uganda. Am J Trop Med Hyg **2017**; 97:1576–80.
 34. Sandeu MM, Abate L, Tchioffo MT, et al. Impact of exposure to mosquito transmission-blocking antibodies on *Plasmodium falciparum* population genetic structure. Infect Genet Evol **2016**; 45:138–44.
 35. Mharakurwa S, Sialumano M, Liu K, Scott A, Thuma P. Selection for chloroquine-sensitive *Plasmodium falciparum* by wild *Anopheles arabiensis* in Southern Zambia. Malar J **2013**; 12:453.
 36. Parobek CM, Parr JB, Brazeau NF, et al. Partner-drug resistance and population substructuring of artemisinin-resistant *Plasmodium falciparum* in Cambodia. Genome Biol Evol **2017**; 9:1673–86.
 37. Björkman AB. Asymptomatic low-density malaria infections: a parasite survival strategy? Lancet Infect Dis **2018**; 18:485–6.
 38. Nguyen TN, von Seidlein L, Nguyen TV, et al. The persistence and oscillations of submicroscopic *Plasmodium falciparum* and *Plasmodium vivax* infections over time in Vietnam: an open cohort study. Lancet Infect Dis **2018**; 18:565–72.
 39. Bousema T, Okell L, Felger I, Drakeley C. Asymptomatic malaria infections: detectability, transmissibility and public health relevance. Nat Rev Microbiol **2014**; 12:833–40.
 40. Paul RE, Packer MJ, Walmsley M, et al. Mating patterns in malaria parasite populations of Papua New Guinea. Science **1995**; 269:1709–11.
 41. Aydemir O, Janko M, Hathaway NJ, et al. Drug-resistance and population structure of *Plasmodium falciparum* across the Democratic Republic of Congo using high-throughput molecular inversion probes. J Infect Dis **2018**; 218:946–55.
 42. Nkhoma SC, Nair S, Cheeseman IH, et al. Close kinship within multiple-genotype malaria parasite infections. Proc Biol Sci **2012**; 279:2589–98.
 43. Wong W, Wenger EA, Hartl DL, Wirth DF. Modeling the genetic relatedness of *Plasmodium falciparum* parasites following meiotic recombination and cotransmission. PLoS Comput Biol **2018**; 14:e1005923.
 44. Lawniczak MK, Eckhoff PA. A computational lens for sexual-stage transmission, reproduction, fitness and kinetics in *Plasmodium falciparum*. Malar J **2016**; 15:487.
 45. Nguitragool W, Mueller I, Kumpitak C, et al. Very high carriage of gametocytes in asymptomatic low-density *Plasmodium falciparum* and *P. vivax* infections in western Thailand. Parasit Vectors **2017**; 10:512.
 46. Rovira-Vallbona E, Contreras-Mancilla JJ, Ramirez R, et al. Predominance of asymptomatic and sub-microscopic infections characterizes the *Plasmodium* gametocyte reservoir in the Peruvian Amazon. PLoS Negl Trop Dis **2017**; 11:e0005674.
 47. Tadesse FG, van den Hoogen L, Lanke K, et al. The shape of the iceberg: quantification of submicroscopic *Plasmodium falciparum* and *Plasmodium vivax* parasitaemia and gametocytaemia in five low endemic settings in Ethiopia. Malar J **2017**; 16:99.
 48. Churcher TS, Trape JF, Cohuet A. Human-to-mosquito transmission efficiency increases as malaria is controlled. Nat Commun **2015**; 6:6054.