Published in final edited form as:

Brain Res. 2020 January 01; 1726: 146502. doi:10.1016/j.brainres.2019.146502.

Increases in compulsivity, inflammation, and neural injury in HIV transgenic rats with escalated methamphetamine self-administration under extended-access conditions

Giordano de Guglielmo^{1,2,*}, Yu Fu^{3,4,*}, Jihuan Chen³, Estefania Larrosa¹, Ivy Hoang³, Tomoya Kawamura³, Irene Lorrai^{3,8}, Barry Zorman⁵, Joseph Bryant⁶, Olivier George², Pavel Sumazin⁵, Celine Lefebvre^{3,7}, Vez Repunte-Canonigo³, Pietro Paolo Sanna³

¹Department of Neuroscience, The Scripps Research Institute, La Jolla, CA, USA

²Department of Psychiatry, University of California San Diego, La Jolla, CA

³Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA, USA

⁴European Bioinformatics Institute (EMBL-EBI), Hinxton, United Kingdom

⁵Department of Pediatrics, Dan L. Duncan Cancer Center, Baylor College of Medicine, Houston, TX, USA

⁶University of Maryland and Institute of Human Virology, Baltimore, MD

⁷Bioinformatics and Computational Biology, Servier, Paris, France

⁸Department of Biomedical Sciences, University of Cagliari, Monserrato, Cagliari, Italy

Abstract

The abuse of stimulants, such as methamphetamine (METH), is associated with treatment non-compliance, a greater risk of viral transmission, and the more rapid clinical progression of immunological and central nervous system human immunodeficiency virus (HIV) disease. The behavioral effects of METH in the setting of HIV remain largely uncharacterized. We used a state-of-the-art paradigm of the escalation of voluntary intravenous drug self-administration in HIV transgenic (Tg) and wildtype rats. The rats were first allowed to self-administer METH under short-access (ShA) conditions, which is characterized by a nondependent and more “recreational” pattern of METH use, and then allowed to self-administer METH under long-access (LgA) conditions, which leads to compulsive (dependent) METH intake. HIV Tg and wildtype rats self-administered equal amounts of METH under ShA conditions. HIV Tg rats under LgA METH self-administration following a 4-week enforced abstinence period to model the intermittent pattern of stimulant abuse in humans, developed greater motivation to self-administer METH and self-administered larger amounts of METH. Impairments in function of the medial prefrontal cortex (mPFC) contribute to compulsive drug and alcohol intake. Gene expression profiling of the mPFC

Correspondence: psanna@scripps.edu, canonigo@scripps.edu.

*Contributed equally

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

in HIV Tg rats with a history of escalated METH self-administration under LgA conditions showed transcriptional evidence of increased inflammation, greater neural injury and impaired aerobic glucose metabolism than wildtype rats that self-administered METH under LgA conditions. The detrimental effects of the interaction between neuroHIV and escalated METH intake on the mPFC are likely key factors in the greater vulnerability to excessive drug intake in the setting of HIV.

Introduction

Human immunodeficiency virus (HIV)-infected individuals have a higher risk of substance abuse, such as methamphetamine, opioids, and alcohol, than the general U.S. population (Cook et al., 2001; Dansak, 1997; Kumar, 2014; Meyerhoff, 2001; Petry, 1999; Samet et al., 2004; Silverstein and Kumar, 2014). Substance abuse is associated with treatment non-compliance, a greater risk of viral transmission, and more rapid clinical progression of HIV disease (Bing et al., 2001; Cadet and Krasnova, 2007; Chana et al., 2006; Galvan et al., 2002; Goodkin et al., 1998; Lucas et al., 2001; Lucas et al., 2002; Lucas et al., 2006; Nath, 2010; Nelson et al., 2002; Scott et al., 2007; Volkow et al., 2007). The abuse of stimulants, such as METH and cocaine, appears to adversely affect cognition in the setting of HIV (Carey et al., 2006; Levine et al., 2006; Nath et al., 2001; Nath, 2010; Scott et al., 2007; Soontornniyomkij et al., 2016). Methamphetamine use is also associated with higher depression scores in HIV-infected individuals (Blackstone et al., 2013; Bousman et al., 2009).

The behavioral effects of METH in the setting of HIV remain largely uncharacterized. The clinical literature suggests that HIV may itself contribute to the perpetuation of METH abuse. HIV patients report using METH to self-medicate HIV-related symptoms (e.g., depression and fatigue) and improve social-cognitive functioning (Gorman, 1996; Homer et al., 2008; Robinson and Rempel, 2006; Semple et al., 2002).

To explore the effects of METH abuse in the setting of HIV, we used voluntary intravenous METH self-administration in HIV transgenic (Tg) rats. Most research to date has focused on the effects of the HIV proteins gp120 and Tat, but evidence indicates that other HIV proteins (e.g., gp41, Nef, and Vpr, among others) also possess neurotoxic potential (Adamson et al., 1999; Jones et al., 2007; Patel et al., 2002; Trillo-Pazos et al., 2000; van de Bovenkamp et al., 2002). Thus, we utilized HIV Tg rats that express multiple HIV products in disease-relevant glial cells, such as microglia and astrocytes, but not in neurons (Reid et al., 2001; Repunte-Canonigo et al., 2014b; Royal et al., 2012).

HIV Tg rats exhibit changes in gene expression that are consistent with HAND in humans (Repunte-Canonigo et al., 2014b; Sanna et al., 2017). HIV Tg rats exhibit deficits in working memory (Repunte-Canonigo et al., 2014b), spatial learning (Vigorito et al., 2007), and performance in a reversal learning task (Lashomb et al., 2009). This is consistent with previous observations that pathological changes that are associated with neuroAIDS, such as synaptodendritic injury, can be induced by HIV proteins in the absence of viral infection and replication (Iskander et al., 2004; Kim et al., 2003; Toggas et al., 1994).

The occasional but limited use of a drug is clinically distinct from escalated drug use, which is characterized by an inability to limit drug intake, the emergence of chronic compulsive drug-seeking behavior, and vulnerability to relapse after cessation (Koob and Le Moal, 2005). Initial and occasional drug use is motivated by a positive emotional state (e.g., euphoria) that is induced by the drug, also described by the process of positive reinforcement (Koob and Le Moal, 2005). Self-administration studies of the neuropharmacological basis of the acute reinforcing effects of drugs of abuse (positive reinforcement) have traditionally used models of limited daily access to drug self-administration (Koob et al., 2004). Using these models, rats and mice with indwelling intravenous catheters are allowed to self-administer drugs of abuse for less than 3 h/day, producing stable levels and patterns of intake. However, addiction is characterized by the loss of control in limiting intake and compulsion to take the drug (Koob and Le Moal, 2005). Animal models of extended daily access to intravenous drug self-administration have been developed for cocaine, heroin, alcohol, nicotine, and METH, which result in escalated levels of intake (Ahmed and Koob, 1998; Cohen et al., 2012; George et al., 2007a; George et al., 2007b; George et al., 2014; Kitamura et al., 2006; Koob et al., 2004; Koob and Le Moal, 2005; Orio et al., 2010).

The present study used intravenous METH self-administration initially under short-access (ShA) conditions, which is characterized by a nondependent “recreational-like” pattern of METH use, followed by METH self-administration under long-access (LgA) conditions, which leads to compulsive (dependent) METH intake. Animals that self-administer METH under LgA conditions exhibit an upward shift of the METH dose-response function, which are indicative of compulsive drug self-administration and working memory impairments (Recinto et al., 2012)(Kitamura et al., 2006; Orio et al., 2010; Wee et al., 2007). Additionally, intracranial self-stimulation (ICSS) reward thresholds gradually become elevated in animals that self-administer METH under LgA conditions and are correlated with an increase in METH intake (Jang et al., 2013). The latter finding indicates that extended access to METH recruits negative reinforcement mechanisms, in which a depressive-like state of withdrawal drives compulsive responding for the drug to relieve the depressive-like state, which is not observed with recreational drug use (Jang et al., 2013; Koob and Le Moal, 2006; Koob and Le Moal, 2005; Koob and Le Moal, 2008; Koob and Volkow, 2010). The model of extended-access self-administration that is used in our laboratory and by others has been shown to recapitulate all seven of the diagnostic criteria for Substance Use Disorder in the *Diagnostic and Statistical Manual of Mental Disorders*, 4th edition (DSM-IV; American Psychiatric Association, 1994), and seven of the 11 criteria in the DSM-5 (George et al., 2014).

Impairments in frontostriatal connectivity are involved in both the progression to compulsive drug intake and neuroHIV (Feil et al., 2010; Ipser et al., 2015; Plessis et al., 2014; Volkow and Morales, 2015), suggesting that METH and HIV can additively promote compulsive drug intake and cognitive impairment. Thus, to better understand the molecular basis of detrimental interactions between HIV and METH, we profiled gene expression from the medial prefrontal cortex (mPFC) in HIV Tg and wildtype (WT) rats with a history of METH self-administration under LgA conditions. Previous studies showed that changes in gene expression that are associated with compulsive cocaine, heroin, and alcohol self-

administration are considerably different from changes in gene expression that are induced by a moderate “recreational-like” pattern of self-administration (Ahmed et al., 2005; Chen et al., 2013; Francesconi et al., 2009; Koob et al., 2004; Repunte-Canonigo et al., 2007; Repunte-Canonigo et al., 2010a; Repunte-Canonigo et al., 2010b; Repunte-Canonigo et al., 2014a; Repunte-Canonigo et al., 2015).

In the present study, HIV Tg rats that self-administered METH under LgA conditions exhibited greater drug seeking and taking than WT rats. The mPFC gene expression analysis showed that HIV Tg rats that self-administered METH under LgA conditions exhibited evidence of increases in neuroinflammation and neurodegeneration, a decrease in the tricarboxylic acid (TCA) cycle, and an increase in glycolysis gene expression.

Collectively, the present results indicate that the combination of neuroHIV and escalated METH intake results in significant additive neurotoxicity in the mPFC. The detrimental effect of neuroHIV and escalated METH intake on the expression of genes that are related to immune system activation and inflammation, neural damage, and metabolism may be significant contributors to neural injury and hypofrontality, which are believed to be key factors in compulsive drug taking and cognitive dysfunction in the HIV setting.

Results

Increase in METH self-administration in HIV transgenic rats

HIV Tg rats were first trained to self-administer METH intravenously under ShA conditions and then under LgA conditions. HIV Tg rats exhibited similar patterns of acquisition and intake under ShA conditions but then greater compulsivity after the escalation of METH intake under LgA conditions. During the pre-escalation phase under ShA conditions (1 h/day access to METH self-administration), mixed factorial analysis of variance (ANOVA), with group as the between-subjects factor and session as the within-subjects factor, indicated no differences in METH intake between HIV Tg and littermate control WT rats during the pre-escalation phase ($F_{11,275} = 3.39$, $p > 0.05$; Fig. 1A).

The rats were then switched to LgA conditions (6 h/day access to METH self-administration), which induced the escalation of METH intake ($p < 0.01$, day 5 vs. day 1 in the WT group; $p < 0.01$, day 10 vs. day 1 in the HIV Tg group; separate one-way ANOVAs followed by Fisher's Least Significant Difference [LSD] *post hoc* test). Mixed factorial ANOVA, with group as the between-subjects factor and session as the within-subjects factor, showed a significant group $\times$ session interaction ($F_{15,330} = 8.48$, $p < 0.001$). Fisher's LSD *post hoc* test showed significant differences in the number of METH infusions between WT and HIV Tg rats between sessions 5 and 8 ($p < 0.01$).

After re-escalation following a 4-week enforced abstinence period, during which METH was unavailable to model the intermittent pattern of stimulant abuse in humans, mixed factorial ANOVA, with group as the between-subjects factor and session as the within-subjects factor, showed a significant group $\times$ session interaction ($F_{7,147} = 3.263$, $p < 0.01$). Fisher's LSD *post hoc* test showed a significant increase in METH intake in HIV Tg rats vs. WT rats starting from session 5 ($p < 0.01$). A progressive-ratio (PR) test was conducted to measure the

motivation for METH intake following the escalation of METH self-administration (before re-escalation). HIV Tg rats exhibited an increase in compulsive-like responding for METH under the PR schedule of reinforcement ($t_{22} = 2.402$, $p < 0.05$).

Overall, HIV Tg rats developed greater compulsivity following the escalation of METH intake under LgA conditions after re-exposure to METH self-administration following a 4-week enforced abstinence period, reflected by greater responding under both fixed-ratio (FR) and PR schedules of reinforcement. These results indicate that HIV interacts with the effects of a history of binge-abstinence patterns of METH intake to increase vulnerability to relapse and motivation for the drug.

Gene expression profiling in the mPFC in HIV Tg and WT rats that self-administered METH under LgA conditions

Functional impairments in the mPFC contribute to compulsive drug intake. Therefore, we profiled gene expression in the mPFC in HIV Tg rats with a history of escalated METH self-administration (Fig. 2). Among the differentially expressed genes were F-Box and WD repeat domain containing 11 (FBXW11; also known as β -transducin repeat containing protein 2 [β TrCP2], homolog of Slimb [HOS]), interferon α (IFN α) inducible protein 27 (IFI27; (Pulliam, 2014), ubiquitin domain containing 2 (UBTD2; (Song et al., 2010), and potassium voltage-gated channel interacting protein 1 (KCNIP1 or KChIP1; (An et al., 2000); Fig. 2A, B). The expression of FBXW11 significantly decreased in HIV Tg rats with a history of escalated METH self-administration (Fig. 2A, B). FBXW11 is an E3 ubiquitin ligase that is implicated in downregulating the interferon type I receptor (IFNAR1) in response to the induction of IFN α (Kumar et al., 2003) and involved in the transcriptional activation of nuclear factor- κ B (NF- κ B) by promoting the degradation of the inhibitor of κ B (I κ B; (Suzuki et al., 2000). The expression of IFI27 significantly increased in HIV Tg rats with a history of escalated METH self-administration (Fig. 2A, B). IFI27 is a gene that mediates antiviral activity in response to neurotropic viruses (Cho et al., 2013) and is a proposed biomarker of monocyte activation (Pulliam, 2014). UBTD2 is a ubiquitin-like protein with an unknown function that can bind ubiquitin, indicating that it may play a role in regulating the ubiquitination of protein substrates (Song et al., 2010). The expression of KCNIP1 significantly increased in HIV Tg rats with a history of escalated METH self-administration (Fig. 2A, B). KCNIP1 is a member of a family of small cytosolic, calcium-binding proteins that were initially identified as subunits of K_v4 potassium channels, which regulate the gating of voltage-gated A-type potassium currents in neuronal tissue and the heart (An et al., 2000). The association between KCNIP1 and K_v4 results in an increase in current, the acceleration of recovery from inactivation, slower inactivation kinetics, and lower neuronal excitability (Gonzalez et al., 2014). KCNIP1 was shown to promote the stronger inhibitory control of firing during sustained activity (Bourdeau et al., 2011). Top differentially expressed genes in HIV Tg rats that self-administered METH under LgA conditions are shown in Fig. 2C. Top differentially expressed pathways are shown in Fig. 2D. Gene expression differences between HIV Tg and WT rats are in part preexisting, consistent with earlier observations (Repunte-Canonigo et al. 2014) and partially attributable to the interaction between HIV and a history of METH intake and re-escalation.

Transcriptional evidence of an increase in neuroinflammation in HIV Tg rats that self-administered METH under LgA conditions

HIV Tg and WT rats that self-administered METH under LgA conditions exhibited a prominent differential expression of pathways related to immune activation and inflammation. These include pathways that are indicative of general immune activation, inflammation, and greater cytokine signaling (Fig. 3A–D). Among the activated pathways were pathways that are related to tumor necrosis factor α (TNF- α) and inflammasome activation (Fig. 3E–G) and the recruitment of specific cytokine signaling pathways, including interleukin-1 β (IL-1 β , a pivotal proinflammatory cytokine that is released following inflammasome activation; Fig. 3D), IL-2, and IL-3 signaling (Fig. 3H, I). IL-3 is associated with active inflammatory processes in the brain (Renner et al., 2016) and has been suggested to induce a neuroprotective phenotype in microglia (Choudhury et al., 2011). Lastly, HIV Tg and WT rats that self-administered METH under LgA conditions exhibited a more pronounced induction of both type I and II IFN signaling (Fig. 3J–L).

Transcriptional evidence of increase in neural injury in HIV Tg rats that self-administered METH under LgA conditions

Differences in the regulation of pathways that are involved in neurodegeneration between HIV Tg and WT rats that self-administered METH under LgA conditions suggest the differential activation of pathogenic mechanisms (Fig. 4). These mechanisms involved the greater induction of pathways that are implicated in neuronal damage, such as genes related to complement (Fig. 4A), amyotrophic lateral sclerosis (Fig. 4B), and the activation of glucocorticoid signaling (Fig. 4C). Escalated METH self-administration interacted with HIV to affect the expression of genes that are involved in the extracellular matrix (Fig. 4D–F) and differential expression of genes related to DNA damage and repair (Fig. 4G–I).

Differences in the regulation of pathways that are involved in metabolism between HIV Tg and WT rats that self-administered METH under LgA conditions were also observed (Fig. 5). Pathway analysis indicated a decrease in genes related to glucose oxidation through the tricarboxylic acid (TCA) cycle and an increase in glycolysis in HIV Tg rats that self-administered METH under LgA conditions compared with METH-self-administering WT rats (Fig. 5).

Pathway analysis also indicated a greater catabolic molecular environment in HIV Tg rats that self-administered METH under LgA conditions (Fig. 6). The gene expression analysis in METH-self-administering HIV Tg rats suggested lower protein synthesis, higher protein degradation (Fig. 4A–C), and impairments in RNA synthesis and processing (Fig. 6D–F).

Discussion

The present study used paradigms of voluntary intravenous METH self-administration under ShA conditions (characterized by a “recreational” pattern of use) followed by LgA conditions (which lead to compulsive/dependent intake; (Kitamura et al., 2006; Orio et al., 2010; Recinto et al., 2012; Wee et al., 2007) in HIV Tg and WT rats. HIV Tg and WT rats self-administered equal amounts of METH under ShA conditions. However, HIV Tg rats

self-administered greater amounts of METH and exhibited greater motivation for METH than WT rats under LgA METH self-administration following a 4-week enforced abstinence period to model the intermittent pattern of stimulant abuse in humans.

HIV Tg rats are a small-animal model of concomitantly low levels of expression of multiple HIV products in disease-relevant cells in the central nervous system, such as microglia and astrocytes, but not in neurons (Reid et al., 2001; Repunte-Canonigo et al., 2014b; Royal et al., 2012), which is a key characteristic of HAND in the setting of viral suppression. HIV Tg rats differ from most of the currently available animal models of HIV that usually focus on a single viral protein. HIV Tg rats exhibit changes in gene expression that are consistent with human neuroHIV (Repunte-Canonigo et al., 2014b; Sanna et al., 2017).

The present model of the escalation of intravenous METH self-administration has not been previously utilized in the HIV/AIDS field. This is an innovative paradigm that was developed at The Scripps Research Institute, which effectively models the negative reinforcement that is associated with escalated drug intake but is not observed with recreational drug use (Ahmed and Koob, 1998; Jang et al., 2013; Koob and Le Moal, 2006; Koob and Le Moal, 2005; Koob and Le Moal, 2008; Koob and Volkow, 2010) and is associated with cognitive impairment and a depressive-like state during withdrawal (Ahmed et al., 2002; Jang et al., 2013; Koob and Kreek, 2007; Koob, 2009; Koob and Volkow, 2010; Recinto et al., 2012). Negative reinforcement is characterized by a depressive-like state during abstinence that drives compulsive drug seeking to relieve the negative emotional state (Ahmed et al., 2002; Jang et al., 2013; Koob and Le Moal, 2006; Koob and Le Moal, 2005; Koob and Le Moal, 2008; Koob and Volkow, 2010). In this paradigm, cognitive deficits that involve impairments in mPFC function emerge during the transition to escalated drug intake and contribute to compulsive drug seeking and taking (Jang et al., 2013; Recinto et al., 2012). Hypofrontality characterizes both compulsive drug intake in dependent individuals and cognitive dysfunction in neuroHIV (Feil et al., 2010; Goldstein and Volkow, 2002; Goldstein and Volkow, 2011; Ipser et al., 2015; Plessis et al., 2014; Volkow et al., 1992a; Volkow et al., 1992b; Volkow and Morales, 2015; Volkow et al., 2015; Wiers et al., 2016), suggesting a neurobiological basis for the additive effects of METH and HIV on compulsive drug intake, as seen in the present study, as well as cognitive impairment.

The identification of factors that promote relapse or resilience is of particular interest (Goldstein and Volkow, 2011). HIV Tg rats in the present study developed greater compulsivity to self-administer METH following the re-escalation (post-abstinence relapse) of METH intake after a 4-week enforced abstinence period, despite acquiring escalated METH self-administration behavior later than their WT littermates. When access to METH self-administration was restored after the 4-week respite, HIV Tg rats exhibited greater METH intake on both FR 8 and PR schedules of reinforcement. Binging patterns of stimulant abuse are prevalent in certain segments of people living with HIV (Harzke et al., 2009; Roy et al., 2017; Wechsberg et al., 2003). Studies in both rats and humans indicate that the motivational impact of drug-associated cues on behavior becomes progressively stronger over periods of repeated abstinence, referred to as the “incubation of craving,” which increases the vulnerability to relapse and represents a challenge to successful treatment (Grimm et al., 2001; Parvaz et al., 2016). Increases in the incubation of cue-

induced craving were found to predict the rate and extent of the re-escalation of cocaine intake after a 4-week period of enforced abstinence in rats (Guillem and Ahmed, 2018). In the present study, HIV Tg and WT littermate controls did not differ in their levels of drug intake before the period of enforced abstinence. Thus, the present results indicate that HIV interacts with the effects of bingeing-abstinence patterns of METH intake to increase the vulnerability to relapse and motivation for the drug.

To investigate the molecular mechanisms that underlie interactions between HIV and METH, we performed gene expression profiling of the mPFC, which has been consistently implicated in craving and the escalation of drug taking (Du et al., 2018; George et al., 2008; Goldstein and Volkow, 2011). Here, we show that the increase in METH self-administration in HIV Tg rats under LgA conditions was associated with an increase in inflammation. We found a global and pronounced increase in the activation of immune and inflammation-related pathways, including pathways that are related to cytokine and both type I and type II IFN signaling, which have been implicated in neural cell injury and cognitive impairment in HIV (Pulliam, 2014).

Inflammation has been associated with the aging brain and degenerative diseases (Mejias et al., 2018; Youm et al., 2013). We observed gene expression evidence of greater neural injury and neurodegeneration in HIV Tg rats vs. WT rats that self-administered METH under LgA conditions. This included pathways that are involved in aging and neurodegenerative diseases (e.g., Alzheimer's disease), including glucocorticoid receptor-regulated genes (Djamshidian and Lees, 2014; Hartmann et al., 1997; Hou et al., 2014; Notarianni, 2013; Vyas and Maatouk, 2013), and pathways that are involved in extracellular matrix deposition (Zhang et al., 2014). Previous studies showed that stimulants, such as cocaine and METH, affect extracellular matrix heparan sulfate content and sulfation levels (Chen et al., 2013; Chen et al., 2017). Heparan sulfate has also been implicated in amyloid- β peptide and tau fibrillization in Alzheimer's disease (Liu et al., 2016).

We observed a reduction of the expression of TCA cycle-related genes, with an increase in glycolysis-related genes in HIV Tg rats with a history of METH self-administration under LgA conditions compared with WT rats that self-administered METH under the same conditions. A decrease in oxidative phosphorylation reflects mitochondrial impairment that leads to less efficient energy metabolism. HIV has been shown to variably affect the TCA cycle in different cell types. The downregulation of TCA proteins has been observed in major blood types of HIV non-progressors (Zhang et al., 2017). The decrease in oxidative phosphorylation may reflect a hypometabolic state that is associated with neurodegenerative alterations in neurons (Butterfield and Halliwell, 2019). Higher lactate levels were seen in the brain in HIV patients in the pre-cART era, attributed to greater anaerobic glycolytic demands of macrophages associated with inflammation (Barker et al., 1995; Simone et al., 1998).

We also observed an apparent increase in the catabolic milieu that was elicited by the interaction between METH and HIV. We observed a reduction of the transcription of genes that are related to RNA processing and a reduction of genes that are related to translation. The dysregulation of translation is emerging as a possible mechanism of

neurodevelopmental disease and neurodegeneration (Darnell and Klann, 2013; Halliday et al., 2017; Hetman and Slomnicki, 2019).

In conclusion, HIV Tg rats self-administered more METH than WT rats under extended-access conditions, leading to an increase in the escalation of METH intake following an intermittent pattern of access to the drug. Dysfunction of the mPFC contributes to compulsivity in the present paradigm. Prefrontal cortex gene expression analyses indicated that HIV Tg rats that escalated their METH intake exhibited transcriptional evidence of increases in neuroinflammation and neurodegeneration and alterations of energy metabolism. These alterations may be key contributors to hypofrontality that contributes to the vulnerability to excessive drug intake in the setting of dependence.

Materials and Methods

Animals

Male HIV Tg rats ($n = 13$) and littermate control WT rats ($n = 14$) on a mixed Fischer/Wistar background (originally on a Fischer background, crossed for three generations with Wistar rats) were housed two per cage on a reverse 12 h/12 h light/dark cycle (lights off at 8:00 AM) in a temperature (20-22°C) and humidity (45-55%) controlled vivarium with *ad libitum* access to tap water and food pellets (PJ Noyes, Lancaster, NH, USA). All of the procedures were conducted in strict adherence to the National Institutes of Health *Guide for the Care and Use of Laboratory Animals* and were approved by the Institutional Animal Care and Use Committee of The Scripps Research Institute. At the time of testing, the rats' body weights ranged between 350 and 400 g.

Intravenous catheterization

The animals were anesthetized by the inhalation of a mixture of isoflurane, and intravenous catheters were aseptically inserted in the right jugular vein using a modified version of a procedure that was described previously (Caine and Koob, 1993; de Guglielmo et al., 2013). The vein was punctured with a 22-gauge needle, and the tubing was inserted and secured inside the vein by tying the vein with suture thread. The catheter assembly consisted of an 18 cm length of Micro-Renathane tubing (0.023 inch inner diameter, 0.037 inch outer diameter; Braintree Scientific, Braintree, MA, USA) that was attached to a guide cannula (Plastics One, Roanoke, VA, USA). The guide cannula was bent at a near right angle, embedded in dental acrylic, and anchored with 2 cm square mesh. The catheter exited through a small incision on the back, and the base was sealed with a small plastic cap and metal cover cap. This design helped keep the catheter base sterile and protected. The catheters were flushed daily with heparinized saline (10 U/ml of heparin sodium; American Pharmaceutical Partners, Schaumburg, IL, USA) in 0.9% bacteriostatic sodium chloride (Hospira, Lake Forest, IL, USA) that contained 20 mg/0.2 ml of the antibiotic Timentin (GlaxoSmithKline, UK).

Drugs

Methamphetamine HCl (National Institute on Drug Abuse, Bethesda, MD, USA) was dissolved in 0.9% saline (Hospira, Lake Forest, IL, USA) and self-administered intravenously at a dose of 0.05 mg/kg/infusion.

Operant training

Self-administration was performed in operant conditioning chambers (Med Associates, St. Albans, VT, USA) that were enclosed in lit, sound-attenuating, ventilated environmental cubicles. The front door and back wall of the chambers were constructed of transparent plastic, and the other walls were opaque metal. Each chamber was equipped with two retractable levers that were located on the front panel. Methamphetamine was delivered through plastic catheter tubing that was connected to an infusion pump. The infusion pump was activated by responses on the right (active) lever. Responses on the left (inactive) lever were recorded but did not have any scheduled consequences. Activation of the pump resulted in the delivery of 0.1 ml of the drug solution. A computer controlled fluid delivery and behavioral data recording.

All of the rats were tested for METH self-administration on a continuous reinforcement (FR1) schedule in three successive phases: pre-escalation, escalation, and re-escalation. Each active lever press resulted in the delivery of one METH dose (0.5 mg/kg/0.1 ml infusion). A 20-s timeout (TO) period followed each METH infusion. During the timeout period, responses on the active lever did not have scheduled consequences. This TO period occurred concurrently with the illumination of a cue light that was located above the active lever to signal delivery of the positive reinforcement.

At the end of the escalation phase, the rats were also tested on a PR schedule of reinforcement. Under these conditions, the response requirements that were necessary to receive a single drug dose increased according to the following equation: $(5e^{[\text{injection numbers} \times 0.2]}) - 5$. This resulted in the following progression of response requirements: 1, 2, 4, 6, 9, 12, 15, 20, 25, 32, 40, 50, 62, 77, 95, 118, 145, 178, 219, 268, etc. The breakpoint was defined as the last ratio that was attained by the rat prior to a 60 min period during which a ratio was not completed.

Pre-escalation phase: ShA.—During this phase, the rats were trained to acquire intravenous METH self-administration. All of the animals were allowed access to an FR1 TO 20 s schedule of METH self-administration with 1 h access to METH for 12 consecutive sessions (0.5 mg/kg/0.1 ml infusion).

Escalation phase. LgA.—This phase began 2 days after the end of the pre-escalation phase. During the escalation phase, drug access was increased to 6 h per LgA session. This phase lasted for 15 consecutive sessions and was followed by a PR test.

Re-escalation phase.—After the first escalation, the animals had no access to METH for 4 weeks. Following these 4 weeks of abstinence, the animals were allowed to self-administer METH in 6 h sessions according to the re-escalation protocol (8 sessions).

Total RNA isolation and RNA sequencing

Microdissected tissue from the mPFC was processed for total RNA isolation using the miRNeasy kit (Qiagen) and Zymo purification kit (Zymo Research). Libraries were prepared with the KAPA mRNA HyperPrep Kit for Illumina sequencing (KAPABiosystems) for mRNA capture with magnetic oligo-dT beads, cDNA synthesis, and library construction and amplification. The Poly-A libraries were subsequently sequenced on an Illumina HiSeq4000 sequencer at 30-million-read target coverage (100 bp paired-end reads).

Gene expression profiling and gene set enrichment analysis

The sequences were first trimmed using Trimmomatic with default setting (version 0.33). All of the samples passed the quality control by fastQC (version 0.11.3). Fastq files were aligned to combined rat (Rnor_6.0) and HIV-1 (NC_001802) transcriptomes on 8/8/2017 using Bowtie2 (Langmead et al., 2009) with default settings. Transcript expression was normalized using RSEM (version 1.3.0; (Li and Dewey, 2011)). The two transcriptomes were combined as described previously (Fu et al., 2019). The rat genome was humanized using the biomaRt package from R. Only genes with the highest counts were kept when multiple rat genes matched to one human gene. Differential expression analysis was performed using the DESeq2 package in R (the Wald method was used). Gene set enrichment analysis (GSEA; (Subramanian et al., 2005) was performed in R for MSigDB curated gene sets but excluding perturbation-based gene sets for a total of 1,452 well-known MSigDB gene sets. Multiple testing adjustment was performed using the False Discovery Rate. Fastq files were deposited in the European Nucleotide Archive project PRJEB33332.

Quantitative PCR validation

Differentially expressed genes in the mPFC were validated in HIV and WT rats with or without a history of METH exposure using the SYBR Green fluorescence detection kit on an iQ5 instrument (Bio-Rad). A set of optimized real-time polymerase chain reaction (PCR) primer assays was designed, and the following sequences were used for the following targets: FBXW11 (AGCCCTGCTTCTGATTTACG [sense] and GAAGACTCTGGGTAGTCGTTTG [antisense]), UBD2 (GGACTATGTGGTGCAGGTTAT [sense] and TAACAATAGAGCGGAGCCAAG [antisense]), KCNIP1 (ACGAATTTCTCGAGTCCTGTC [sense] and GCCAGTGTCTCAGTTACAT [antiSense]), IFI27 (CAGCCAAGCTGATGTCTGT [sense] and AGGACCTTAGATGTCATGGAGA [antisense]), and β -actin (AGATTACTGCCCTGGCTCCT [sense] and CAGTGAGGCCAGGATAGAGC [antisense]). Gene expression was normalized to β -actin and analyzed based on the CT method.

Acknowledgements

Supported by National Institutes of Health grants DA041750, DA043268, DA046170, DA046204, and DA048882. The authors would like to thank Michael Arends for assistance with manuscript preparation.

References

- Adamson DC, Kopnisky KL, Dawson TM, Dawson VL, 1999 Mechanisms and structural determinants of HIV-1 coat protein, gp41-induced neurotoxicity. *J Neurosci.* 19, 64–71. [PubMed: 9870939]
- Ahmed SH, Koob GF, 1998 Transition from moderate to excessive drug intake: change in hedonic set point. *Science.* 282, 298–300. [PubMed: 9765157]
- Ahmed SH, Kenny PJ, Koob GF, Markou A, 2002 Neurobiological evidence for hedonic allostasis associated with escalating cocaine use. *Nat Neurosci.* 5, 625–6. [PubMed: 12055635]
- Ahmed SH, Lutjens R, van der Stap LD, Lekic D, Romano-Spica V, Morales M, Koob GF, Repunte-Canonigo V, Sanna PP, 2005 Gene expression evidence for remodeling of lateral hypothalamic circuitry in cocaine addiction. *Proc Natl Acad Sci U S A.* 102, 11533–8. [PubMed: 16076954]
- An WF, Bowlby MR, Betty M, Cao J, Ling HP, Mendoza G, Hinson JW, Mattsson KI, Strassle BW, Trimmer JS, Rhodes KJ, 2000 Modulation of A-type potassium channels by a family of calcium sensors. *Nature.* 403, 553–6. [PubMed: 10676964]
- Barker PB, Lee RR, McArthur JC, 1995 AIDS dementia complex: evaluation with proton MR spectroscopic imaging. *Radiology.* 195, 58–64. [PubMed: 7892496]
- Bing EG, Burnam MA, Longshore D, Fleishman JA, Sherbourne CD, London AS, Turner BJ, Eggan F, Beckman R, Vitiello B, Morton SC, Orlando M, Bozzette SA, Ortiz-Barron L, Shapiro M, 2001 Psychiatric disorders and drug use among human immunodeficiency virus-infected adults in the United States. *Arch Gen Psychiatry.* 58, 721–8. [PubMed: 11483137]
- Blackstone K, Iudicello JE, Morgan EE, Weber E, Moore DJ, Franklin DR, Ellis RJ, Grant I, Woods SP, Translational Methamphetamine ARCG, 2013 Human immunodeficiency virus infection heightens concurrent risk of functional dependence in persons with long-term methamphetamine use. *J Addict Med.* 7, 255–63. [PubMed: 23648641]
- Bourdeau ML, Laplante I, Laurent CE, Lacaille JC, 2011 KChIP1 modulation of Kv4.3-mediated A-type K(+) currents and repetitive firing in hippocampal interneurons. *Neuroscience.* 176, 173–87. [PubMed: 21129448]
- Bousman CA, Cherner M, Ake C, Letendre S, Atkinson JH, Patterson TL, Grant I, Everall IP, 2009 Negative mood and sexual behavior among non-monogamous men who have sex with men in the context of methamphetamine and HIV. *J Affect Disord.* 119, 84–91. [PubMed: 19419773]
- Brew BJ, 2004 Evidence for a change in AIDS dementia complex in the era of highly active antiretroviral therapy and the possibility of new forms of AIDS dementia complex. *Aids.* 18 Suppl 1, S75–8. [PubMed: 15075501]
- Butterfield DA, Halliwell B, 2019 Oxidative stress, dysfunctional glucose metabolism and Alzheimer disease. *Nat Rev Neurosci.* 20, 148–160. [PubMed: 30737462]
- Cadet JL, Krasnova IN, 2007 Interactions of HIV and methamphetamine: cellular and molecular mechanisms of toxicity potentiation. *Neurotox Res.* 12, 181–204. [PubMed: 17967742]
- Caine SB, Koob GF, 1993 Modulation of cocaine self-administration in the rat through D-3 dopamine receptors. *Science.* 260, 1814–6. [PubMed: 8099761]
- Carey C, Woods S, Rippeth J, Gonzalez R, Heaton R, Grant I, 2006 Additive deleterious effects of methamphetamine dependence and immunosuppression on neuropsychological functioning in HIV infection. *AIDS Behav.* 3, 10, 185–90. [PubMed: 16477511]
- Chana G, Everall I, Crews L, Langford D, Adame A, Grant I, Cherner M, Lazzaretto D, Heaton R, Ellis R, Masliah E, Group H, 2006 Cognitive deficits and degeneration of interneurons in HIV+ methamphetamine users. *Neurology.* 67, 1486–9. [PubMed: 17060582]
- Chen J, Repunte-Canonigo V, Kawamura T, Lefebvre C, Shin W, Howell LL, Hemby SE, Harvey BK, Califano A, Morales M, Koob GF, Sanna PP, 2013 Hypothalamic proteoglycan syndecan-3 is a novel cocaine addiction resilience factor. *Nat Commun.* 4, 1955. [PubMed: 23736082]
- Chen J, Kawamura T, Sethi MK, Zaia J, Repunte-Canonigo V, Sanna PP, 2017 Heparan sulfate: Resilience factor and therapeutic target for cocaine abuse. *Sci Rep.* 7, 13931. [PubMed: 29066725]
- Cho H, Proll SC, Szretter KJ, Katze MG, Gale M Jr., Diamond MS, 2013 Differential innate immune response programs in neuronal subtypes determine susceptibility to infection in the brain by positive-stranded RNA viruses. *Nat Med.* 19, 458–64. [PubMed: 23455712]

- Choudhury ME, Sugimoto K, Kubo M, Nagai M, Nomoto M, Takahashi H, Yano H, Tanaka J, 2011 A cytokine mixture of GM-CSF and IL-3 that induces a neuroprotective phenotype of microglia leading to amelioration of (6-OHDA)-induced Parkinsonism of rats. *Brain Behav.* 1, 26–43. [PubMed: 22398979]
- Cohen A, Koob GF, George O, 2012 Robust escalation of nicotine intake with extended access to nicotine self-administration and intermittent periods of abstinence. *Neuropsychopharmacology.* 37, 2153–60. [PubMed: 22549121]
- Cook RL, Sereika SM, Hunt SC, Woodward WC, Erlen JA, Conigliaro J, 2001 Problem drinking and medication adherence among persons with HIV infection. *J Gen Intern Med.* 16, 83–8. [PubMed: 11251758]
- Cysique LA, Maruff P, Brew BJ, 2004 Prevalence and pattern of neuropsychological impairment in human immunodeficiency virus-infected/acquired immunodeficiency syndrome (HIV/AIDS) patients across pre- and post-highly active antiretroviral therapy eras: a combined study of two cohorts. *J Neurovirol.* 10, 350–7. [PubMed: 15765806]
- Dansak DA, 1997 Medical use of recreational drugs by AIDS patients. *J Addict Dis.* 16, 25–30.
- Darnell JC, Klann E, 2013 The translation of translational control by FMRP: therapeutic targets for FXS. *Nat Neurosci.* 16, 1530–6. [PubMed: 23584741]
- de Guglielmo G, Cipitelli A, Somaini L, Gerra G, Li H, Stopponi S, Ubaldi M, Kallupi M, Ciccocioppo R, 2013 Pregabalin reduces cocaine self-administration and relapse to cocaine seeking in the rat. *Addict Biol.* 18, 644–53. [PubMed: 22734646]
- Djamshidian A, Lees AJ, 2014 Can stress trigger Parkinson's disease? *J Neurol Neurosurg Psychiatry.* 85, 878–81. [PubMed: 24259593]
- Du C, Volkow ND, You J, Park K, Allen CP, Koob GF, Pan Y, 2018 Cocaine-induced ischemia in prefrontal cortex is associated with escalation of cocaine intake in rodents. *Mol Psychiatry.*
- Everall I, Vaida F, Khanlou N, Lazzaretto D, Achim C, Letendre S, Moore D, Ellis R, Cherne M, Gelman B, Morgello S, Singer E, Grant I, Masliah E, 2009 Cliniconeuropathologic correlates of human immunodeficiency virus in the era of antiretroviral therapy. *J Neurovirol.* 1–11. [PubMed: 19462266]
- Feil J, Sheppard D, Fitzgerald PB, Yucel M, Lubman DI, Bradshaw JL, 2010 Addiction, compulsive drug seeking, and the role of frontostriatal mechanisms in regulating inhibitory control. *Neurosci Biobehav Rev.* 35, 248–75. [PubMed: 20223263]
- Francesconi W, Berton F, Repunte-Canonigo V, Hagihara K, Thurbon D, Lekic D, Specio SE, Greenwell TN, Chen SA, Rice KC, Richardson HN, O'Dell LE, Zorrilla EP, Morales M, Koob GF, Sanna PP, 2009 Protracted withdrawal from alcohol and drugs of abuse impairs long-term potentiation of intrinsic excitability in the juxtacapsular bed nucleus of the stria terminalis. *J Neurosci.* 29, 5389–401. [PubMed: 19403807]
- Fu Y, Zorman B, Sumazin P, Sanna PP, Repunte-Canonigo V, 2019 Epitranscriptomics: Correlation of N6-methyladenosine RNA methylation and pathway dysregulation in the hippocampus of HIV transgenic rats. *PloS one.* 14, e0203566. [PubMed: 30653517]
- Galvan FH, Bing EG, Fleishman JA, London AS, Caetano R, Burnam MA, Longshore D, Morton SC, Orlando M, Shapiro M, 2002 The prevalence of alcohol consumption and heavy drinking among people with HIV in the United States: results from the HIV Cost and Services Utilization Study. *J Stud Alcohol.* 63, 179–86. [PubMed: 12033694]
- George O, Ghosland S, Azar MR, Cottone P, Zorrilla EP, Parsons LH, O'Dell LE, Richardson HN, Koob GF, 2007a CRF-CRF1 system activation mediates withdrawal-induced increases in nicotine self-administration in nicotine-dependent rats. *Proc Natl Acad Sci U S A.* 104, 17198–203. [PubMed: 17921249]
- George O, Mandyam CD, Wee S, Koob GF, 2007b Extended Access to Cocaine Self-Administration Produces Long-Lasting Prefrontal Cortex-Dependent Working Memory Impairments. *Neuropsychopharmacology.*
- George O, Mandyam CD, Wee S, Koob GF, 2008 Extended access to cocaine self-administration produces long-lasting prefrontal cortex-dependent working memory impairments. *Neuropsychopharmacology.* 33, 2474–82. [PubMed: 18033234]

- George O, Koob GF, Vendruscolo LF, 2014 Negative reinforcement via motivational withdrawal is the driving force behind the transition to addiction. *Psychopharmacology (Berl)*.
- Goldstein RZ, Volkow ND, 2002 Drug addiction and its underlying neurobiological basis: neuroimaging evidence for the involvement of the frontal cortex. *Am J Psychiatry*. 159, 1642–52. [PubMed: 12359667]
- Goldstein RZ, Volkow ND, 2011 Dysfunction of the prefrontal cortex in addiction: neuroimaging findings and clinical implications. *Nat Rev Neurosci*. 12, 652–69. [PubMed: 22011681]
- Gonzalez WG, Pham K, Miksovska J, 2014 Modulation of the voltage-gated potassium channel (Kv4.3) and the auxiliary protein (KChIP3) interactions by the current activator NS5806. *J Biol Chem*. 289, 32201–13. [PubMed: 25228688]
- Goodkin K, Shapshak P, Metsch LR, McCoy CB, Crandall KA, Kumar M, Fujimura RK, McCoy V, Zhang BT, Reyblat S, Xin KQ, Kumar AM, 1998 Cocaine abuse and HIV-1 infection: epidemiology and neuropathogenesis. *J Neuroimmunol*. 83, 88–101. [PubMed: 9610677]
- Gorman M, 1996 Speed use and HIV transmission. *Focus*. 11, 4–6.
- Grimm JW, Hope BT, Wise RA, Shaham Y, 2001 Neuroadaptation. Incubation of cocaine craving after withdrawal. *Nature*. 412, 141–2. [PubMed: 11449260]
- Guillem K, Ahmed SH, 2018 Incubation of Accumbal Neuronal Reactivity to Cocaine Cues During Abstinence Predicts Individual Vulnerability to Relapse. *Neuropsychopharmacology*. 43, 1059–1065. [PubMed: 28920590]
- Halliday M, Radford H, Zents KAM, Molloy C, Moreno JA, Verity NC, Smith E, Ortori CA, Barrett DA, Bushell M, Mallucci GR, 2017 Repurposed drugs targeting eIF2 α -P-mediated translational repression prevent neurodegeneration in mice. *Brain*. 140, 1768–1783. [PubMed: 28430857]
- Hartmann A, Veldhuis JD, Deuschle M, Standhardt H, Heuser I, 1997 Twenty-four hour cortisol release profiles in patients with Alzheimer's and Parkinson's disease compared to normal controls: ultradian secretory pulsatility and diurnal variation. *Neurobiol Aging*. 18, 285–9. [PubMed: 9263193]
- Harzke AJ, Williams ML, Bowen AM, 2009 Binge use of crack cocaine and sexual risk behaviors among African-American, HIV-positive users. *AIDS Behav*. 13, 1106–18. [PubMed: 18758935]
- Heaton RK, Franklin DR, Ellis RJ, McCutchan JA, Letendre SL, Leblanc S, Corkran SH, Duarte NA, Clifford DB, Woods SP, Collier AC, Marra CM, Morgello S, Mindt MR, Taylor MJ, Marcotte TD, Atkinson JH, Wolfson T, Gelman BB, McArthur JC, Simpson DM, Abramson I, Gamst A, Fennema-Notestine C, Jernigan TL, Wong J, Grant I, 2011 HIV-associated neurocognitive disorders before and during the era of combination antiretroviral therapy: differences in rates, nature, and predictors. *J Neurovirol*. 17, 3–16. [PubMed: 21174240]
- Hetman M, Slomnicki LP, 2019 Ribosomal biogenesis as an emerging target of neurodevelopmental pathologies. *J Neurochem*. 148, 325–347. [PubMed: 30144322]
- Homer BD, Solomon TM, Moeller RW, Mascia A, DeRaleau L, Halkitis PN, 2008 Methamphetamine abuse and impairment of social functioning: a review of the underlying neurophysiological causes and behavioral implications. *Psychol Bull*. 134, 301–10. [PubMed: 18298273]
- Hou G, Tian R, Li J, Yuan TF, 2014 Chronic stress and Parkinson's disease. *CNS Neurosci Ther*. 20, 1–2. [PubMed: 24341934]
- Ipser JC, Brown GG, Bischoff-Grethe A, Connolly CG, Ellis RJ, Heaton RK, Grant I, Translational Methamphetamine ARCG, 2015 HIV infection is associated with attenuated frontostriatal intrinsic connectivity: a preliminary study. *J Int Neuropsychol Soc*. 21, 203–13. [PubMed: 25824201]
- Iskander S, Walsh KA, Hammond RR, 2004 Human CNS cultures exposed to HIV-1 gp120 reproduce dendritic injuries of HIV-1-associated dementia. *J Neuroinflammation*. 1, 7. [PubMed: 15285795]
- Jang CG, Whitfield T, Schulteis G, Koob GF, Wee S, 2013 A dysphoric-like state during early withdrawal from extended access to methamphetamine self-administration in rats. *Psychopharmacology (Berl)*. 225, 753–63. [PubMed: 23007601]
- Jones GJ, Barsby NL, Cohen EA, Holden J, Harris K, Dickie P, Jhamandas J, Power C, 2007 HIV-1 Vpr causes neuronal apoptosis and in vivo neurodegeneration. *J Neurosci*. 27, 3703–11. [PubMed: 17409234]

- Kim BO, Liu Y, Ruan Y, Xu ZC, Schantz L, He JJ, 2003 Neuropathologies in transgenic mice expressing human immunodeficiency virus type 1 Tat protein under the regulation of the astrocyte-specific glial fibrillary acidic protein promoter and doxycycline. *Am J Pathol.* 162, 1693–707. [PubMed: 12707054]
- Kitamura O, Wee S, Specio SE, Koob GF, Pulvirenti L, 2006 Escalation of methamphetamine self-administration in rats: a dose-effect function. *Psychopharmacology (Berl).* 186, 48–53. [PubMed: 16552556]
- Koob G, Le Moal M, 2006 Psychostimulants. In *Neurobiology of Addiction*. Vol., ed.^eds. Academic Press, Amsterdam, San Diego, pp. 69–120.
- Koob G, Kreek MJ, 2007 Stress, dysregulation of drug reward pathways, and the transition to drug dependence. *Am J Psychiatry.* 164, 1149–59. [PubMed: 17671276]
- Koob GF, Ahmed SH, Boutrel B, Chen SA, Kenny PJ, Markou A, O'Dell LE, Parsons LH, Sanna PP, 2004 Neurobiological mechanisms in the transition from drug use to drug dependence. *Neurosci Biobehav Rev.* 27, 739–49. [PubMed: 15019424]
- Koob GF, Le Moal M, 2005 Plasticity of reward neurocircuitry and the 'dark side' of drug addiction. *Nat Neurosci.* 8, 1442–4. [PubMed: 16251985]
- Koob GF, Le Moal M, 2008 Addiction and the brain antireward system. *Annu Rev Psychol.* 59, 29–53. [PubMed: 18154498]
- Koob GF, 2009 Neurobiological substrates for the dark side of compulsivity in addiction. *Neuropharmacology.* 56 Suppl 1, 18–31. [PubMed: 18725236]
- Koob GF, Volkow ND, 2010 Neurocircuitry of addiction. *Neuropsychopharmacology.* 35, 217–38. [PubMed: 19710631]
- Kumar A, 2014 Editorial: Impact of alcohol on HIV related issues in human population or model system. *Curr HIV Res.* 12, 233. [PubMed: 25142713]
- Kumar KG, Tang W, Ravindranath AK, Clark WA, Croze E, Fuchs SY, 2003 SCF(HOS) ubiquitin ligase mediates the ligand-induced down-regulation of the interferon-alpha receptor. *EMBO J.* 22, 5480–90. [PubMed: 14532120]
- Langmead B, Trapnell C, Pop M, Salzberg SL, 2009 Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* 10, R25. [PubMed: 19261174]
- Lashomb AL, Vigorito M, Chang SL, 2009 Further characterization of the spatial learning deficit in the human immunodeficiency virus-1 transgenic rat. *J Neurovirol.* 15, 14–24. [PubMed: 19085205]
- Levine AJ, Hardy DJ, Miller E, Castellon SA, Longshore D, Hinkin CH, 2006 The effect of recent stimulant use on sustained attention in HIV-infected adults. *J Clin Exp Neuropsychol.* 28, 29–42. [PubMed: 16448974]
- Li B, Dewey CN, 2011 RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. *BMC bioinformatics.* 12, 323. [PubMed: 21816040]
- Liu CC, Zhao N, Yamaguchi Y, Cirrito JR, Kanekiyo T, Holtzman DM, Bu G, 2016 Neuronal heparan sulfates promote amyloid pathology by modulating brain amyloid-beta clearance and aggregation in Alzheimer's disease. *Sci Transl Med.* 8, 332ra44.
- Lucas GM, Cheever LW, Chaisson RE, Moore RD, 2001 Detrimental effects of continued illicit drug use on the treatment of HIV-1 infection. *J Acquir Immune Defic Syndr.* 27, 251–9. [PubMed: 11464144]
- Lucas GM, Gebo KA, Chaisson RE, Moore RD, 2002 Longitudinal assessment of the effects of drug and alcohol abuse on HIV-1 treatment outcomes in an urban clinic. *Aids.* 16, 767–74. [PubMed: 11964533]
- Lucas GM, Griswold M, Gebo KA, Keruly J, Chaisson RE, Moore RD, 2006 Illicit drug use and HIV-1 disease progression: a longitudinal study in the era of highly active antiretroviral therapy. *Am J Epidemiol.* 163, 412–20. [PubMed: 16394200]
- McArthur JC, 2004 HIV dementia: an evolving disease. *J Neuroimmunol.* 157, 3–10. [PubMed: 15579274]
- Mejias NH, Martinez CC, Stephens ME, de Rivero Vaccari JP, 2018 Contribution of the inflammasome to inflamming. *J Inflamm (Lond).* 15, 23. [PubMed: 30473634]

- Meyerhoff DJ, 2001 Effects of alcohol and HIV infection on the central nervous system. *Alcohol Res Health*. 25, 288–98. [PubMed: 11910707]
- Nath A, Maragos WF, Avison MJ, Schmitt FA, Berger JR, 2001 Acceleration of HIV dementia with methamphetamine and cocaine. *J Neurovirol*. 7, 66–71. [PubMed: 11519485]
- Nath A, 2010 Human immunodeficiency virus-associated neurocognitive disorder: pathophysiology in relation to drug addiction. *Ann N Y Acad Sci*. 1187, 122–8. [PubMed: 20201849]
- Nelson KE, Galai N, Safaeian M, Strathdee SA, Celentano DD, Vlahov D, 2002 Temporal trends in the incidence of human immunodeficiency virus infection and risk behavior among injection drug users in Baltimore, Maryland, 1988–1998. *Am J Epidemiol*. 156, 641–53. [PubMed: 12244033]
- Neuenburg JK, Brodt HR, Herndier BG, Bickel M, Bacchetti P, Price RW, Grant RM, Schlote W, 2002 HIV-related neuropathology, 1985 to 1999: rising prevalence of HIV encephalopathy in the era of highly active antiretroviral therapy. *J Acquir Immune Defic Syndr*. 31, 171–7. [PubMed: 12394795]
- Notarianni E, 2013 Hypercortisolemia and glucocorticoid receptor-signaling insufficiency in Alzheimer's disease initiation and development. *Curr Alzheimer Res*. 10, 714–31. [PubMed: 23906001]
- Orio L, Wee S, Newman AH, Pulvirenti L, Koob GF, 2010 The dopamine D3 receptor partial agonist CJB090 and antagonist PG01037 decrease progressive ratio responding for methamphetamine in rats with extended-access. *Addict Biol*. 15, 312–23. [PubMed: 20456290]
- Parvaz MA, Moeller SJ, Goldstein RZ, 2016 Incubation of Cue-Induced Craving in Adults Addicted to Cocaine Measured by Electroencephalography. *JAMA Psychiatry*. 73, 1127–1134. [PubMed: 27603142]
- Patel CA, Mukhtar M, Harley S, Kulkosky J, Pomerantz RJ, 2002 Lentiviral expression of HIV-1 Vpr induces apoptosis in human neurons. *J Neurovirol*. 8, 86–99. [PubMed: 11935461]
- Petry NM, 1999 Alcohol use in HIV patients: what we don't know may hurt us. *Int J STD AIDS*. 10, 561–70. [PubMed: 10492422]
- Plessis SD, Vink M, Joska JA, Koutsilieri E, Stein DJ, Emsley R, 2014 HIV infection and the fronto-striatal system: a systematic review and meta-analysis of fMRI studies. *AIDS*. 28, 803–11. [PubMed: 24300546]
- Pulliam L, 2014 Cognitive consequences of a sustained monocyte type 1 IFN response in HIV-1 infection. *Curr HIV Res*. 12, 77–84. [PubMed: 24862334]
- Recinto P, Samant AR, Chavez G, Kim A, Yuan CJ, Soleiman M, Grant Y, Edwards S, Wee S, Koob GF, George O, Mandym CD, 2012 Levels of neural progenitors in the hippocampus predict memory impairment and relapse to drug seeking as a function of excessive methamphetamine self-administration. *Neuropsychopharmacology*. 37, 1275–87. [PubMed: 22205547]
- Reid W, Sadowska M, Denaro F, Rao S, Foulke J Jr., Hayes N, Jones O, Doodnauth D, Davis H, Sill A, O'Driscoll P, Huso D, Fouts T, Lewis G, Hill M, Kamin-Lewis R, Wei C, Ray P, Gallo RC, Reitz M, Bryant J, 2001 An HIV-1 transgenic rat that develops HIV-related pathology and immunologic dysfunction. *Proc Natl Acad Sci U S A*. 98, 9271–6. [PubMed: 11481487]
- Renner K, Hellerbrand S, Hermann F, Riedhammer C, Talke Y, Schiechl G, Rodriguez Gomez M, Kutzi S, Halbritter D, Goebel N, Bruhl H, Weissert R, Mack M, 2016 IL-3 promotes the development of experimental autoimmune encephalitis. *JCI Insight*. 1, e87157. [PubMed: 27734026]
- Repunte-Canonigo V, Lütjens R, van der Stap LD, Sanna PP, 2007 Increased expression of protein kinase A inhibitor alpha (PKI-alpha) and decreased PKA-regulated genes in chronic intermittent alcohol exposure. *Brain Res*. 1138, 48–56. [PubMed: 17270154]
- Repunte-Canonigo V, Berton F, Cottone P, Reifel-Miller A, Roberts AJ, Morales M, Francesconi W, Sanna PP, 2010a A potential role for adiponectin receptor 2 (AdipoR2) in the regulation of alcohol intake. *Brain Res*. 1339, 11–7. [PubMed: 20380822]
- Repunte-Canonigo V, van der Stap LD, Chen J, Sabino V, Wagner U, Zorrilla EP, Schumann G, Roberts AJ, Sanna PP, 2010b Genome-wide gene expression analysis identifies K-ras as a regulator of alcohol intake. *Brain Res*. 1339, 1–10. [PubMed: 20388501]

- Repunte-Canonigo V, Chen J, Lefebvre C, Kawamura T, Kreifeldt M, Basson O, Roberts AJ, Sanna PP, 2014a MeCP2 regulates ethanol sensitivity and intake. *Addict Biol.* 19, 791–9. [PubMed: 23448145]
- Repunte-Canonigo V, Lefebvre C, George O, Kawamura T, Morales M, Koob GF, Califano A, Masliah E, Sanna PP, 2014b Gene expression changes consistent with neuroAIDS and impaired working memory in HIV-1 transgenic rats. *Mol Neurodegener.* 9, 26. [PubMed: 24980976]
- Repunte-Canonigo V, Shin W, Vendruscolo LF, Lefebvre C, Koob GF, Califano A, Sanna PP, 2015 Identifying candidate drivers of alcohol dependence-induced excessive drinking by assembly and interrogation of brain-specific regulatory networks. *Genome Biology.* 16:68 DOI: 10.1186/s13059-015-0593-5 [PubMed: 25886852]
- Robinson L, Rempel H, 2006 Methamphetamine use and HIV symptom self-management. *J Assoc Nurses AIDS Care.* 17, 7–14.
- Roy E, Arruda N, Jutras-Aswad D, Berbiche D, Perreault M, Bertrand K, Dufour M, Bruneau J, 2017 Examining the link between cocaine bingeing and individual, social and behavioral factors among street-based cocaine users. *Addict Behav.* 68, 66–72. [PubMed: 28103534]
- Royal W 3rd, Zhang L, Guo M, Jones O, Davis H, Bryant JL, 2012 Immune activation, viral gene product expression and neurotoxicity in the HIV-1 transgenic rat. *J Neuroimmunol.* 247, 16–24. [PubMed: 22503372]
- Sacktor N, Robertson K, 2014 Evolving clinical phenotypes in HIV-associated neurocognitive disorders. *Curr Opin HIV AIDS.* 9, 517–20. [PubMed: 25203640]
- Samet JH, Phillips SJ, Horton NJ, Traphagen ET, Freedberg KA, 2004 Detecting alcohol problems in HIV-infected patients: use of the CAGE questionnaire. *AIDS Res Hum Retroviruses.* 20, 151–5. [PubMed: 15018702]
- Sanna PP, Repunte-Canonigo V, Masliah E, Lefebvre C, 2017 Gene expression patterns associated with neurological disease in human HIV infection. *PLoS One.* 12, e0175316. [PubMed: 28445538]
- Scott JC, Woods SP, Matt GE, Meyer RA, Heaton RK, Atkinson JH, Grant I, 2007 Neurocognitive effects of methamphetamine: a critical review and meta-analysis. *Neuropsychol Rev.* 17, 275–97. [PubMed: 17694436]
- Semple S, Patterson T, Grant I, 2002 Motivations associated with methamphetamine use among HIV+ men who have sex with men. *J Subst Abuse Treat.* 22, 149–56. [PubMed: 12039618]
- Silverstein PS, Kumar A, 2014 HIV-1 and alcohol: interactions in the central nervous system. *Alcohol Clin Exp Res.* 38, 604–10. [PubMed: 24134164]
- Simone IL, Federico F, Tortorella C, Andreula CF, Zimatore GB, Giannini P, Angarano G, Lucivero V, Picciola P, Carrara D, Bellacosa A, Livrea P, 1998 Localised 1H-MR spectroscopy for metabolic characterisation of diffuse and focal brain lesions in patients infected with HIV. *J Neurol Neurosurg Psychiatry.* 64, 516–23. [PubMed: 9576546]
- Song AX, Zhou CJ, Guan X, Sze KH, Hu HY, 2010 Solution structure of the N-terminal domain of DC-Ubp/UBTD2 and its interaction with ubiquitin. *Protein Sci.* 19, 1104–9. [PubMed: 20440844]
- Soontornniyomkij V, Kesby JP, Morgan EE, Bischoff-Grethe A, Minassian A, Brown GG, Grant I, Translational Methamphetamine ARCG, 2016 Effects of HIV and Methamphetamine on Brain and Behavior: Evidence from Human Studies and Animal Models. *J Neuroimmune Pharmacol.* 11, 495–510. [PubMed: 27484318]
- Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP, 2005 Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A.* 102, 15545–50. [PubMed: 16199517]
- Suzuki H, Chiba T, Suzuki T, Fujita T, Ikenoue T, Omata M, Furuichi K, Shikama H, Tanaka K, 2000 Homodimer of two F-box proteins betaTrCP1 or betaTrCP2 binds to IkappaBalpha for signal-dependent ubiquitination. *J Biol Chem.* 275, 2877–84. [PubMed: 10644755]
- Toggas SM, Masliah E, Rockenstein EM, Rall GF, Abraham CR, Mucke L, 1994 Central nervous system damage produced by expression of the HIV-1 coat protein gp120 in transgenic mice. *Nature.* 367, 188–93. [PubMed: 8114918]

- Tozzi V, Balestra P, Serraino D, Bellagamba R, Corpolongo A, Piselli P, Lorenzini P, Visco-Comandini U, Vlassi C, Quartuccio ME, Giulianelli M, Noto P, Galgani S, Ippolito G, Antinori A, Narciso P, 2005 Neurocognitive impairment and survival in a cohort of HIV-infected patients treated with HAART. *AIDS Res Hum Retroviruses*. 21, 706–13. [PubMed: 16131310]
- Trillo-Pazos G, McFarlane-Abdulla E, Campbell IC, Pilkington GJ, Everall IP, 2000 Recombinant nef HIV-IIIB protein is toxic to human neurons in culture. *Brain Res*. 864, 315–26. [PubMed: 10802040]
- van de Bovenkamp M, Nottet HS, Pereira CF, 2002 Interactions of human immunodeficiency virus-1 proteins with neurons: possible role in the development of human immunodeficiency virus-1-associated dementia. *Eur J Clin Invest*. 32, 619–27. [PubMed: 12190962]
- Vigorito M, LaShomb AL, Chang SL, 2007 Spatial learning and memory in HIV-1 transgenic rats. *J Neuroimmune Pharmacol*. 2, 319–28. [PubMed: 18040850]
- Volkow ND, Hitzemann R, Wang GJ, Fowler JS, Burr G, Pascani K, Dewey SL, Wolf AP, 1992a Decreased brain metabolism in neurologically intact healthy alcoholics. *Am J Psychiatry*. 149, 1016–22. [PubMed: 1636801]
- Volkow ND, Hitzemann R, Wang GJ, Fowler JS, Wolf AP, Dewey SL, Handlesman L, 1992b Long-term frontal brain metabolic changes in cocaine abusers. *Synapse*. 11, 184–90. [PubMed: 1636149]
- Volkow ND, Wang GJ, Fowler JS, Telang F, Jayne M, Wong C, 2007 Stimulant-induced enhanced sexual desire as a potential contributing factor in HIV transmission. *Am J Psychiatry*. 164, 157–60. [PubMed: 17202559]
- Volkow ND, Morales M, 2015 The Brain on Drugs: From Reward to Addiction. *Cell*. 162, 712–25. [PubMed: 26276628]
- Volkow ND, Wang GJ, Shokri Kojori E, Fowler JS, Benveniste H, Tomasi D, 2015 Alcohol decreases baseline brain glucose metabolism more in heavy drinkers than controls but has no effect on stimulation-induced metabolic increases. *J Neurosci*. 35, 3248–55. [PubMed: 25698759]
- Vyas S, Maatouk L, 2013 Contribution of glucocorticoids and glucocorticoid receptors to the regulation of neurodegenerative processes. *CNS Neurol Disord Drug Targets*. 12, 1175–93. [PubMed: 24040816]
- Wechsberg WM, Lam WK, Zule W, Hall G, Middlesteadt R, Edwards J, 2003 Violence, homelessness, and HIV risk among crack-using African-American women. *Subst Use Misuse*. 38, 669–700. [PubMed: 12747401]
- Wee S, Wang Z, Woolverton WL, Pulvirenti L, Koob GF, 2007 Effect of aripiprazole, a partial dopamine D2 receptor agonist, on increased rate of methamphetamine self-administration in rats with prolonged session duration. *Neuropsychopharmacology*. 32, 2238–47. [PubMed: 17327886]
- Wiers CE, Shokri-Kojori E, Wong CT, Abi-Dargham A, Demiral SB, Tomasi D, Wang GJ, Volkow ND, 2016 Cannabis Abusers Show Hypofrontality and Blunted Brain Responses to a Stimulant Challenge in Females but not in Males. *Neuropsychopharmacology*. 41, 2596–605. [PubMed: 27156854]
- Youm YH, Grant RW, McCabe LR, Albarado DC, Nguyen KY, Ravussin A, Pistell P, Newman S, Carter R, Laque A, Munzberg H, Rosen CJ, Ingram DK, Salbaum JM, Dixit VD, 2013 Canonical Nlrp3 inflammasome links systemic low-grade inflammation to functional decline in aging. *Cell Metab*. 18, 519–32. [PubMed: 24093676]
- Zhang GL, Zhang X, Wang XM, Li JP, 2014 Towards understanding the roles of heparan sulfate proteoglycans in Alzheimer's disease. *Biomed Res Int*. 2014, 516028. [PubMed: 25157361]
- Zhang LL, Zhang ZN, Wu X, Jiang YJ, Fu YJ, Shang H, 2017 Transcriptomic meta-analysis identifies gene expression characteristics in various samples of HIV-infected patients with nonprogressive disease. *J Transl Med*. 15, 191. [PubMed: 28899396]

- The abuse of stimulants, such as methamphetamine (METH), is associated with treatment non-compliance, a greater risk of viral transmission, and more rapid clinical progression of immunological and central nervous system (CNS) human immunodeficiency virus (HIV) disease.
- We used a state-of-the-art paradigm of escalated voluntary intravenous drug self-administration in HIV transgenic (Tg) and wildtype (WT) rats. The rats were first allowed to self-administer METH under short-access (ShA) conditions, which is characterized by a nondependent and more “recreational” pattern of METH use, and then under long-access (LgA) conditions, which leads to compulsive (dependent) METH intake.
- Compulsivity in METH abuse appears to be driven by medial prefrontal cortex (mPFC) dysfunction. Gene expression profiling of the mPFC in HIV Tg rats with a history of escalated METH self-administration under LgA conditions showed evidence of greater inflammation than WT rats that self-administered METH under LgA conditions, greater neural injury, lower aerobic and higher anaerobic glucose metabolism, and impairments in protein synthesis, a potential mechanism of neurodegeneration.
- These alterations may be key contributors to neural injury and hypofrontality that are believed to underlie cognitive dysfunction in HIV and greater vulnerability to excessive drug intake in the setting of dependence.

Fig. 1. HIV Tg rats exhibited greater compulsivity than WT rats for METH self-administration under long-access conditions.

(A) HIV Tg and littermate control wildtype (WT) rats were trained to self-administer methamphetamine (METH) under short-access conditions (ShA) (1 h/day) on a fixed-ratio 1 schedule of reinforcement. (B) The rats were then switched to long-access conditions (LgA) (6 h/day), resulting in the escalation of METH intake ($^{##}p < 0.01$, day 5 vs. day 1 in the WT group; $^{$$}p < 0.01$, day 10 vs. day 1 in the HIV Tg group). Significant differences were detected in the number of METH infusions between WT and HIV Tg rats between sessions 5 and 8 ($^{++}p < 0.01$). After re-escalation, a significant increase in METH intake was detected in HIV Tg rats vs. WT rats starting from session 5 ($^{**}p < 0.01$). (C) Following the escalation of METH self-administration (before re-escalation), the rats were tested under a progressive-ratio schedule ($^{*}p < 0.05$, vs. WT). HIV Tg rats that self-administered METH under LgA conditions exhibited an increase in responding, reflected by increases in the number of infusions and breakpoints on the PR schedule of reinforcement.

Fig. 2. Differential gene expression in HIV Tg rats and WT rats that self-administered METH under long-access conditions.

(A) Volcano plot of changes in gene expression in the medial prefrontal cortex (mPFC) in HIV Tg rats that self-administered METH under LgA conditions compared with METH-self-administering WT rats. The plots show significance (Log_{10} of p value) vs. fold-change (Log_2) on the y and x axes, respectively. (B) RT-PCR validation of representative differentially expressed genes indicated in (A), including F-Box and WD repeat domain containing 11 (FBXW11), IFN α inducible protein 27 (IFI27), ubiquitin domain containing 2

(UBTD2), and potassium voltage-gated channel interacting protein 1 (KCNIP1). $**p < 0.01$, $***p < 0.0001$. Hm, HIV Tg rats that self-administered METH under LgA conditions; Wm, WT rats that self-administered METH under the same conditions. (C) Heatmap that depicts regulation of the top differentially expressed genes in HIV *vs.* WT (HIV_WT) and HIV *vs.* WT self-administering METH (Hm_Wm). (D) Heatmap of the top differentially expressed pathways indicated by the Gene Set Enrichment Analysis (GSEA) algorithm (Subramanian et al., 2005). NES, normalized enrichment score (Subramanian et al., 2005); Hm_HIV = HIV Tg rats that self-administered METH *vs.* HIV Tg rats; Hm_Wm = HIV Tg *vs.* WT rats that self-administered METH; Wm_W = WT rats that self-administered METH *vs.* WT rats. As further expanded in the next figures, differentially regulated pathways included pathways that are involved in inflammation and immune activation, energy metabolism, protein and RNA metabolism, and neurodegeneration.

Fig. 3. Transcriptional evidence of an increase in neuroinflammation in HIV Tg rats that self-administered METH under long-access conditions.

Representative pathways are indicative of (A, B) broad immune activation and inflammation and (C, D) cytokine signaling, including (E) tumor necrosis factor α (TNF- α), (F) inflammasome activation, (G-I) IL-1, IL-2, and IL-3 activation, and (J-L) type I and II interferon (IFN) signaling, in the mPFC in HIV Tg rats that self-administered METH under LgA conditions (Hm) compared with METH-self-administering WT rats (Wm).

Fig. 4. Gene expression changes indicative of increase in neural damage in HIV Tg rats that self-administered METH under long-access conditions.

In the mPFC of HIV Tg rats that self-administered METH under LgA conditions (Hm) compared with METH-self-administering WT rats (Wm), increases were observed in the transcription of (A) complement-related genes; (B) genes that are related to neurodegeneration in amyotrophic lateral sclerosis; (C) genes that are related to glucocorticoid signaling; increases were also seen in the differential expression of pathways that are involved in (D-F) the extracellular matrix and (G-I) DNA damage and repair.

Fig. 5. Differential regulation of pathways that are involved in glucose metabolism between HIV Tg rats and WT rats that self-administered METH under long-access conditions.

(A-C) Decrease in the expression of genes that are related to glucose oxidation through the tricarboxylic acid (TCA) cycle (A) and increase in the expression of genes that are related to glycolysis (B) and the metabolism of pyruvate (C), the end product of glycolysis, in the mPFC in HIV Tg rats that self-administered METH under LgA conditions (Hm) compared with METH-self-administering WT rats (Wm). A reduction of oxidative phosphorylation through the TCA cycle may reflect a hypometabolic state that can contribute to neural damage and mPFC dysfunction.

Fig. 6. Catabolic molecular environment in HIV Tg rats that self-administered METH under long-access conditions.

(A-F) Transcriptional evidence of impairments in metabolism of proteins and proteostasis (A, B), lower protein synthesis (C) and alterations of RNA processing (D-F) in the mPFC in HIV Tg rats that self-administered METH under LgA conditions (Hm) compared with METH-self-administering WT rats (Wm).