Published in final edited form as:

Public Health Nutr. 2020 June ; 23(8): 1320–1328. doi:10.1017/S1368980019004099.

Shift in body fat distribution from lower body to upper body among urban Colombian women, 1988-89 – 2007-08

Richard L Bender¹, Traci A Bekelman², Paul A Sandberg³, Darna L Dufour¹, Julio C Reina⁴

¹Department of Anthropology, University of Colorado Boulder, Boulder, CO 80309, USA;

²Department of Pediatrics, Section of Nutrition, University of Colorado Anschutz Medical Campus, Aurora, CO 80045, USA;

³Center for the Advanced Study of Human Paleobiology, The George Washington University, Washington, DC 20052, USA;

⁴Departamento de Pediatría, Universidad del Valle and Centro Médico Imbanaco de Cali, Colombia

Abstract

Objective: Body fat distribution may be a stronger predictor of metabolic risk than body mass index (BMI). Yet, few studies have investigated secular changes in body fat distribution in middle-income countries or how those changes vary by socioeconomic status (SES). This study evaluated change in body fat distribution by SES in Colombia, a middle-income country where BMI is increasing rapidly.

Design: We applied factor analysis to previously published data to assess secular changes in adiposity and body fat distribution in cross-sectional samples of urban Colombian women. Anthropometry was used to assess weight, height, and skinfolds (biceps, triceps, subscapular, suprailiac, thigh, calf).

Setting: Cali, Colombia.

Participants: Women (18–44 years) in 1988–89 (*n* 1,533) and 2007–09 (*n* 577) from three SES groups.

Results: We identified an overall adiposity factor, which increased between 1988–89 and 2007–08 in all SES groups, particularly in the middle SES group. We also identified arm, leg, and trunk

Corresponding Author: Richard L Bender, Department of Anthropology, 1350 Pleasant St, 233 UCB, Boulder, CO 80309, Richard.Bender@colorado.edu, Telephone: 303-492-2547, Fax: 303-492-1971.

Authorship

R.L.B., D.L.D., P.A.S., and T.A.B. conceptualized the project. J.C.R. and D.L.D. supervised the data collection. R.L.B. and P.A.S. analysed the data. R.L.B. and D.L.D. drafted the manuscript. T.A.B. and P.A.S. provided substantive feedback on the manuscript. All authors contributed to data interpretation and critically reviewed the final manuscript.

Conflict of Interest

None

Ethical Standards Disclosure

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human participants were approved by the Institutional Review Board of the University of Colorado Boulder and the Comité de Investigaciones y Ética en Investigación del Centro Médico Imbanaco in Cali. Written informed consent was obtained from all participants.

adiposity factors. In all SES groups, leg adiposity decreased, while trunk and arm adiposity increased.

Conclusions: Factor analysis highlighted three trends that were not readily visible in the BMI data and variable-by-variable analysis of skinfolds: (1) overall adiposity increased between time periods in all SES groups; (2) the adiposity increase was driven by a shift from lower body to upper body adiposity; (3) the increase in adiposity was greatest in the middle SES group. Factor analysis provides novel insights into secular changes and socioeconomic variation in body fat distribution during a period of rapid economic development in a middle-income country.

Keywords

Adiposity; anthropometry; socioeconomic status; Colombia; factor analysis; secular shifts

Introduction

The socioeconomic transitions occurring in middle-income countries are accompanied by increases in obesity and related co-morbidities like diabetes and cardiovascular disease^(1–3). Obesity is typically assessed in terms of body mass index (BMI), a measure of weight for height which is used as a proxy for adiposity but is not a measure of body composition or body fat distribution. In recent years, however, it has become increasingly clear that the distribution of body fat may be a stronger predictor of metabolic and cardiovascular risk than BMI or total adiposity alone⁽⁴⁾.

Specifically, studies have shown that adiposity on the trunk of the body versus on the periphery (e.g., limbs) exerts an independent, and sometimes opposite effect on cardiometabolic outcomes and mortality^(5,6). Indeed, lower body (legs) adiposity appears to be protective against cardiometabolic risk relative to upper body adiposity, when controlled for total adiposity^(7,8). Despite the critical role of body fat distribution in understanding chronic disease risk, the possibility that the increases in obesity in middle-income countries are also accompanied by changes in body fat distribution has received little attention. Additionally, given the well-established relationship between socioeconomic status (SES) and BMI observed in middle-income countries^(9–11), it is also possible that body fat distribution, and its concomitant health implications, may differ according to SES in these settings.

The most commonly used indicators of body fat distribution in population-based studies in resource-limited settings, such as middle-income countries, are body circumferences and measures of subcutaneous skinfold thickness⁽¹²⁾, which can provide discrete data points at 12 or more different locations on the upper and lower body. These multiple measures of adiposity can be difficult to interpret since the individual measures all provide unique information but are generally correlated with each other, and researchers must consider several differences and comparisons simultaneously. This is problematic for sets of related variables such as skinfold measures, which are treated as providing independent information (e.g., adiposity at different bodily sites), yet also as measuring aspects of a single underlying source of variation (e.g., overall adiposity). For example, ratios of individual skinfolds, such as the subscapular-triceps ratio, are used as measures of body fat distribution. At the same

time, individual skinfold measures are also entered into equations for predicting total body fat⁽¹³⁾, or are simply added together to calculate variables such as the sum of 4 or 6 skinfolds. Here, we show how a multivariate statistical approach – namely, factor analysis – can aid in the interpretation of secular changes in body fat distribution and overall adiposity, while minimizing the problems associated with an individual variable-by-variable analysis.

Factor analysis is a multivariate approach that describes a set of observed variables in terms of a smaller number of latent (i.e., unobserved) variables or factors. These factors are constructed as linear combinations of the observed variables that account for the maximum amount of variability in the original data. Factor analysis can be used to reduce the dimensionality of a dataset, that is, to include fewer variables in an analysis while maintaining most of the information contained in the original dataset. Factor analysis is also used to interpret the shared variation in a dataset in terms of underlying constructs that may not be directly measurable. In the present context, this approach allows us to analyse all skinfold measures simultaneously, parsing the unique contributions of each skinfold from the shared variability among all skinfolds. The approach also provides the opportunity to explore the pattern of adiposity that may not be readily apparent when interpreting individual skinfold measures.

In this analysis we use previously-published data from two separate cross-sectional anthropometric surveys of women in the city of Cali, Colombia⁽¹⁴⁾. The surveys bracketed a 20-year period (1988–89 and 2007–08) during which Colombia showed rapid economic development⁽¹⁵⁾. The anthropometric surveys documented increases in body weight as well as stature, but no significant change in mean BMI, except for a small increase in the lowest SES group. In approximately the same period of time (1995–2005) there was a 40% increase in the burden of chronic disease in Colombia, of which cardiovascular disease was a leading contributor⁽¹⁶⁾. Indeed, given this shift in the disease burden, the lack of clear increases in BMI found in the original study⁽¹⁴⁾ was surprising in light of the expected link between obesity and chronic disease^(1–3). Thus, using a novel analytic approach to understand changes in both overall adiposity and in body fat distribution in a middle-income country during a period of economic development and increased chronic disease burden may help to inform public health practices in similar contexts.

To more precisely understand changes in adiposity during a period of rapid economic transition in Colombia, the current paper focuses explicitly on shifts in adiposity and body fat distribution between time periods in the Cali dataset. These are the only data of which we are aware that provide skinfold measurements in the lower as well as the upper body, and hence can document secular changes in overall adiposity as well as in whole-body fat distribution.

Although these data are from 1988–1989 and 2007–2008, they are relevant to current public health practice for two primary reasons. First, understanding changes in body fat distribution and overall adiposity over time is critical for identifying modifiable targets for interventions to reduce the burden of chronic disease in middle-income countries, and few studies provide comparable skinfold data at multiple time points. Second, unfavorable shifts in body fat distribution were not readily apparent in the original analysis of Colombian BMI data and

skinfold measures using more traditional analytic methods⁽¹⁴⁾. Studies that use such methods to assess changes in adiposity between time points may similarly fail to identify or underestimate population-level shifts in body fat distribution that have important implications for public health practice. The novel analytic approach used in the current paper may be applicable to the broader public health and nutrition communities because it allows for better interpretation of multiple, interrelated biological variables. Our goals are to a) use factor analysis to better understand the secular changes in overall adiposity and body fat distribution in Cali women and b) determine if these changes vary by SES.

Methods

Study participants and data collection methods are discussed in detail in the original analysis⁽¹⁴⁾ and are summarized briefly here. Participants in both the 1988–89 (n 1,553) and 2007–08 (n 577) studies were distinct cross-sectional samples of two different groups of women aged 18–44 living in the city of Cali. In both studies, a socioeconomically diverse sample of participants was recruited as a convenience sample from medical clinics (staff and non-patients), private schools, a government institute, and a health club. Only women who self-reported as healthy and not pregnant or lactating were enrolled in the studies.

In both the 1988–89 and the 2007–08 studies, the SES of participants was defined in terms of their residential address using the classification system of the city of Cali^(17,18). In this system, each city block is classified into one of six *estratos* (strata) based on the external appearance of the houses, access to municipal services, and the condition of the streets. The six *estratos* are compressed into three SES groups for analysis: lower, *estratos* one and two; middle, *estratos* three and four; higher, *estratos* five and six.

In both the 1988–89 and 2007–08 studies, anthropometric measurements were taken by the same experienced technician following standardized procedures⁽¹⁹⁾. Height (m) was measured with a Harpenden stadiometer, and weight (kg) with a digital Seca scale. Skinfold thicknesses (mm) were measured in triplicate with a Lange skinfold calliper at six sites: triceps, biceps, subscapular, suprailiac, mid-thigh, and medial calf. Waist circumference (WC; cm) was measured in duplicate with a flexible steel tape. BMI was calculated as weight (kg)/height (m²).

Statistical analysis

We applied a factor analysis to the six skinfold variables reported in the original analysis⁽¹⁴⁾. While previous researchers have employed multivariate methods to analyse sets of skinfold measurements, most of these studies date from the 1990s, and most used principal components analysis rather than factor analysis^(20–22). Principal components analysis partitions the total variability in the dataset, including unique and error variability, but factor analysis partitions shared variability only⁽²³⁾. Therefore, factor analysis is better-suited to our goal of identifying and interpreting latent variables in the dataset.

We used the maximum likelihood method of factor extraction. This allowed us to subsequently use Bartlett's test for sphericity to test for the sufficiency of the number of factors extracted from the pooled data⁽²⁴⁾. Skinfold data from the 1988–89 and 2007–08

studies were pooled to allow for the extraction of common factors that could be directly compared between the two time periods. Following the initial factor extraction, the varimax method was used for orthogonal rotation of factors. This method aids the interpretation of factors by maximizing the variance accounted for by each factor, while keeping the factors orthogonal (i.e., uncorrelated) and without altering the multivariate relationship among the individual data points^(25,26). We report factor loadings, i.e., the correlation between an initial skinfold variable and a factor, for each extracted factor both before and after varimax rotation. The strength of the factor loadings was used to interpret and label the factors in biological terms. We placed our factors' labels in italics and quotation marks (e.g., “*overall adiposity*”) to emphasize that the labels reflect our own interpretations of the data.

For both the un-rotated and rotated factors, we calculated individual factor scores for each subject, which are interpretable as standard scores (i.e., *z*-scores) in units of standard deviations with a mean of zero. Subsequently, we used independent-samples *t*-tests with Tukey-Kramer correction for multiple comparisons to compare mean age, height, weight, and factor scores between time periods within each SES group. Pearson's χ^2 test was used to assess the distribution of SES groups by time period, with Cramér's *V* as the measure of association. Correlation analysis (Pearson's *r*) and ordinary least squares regression were used to assess the associations among age, height, weight, waist circumference and factor scores. We used analysis of covariance (ANCOVA) to compare mean factor scores between time periods while controlling for the potential confounding effects of age and height. Factor analyses and statistical tests were conducted in JMP Pro 12 with statistical significance set at $P < 0.05$ (two-sided). Figures were created in SigmaPlot 12.5.

Results

Sample characteristics

Sample sizes by SES group and time period are shown in Table 1. There was a significant association between SES and time period, but the measure of association was weak.

Mean age, height, and weight of women by SES group in both time periods are shown in Table 2. For all SES groups, the 2007–08 sample was significantly older, taller, and heavier. Also, in both time periods, age was positively correlated with weight and skinfold measures (data not shown).

Factor loadings

The factor analysis resulted in 3 factors extracted from the pooled 1988–89 and 2007–08 data (Bartlett's test of sphericity, $\chi^2 = 326.0$, $df = 4$, $P < 0.001$), together accounting for 72.2% of the total variation in the skinfold data. We first present the loadings for the 3 factors before rotation (Table 3). We interpret the first of these factors, which accounts for 56.6% of total variation and is loaded positively on all 6 skinfold variables, as an “*overall adiposity*” factor.

Factor loadings for the varimax-rotated solution are shown in Table 4. Here, each of the 3 extracted factors accounts for between 23%–26% of total variation. Factor 1 loads strongly on the triceps and biceps skinfolds, with moderate loadings on the subscapular and

suprailiac skinfolds. Factor 2 loads strongly on the mid-thigh and medial calf skinfolds, with a moderate loading on the triceps skinfold. Finally, Factor 3 loads strongly on the subscapular and suprailiac skinfolds, with moderate loadings on the biceps and triceps skinfolds.

We interpret the 3 rotated factors as follows. Factor 1 is an arm adiposity factor (“*arms*”), reflecting adiposity mainly in the biceps and triceps depots of the arms, with some contribution from the subscapular and suprailiac depots of the trunk. Factor 2 is a leg adiposity factor (“*legs*”), reflecting adiposity in the mid-thigh and medial calf depots of the legs. Factor 3 is a trunk adiposity factor (“*trunk*”), reflecting adiposity mainly in the subscapular and suprailiac depots of the trunk, with some contribution from the biceps and triceps depots of the arms.

Correlations among factors and anthropometric variables

The “*overall adiposity*” factor was positively correlated with weight in all SES groups in both time periods ($0.68 \leq r \leq 0.82$, $P < 0.001$ in all cases; Figure 1), but not significantly correlated with height in any SES group or time period ($0.02 \leq |r| \leq 0.09$, $P > 0.05$ in all cases; Figure 1). Additionally, the “*overall adiposity*” factor was positively correlated with WC in all SES groups and time periods ($0.68 \leq r \leq 0.82$, $P < 0.001$ in all cases; Figure 1). Finally, all three of the rotated factors (“*arms*,” “*legs*,” and “*trunk*”) were positively correlated with WC in all SES groups and time periods ($0.33 \leq r \leq 0.69$; $P < 0.001$ in all cases; data not shown).

Changes between time periods within SES groups

Table 5 and Figure 2 show changes in means of the factor scores for the extracted adiposity factors (“*overall*,” “*arms*,” “*legs*,” and “*trunk*”) between time periods in each SES group. Mean “*overall adiposity*” factor scores increased significantly from 1988–89 to 2007–08 in all SES groups, with the smallest increase in the higher SES group and the largest increase in the middle SES group. Means of “*arms*” and “*trunk*” factor scores also increased significantly between time periods in all SES groups. For the “*arms*” factor, the smallest increase was observed in the higher SES group, with a lesser (but equal) increase in the lower SES and middle SES groups. For the “*trunk*” factor, the smallest increase was in the lower SES group, with the largest increase in the middle SES group. Lastly, mean “*legs*” factor scores decreased significantly between 1988–89 and 2007–08 in all SES groups, with the smallest decrease in the lower SES group and a greater (but equal) decrease in the middle SES and higher SES groups.

To control for the potential confounding effects of age and height on adiposity factor scores, we repeated these comparisons with ANCOVA, using age and height as covariates. In all cases, the main effect of time period on adiposity factor scores remained significant at $P = 0.009$.

Discussion

Our goals in this paper were to a) better understand the secular changes in body fat distribution in Cali women and b) determine if these changes vary by SES. To accomplish

that, we applied a multivariate method that is particularly well-suited to the analysis of intercorrelated biological variables in a variety of populations: factor analysis. The re-analysis highlighted three features of the original data set that were not readily apparent in the variable-by-variable analysis of discrete skinfold measurements or BMI.

First, our initial factor analysis provided a measure of “*overall adiposity*” that increased significantly between time periods in all SES groups. The magnitude of the increases in “*overall adiposity*” scores, between +0.54 and +0.86 standard deviations, suggests a substantial increase in the aspect of body fatness measured by this factor. This measure was highly correlated with body weight, but not height, an observation that increased our confidence that the “*overall adiposity*” factor was not tracking overall body size, but rather a dimension of body weight that was independent of height – i.e., adiposity. The overall increases in adiposity were not evident when BMI was used as a proxy for adiposity in the original study⁽¹⁴⁾: mean BMI only increased significantly in the lower SES group, and that increase was small (1.5 kg/m² over 20 years).

The relative lack of changes in BMI stems from the fact that although weight increased between time periods in all SES groups, so did height. Thus, the effect of the increases in weight were attenuated by a concurrent secular change in height. Our reanalysis suggests that adiposity did in fact increase in this sample over time, even though the most commonly-used indicator of obesity – BMI – did not. Unlike BMI, our “*overall adiposity*” factor was not correlated with height, and the changes in “*overall adiposity*” scores between time periods remained statistically significant after controlling for height and age. Therefore, the change in “*overall adiposity*” between 1988–89 and 2007–08 is unlikely to be an artefact of increased height between these time periods. Also, although the SES distribution differed in the 1988–1989 and 2007–08 samples, it is unlikely that these differences introduced bias for two reasons: the magnitude of the difference in distribution is small, and our analysis focuses on changes within each SES group rather than on SES itself as a driver of adiposity.

Furthermore, the “*overall adiposity*” factor was positively correlated with WC in all SES groups and both time periods. WC is a commonly-measured variable in nutritional epidemiology due to its strong association with numerous health risks including hypertension, type 2 diabetes mellitus, and cardiovascular disease (CVD)^(27–31). Thus, the association between our “*overall adiposity*” factor and WC further suggests that the “*overall adiposity*” factor may be a biologically-meaningful indicator of health risks.

Second, following varimax rotation, our factor analysis distinguished a pattern of body fat distribution in which the increases in adiposity were restricted to the upper body, i.e., “*trunk*” and “*arms*”; adiposity actually decreased in the lower body, i.e., “*legs*.” These changes were generally consistent with the changes in individual skinfolds originally reported⁽¹⁴⁾. However, given the opposite direction of adiposity changes in the upper and lower body, it was difficult to judge from the original variable-by-variable analysis if overall adiposity had increased. Hence, our factor analytic approach more clearly demonstrated that the general adiposity increase observed in this population was driven by an increase in upper body fatness specifically. The magnitude of the changes suggests a meaningful shift in the aspects of adiposity measured by these factors. Overall, the smallest change was the increase

in “*trunk*” adiposity in the lower SES group (+0.21 standard deviations), while the greatest change was the “*arms*” adiposity increase in both the lower and middle SES groups (+1.15 standard deviations). The potential health implications of this increase in upper body fatness in conjunction with decreased leg fatness are unclear.

The distinctions in the pattern of subcutaneous fat distribution we found are in reasonable agreement with results from studies conducted in the 1970s, 80s, and 90s using multivariate techniques (principal components analysis and factor analysis) to analyse skinfold data in a variety of populations. For example, principal components of overall fatness, trunk vs. extremity fatness, and upper vs. lower body fatness were uncovered in adults and children in a re-analysis of 44 samples of anthropometric data^(32,33). Indices of trunk vs. extremity fatness were also reported among Polish adults⁽³⁴⁾ and among Mexican-American children⁽²⁰⁾. These previous studies of skinfold data, having uncovered patterns of body fat distribution similar to those in the current study, provide a good basis for between-group comparisons and for our cross-sectional examination of changes between time periods.

Third, in our reanalysis we found that the increase in “*overall adiposity*” and in “*trunk*” adiposity specifically was greatest in the middle SES group. This was not evident in the original analysis using WC and waist hip ratio (WHR) as indices of truncal fat deposition: WC increased in all groups, but WHR only increased in the lower and higher SES groups⁽¹⁴⁾. Hence, the assumption was that the middle SES group had experienced less of an increase in truncal body fat than the other two groups. Since the “*trunk*” factor was positively correlated with WC, the “*trunk*” adiposity data suggest that the greatest increase in adiposity-related health risk occurred within the middle SES group. Thus, the factor analysis approach revealed a potential health risk that was not readily apparent in the original variable-by-variable analysis of individual skinfold measures⁽¹⁴⁾.

Comparable data on secular changes in body fat distribution in reproductive age women are limited, and most focus on WC and/or WHR as indices of truncal adiposity; only a few studies report skinfold data. Reports from developing countries during periods of significant economic change are of particular interest. For example, an increase in WC was found in 20-year old female students in Slovenia over a 30-year period (1964–5 to 2008)⁽³⁵⁾. It is noteworthy that this change occurred in the absence of a change in BMI. A 12-year (1999 to 2012) increase in WC was also reported in Mexican women of reproductive age⁽³⁶⁾. It was accompanied by an increase in BMI, but interpreted as an increase in truncal adiposity, because the increases in WC were greater than expected for a given BMI. Data for reproductive age women in developed countries (USA, Mexico, Canada, Sweden, Finland) in the past ~50 years also provide evidence of an increase in WC in the absence of a change in mean BMI, or an increase in WC greater than expected for the change in BMI^(36–39). In all cases, the WC increases are interpreted as changes in the distribution of body fat to a more truncal pattern, but since the data are limited to the upper body, they are incomplete measures of fat distribution. However, observations that hip circumferences were smaller, or unchanged, in the more recent cohorts of Scandinavian women^(38,39) suggest that fat was being preferentially deposited in the upper body rather than the lower. Several studies also measured skinfolds and reported secular increases in subcutaneous fatness^(35,37,39), but all of

the skinfolds were on the upper body, or reported as the sum of skinfolds, and hence limit our understanding of possible changes in fat distribution over the whole body.

The upper-versus lower-body sites of fat deposition matter in terms of health outcomes. It is generally accepted that higher levels of abdominal and trunk fat are associated with greater obesity related co-morbidities, and lower-body adipose tissue depots are thought to be associated with decreased CVD risk⁽⁴⁾. For example, a number of studies using imaging technologies like dual energy x-ray absorptiometry (DXA) have shown that after controlling for total adiposity, higher leg adiposity is associated with lower CVD risk in USA and British women over a wide age range (18–64 years)^(5,8), pre-menopausal European women⁽⁴⁰⁾, and even severely obese women⁽⁴¹⁾. Indeed, the association is stronger in more obese women.

The health significance of fat depots on the arms is less clear. Arm fat has been reported to be associated with greater CVD risk in USA adult women⁽⁵⁾. In contrast, other studies have found no association of arm fat with increased risk of type-2 diabetes mellitus or CVD in adult women^(8,40). Reasons for these disparate results are not readily apparent but could be related to participant age: abdominal fat deposition increases post-menopause, and arm fat also increases until age 60–70 years^(42,43).

In Cali, the changes in body fat distribution between 1988–89 and 2007–08 occurred in the context of an ongoing period of economic development, but causal linkages are not readily apparent. The general assumption is that nutrition transitions in middle-income countries such as Colombia involve changes in lifestyle, especially in diet and physical activity⁽⁴⁴⁾. Results of research attempting to link changes in lifestyle to both overall adiposity and body fat distribution are contradictory. Studies of Scandinavian women have not been able to link increases in WC to lifestyle changes (alcohol consumption, smoking, physical activity) between 1980–81 and 1992–93^(38,39). In contrast, increases in WC in USA women between 1988 and 2010 have been linked to decreases in self-reported physical activity⁽⁴⁵⁾, although other studies have not demonstrated a strong link between physical activity and abdominal (specifically visceral) adiposity^(46,47).

The strengths of the present study, a reanalysis of secular changes in adiposity in urban women in three SES groups in a developing country, include the use of factor analysis to understand intercorrelated biological variables and the potential for the application of this method to other populations. Factor analysis allowed us to analyse all skinfold measures simultaneously and helped to reveal the underlying patterns within the body fat distribution data, suggesting avenues for future population health monitoring. Public health and nutrition scientists interested in body fat distribution can benefit from using factor analysis to interpret skinfold data, especially for assessing secular changes, because typical analytic approaches may fail to identify or underestimate important population-changes in human biology.

Could factor analysis also be useful in analysing more direct measures of body fat, such as those provided by imaging technologies like DXA scans? Presumably yes, because a number of studies have demonstrated good agreement between estimates of total and extremity fat mass derived from skinfolds and their DXA analogues^(12,48,49). Further,

analyses of intercorrelated biological variables such as nutritional biomarkers, biological measures of stress, and mental health outcomes, could also benefit from factor analysis, a low cost analytical approach that is easily implemented in current statistical packages.

Another strength of our study is the use of anthropometric data from reasonably large samples, collected by the same technician using the same equipment and techniques which reduced measurement error, and at time points that bracketed a period of significant economic change. Further, skinfold data from different time periods within the same setting are scarce among Latin American countries, and most studies rely solely on BMI to assess secular changes in adiposity^(10,11). We also recognize the limitations of the data. The samples were cross-sectional convenience samples from one urban area. We were not able to assess the lifestyle changes which may have been associated with the changes in body fat distribution. Additional studies incorporating multiple measures of body fat distribution over time are needed to fully understand the transition in body composition in this population.

Conclusions

We used factor analysis to re-examine secular changes in adiposity and body fat distribution among women in Cali, Colombia between 1988–89 and 2007–08, a period of rapid economic development and increased chronic disease burden in this middle-income country. Factor analysis provides a parsimonious method of analysing large datasets of intercorrelated anthropometric variables, such as skinfolds. We found that this method highlighted three trends that were not readily visible in the original analysis of the skinfold data⁽¹⁴⁾. First, overall adiposity increased between time periods in all three SES groups, a change that was not apparent in the original analysis using BMI as a proxy. Second, the increase in adiposity was greatest in the middle SES group. Third, the overall adiposity increase was driven by a shift in adiposity from the lower body to the upper body, a shift associated with increased health risks. Factor analysis provided novel insights into secular changes and socioeconomic variation in body fat distribution during a period of rapid economic development in Colombia.

Acknowledgments

The authors thank the women who volunteered for the studies for their patience and cooperation. The authors also express their gratitude to Betty de Orozco and Yazmin Villota for their invaluable assistance with the data collection, to Coldeportes del Valle for their logistical support, and to Kathryn M. Olszowy for collaboration in analysing the original dataset. The authors thank the directors of the following facilities for their assistance in recruiting participants: Centro Médico Imbanaco de Cali; Hospital Carlos Holmes Trujillo; Instituto de Seguros Sociales; FENALCO; Colegio Bolívar; Colegio Gimnasio de Colina; Gimnasio Estamina. Additionally, the authors thank three anonymous reviewers for their helpful comments and suggestions.

Financial Support

The original study was supported by the National Science Foundation (D.L.D., grant number BCS 0742959) and the National Institutes of Health (D.L.D., grant number 5-R22-DK39734). T.A.B. receives salary support from NIH T32 DK007658–27. This research received no specific grant from any funding agency, commercial or not-for-profit sectors. The National Science Foundation and the National Institutes of Health had no role in the design, analysis or writing of this article.

References

1. Monteiro CA, Moura EC, Conde WL et al. (2004) Socioeconomic status and obesity in adult populations of developing countries: a review. *Bull World Health Org* 82, 940–946. [PubMed: 15654409]
2. Prentice AM (2006) The emerging epidemic of obesity in developing countries. *Int J Epidemiol* 35, 93–99. [PubMed: 16326822]
3. Subramanian SV, Perkins JM, Özaltin E et al. (2011) Weight of nations: a socioeconomic analysis of women in low- to middle-income countries. *Am J Clin Nutr* 93, 413–421. [PubMed: 21068343]
4. Goossens GH (2017) The metabolic phenotype in obesity: fat mass, body fat distribution, and adipose tissue function. *Obes Facts* 10, 207–215. [PubMed: 28564650]
5. Hu G, Bouchard C, Bray GA et al. (2011) Trunk versus extremity adiposity and cardiometabolic risk factors in white and African American adults. *Diabetes Care* 34, 1415–1418. [PubMed: 21505210]
6. Lee JJ, Pedley A, Therkelsen KE et al. (2017) Upper body subcutaneous fat is associated with cardiometabolic risk factors. *Am J Med* 130, 958–966. [PubMed: 28238696]
7. Okura T, Nakata Y, Yamabuki K et al. (2004) Regional body composition changes exhibit opposing effects on coronary heart disease risk factors. *Arterioscler Thromb Vasc Biol* 24, 923–929. [PubMed: 15016639]
8. Vasan SK, Osmond C, Canoy D et al. (2018) Comparison of regional fat measurements by dual-energy X-ray absorptiometry and conventional anthropometry and their association with markers of diabetes and cardiovascular disease risk. *Int J Obes* 42, 850–857.
9. Neuman M, Finlay JE, Davey Smith G et al. (2011) The poor stay thinner: stable socioeconomic gradients in BMI among women in lower- and middle-income countries. *Am J Clin Nutr* 94, 1348–1357. [PubMed: 21993437]
10. Kasper NM, Herrán OF & Villamor E (2014) Obesity prevalence in Colombian adults is increasing fastest in lower socio-economic status groups and urban residents: results from two nationally representative surveys. *Public Health Nutr* 17, 2398–2406. [PubMed: 24476690]
11. Yamada G, Jones-Smith JC, Castillo-Salgado C et al. (2019) Differences in magnitude and rates of change in BMI distributions by socioeconomic and geographic factors in Mexico, Colombia and Peru, 2005–2010. *Eur J Clin Nutr*. 10.1038/s41430-019-0479-7.
12. Scafoglieri A, Tresignie J, Provyn S et al. (2013) Accuracy and concordance of anthropometry for measuring regional fat distribution in adults aged 20–55 years. *Am J Hum Biol* 25, 63–70. [PubMed: 23132592]
13. Durnin JVGA & Womersley J (1974) Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged 16 to 72 years. *Br J Nutr* 32, 77–97. [PubMed: 4843734]
14. Olszowy KM, Dufour DL, Bender RL et al. (2012) Socioeconomic status, stature, and obesity in women: 20-year trends in urban Colombia. *Am J Hum Biol* 24, 602–610. [PubMed: 22513997]
15. World Bank (2009) World Bank Country and Lending Groups. <http://go.worldbank.org/K2CKM78CC0> (accessed November 2018).
16. Webster PC (2012) Health in Colombia; the chronic disease burden. *CMAJ* 184, E293–E294. [PubMed: 22392946]
17. EMCALI (1982) La estructura tarifaria actual. Empresas Municipales de Cali (EMCALI).
18. Reyes RT, Ponce de Leon SJC, Suarez Becerra CH et al. (2007) Cali en Cifras. Alcaldía de Santiago de Cali, Colombia http://planeacion.cali.gov.co/Publicaciones/Cali_en_Cifras/Caliencifras2007.pdf (accessed October 2011).
19. Lohman TG, Roche AF & Martorell R (1988) Anthropometric standardization reference manual. Champaign, IL: Human Kinetics Books.
20. Baumgartner RN, Roche AF, Guo S et al. (1990) Fat patterning and centralized obesity in Mexican-American children in the Hispanic Health and Nutrition Examination Survey (HHANES 1982–1984). *Am J Clin Nutr* 51, 936S–943S.

21. Georges E, Mueller WH & Wear ML (1991) Body fat distribution: associations with socioeconomic status in the Hispanic Health and Nutrition Examination Survey. *Am J Hum Biol* 3, 489–501. [PubMed: 28597473]
22. Rebato E, Salces I, San Martin L et al. (1998) Fat distribution in relation to sex and socioeconomic status in children 4–19 years. *Am J Hum Biol* 10, 799–806. [PubMed: 28561408]
23. Bryant FB & Yarnold PR (1995) Principal-components analysis and exploratory and confirmatory factor analysis In *Reading and Understanding Multivariate Statistics*, pp. 99–136 G Grimm G and Yarnold PR, editors]. Washington, DC: American Psychological Association.
24. Tobias S & Carlson JE (1969) Brief report: Bartlett's test of sphericity and chance findings in factor analysis. *Multivar Behav Res* 4, 375–377.
25. Kaiser HF (1958) The varimax criterion for analytic rotation in factor analysis. *Psychometrika* 23, 187–200.
26. Abdi H (2003) Factor rotations in factor analysis In *Encyclopedia of Social Science Research Methods* pp. 1–8. [Lewis-Beck M, Bryman A and Futing T, editors]. Thousand Oaks, CA: Sage.
27. Janssen I, Katzmarzyk PT & Ross R (2004) Waist circumference and not body mass index explains obesity-related health risk. *Am J Clin Nutr* 79, 379–384. [PubMed: 14985210]
28. Wahrenberg H, Hertel K, Leijonhufvud BM et al. (2005) Use of waist circumference to predict insulin resistance: retrospective study. *Br Med J* 330, 1363–1364. [PubMed: 15833749]
29. Katzmarzyk PT, Janssen I, Ross R et al. (2006) The importance of waist circumference in the definition of metabolic syndrome: prospective analyses of mortality in men. *Diabetes Care* 29, 404–409. [PubMed: 16443895]
30. Klein S, Allison DB, Heymsfield SB et al. (2007) Waist circumference and cardiometabolic risk: a consensus statement from Shaping America's Health: Association for Weight Management and Obesity Prevention; NAASO, The Obesity Society; the American Society for Nutrition; and the American Diabetes Association. *Obesity* 15, 1061–1067. [PubMed: 17495180]
31. Balkau B, Deanfield JE, Després JP et al. (2007) International Day for the Evaluation of Abdominal Obesity (IDEA): a study of waist circumference, cardiovascular disease, and diabetes mellitus in 168,000 primary care patients in 63 countries. *Circulation* 116, 1942–1951. [PubMed: 17965405]
32. Mueller MH & Reid RM (1979) A multivariate analysis of fatness and relative fat patterning. *Am J Phys Anthropol* 50, 199–208. [PubMed: 443356]
33. Mueller MH & Wohlleb JC (1981) Anatomical distribution of subcutaneous fat and its description by multivariate methods: How valid are principle components? *Am J Phys Anthropol* 54, 25–35. [PubMed: 7234976]
34. Badora G (1975) The distribution of subcutaneous fat tissue in young men and women. *Studies in Physical Anthropology* 1, 91–108.
35. Kotnik KJ & Golja P (2012) Changes in body composition of University students in a country in socio-economic transition. *Anthrop Anz* 69, 261–271.
36. Albrecht SS, Barquera S & Popkin BM (2014) Exploring secular changes in the association between BMI and waist circumference in Mexican-origin and White women: A comparison of Mexico and the United States. *Am J Hum Biol* 26, 627–634. [PubMed: 24917415]
37. Janssen I, Shields M, Craig CI et al. (2012) Changes in the obesity phenotype within Canadian children and adults, 1981 to 2007–2009. *Obesity* 20, 916–919. [PubMed: 21566567]
38. Lahti-Koski M, Pietinen P, Männistö S et al. (2000) Trends in waist-to-hip ratio and its determinants in adults in Finland from 1987–1997. *Am J Clin Nutr* 72, 1436–44. [PubMed: 11101468]
39. Lissner L, Björkelund C, Heitmann BL et al. (1998) Secular increases in waist-hip ratio among Swedish women. *Int J Obes* 22, 1116–1120.
40. Sánchez-López M, Ortega FB, Moya-Martínez P et al. (2013) Leg fat might be more protective than arm fat in relation to lipid profile. *Eur J Nutr* 52, 489–495. [PubMed: 22484387]
41. Faloia E, Tirabassi G, Canibus P et al. (2009) Protective effect of leg fat against cardiovascular risk factors in obese premenopausal women. *Nutr Metab Cardiovasc Dis* 19, 39–44. [PubMed: 18456482]

42. Svendsen OL, Hassager C & Christiansen C (1995) Age- and menopause-associated variation in body composition and fat distribution in healthy women as measured by dual-energy x-ray absorptiometry. *Metabolism* 44, 369–373. [PubMed: 7885283]
43. Shimokata H, Tobin JD, Muller DC et al. (1989) Studies in the distribution of body fat: I Effects of age, sex, and obesity. *J Gerontol* 44, M66–M73. [PubMed: 2921472]
44. Popkin BM (1994) The nutrition transition in low-income countries: an emerging crisis. *Nutr Rev* 52, 285–298. [PubMed: 7984344]
45. Ladabaum U, Mannalithara A, Myer PA et al. (2014) Obesity, abdominal obesity, physical activity, and caloric intake in U.S. Adults: 1988–2010. *Am J Med* 127, 717–727. [PubMed: 24631411]
46. Thompson D, Karpe F, Lafontan M et al. (2012) Physical activity and exercise in the regulation of human adipose tissue physiology. *Physiol Rev* 92, 157–191. [PubMed: 22298655]
47. Karpe F & Pinnick KE (2015) Biology of upper-body and lower-body adipose tissue – link to whole-body phenotypes. *Nat Rev Endocrinol* 11, 90–100. [PubMed: 25365922]
48. Demura S, Sato S & Noguchi T (2005) Prediction of segmental percent fat using anthropometric variables. *J Sports Med Phys Fitness* 45, 518–523. [PubMed: 16446684]
49. Holmes JD, Andrews DM, Durkin JL et al. (2005) Predicting in vivo soft tissues masses of the lower extremity using segment anthropometric measures and DXA. *J Appl Biomech* 21, 371–382. [PubMed: 16498182]

Fig. 1.

Relationship among "overall adiposity" score, weight (kg), height (m), and waist circumference (cm) within each time period, with ordinary least squares regression lines. Correlations in both time periods are statistically significant between "overall adiposity" score and weight ($r > 0.743$, $P < 0.001$ in both cases) and between "overall adiposity" score and waist circumference ($r > 0.717$, $P < 0.001$ in both cases). Correlations between "overall adiposity" score and height are not significant in either time period ($|r| < 0.042$, $P = 0.275$ in both cases).

Fig. 2.

Change between time periods in mean scores for the “overall adiposity,” “arms,” “legs,” and “trunk” factors. Filled squares, lower socioeconomic status (SES); shaded circles, middle SES; open triangles, higher SES.

TABLE 1

Comparison (χ^2) of sample sizes by socioeconomic status (SES) groups between time periods.

	1988–89		2007–08	
	<i>n</i>	%	<i>n</i>	%
Higher SES	454	29.2	93	16.1
Middle SES	586	37.7	267	46.3
Lower SES	513	33.0	217	37.6
χ^2 (df)	38.41 (2)			
<i>P</i>	< 0.001			
Cramér's V	0.134			

TABLE 2

Comparisons (*t*-tests) of age, height, and weight between time periods within socioeconomic status (SES) groups.

	1988–89		2007–08			
Variable	Mean	SD	Mean	SD	<i>t</i> (df)	<i>P</i>
Age (years)						
Higher SES	29.0	7.0	32.2	8.4	3.94 (545)	< 0.001
Middle SES	28.6	7.0	32.3	7.4	7.01 (851)	< 0.001
Lower SES	29.9	7.1	31.5	7.9	2.70 (728)	0.007
Height (m)						
Higher SES	1.58	0.06	1.61	0.07	4.61 (545)	< 0.001
Middle SES	1.53	0.06	1.59	0.06	12.64 (851)	< 0.001
Lower SES	1.53	0.06	1.56	0.06	6.92 (728)	< 0.001
Weight (kg)						
Higher SES	56.5	7.5	59.4	9.1	3.29 (545)	0.001
Middle SES	55.1	7.6	61.3	10.3	9.92 (851)	< 0.001
Lower SES	57.0	9.9	63.7	12.3	7.80 (728)	< 0.001

TABLE 3

Factor loadings for initial un-rotated solution. 3 factors extracted, accounting for 72.2% of variation. Darker text corresponds to stronger loadings.

Variable	Factor loadings (un-rotated)		
	Factor 1	Factor 2	Factor 3
Triceps	0.87	−0.06	−0.24
Biceps	0.85	−0.24	−0.15
Subscapular	0.87	−0.10	0.26
Suprailiac	0.73	−0.03	0.29
Mid-thigh	0.61	0.59	0.02
Medial calf	0.51	0.52	−0.10
<i>Variance explained</i>	56.6%	11.6%	4.0%

TABLE 4

Factor loadings after varimax rotation. 3 factors extracted, accounting for 72.2% of variation. Darker text corresponds to stronger loadings.

Variable	Factor loadings (varimax)		
	Factor 1	Factor 2	Factor 3
Triceps	0.76	0.35	0.35
Biceps	0.76	0.17	0.44
Subscapular	0.47	0.25	0.74
Suprailiac	0.33	0.26	0.67
Mid-thigh	0.16	0.79	0.27
Medial calf	0.20	0.70	0.13
<i>Variance explained</i>	25.8%	23.2%	23.2%

TABLE 5

Comparisons (*t*-tests) of adiposity factor scores between time periods within socioeconomic status (SES) groups.

Factor	1988–89 factor scores		2007–08 factor scores		Difference	<i>t</i> (df)	<i>P</i> *
	Mean	SD	Mean	SD			
“Overall Adiposity“							
Higher SES	−0.33	0.76	0.21	0.96	+0.54	5.93 (545)	< 0.001
Middle SES	−0.23	0.79	0.63	1.04	+0.86	13.36 (851)	< 0.001
Lower SES	−0.09	0.91	0.66	1.16	+0.75	9.29 (728)	< 0.001
“Arms“							
Higher SES	−0.13	0.55	0.70	0.72	+0.83	12.52 (545)	< 0.001
Middle SES	−0.32	0.63	0.82	0.80	+1.15	22.52 (851)	< 0.001
Lower SES	−0.39	0.75	0.76	0.94	+1.15	17.50 (728)	< 0.001
“Legs“							
Higher SES	0.03	0.80	−0.55	0.69	−0.58	−6.51 (545)	< 0.001
Middle SES	0.29	0.86	−0.29	0.72	−0.58	−9.53 (851)	< 0.001
Lower SES	0.03	0.86	−0.33	0.87	−0.36	−5.20 (728)	< 0.001
“Trunk“							
Higher SES	−0.42	0.69	−0.04	0.80	+0.38	4.67 (545)	< 0.001
Middle SES	−0.24	0.67	0.33	0.87	+0.56	10.33 (851)	< 0.001
Lower SES	0.28	0.80	0.49	0.79	+0.21	3.24 (728)	0.001

* Tukey-Kramer adjustment for multiple comparisons