Published in final edited form as:

J Am Chem Soc. 2020 March 04; 142(9): 4390–4399. doi:10.1021/jacs.9b13173.

Multicomponent Pyrazole Synthesis from Alkynes, Nitriles, and Titanium Imido Complexes via Oxidatively Induced N–N Bond Coupling

Adam J. Pearce,

Department of Chemistry, University of Minnesota–Twin Cities, Minneapolis, Minnesota 55455, United States

Robin P. Harkins,

Department of Chemistry, University of Minnesota–Twin Cities, Minneapolis, Minnesota 55455, United States

Benjamin R. Reiner,

Department of Chemistry, University of Minnesota–Twin Cities, Minneapolis, Minnesota 55455, United States

Alexander C. Wotal,

Department of Chemistry, University of Minnesota–Twin Cities, Minneapolis, Minnesota 55455, United States

Rachel J. Dunscomb,

Department of Chemistry, University of Minnesota–Twin Cities, Minneapolis, Minnesota 55455, United States

Ian A. Tonks

Department of Chemistry, University of Minnesota–Twin Cities, Minneapolis, Minnesota 55455, United States

Abstract

Pyrazoles are an important class of heterocycles found in a wide range of bioactive compounds and pharmaceuticals. Pyrazole synthesis often requires hydrazine or related reagents where an intact N–N bond is conservatively installed into a pyrazole precursor fragment. Herein, we report the multicomponent oxidative coupling of alkynes, nitriles, and Ti imido complexes for the synthesis of multisubstituted pyrazoles. This modular method avoids potentially hazardous reagents like hydrazine, instead forming the N–N bond in the final step via oxidation-induced coupling on Ti. The mechanism of this transformation has been studied in-depth through stoichiometric reactions of the key diazatitanacyclohexadiene intermediate, which can be accessed

Corresponding Author: itonks@umn.edu.

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/jacs.9b13173>

The authors declare no competing financial interest.

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.9b13173>.

Full experimental and computational details, including Figures S1–S109 and Tables S1 and S2 (PDF) CIF files for 5a, 6a, 7a, 7b, 9a, 10a, and S1 (ZIP)

via multicomponent coupling of Ti imidos with nitriles and alkynes, ring opening of 2-imino-2*H*-azirines, or direct metalation of 4-azadiene-1-amine derivatives. The critical transformation in this reaction is the 2-electron oxidation-induced N–N coupling on Ti. This is a rare example of formal N–N coupling on a metal center, which likely occurs through an electrocyclic mechanism analogous to a Nazarov cyclization. Conveniently, these 2-electron-oxidized diazatitanacyclohexadiene intermediates can be accessed via disproportionation of the 1-electron-oxidized species, which allows utilization of weak oxidants such as TEMPO

Graphical Abstract

INTRODUCTION

Pyrazoles are an important class of N–N-bond-containing heterocycles found in flagship pharmaceuticals such as Eliquis, Viagra, Celebrex, and Acomplia.^{1–3} Multisubstituted pyrazoles are most commonly synthesized via the Knorr condensation of hydrazines onto 1,3-diketones^{4–6} or through a 1,3-dipolar cycloaddition of hydrazones with alkynes or electron-deficient alkenes after autoxidation.^{7–15} Direct N–N bond formation is an attractive alternative to these classical methods, as it would remove the need to use potentially toxic and/or explosive hydrazine reagents. However, N–N bond-forming methods are limited, particularly those proceeding through formal reductive elimination at a transition metal center. This is due to the unfavorable thermodynamics of N–N bond reductive elimination, which results in a weak N–N bond at the expense of two strong metal–N bonds. Despite this challenge, there are a few elegant examples of late transition metal facilitated N–N bond reductive eliminations: (1) Cu-catalyzed or -mediated synthesis of pyrazoles from nitriles and enamines (Figure 1a),^{16,17} as well as a few related transformations^{18–22} and (2) a stoichiometric example featuring a binuclear Ni complex which couples triazabicyclodecene ligands after oxidation with hypervalent iodine reagent PhICl₂ (Figure 1b).²³ In the pyrazole examples, the aromaticity gained provides driving force to overcome the otherwise weak N–N bond (approximately 24 kcal/mol for the parent pyrazole²⁴). In the binuclear Ni example, generation of high-valent metal centers makes reductive elimination more favorable. Oxidatively induced reductive elimination is a common strategy employed in late-transition-

metal organometallic chemistry to overcome otherwise challenging couplings.^{25–32} Moreover, increasing the oxidation state of the system as a whole has been shown to increase the electrophilicity of the *ligands* toward nucleophilic coupling.³³ Other aerobically triggered reductive eliminations have been reported for the synthesis of hydrazines and azoarenes,³⁴ and there is also an example of air oxidation of a d⁰ yttrium triamide complex that liberates aminyl radicals competent for N–N coupling.³⁵

Our group has produced several examples of early transition metal-catalyzed oxidative aminations of alkynes for the synthesis of pyrroles,^{36–40} which proceeds through an electrocyclic reductive elimination to form a C–N bond from an azatitanacyclohexadiene intermediate (Figure 1c, bottom right).⁴¹ We envisioned that reaction of an analogous diazatitanacyclohexadiene^{42,43}—formed from the reaction between a nitrile and the common azatitanacyclobutene intermediate^{44–54}—could result in a formal N–N bond reductive elimination to produce multisubstituted pyrazoles, either through catalytic turnover with azobenzene as an oxidant or by further reaction with an exogenous oxidant (Figure 1c, bottom left). Herein, we report an example of oxidation-induced N–N bond reductive elimination, yielding pyrazoles from diazatitanacyclohexadienes (Figure 1c, top). These diazatitanacyclohexadienes can be generated from the coupling of an alkyne and a nitrile with a Ti imido complex, providing a practical and simple multicomponent [2+2+1] route to pyrazoles. Mechanistic studies indicate that N–N coupling is triggered by 2-electron oxidation of the diazatitana- cyclohexadiene, which generates an intermediate that can undergo electrocyclic ring closure in analogy to a Nazarov-type cyclization.

RESULTS AND DISCUSSION

Multicomponent Coupling for the Synthesis of Pyrazoles

Initial catalytic multicomponent coupling attempts of azobenzene, alkynes, and nitriles catalyzed by [py₂TiCl₂(NPh)]₂, were met with limited success. For example, reaction of *p*-tolunitrile, 3-hexyne, and azobenzene for 20 h at 145 °C yields 16% of the desired 4,5-diethyl-*N*-phenyl-3-*p*-tolylpyrazole (1) in addition to 57% *N*-phenyl-3-hexanimine (2) and 9% 2,3,4,5-tetraethyl-*N*-phenylpyrrole (3) (Figure 2). Although the yield of 1 is lower than the catalyst loading, we were able to demonstrate catalytic turnover using a labeled azobenzene, ArNNAr (Ar = *p*-(CF₃O)C₆H₄), which showed incorporation of both Ph (from the precatalyst) and Ar into the pyrazole product (Table S2).

A potential mechanism for the multicomponent catalytic reaction was computed in analogy to our previous multicomponent synthesis of pyrroles (Figure 3). Computations indicate that the key intermediate diazatitanacyclohexadiene **IM5-pyz** is very stable ($G = -17.4$ kcal/mol relative to py₃TiCl₂(NPh)) and the barrier to N–N reductive elimination (TS3-pyz 47.1 kcal/mol from **IM5-pyz**) is overwhelmingly high. In fact, ¹H NMR monitoring of the reaction shows the formation of a new species consistent with the structure of **IM5-pyz** at 90% yield relative to catalyst within 10 min, suggesting the stability of **IM5-pyz** is inhibitive for productive catalysis. While the formation of 1a demonstrates that diazatitanacyclohexadiene **IM5-pyz** or another analogous species is capable of N–N coupling, we suspect that the high barrier to ring closure likely facilitates competitive acid/base chemistry to occur, ultimately leading to hydroamination. In fact, protonated

diazatitanacycles crystallize from the reaction mixture (see Supporting Information). Thus, unlike previous examples of Ti-catalyzed nitrene transfer, PhNNPh may not be a strong enough oxidant to drive productive N–N bond-forming catalysis.

Diazatitanacyclohexadiene Synthesis

We next synthesized several diazatitanacyclohexadiene model complexes in order to study potential routes to productive N–N bond formation with other oxidants beyond PhNNPh (Figure 4). Protonolysis of $\text{TiCl}_2(\text{NMe}_2)_2$ with a 4-azadiene-1-amine (H_2ADA) (**4a**) gives $(\text{HNMe}_2)\text{TiCl}_2(\text{ADA}^{\text{Ph}})$ (**5a**) in 46% isolated yield. Reaction of H_2ADA (**4a**, **4b**) with $\text{TiCl}_4(\text{THF})_2$ in THF followed by the addition of 2 equiv of LiHMDS (HMDS = hexamethyldisilazide) produces the dinuclear species $[\text{TiCl}_2(\text{ADA})]_2$ (**6a**, **6b**), which is bridged through the imido ligands. Treatment of **6a** and **6b** with pyridine generates $\text{py}_2\text{TiCl}_2(\text{ADA})$ (**7a**, **7b**) in 30% crystallized yield. The X-ray crystal structures for **5a** and **7a** are shown in Figure 4. **5a**, **7a**, and **7b** all have similar solid-state metrics and contain a titanium vinylimido with a pendent imine donor. For example, **7a** has a Ti1–N1 distance of 1.768(1) Å, indicative of Ti–N imido multiple bonding, and a Ti1–N2 distance of 2.126(1) Å, indicative of a Ti–N imine dative bond. These bond lengths are consistent with the computed values for **IM5-pyz** (Figure 3, bottom), where Ti1–N1 and Ti–N2 were computed to be 1.71701 and 2.10562 Å, respectively. Further reactions were carried out with **7a** and **7b** because the HNMe_2 ligand in **5a** can potentially participate in deleterious acid/base reactions.

Diazatitanacyclohexadiene Oxidation

Although **7a** and **7b** are both Ti^{IV} , we hypothesized that ligand-based oxidation could increase the electrophilicity of N_α and promote N–N coupling in analogy to late transition metal-based oxidations that induce reductive elimination. In fact, several oxidants promote facile, room temperature N–N coupling with **7a** and **7b** (Table 1). Reaction of **7a** and **7b** with 2 equiv of TEMPO resulted in 80% isolated yield of the N–N coupled pyrazole product **1a** or **1b** after 2 h (Table 1, entry 1), respectively. When the reaction is performed with 1 equiv of TEMPO, a 1:1 mixture of unreacted **7a** and pyrazole **1a** is observed, which is indicative of a net 2-electron stoichiometry requirement for the reaction (Table 1, entry 2). Likewise, hypervalent iodine reagents PhICl_2 , $\text{PhI}(\text{TFA})_2$, and $\text{PhI}(\text{OAc})_2$ all resulted in good yields of **1a** or **1b** (Table 1, entries 3–5). Interestingly, oxidation of **7a** by $\text{PhI}(\text{OAc})_2$ (Table 1, entry 5) results in an initial mixture of **1a** with a kinetic azirine product (**8a**), which slowly converts to **1a** over time. Intermediacy of **8a** could indicate potential free nitrene reactivity^{55–57} or transmetalation to I, but control experiments rule out these possibilities (Figures S90 and S96). The intermediacy of **8a** was not evident in any other reaction. Other oxidants also engender productive N–N coupling, including FcPF_6 ($\text{Fc} = \text{Cp}_2\text{Fe}^+$) and even O_2 , albeit in lower yields. Prolonged exposure of **7a** to PhNNPh even at elevated temperatures (up to 145 °C) did not furnish tractable amounts of **1a**, consistent with early catalytic attempts. Thermolysis of **7a** in the absence of any oxidant also only led to trace amounts of **1a**. N–N bond coupling was not observed when reacting with several common oxidants such as I_2 , NBS, NOBF_4 , and Me_3NO , instead leading to intractable mixtures.

***In Situ* Multicomponent Pyrazole Synthesis**

One-pot multicomponent synthesis via alkynes and nitriles is an attractive, unrealized, and simple route to pyrazoles. Given that diazatitanacycles have been observed as long-lived intermediates in our initial catalytic reaction attempts, it seemed plausible that they could be generated *in situ* by reaction of an alkyne and nitrile with a Ti imido and then oxidized to generate the desired pyrazole product in a one-pot procedure (Table 2). Indeed, diazatitanacycles analogous to **7a** and **7b** can be synthesized in 81% *in situ* yield (¹H NMR) by reaction of [py₂TiCl₂(NPh)]₂ with an excess of *p*-tolunitrile (6 equiv; 3 equiv with respect to [Ti]) and 3-hexyne (6 equiv) in C₆D₅Br after 1 h at 115 °C (Table 2, entry 1). C₆D₅Br solvent was chosen because it is an affordable-to-access polar NMR solvent, and was successful for our previous [2+2+1] pyrrole cyclizations.⁴¹ Addition of TEMPO after diazatitanacyclohexadiene formation, followed by heating at 50 °C for 2 h affords pyrazole **1** in 75% yield with respect to [py₂TiCl₂(NPh)]₂ (98% yield with respect to **IM5-pyz**). Based off of this result, comparative reactions were next carried out at a 3:3:1 and 1:1:1 stoichiometry of benzonitrile, 3-hexyne, and Ti=NR (Table 2, entries 2 and 3). While the 1:1:1 stoichiometry results in a lower conversion to the metallacyclic intermediate (Figures S54 and S57), **1b** is still generated in 56% overall yield with respect to alkyne (Table 2, entry 3). This reaction could be performed at 1 mmol scale, resulting in 65% yield of **1b** (see Supporting Information). Inferring that the alkyne and nitrile fragments may often be the more precious components of this reaction, we next briefly explored the scope of this reaction at the 1:1:1 stoichiometry. In all cases, the desired pyrazole products were furnished in moderate to good yield. This method can tolerate both electron-rich and electron-poor nitriles (entries 3–5). *p*-(Trifluoromethyl)benzonitrile affects a slightly lower yield (49%) which is explained by detectable amounts of titanacycle decomposition (entry 5). Alkyl nitriles (entries 6–8) are also effective coupling partners in the reaction; however, the yield of diazatitanacycle oxidation is poor when coupling acetonitrile to 1-phenyl-1-propyne (entry 8). 1-Phenyl-1-propyne couples with both acetonitrile and benzonitrile, affording the 5-methyl-4-phenyl regioisomers selectively (entries 8 and 9); notably, 5-alkyl-3-arylpyrazoles are challenging to access via Knorr pyrazole synthesis due to the regioselectivity of the hydrazine condensation step. Tetraarylpyrazoles, which have shown the enhancement of aggregation-induced emission,^{58,59} are accessible in 37% *in situ* yield for pyrazole **1j** (entry 11) or 42% isolated yield when diphenylacetylene (**1k**) is used (see Supporting Information). This scope demonstrates that multisubstituted pyrazoles can be accessed directly and generally in a single step through multicomponent coupling of Ti imidos, while avoiding prerequisite hydrazine or hydrazine-derived coupling partners. Conveniently, [py₂TiCl₂(NPh)]₂ and derivatives thereof can be prepared in a single pot from TiCl₄(THF)₂, and demonstrated benchtop-compatible Ti imido generation.⁶⁰ Thus, access to these multicomponent coupling reactions does not require specialized equipment or protocols.

Mechanistic Studies of N–N Coupling

N–N coupling on Ti could plausibly occur through one (or more) of three possible oxidation states—either the “default” oxidation state, 1-electron-oxidized, or 2-electron-oxidized (Figure 5, top). In order to determine the system oxidation state that triggers N–N coupling

on Ti, we sought to conduct N–N coupling reactions with the entire redox series. Previous work in our group has shown that $\text{Cp}_2\text{Ti}(\text{BTMSA})$ ($\text{BTMSA} = \text{bis}(\text{trimethylsilyl})\text{acetylene}$)^{61–65} rapidly ring-opens 2*H*-azirines at the C–N bond.⁶⁶ Thus, we hypothesized 2-imino-2*H*-azirine **8a**—an isomer of pyrazole **1a**—could react with a Ti^{II} source to give a diazatitanacyclohexadiene in the “default” oxidation state. By extension of this logic, reacting 2-imino-2*H*-azirines with Ti^{III} and Ti^{IV} precursors could allow for interrogation of 1- and 2-electron-oxidized diazatitana- cyclohexadiene species, respectively.

The reaction of Ti^{II} , Ti^{III} , and Ti^{IV} precursors with the 2-imino-2*H*-azirine **8a** is shown in Figure 5. Reaction of **8a** and $\text{Cp}_2\text{Ti}(\text{BTMSA})$ rapidly gives the expected diazatitanacyclohexadiene $\text{Cp}_2\text{Ti}(\text{ADA}^{\text{Ph}})$ (**9a**) in 89% yield (Figure 5, reaction II). Interestingly, XRD analysis of **9a** (Figure 6) shows a Ti1–N1 bond distance of 1.857(1) Å and a Ti1–N2 distance of 2.121(1) Å, indicating an iminyl-enamide bonding description for this complex, contrasting the vinylimidoimine resonance form in complexes **5a**, **7a**, and **7b** (Figure 4). Consistent with earlier results with **7a** and **7b**, no N–N coupling occurs from **9a** in this oxidation state.

Having shown that Ti^{II} reacts cleanly with **8a** to give diazatitanacyclohexadiene **9a**, we moved to investigate the *in situ* formation of a diazatitanacycle radical via radical ring opening of **8a** by $\text{TiCl}_3(\text{THF})_3$ (Figure 5, reaction III). A benzene solution of **8a** reacts instantaneously with a benzene slurry of $\text{TiCl}_3(\text{THF})_3$ yielding a 1:1 mixture of pyrazole **1a** and presumably the “default” oxidation state diazatitanacycle, $(\text{THF})_x\text{TiCl}_2(\text{ADA}^{\text{Ph}})$ (**7a-0e⁻**). Addition of 2 equiv of pyridine generates **7a** with the gradual precipitation of a yellow solid presumed to be $(\text{py})_2\text{TiCl}_4$. The formation of equal parts pyrazole **1a**, **7a**, and TiCl_4 strongly suggests the initial formation of diazatitanacycle radical $\text{TiCl}_3(\text{ADA}^{\text{Ph}})$ (**7a-1e⁻**), followed by disproportionation into the “default” oxidation state and the 2-electron-oxidized species **7a-2e⁻**, which then couples to form pyrazole.

Given the disproportionation reactivity with Ti^{III} , we then tested the reactivity of 2-imino-2*H*-azirines with Ti^{IV} (Figure 5, reaction IV). Treatment of azirine **8a** with $\text{TiCl}_4(\text{THF})_2$ generates an unidentified intermediate species, which upon heating to 50 °C results in conversion to **1a** as the major product. Alper demonstrated that TiCl_4 will ring open 2*H*-azirines, in which γ -chlorination occurs after C–N bond cleavage.⁶⁷ By analogy, 2-imino-2*H*-azirines likely undergo similar reactivity to form **7a-Cl-2e⁻**, which can then undergo N–N coupling upon chloride loss. Consistent with this mechanism, there are examples of late transition metal-catalyzed 2-acyl-2*H*-azirine or 2-imino-2*H*-azirine rearrangements to isoxazoles or pyrazoles, respectively, that may proceed through similar vinyl-nitrenoid intermediates.^{68–71}

Reactivity of $\text{Cp}_2\text{Ti}(\text{ADA}^{\text{Ph}})$ (**9a**) with PhICl_2

The 2*H*-azirine ring-opening redox series in Figure 5 strongly suggests that N–N coupling to form pyrazoles occurs through the 2-electron-oxidized diazatitanacycle of the form **7a-2e⁻**. We next aimed to synthesize analogues of the **7a-2e⁻** intermediate and observe their direct N–N coupling reactivity. We hoped that the stability imbued by the titanocene moiety in **9a** would allow for isolation of an intermediate in the PhICl_2 oxidations of

diazatitanacyclohexadienes. Mixing a dark claret C₆D₆ solution of **9a** with PhICl₂ at room temperature immediately furnishes a bright yellow solution, yielding **10a** quantitatively (Figure 6), which exists as two isomers. The solid-state structure of **10a** (Figure 6) reveals chlorination of Ti and the C_γ position of the diazatitanacyclohexadiene ring. **10a** is a “2-electron-oxidized” organic moiety, similar to the proposed azirine ring-opened product from TiCl₄(THF)₂ **7a-Cl-2e⁻** (Figure 5, reaction IV) and Alper’s earlier observations of γ-chlorination. However, steric encumbrance imposed by the Cp rings in **10a** likely prevents imine coordination to Ti. Interestingly, **9a** does not react with TEMPO, which may indicate that reaction of TEMPO with **7a** initially requires coordination to Ti.

Refluxing **10a** in C₆D₆ for 22 h gives pyrazole **1a** and Cp₂TiCl₂ as the major products (Figure 6), consistent with 2-imino-2*H*-azirine ring-opening by TiCl₄(THF)₂. The slower rate of decomposition of **10a** into pyrazole compared to the reaction of **8a** and TiCl₄ is likely due to the hindered approach of the imine during N–N coupling. Further, cyclic voltammetry of **9a** features a quasi-reversible 1-electron oxidation at –0.55 mV vs Fc^{0/+} which exhibits fully reversible behavior at scan rates > 200 mV/s (Figure 7). A linear dependence of *i*_{pa} on $v^{1/2}$ was observed, consistent with a mass-transfer-limited process. This suggests that the quasi-reversibility of **9a** oxidation at low scan rates arises from a chemical event coupled to oxidation, which, based on the 2-imino-2*H*-azirine reactivity above, is likely related to disproportionation of the 1-electron-oxidized Cp₂Ti(ADA^{Ph•}) en route to **1a** formation. Notably, no other current deflection is observed in the anodic sweep before oxidation of the supporting electrolyte at ~700 mV.

Discussion of the Mechanism of Oxidatively Induced N–N Coupling Mediated by Ti

We propose the mechanism for N–N coupling in Figure 8 based on the stoichiometric oxidations of **7a,b**, the 2*H*-azirine **8a** ring-opening reactivity with various Ti oxidation states, and the reactivity and electrochemistry of **9a**. First, multicomponent coupling of a Ti imido with an alkyne and nitrile via the previously established [2+2]/insertion mechanism^{44,49} gives diazatitanacycle **A**, arbitrarily considered to be in the “default” oxidation state of this system. As demonstrated in the series of redox experiments above (Figure 5, reaction II), **A** can also be accessed via ring-opening of 2-imino-2*H*-azirines by a Ti^{II} precursor such as Cp₂Ti(BTMSA). The barrier for N–N reductive elimination from **A** is too high, facilitating entry into competitive decomposition pathways. Single-electron oxidation of **A** leads to the 1-electron-oxidized diazatitanacycle **B**, which is unstable. From **B**, there are two possible reaction pathways: (1) disproportionation to **A** and 2-electron-oxidized **C**, which undergoes rapid N–N coupling to form **1** and a TiX₄ coproduct, or (2) further oxidation to **D**, as seen in the oxidation of **9a** to **10a** by PhICl₂. The reaction of **8a** with TiCl₃(THF)₃, which results in 50% N–N coupled **1a** and 50% of the “default” oxidation state **7a** (Figure 5, reaction III), provides evidence for disproportionation. Further, weak oxidants such as TEMPO or FcPF₆ result in product formation despite an inability to oxidize **7a** by two electrons, indicating that a disproportionation mechanism is likely in play.

Although **D** is also 2-electron-oxidized like **C**, C–X dissociation to **C** is likely necessary for rapid N–N coupling to occur. Evidence for this can be seen in several reactions. For example, reaction of TiCl₄(THF)₂ with **8a** (Figure 5, reaction IV) results in full conversion

to **1a**, but the reaction is slower than that of $\text{TiCl}_3(\text{THF})_3$ and requires heat to drive it to completion. Based on work from Alper, reaction IV likely generates an intermediate analogous to **D**. Thus, the major difference between the oxidized diazatitanacycles in reactions III and IV is the presence of an extra equivalent of Cl^- in IV; indicating that dissociation of halide from **D** to make the carbocationic diazatitanacyclic **C** is likely kinetically relevant. Further evidence for the role of X^- dissociation from **D** may also be seen in the overall yields with hypervalent iodine oxidants presented in Table 1: reactions with anion equivalents that would be poor leaving groups (e.g., OAc^-) generally result in lower yields than more weakly coordinating anion equivalents and/or better leaving groups (Cl^- , O_2CCF_3^-).

Based on the above evidence, the 2-electron-oxidized diazatitanacyclic **C** is likely responsible for rapid N–N bond formation, which depending on oxidant strength can be accessed either through disproportionation of a 1-electron-oxidized species or directly through 2-electron oxidation. We have previously established in related [2+2+1] pyrrole syntheses that C–N bond reductive elimination from a 6-membered azatitanacyclic proceeds through an electrocyclic mechanism. Here, oxidation of the diazatitanacyclic should make the α -N significantly more electrophilic, prompting ringclosure in analogy to a Nazarov-like electrocyclization (Figure 9). The kinetic slowdown in the presence of coordinating anions is further evidence for an electrocyclic mechanism: intermediate **C**, with full π -conjugation, undergoes rapid reaction, whereas intermediate **D** lacks the conjugated π -system requisite for electrocyclization.

CONCLUSION

In summary, we have shown that simple titanium imido complexes are capable of selective coupling with alkynes and nitriles to generate diazatitanacyclohexadienes, which can be oxidized under mild conditions to generate pyrazoles in good yield. By exploring an electrochemical series of 2-imino-2*H*-azirine ring openings with Ti^{II} , Ti^{III} , and Ti^{IV} , we have demonstrated that Ti-mediated N–N coupling occurs from 2-electron-oxidized diazatitanacyclohexadienes via a potentially electrocyclic process. While oxidatively induced reductive elimination has been extensively explored, the concept of d^0 transition metals mediating oxidatively induced couplings through ligand-centered oxidations may be a general strategy for accessing new reactivity and myriad new types of metal-mediated bond couplings, particularly via these types of electrocyclic pathways.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We acknowledge the Minnesota Supercomputing Institute (MSI) at the University of Minnesota for providing resources that contributed to the results reported within this paper. A.J.P. would like to thank Victor Young Jr. for assistance with XRD analysis, as well as Evan P. Beaumier and Xin Yi See for enlightening discussions and generously sharing reagents.

Funding

Financial support was provided by the National Institutes of Health (1R35GM119457) and the Alfred P. Sloan Foundation (L.A.T. is a 2017 Sloan Fellow). Instrumentation for the University of Minnesota Chemistry NMR facility was supported through a grant through the National Institutes of Health (S10OD011952). X-ray diffraction experiments were performed with a diffractometer purchased through a grant from NSF/MRI (1229400) and the University of Minnesota.

REFERENCES

- (1). Ansari A; Ali A; Asif M; Shamsuzzaman S Review: Biologically Active Pyrazole Derivatives. *New J. Chem.* 2017, 41, 16–41.
- (2). Karrouchi K; Radi S; Ramli Y; Taoufik J; Mabkhot Y; Al- Aizari F; Ansar M Synthesis and Pharmacological Activities of Pyrazole Derivatives: A Review. *Molecules* 2018, 23, 134.
- (3). Fustero S; Sánchez-Roselló M; Barrio P; Simón-Fuentes A From 2000 to Mid-2010: A Fruitful Decade for the Synthesis of Pyrazoles. *Chem. Rev.* 2011, 111, 6984–7034. [PubMed: 21806021]
- (4). Knorr L Einwirkung von Acetessigester Auf Phenylhydrazin. *Ber. Dtsch. Chem. Ges.* 1883, 16, 2597–2599.
- (5). Silva VLM; Silva AMS; Pinto DCGA; Cavaleiro JAS; Elguero J 3(5)-(2-Hydroxyphenyl)-5(3)-Styrylpyrazoles: Synthesis and Diels-Alder Transformations. *Eur. J. Org. Chem.* 2004, 2004, 4348–4356.
- (6). Heller ST; Natarajan SR 1,3-Diketones from Acid Chlorides and Ketones: A Rapid and General One-Pot Synthesis of Pyrazoles. *Org. Lett.* 2006, 8, 2675–2678. [PubMed: 16774229]
- (7). Snider BB; Conn RSE; Sealfon S Reactions of Phenylhydrazones with Electron-Deficient Alkenes. *J. Org. Chem.* 1979, 44, 218–221.
- (8). Gómez Guillén M; Conde Jiménez JL Reaction of D-Galactose Phenylhydrazone with Nitroalkenes: Synthesis of Pentahydroxypentylpyrazoles. *Carbohydr. Res.* 1988, 180, 1–17.
- (9). Gómez-Guillén M; Hans-Hans F; Simon JML; Martín-Zamora ME New Pentahydroxypentylpyrazoles from the Reactions of D-Mannose and d-Galactose Methylhydrazones with Nitroalkenes. *Carbohydr. Res.* 1989, 189, 349–358.
- (10). Arrieta A; Ramón Carrillo J; Cossío FP; Díaz-Ortiz A; JoséGómez-Escalonilla M; de la Hoz A; Langa F; Moreno A Efficient Tautomerization Hydrazone-Azomethine Imine under Microwave Irradiation. Synthesis of [4,3'] and [5,3']Bipyrazoles. *Tetrahedron* 1998, 54, 13167–13180.
- (11). Deng X; Mani NS Reaction of N-Monosubstituted Hydrazones with Nitroolefins: A Novel Regioselective Pyrazole Synthesis. *Org. Lett.* 2006, 8, 3505–3508. [PubMed: 16869646]
- (12). Tang M; Wang Y; Wang H; Kong Y Aluminum Chloride Mediated Reactions of N-Alkylated Tosyl-hydrazones and Terminal Alkynes: A Regioselective Approach to 1,3,5-Trisubstituted Pyrazoles. *Synthesis* 2016, 48, 3065–3076.
- (13). Kong Y; Tang M; Wang Y Regioselective Synthesis of 1,3,5-Trisubstituted Pyrazoles from N -Alkylated Tosylhydrazones and Terminal Alkynes. *Org. Lett.* 2014, 16, 576–579. [PubMed: 24432814]
- (14). Ma C; Li Y; Wen P; Yan R; Ren Z; Huang G Copper(I)-Catalyzed Synthesis of Pyrazoles from Phenylhydrazones and Dialkyl Ethylenedicarboxylates in the Presence of Bases. *Synlett* 2011, 2011, 1321–1323.
- (15). Deng X; Mani NS Regioselective Synthesis of 1,3,5-Tri- and 1,3,4,5-Tetrasubstituted Pyrazoles from N -Arylhidrazones and Nitroolefins. *J. Org. Chem.* 2008, 73, 2412–2415. [PubMed: 18278943]
- (16). Neumann JJ; Suri M; Glorius F Efficient Synthesis of Pyrazoles: Oxidative C-C/N-N Bond-Formation Cascade. *Angew. Chem., Int. Ed.* 2010, 49, 7790–7794.
- (17). Suri M; Jousseau T; Neumann JJ; Glorius F An Efficient Copper-Catalyzed Formation of Highly Substituted Pyrazoles Using Molecular Oxygen as the Oxidant. *Green Chem.* 2012, 14, 2193.
- (18). Ueda S; Nagasawa H Facile Synthesis of 1,2,4-Triazoles via a Copper-Catalyzed Tandem Addition-Oxidative Cyclization. *J. Am. Chem. Soc.* 2009, 131, 15080–15081. [PubMed: 19799379]

- (19). Tang X; Huang L; Yang J; Xu Y; Wu W; Jiang H Practical Synthesis of Pyrazoles via a Copper-Catalyzed Relay Oxidation Strategy. *Chem. Commun.* 2014, 50, 14793–14796.
- (20). Chen B; Zhu C; Tang Y; Ma S Copper-Mediated Pyrazole Synthesis from 2,3-Alkenoates or 2-Alkynoates, Amines and Nitriles. *Chem. Commun.* 2014, 50, 7677.
- (21). Dai G; Yang L; Zhou W Copper-Catalyzed Oxidative Dehydrogenative N-N Bond Formation for the Synthesis of N, N'-Diarylindazol-3-Ones. *Org. Chem. Front* 2017, 4, 229–231.
- (22). Ryan MC; Kim YJ; Gerken JB; Wang F; Aristov MM; Martinelli JR; Stahl SS Mechanistic Insights into Copper-Catalyzed Aerobic Oxidative Coupling of N–N Bonds. *Chem. Sci.* 2020, 11, 1170–1175.
- (23). Diccianni JB; Hu C; Diao T N-N Bond Forming Reductive Elimination via a Mixed-Valent Nickel(n)-Nickel(IH) Intermediate. *Angew. Chem., Int Ed.* 2016, 55, 7534–7538.
- (24). Cyra ski MK Energetic Aspects of Cyclic Pi-Electron Delocalization: Evaluation of the Methods of Estimating Aromatic Stabilization Energies. *Chem. Rev.* 2005, 105, 3773–3811. [PubMed: 16218567]
- (25). Bour JR; Camasso NM; Sanford MS Oxidation of Ni(II) to Ni(IV) with Aryl Electrophiles Enables Ni-Mediated Aryl-CF₃ Coupling. *J. Am. Chem. Soc.* 2015, 137, 8034. [PubMed: 26079544]
- (26). Lau W; Huffman JC; Kochi JK Electrochemical Oxidation-Reduction of Organometallic Complexes. Effect of the Oxidation State on the Pathways for Reductive Elimination of Dialkyliron Complexes. *Organometallics* 1982, 1, 155–169.
- (27). Kim J; Shin K; Jin S; Kim D; Chang S Oxidatively Induced Reductive Elimination: Exploring the Scope and Catalyst Systems with Ir, Rh, and Ru Complexes. *J. Am. Chem. Soc.* 2019, 141, 4137–4146. [PubMed: 30762365]
- (28). Lanci MP; Remy MS; Kaminsky W; Mayer JM; Sanford MS Oxidatively Induced Reductive Elimination from (tBu₂Bpy)Pd(Me)₂: Palladium(IV) Intermediates in a One-Electron Oxidation Reaction. *J. Am. Chem. Soc.* 2009, 131, 15618–15620. [PubMed: 19824643]
- (29). Kraatz H-B; van der Boom ME; Ben-David Y; Milstein D Reaction of Aryl Iodides with (PCP)Pd(II)-Alkyl and Aryl Complexes: Mechanistic Aspects of Carbon-Carbon Bond Formation. *Isr. J. Chem.* 2001, 41, 163–172.
- (30). Watson MB; Rath NP; Mirica LM Oxidative C–C Bond Formation Reactivity of Organometallic Ni(II), Ni(III), and Ni(IV) Complexes. *J. Am. Chem. Soc.* 2017, 139, 35–38. [PubMed: 27997181]
- (31). Tsou TT; Kochi JK Mechanism of Biaryl Synthesis with Nickel Complexes. *J. Am. Chem. Soc.* 1979, 101, 7547–7560.
- (32). Tsou TT; Kochi JK Reductive Coupling of Organometals Induced by Oxidation. Detection of Metastable Paramagnetic Intermediates. *J. Am. Chem. Soc.* 1978, 100, 1634–1635.
- (33). Camasso NM; Sanford MS Design, Synthesis, and CarbonHeteroatom Coupling Reactions of Organometallic Nickel(IV) Complexes. *Science* 2015, 347, 1218–1220. [PubMed: 25766226]
- (34). Kinoshita K On the Mechanism of Oxidation by Cuprous Chloride, Pyridine and Air. I. The Properties of the Reaction. *Bull. Chem. Soc. Jpn.* 1959, 32, 777–780.
- (35). Zhang L; Xia J; Li Q; Li X; Wang S Fast Synthesis of Hydrazine and Azo Derivatives by Oxidation of Rare-Earth-Metal-Nitrogen Bonds. *Organometallics* 2011, 30, 375–378.
- (36). Pearce AJ; See XY; Tonks IA Oxidative Nitrene Transfer from Azides to Alkynes via Ti(II)/Ti(IV) Redox Catalysis: Formal [2+2+1] Synthesis of Pyrroles. *Chem. Commun.* 2018, 54, 6891–6894.
- (37). Gilbert ZW; Hue RJ; Tonks IA Catalytic Formal [2+2+1] Synthesis of Pyrroles from Alkynes and Diazenes via TiII/TiIV Redox Catalysis. *Nat. Chem* 2016, 8, 63–68. [PubMed: 26673265]
- (38). Chiu H-C; Tonks IA Trimethylsilyl-Protected Alkynes as Selective Cross-Coupling Partners in Titanium-Catalyzed [2+2+1] Pyrrole Synthesis. *Angew. Chem., Int. Ed.* 2018, 57, 6090–6094.
- (39). Chiu H-C; See XY; Tonks IA Dative Directing Group Effects in Ti-Catalyzed [2+2+1] Pyrrole Synthesis: Chemo- and Regioselective Alkyne Heterocoupling. *ACS Catal* 2019, 9, 216–223. [PubMed: 31768294]

- (40). Kawakita K; Beaumier EP; Kakiuchi Y; Tsurugi H; Tonks IA; Mashima K Bis(Imido)Vanadium(V)-Catalyzed [2+2+1] Coupling of Alkynes and Azobenzenes Giving Multisubstituted Pyrroles. *J. Am. Chem. Soc.* 2019, 141, 4194–4198. [PubMed: 30731038]
- (41). Davis-Gilbert ZW; Wen X; Goodpaster JD; Tonks IA Mechanism of Ti-Catalyzed Oxidative Nitrene Transfer in [2+2+1] Pyrrole Synthesis from Alkynes and Azobenzene. *J. Am. Chem. Soc.* 2018, 140, 7267–7281. [PubMed: 29763560]
- (42). Doxsee KM; Farahi JB; Hope H Synthesis and Formal [4 + 2] Cycloaddition Reactions of Vinylimido Complexes of Titanocene. *J. Am. Chem. Soc.* 1991, 113, 8889–8898.
- (43). Doxsee KM; Farahi JB Reductive Coupling of Nitriles via Formal [2+2] Cycloadditions to the Titanium–Carbon Double Bond. *J. Am. Chem. Soc.* 1988, 110, 7239–7240.
- (44). Schofield AD; Nova A; Selby JD; Schwarz AD; Clot E; Mountford P Reaction Site Diversity in the Reactions of Titanium Hydrazides with Organic Nitriles, Isonitriles and Isocyanates: Ti = Na Cycloaddition, Ti = Na Insertion and Na–N β Bond Cleavage. *Chem. - Eur. J* 2011, 17, 265–285. [PubMed: 21207623]
- (45). Manßen M; De Graaff S; Meyer MF; Schmidtman M; Beckhaus R Direct Access to Titanocene Imides via Bis(η^5 : η^1 -Penta-Fulvene)Titanium Complexes and Primary Amines. *Organometallics* 2018, 37, 4506–4514.
- (46). Groom LR; Schwarz AD; Nova A; Clot E; Mountford P Synthesis and Reactions of a Cyclopentadienyl-Amidinate Titanium Tert-Butoxyimido Compound. *Organometallics* 2013, 32, 7520–7539.
- (47). Pugh SM; Trösch DJM; Wilson DJ; Bashall A; Cloke FGN; Gade LH; Hitchcock PB; McPartlin M; Nixon JF; Mountford P Cycloaddition Reactions of the Titanium Imide [Ti(NBut)(MeC(2-C5H4N)(CH2NSiMe3)2)(Py)] with ButCP and MeCN. *Organometallics* 2000, 19, 3205–3210.
- (48). Tiong PJ; Groom LR; Clot E; Mountford P Synthesis, Bonding and Reactivity of a Terminal Titanium Alkylidene Hydrazido Compound. *Chem. - Eur. J* 2013, 19, 4198–4216. [PubMed: 23401149]
- (49). McGrane PL; Jensen M; Livinghouse T Intramolecular [2+2] Cycloadditions of Group IV Metal-Imido Complexes. Applications to the Synthesis of Dihydropyrrole and Tetrahydropyridine Derivatives. *J. Am. Chem. Soc.* 1992, 114, 5459–5460.
- (50). Nguyen TT; Kortman GD; Hull KL Synthesis, Cycloaddition, and Cycloreversion Reactions of Mononuclear Titanocene-Oxo Complexes. *Organometallics* 2016, 35, 1713–1725.
- (51). Barnea E; Majumder S; Staples RJ; Odom AL One-Step Route to 2,3-Diaminopyrroles Using a Titanium-Catalyzed Four-Component Coupling. *Organometallics* 2009, 28, 3876–3881.
- (52). Cao C; Shi Y; Odom AL A Titanium-Catalyzed Three- Component Coupling To Generate α,β -Unsaturated β -Iminoamines. *J. Am. Chem. Soc.* 2003, 125, 2880–2881. [PubMed: 12617647]
- (53). Odom AL; McDaniel TJ Titanium-Catalyzed Multicomponent Couplings: Efficient One-Pot Syntheses of Nitrogen Heterocycles. *Acc. Chem. Res.* 2015, 48, 2822–2833. [PubMed: 26295382]
- (54). Basuli F; Aneetha H; Huffman JC; Mindiola DJ A Fluorobenzene Adduct of Ti(IV), and Catalytic Carboamination to Prepare α,β -Unsaturated Imines and Triaryl-Substituted Quinolines. *J. Am. Chem. Soc.* 2005, 127, 17992–17993. [PubMed: 16366536]
- (55). Padwa A; Smolanoff J; Tremper A Photochemical Transformations of Small Ring Heterocyclic Systems. LXV. Intramolecular Cycloaddition Reactions of Vinyl-Substituted 2H-Azirines. *J. Am. Chem. Soc.* 1975, 97, 4682–4691.
- (56). Nishiwaki T; Fujiyama F Studies on Heterocyclic Chemistry. Part XIII. Cleavage of 5-Benzyl-Amino-Oxazoles, Photoproducts of N-Benzyl-2H-Azirine-2-Carboxamides, by Dialkyl Phosphite. *J. Chem. Soc., Perkin Trans. 1* 1972, 1456.
- (57). Nishiwaki T; Nakano A; Matsuoka H Studies on Heterocyclic Chemistry. Part VII. Thermally Induced Dimerization of 5-Aminoisoxazoles and 2H-Azirines and Photochemistry of 5-Aminoisoxazoles. *J. Chem. Soc. C* 1970, 1825.
- (58). Mukherjee S; Salini PS; Srinivasan A; Peruncheralathan S AIEE Phenomenon: Tetraaryl vs. Triaryl Pyrazoles. *Chem. Commun.* 2015, 51, 17148–17151.

- (59). Mukundam V; Kumar A; Dhanunjayarao K; Ravi A; Peruncheralathan S; Venkatasubbaiah K Tetraaryl Pyrazole Polymers: Versatile Synthesis, Aggregation Induced Emission Enhancement and Detection of Explosives. *Polym. Chem.* 2015, 6, 7764–7770.
- (60). Davis-Gilbert ZW; Kawakita K; Blechschmidt DR; Tsurugi H; Mashima K; Tonks IA In Situ Catalyst Generation and Benchtop-Compatible Entry Points for TiII/TiIV Redox Catalytic Reactions. *Organometallics* 2018, 37, 4439–4445. [PubMed: 31802785]
- (61). Rosenthal U; Pellny PM; Kirchbauer FG; Burlakov VV What Do Titano- and Zirconocenes Do with Dienes and Polyynes? *Acc. Chem. Res.* 2000, 33, 119–129.
- (62). Rosenthal U; Burlakov VV; Arndt P; Baumann W; Spannenberg A The Titanocene Complex of Bis(Trimethylsilyl)-Acetylene: Synthesis, Structure, and Chemistry. *Organometallics* 2003, 22, 884–900.
- (63). Burlakov VV; Polyakov AV; Yanovsky AI; Struchkov YT; Shur VB; Vol'pin ME; Rosenthal U; Gorls H Novel Acetylene Complexes of Titanocene and Permethyltitanocene without Additional Ligands. Synthesis Spectral Characteristics and X-Ray Diffraction Study. *J. Organomet. Chem.* 1994, 476, 197–206.
- (64). Peulecke N; Lefebvre C; Ohff A; Baumann W; Tillack A; Kempe R; Burlakov VV; Rosenthal U Ansa-Titanocene and -Zirconocene η^2 -Alkyne Complexes -Synthesis, Spectral Characteristics, and X-Ray Crystal Structure. *Chem. Ber.* 1996, 129, 959–962.
- (65). Ohff A; Pulst S; Lefebvre C; Peulecke N; Arndt P; Burlakov VV; Rosenthal U Unusual Reactions of Titanocene- and Zirconocene-Generating Complexes. *Synlett* 1996, 1996, 111–118.
- (66). Desnoyer AN; See XY; Tonks IA Diverse Reactivity of Diazatitanacyclohexenes: Coupling Reactions of 2H-Azirines Mediated by Titanium(II). *Organometallics* 2018, 37, 4327–4331.
- (67). Alper H; Prickett JE; Wollowitz S Intermolecular and Intramolecular Cycloaddition Reactions of Azirines by Group 6 Metal Carbonyls and by Titanium Tetrachloride. *J. Am. Chem. Soc.* 1977, 99, 4330–4333.
- (68). Li X; Du Y; Liang Z; Li X; Pan Y; Zhao K Simple Conversion of Enamines to 2H-Azirines and Their Rearrangements under Thermal Conditions. *Org. Lett.* 2009, 11, 2643–2646. [PubMed: 19438258]
- (69). Pinho e Melo TMVD; Lopes CSJ; Rocha Gonsalves AM; Storrs RC Reactivity of 2-Halo-2H-Azirines. Part II. Thermal Ring Expansion Reactions: Synthesis of 4-Haloisoxazoles. *Synthesis* 2002, 2002, 605–608.
- (70). Padwa A; Stengel T Transition Metal Catalyzed Ring Opening Reactions of 2-Phenyl-3-Vinyl Substituted 2H-Azirines. *Tetrahedron Lett.* 2004, 45, 5991–5993.
- (71). Alper H; Sakakibara T An Interesting Azirine Induced Reaction of the Cyclopentadienyliron Dicarbonyl Dimer. *Can. J. Chem.* 1979, 57, 1541–1543.

Figure 1.

(a) Cu-catalyzed oxidation-induced formal N–N reductive elimination for the synthesis of pyrazoles from electron-poor enamines and nitriles. (b) Ni-mediated N–N bond reductive elimination. (c) Formal [2+2+1] multicomponent synthesis of pyrazoles via oxidation-induced N–N coupling, and a proposed catalytic cycle for pyrazoles compared to the previously reported [2+2+1] synthesis of pyrroles.

Figure 2. Multicomponent coupling of p-tolunitrile, 3-hexyne, and azobenzene catalyzed by $\text{py}_2\text{TiCl}_2(\text{NPh})$.

Figure 3.

Computed cycle for pyrazole formation via direct N–N bond coupling (M06/6–311G(d,p)/SMD, 145 °C, $\text{C}_6\text{D}_5\text{Br}$) compared to C–N bond formation in the analogous multicomponent pyrrole synthesis (M06/6–311G(d,p)/SMD, 115 °C, PhCF_3) by Ti imido catalysts. Pyrrole computations and intermediate labels are adapted from ref 37. All free energies are referenced to $\text{py}_3\text{TiCl}_2(\text{NPh}) = 0.0 \text{ kcal/mol}$.

Figure 4. Syntheses of diazatitanacyclohexadienes **5a**, **7a**, and **7b** and ORTEP diagrams of **5a** and **7a**. Thermal ellipsoids are drawn at 50% probability and hydrogen atoms omitted for clarity. Full crystallographic data for **5a**, **7a**, and **7b** is available in the Supporting Information.

Figure 5.
Reaction of **8a** with a series of Ti complexes spanning Ti^{II}, Ti^{III}, and Ti^{IV} oxidation states.

Figure 6. Oxidation of **9a** to **10a** by PhICl_2 followed by thermal elimination of Cp_2TiCl_2 to give **1a**, and ORTEP diagrams of **9a** and **10a**. Thermal ellipsoids are drawn at 50% probability, and hydrogen atoms are omitted for clarity. One crystallographically independent molecule of **10a** and a solvent Et_2O have also been omitted. Full crystallographic data is available in the Supporting Information, including CIF files.

Figure 7. Cyclic voltammogram of **9a** (DCM solvent, 3.5 mM **9a**, 0.1 M NBu₄BPh₄) at scan rates 1–200 mV/s (red-blue). Inset displays i_{pa}/i_{pc} as a function of scan rate.

Figure 8.
Proposed mechanism for pyrazole formation from diazatitanacyclohexadienes.

Figure 9. Analogy between electrocyclic N–N coupling and the electrocyclic Nazarov cyclization of dienones.

Table 1.

Oxidation of **7a** with Various Oxidants^a

entry	oxidant	time (h)	<i>T</i> (°C)	yield (%)
1	TEMPO (2 equiv)	2	50	87(80 ^b)
2	TEMPO (1 equiv)	2	50	40
3 ^c	PhICl ₂	3	rt	70
4 ^c	PhI(O ₂ CCF ₃) ₂	3	rt	76
5 ^c	PhI(OAc) ₂	3	rt	44
6	FcPF ₆ (2 equiv)	16	rt	32
7	FcPF ₆ (2 equiv)	70	rt	50
8	O ₂ (1 atm)	3	60	17

^a Conditions: **7a** (0.02 M), oxidant (1–2 equiv), 0.5 mL of benzene. Yields are determined *in situ* via ¹H NMR against internal TMB standard.^b Isolated yield.^c Yields are determined via ¹H NMR against internal TCE standard added after the reaction.

Table 2.

Substrate Scope of One-Pot *In Situ* Pyrazole Synthesis from Alkynes and Nitriles^a

entry	R ¹	R ²	R ³	product	yield (%)
1 ^b	4-Me-C ₆ H ₄	Et	Et	1	75 ^c
2 ^b	Ph	Et	Et	1b	79 ^c
3	Ph	Et	Et	1b	56
4	4-MeO-C ₆ H ₄	Et	Et	1c	52
5	4-CF ₃ -C ₆ H ₄	Et	Et	1d	49
6	<i>i</i> Pr	Et	Et	1e	54
7	Me	Et	Et	1f	43
8	Me	Ph	Me	1g	28
9	Ph	Ph	Me	1h	54
10	Ph	Me	Me	1i	52
11	Ph	4- <i>t</i> -BuC ₆ H ₄	4- <i>t</i> -BuC ₆ H ₄	1j	37

^a Conditions: 0.025 mmol of [py₂TiCl₂(NPh)]₂, 0.05 mmol (2 equiv) of alkyne, 0.05 mmol (2 equiv) of nitrile, 0.5 mL of PhBr, 115 °C, 4h; yields with respect to alkyne are determined *in situ* via ¹H NMR against internal 1,3,5-trimethoxybenzene standard.

^b Conditions: 0.025 mmol of [py₂TiCl₂(NPh)]₂, 0.15 mmol (6 equiv) of alkyne, 0.15 mmol (6 equiv) of nitrile, 0.5 mL of PhBr, 115 °C, 1 h.

^c Yields reported with respect to [py₂TiCl₂(NPh)]₂, which is the limiting reagent under condition B.