Published in final edited form as:

Inorg Chem. 2019 April 15; 58(8): 5042–5050. doi:10.1021/acs.inorgchem.9b00102.

Spectroscopic and $^1\text{O}_2$ Sensitization Characteristics of a Series of Isomeric $\text{Re}(\text{bpy})(\text{CO})_3\text{Cl}$ Complexes Bearing Pendant BODIPY Chromophores

Andrea M. Potocny, Justin J. Teesdale, Alize Marangoz, Glenn P. A. Yap, Joel Rosenthal*

Department of Chemistry and Biochemistry, University of Delaware, Newark, Delaware 19716

Abstract

Two new $\text{Re}(\text{I})$ bipyridyltricarboxylchloride complexes, $\text{Re}(\text{BB3})(\text{CO})_3\text{Cl}$, and $\text{Re}(\text{BB4})(\text{CO})_3\text{Cl}$, featuring BODIPY groups appended to the 5,5'-, or 6,6'- positions of the bipyridine ligand respectively, have been synthesized as structurally isomeric compliments to a previously reported 4,4'-substituted homologue, $\text{Re}(\text{BB2})(\text{CO})_3\text{Cl}$. X-ray crystal structures of the compounds show that the 4,4'-, 5,5'- and 6,6'- substitution patterns place the BODIPY groups at progressively shorter distances of 9.43 Å, 8.39 Å and 5.56 Å, respectively, from the complexes' Re centers. The photophysical properties of the isomeric complexes were investigated to ascertain the manner in which the heavy rhenium atom might induce intersystem crossing of the pendant BODIPY moieties positioned at progressively shorter through-space distances. Electronic absorption spectroscopy revealed that the three metal complexes retain the strong visible absorption features characteristic of the bpy-BODIPY (**BB2–BB4**) ligands, however, the fluorescence of the parent borondipyrromethane appended ligands is attenuated by more than an order of magnitude in $\text{Re}(\text{BB2})(\text{CO})_3\text{Cl}$ and $\text{Re}(\text{BB3})(\text{CO})_3\text{Cl}$, and by more than two orders of magnitude in $\text{Re}(\text{BB4})(\text{CO})_3\text{Cl}$. Furthermore, phosphorescence from $\text{Re}(\text{BB4})(\text{CO})_3\text{Cl}$ is observed under a nitrogen atmosphere, consistent with highly efficient ISC to the triplet-excited state. Singlet oxygen sensitization studies confirm that all three complexes produce singlet oxygen with quantum yields that increase as the distance of the BODIPY groups to the heavy rhenium center is decreased. The trends observed across the series of rhenium complexes with respect to emission and $^1\text{O}_2$ sensitization properties can be rationalized in terms of the varied distal separation between the metal center and BODIPY groups in each system.

Graphical Abstract

*Corresponding Author : joelr@udel.edu. Tel: 302-831-0716.

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

ASSOCIATED CONTENT

Supporting Information. Crystallographic and spectroscopic data. This material is available free of charge via the Internet at <http://pubs.acs.org>. Crystallographic data are also available from the Cambridge Crystallographic Data Centre (CCDC 1879091 & 1879092).

The authors declare no competing financial interest.

A set of isomeric Re(I)bipyridyltricarboxylchloride complexes bearing BODIPY groups at the 4,4'-, 5,5'-, or 6,6'- positions of the bpy ligand are presented. Crystallography revealed that the 4,4'-, 5,5'-, and 6,6'-, substitution patterns place the BODIPYS at progressively shorter distances of 9.43 Å, 8.39 Å and 5.56 Å from the complexes' Re centers. The complexes' photophysics vary as a function of the distal separation between the Re centers and BODIPY moieties.

Keywords

singlet oxygen; boron dipyrromethene; BODIPY; photophysical properties; rhenium bipyridyltricarboxyl chloride

INTRODUCTION

Boron dipyrromethenes (BODIPYs) are well-known for their strong absorption of visible light, fluorescence quantum yields that can approach unity, and good photostability.^{1,2} These characteristics have inspired studies of BODIPY-based compounds for applications as light harvesters in dye-sensitized solar cells,³ as laser dyes⁴ and as fluorescent probes for biomolecules, pH, certain cations, anions or gases, reactive oxygen or nitrogen species, redox potential, and viscosity among other stimuli.^{5–7} While population of the triplet excited-state manifold of traditional BODIPY compounds is typically negligible, other photosensitizers that support a more extensive triplet photochemistry have found uses in triplet–triplet annihilation upconversion,^{8,9} photocatalysis,¹⁰ and photodynamic therapy.¹¹ The ease with which the BODIPY framework can be synthetically modified^{12–14} has buoyed efforts aimed at enabling the use of this popular class of luminophore for applications that require the photochemical generation of reactive triplet intermediates.¹⁵

A common strategy for transforming conventional fluorophores into triplet photosensitizers exploits the heavy atom effect to promote intersystem crossing (ISC) through enhanced spin orbit coupling.¹¹ Accordingly, numerous halogenated BODIPY derivatives featuring bromine or iodine substituents directly attached to, or in π -conjugation with the pyrrolic backbone have been synthesized and reported to efficiently access the triplet state.^{16–21} Ligands bearing BODIPY moieties have also been complexed with heavy transition metals such as Ru(II),^{22–24} Pt(II),^{25–29} Ir(III)^{30–32} or Re(I)^{33–35} to deliver systems that exhibit lower fluorescence quantum yields and, in some cases, detectable phosphorescence from the BODIPY triplet state following photoexcitation. Along these lines, we have reported a 2,2'-bipyridine ligand functionalized with two BODIPY substituents at the 4- and 4'-positions

(BB2),³⁶ and subsequently incorporated this ligand into a rhenium(I)tricarbonylchloride complex to form **Re(BB2)(CO)₃Cl** (Scheme 1).³⁷ In related work, we synthesized the analogous 5,5'-substituted 2,2'-bpyBODIPY ligand (**BB3**), which had been introduced previously in the literature using a slightly different synthetic strategy,³⁸ as well as the homologous 6,6'-substituted 2,2'-bpyBODIPY ligand (**BB4**).³⁹ In this study, **BB3** and **BB4** have also been incorporated into rhenium(I)tricarbonylchloride complexes (**Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl**, respectively) to complete the set of three-homologous complexes shown in Scheme 1.

For each of the isomeric Re complexes of Scheme 1, the positions at which the BODIPY units are attached to the bipyridine scaffolds ultimately dictates the chromophores' proximities (~5–10 Å) to the rhenium center. Accordingly, we anticipated that the photophysics of the pendant BODIPY units might be perturbed by varying degrees for each Re complex, and conducted steady state absorption and emission studies to probe how altering the (Re•••BODIPY) separation influences the photophysical properties of the three isomers. The absorption profiles of the complexes are dominated by the strong BODIPY features in the visible region while the emission spectra reveal evidence of greater spin-orbit coupling when the BODIPY groups are positioned closer to the metal center. Additionally, **Re(BB2)(CO)₃Cl**, **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** all show the ability to sensitize ¹O₂, but **Re(BB4)(CO)₃Cl**, which has the shortest average Re•••BODIPY separation, photosensitizes the formation of ¹O₂ with the highest quantum yield across the series. These constructs are notable as they form a structurally characterized set of isomeric homologues that add to the small number of BODIPY compounds tethered to a rhenium heavy atom to induce a triplet photochemistry,^{33–35} and as new examples of rhenium(I) complexes capable of generating ¹O₂ upon irradiation with wavelengths of light longer than 500 nm.^{11,33,40,41}

EXPERIMENTAL SECTION

General Materials and Methods.

Reactions requiring an inert atmosphere were carried out under positive pressure of N₂ using flasks fitted with Suba-Seal rubber septa and standard schlenk techniques. Air and moisture sensitive reagents were transferred using standard syringe or cannula techniques. Reagents and solvents were purchased from Sigma Aldrich, Acros, Fisher, Strem, or Cambridge Isotopes Laboratories. **Re(BB2)(CO)₃Cl** and the **BB3** and **BB4** ligands were prepared according to published procedures.^{36,37,39} Solvents used for synthesis were of reagent grade or better. Anhydrous solvents were dried by passage through activated alumina and stored over 4 Å molecular sieves prior to use. Column chromatography was performed with 40–63 μm silica gel from Silicycle.

Compound Characterization.

¹H NMR and ¹³C NMR spectra were recorded at 25 °C on a Bruker 400 MHz spectrometer with a cryogenic QNP probe. Proton spectra are referenced to the residual proton resonance of the deuterated solvent (CDCl₃ = δ 7.26) and carbon spectra are referenced to the carbon resonances of the solvent (CDCl₃ = δ 77.16).⁴² All chemical shifts are reported using the standard δ notation in parts-per-million, and positive chemical shifts are to higher frequency

from the given reference. High-resolution mass spectrometry analyses were performed by the University of Delaware Mass Spectrometry Laboratory in the Department of Chemistry and Biochemistry.

Rhenium[5,5'-bis(BODIPY)-2,2'-bipyridine](CO)₃Cl (Re(BB3)(CO)₃Cl).

5,5'-bis(BODIPY)-2,2'-bipyridine (**BB3**) (84 mg, 0.11 mmol) and rhenium(I) pentacarbonylchloride (40 mg, 0.11 mmol) were combined in 50 mL of 1,2-dichloroethane. The resulting solution was heated to reflux while being stirred for 20 h under nitrogen and underwent a change in color from orange to dark red. The solvent was removed under reduced pressure, and the crude product was purified on silica by eluting with 5% hexanes in CH₂Cl₂ followed by straight CH₂Cl₂. The product was obtained as the second band immediately following a band of unreacted ligand. The desired product was obtained as 41 mg of a dark red powder (Yield = 35%). ¹H NMR (400 MHz, CDCl₃, 25 °C) δ /ppm: 9.04 (d, *J* = 1.7 Hz, 2H), 8.42 (d, *J* = 8.2 Hz, 2H), 8.11 (dd, *J* = 8.2, 1.9 Hz, 2H), 2.57 (s, 12H), 2.39 – 2.28 (m, 8H), 1.47 (s, 6H), 1.46 (s, 6H), 1.02 (dt, *J* = 10.8, 7.6 Hz, 12H). ¹³C{¹H} NMR (101 MHz, CDCl₃, 25 °C) δ /ppm: 196.02, 156.88, 155.35, 154.99, 152.40, 139.30, 138.18, 136.22, 134.44, 133.80, 130.64, 130.50, 129.89, 123.13, 29.50, 16.92, 16.89, 14.41, 14.29, 13.45, 13.01, 12.63, 12.51. HR-ESI-MS: [M+H]⁺ *m/z*: calcd for C₄₇H₅₁B₂ClN₆F₄O₃Re, 1067.3391; found, 1067.3428.

Rhenium[6,6'-bis(BODIPY)-2,2'-bipyridine](CO)₃Cl (Re(BB4)(CO)₃Cl).

6,6'-bis(BODIPY)-2,2'-bipyridine (**BB4**) (84 mg, 0.11 mmol) and rhenium(I) pentacarbonylchloride (40 mg, 0.11 mmol) were combined in 50 mL of 1,2-dichloroethane. The resulting solution was heated to reflux while being stirred for 20 h under nitrogen and underwent a change in color from orange to dark red. The solvent was removed under reduced pressure, and the crude product was purified on silica by eluting 25% hexanes in CH₂Cl₂ followed by 1% methanol in CH₂Cl₂. The desired product was obtained as 75 mg of a dark red powder (Yield = 64%). ¹H NMR (400 MHz, CDCl₃, 25 °C) δ /ppm: 8.46 (d, *J* = 8.2 Hz, 2H), 8.20 (t, *J* = 7.9 Hz, 2H), 7.72 (d, *J* = 7.7 Hz, 2H), 2.52 (s, 12H), 2.28 (q, *J* = 7.5 Hz, 4H), 2.20 (q, *J* = 7.6 Hz, 4H), 1.65 (s, 6H), 1.07 (s, 6H), 0.95 (t, *J* = 7.5 Hz, 6H), 0.87 (t, *J* = 7.5 Hz, 6H). ¹³C{¹H} NMR (101 MHz, CDCl₃, 25 °C) δ /ppm: 192.04, 187.33, 158.68, 158.09, 158.04, 155.29, 139.29, 138.93, 135.70, 134.50, 133.86, 133.13, 131.18, 129.94, 128.16, 124.26, 17.26, 17.12, 14.65, 14.25, 13.53, 13.09, 12.83, 11.36. HR-ESI-MS: [M+H]⁺ *m/z*: calcd for C₄₇H₅₁B₂ClN₆F₄O₃Re, 1067.3391; found, 1067.3435.

X-ray Structural Analysis.

Crystals of **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** were obtained via slow evaporation of saturated solutions of the respective complexes in 1,2-dichloroethane. Crystals were mounted on a glass fiber using viscous oil and cooled to the data collection temperature. Data were collected on a Bruker-AXS APEX CCD diffractometer with graphite-monochromated Mo-Kα radiation (λ = 0.71073 Å). Unit cell parameters were obtained from 60 data frames, 0.3° ω, from three different sections of the Ewald sphere. The systematic absences in the diffraction data are uniquely consistent with Pmn2₁/n for **Re(BB4)(CO)₃Cl** with the Flack parameter refined to virtually nil, indicating the true hand

of the data had been determined. For **Re(BB3)(CO₃)Cl**, no symmetry higher than triclinic was observed and solution in the centrosymmetric space group option yielded chemically reasonable and computationally stable results of refinement. The data-sets were treated with SADABS absorption corrections based on redundant multi-scan data (Sheldrick, G., Bruker-AXS Inc, Madison, WI, 2001). For **Re(BB4)(CO₃)Cl**, the compound molecule was located on a mirror plane. Molecules of 1,2-dichloroethane solvent were located cocrystallized in a 1:1 ratio with the **Re(BB4)(CO₃)Cl** complex with the solvent disordered at a mirror plane. For **Re(BB3)(CO₃)Cl**, the pyrrole(C)–ethyl(C) bonds were restrained to be similar with one ethyl group found disordered in two positions having a refined site occupancy of 60/40 with the disordered chemically equivalent atoms constrained with equal atomic displacement parameters and treated with three-dimensional rigid-bond displacement restraints. The structures were solved using direct methods and refined with full-matrix, least-squares procedures on F^2 .⁴³ All non-hydrogen atoms were refined with anisotropic displacement parameters. All other hydrogen atoms were treated as idealized contributions. Structure factors are contained in the SHELXTL program library.⁴⁴ CIFs are deposited under CCDC depositary numbers 1879091 and 1879092.

Steady-State Spectroscopy Experiments.

UV-visible absorbance spectra were collected at room temperature on a StellarNet Silver-NoVA CCD array UV-vis spectrometer using quartz cuvettes (6Q) with a 1.0 cm pathlength from Firefly Scientific. Absorption spectra of **BB2**, **BB4**, **Re(BB2)(CO₃)Cl**, **Re(BB3)(CO₃)Cl**, and **Re(BB4)(CO₃)Cl** were collected in N,N-dimethylformamide at concentrations of 1.0, 2.0, 3.0, 4.0 and 5.0 μM . Absorption spectra of **BB3** were collected in 10% CH_2Cl_2 in N,N-dimethylformamide at concentrations of 1.0, 2.0, 3.0, 4.0 and 5.0 μM .

Emission and excitation spectra were recorded on an automated Photon Technology International (PTI) QuantaMaster 40 fluorometer equipped with a 75-W Xenon arc lamp, an LPS-220B lamp power supply and a Hamamatsu R2658 photomultiplier tube. All samples were prepared in screw cap quartz cuvettes of 1.0 cm path length from Starna Cells, Inc. Solutions of **BB2** (7.0 μM), **BB4** (6.0 μM), **Re(BB2)(CO₃)Cl** (8.0 μM), **Re(BB3)(CO₃)Cl** (9.0 μM) and **Re(BB4)(CO₃)Cl** (10.0 μM) in N,N-dimethylformamide, and a solution of **BB3** (6.0 μM) in 10% CH_2Cl_2 in N,N-dimethylformamide were prepared in a nitrogen-filled glovebox such that the absorbance values of all samples at $\lambda = 500 \text{ nm}$ ranged between 2.60×10^{-1} and 3.15×10^{-1} (1 cm pathlength). Samples for emission experiments were excited at $\lambda_{\text{ex}} = 500 \text{ nm}$, and emission was monitored from $\lambda_{\text{em}} = 515\text{--}950 \text{ nm}$ using a step size of 1 nm and an integration time of 0.25 seconds. Emission spectra were also measured following exposure of each of the samples to air. All reported spectra are the average of five individual acquisitions. Excitation spectra recorded for **Re(BB4)(CO₃)Cl** were monitored at either 575 nm or 828 nm, while varying the excitation wavelength from 420–650 nm.

A solution of $[\text{Ru}(\text{bpy})_3](\text{PF}_6)_2$ in air-saturated acetonitrile was prepared at a concentration of $\sim 120 \mu\text{M}$ such that its absorbance at $\lambda = 500 \text{ nm}$ fell between 2.60×10^{-1} and 3.15×10^{-1} for use as a luminescence reference ($\Phi_{\text{ref}} = 0.018$).⁴⁵ Emission quantum yields were calculated using the expression below:

$$\Phi_s = \Phi_{ref} \left(\frac{I_s}{I_{ref}} \right) \left(\frac{A_{ref}}{A_s} \right) \left(\frac{\eta_s}{\eta_{ref}} \right)^2$$

where Φ_s and Φ_{ref} are the emission quantum yield of the sample and the reference, respectively, I_s and I_{ref} are the integrated emission intensities of the sample and reference, respectively, A_s and A_{ref} are the measured absorbances of the sample and reference at the excitation wavelength, and η_s and η_{ref} are the refractive indices of the solvents used for the sample and reference, respectively.

Determination of Singlet Oxygen Sensitization Quantum Yields.

$^1\text{O}_2$ production was quantified by monitoring the fluorescence quenching of the $^1\text{O}_2$ trapping agent, 1,3-diphenylisobenzofuran (DPBF).^{46,47} Measurements were carried out on an automated Photon Technology International (PTI) QuantaMaster 40 fluorometer equipped with a 75-W Xenon arc lamp, an LPS-220B lamp power supply and a Hamamatsu R2658 photomultiplier tube using quartz cuvettes (6Q) of 1.0 cm path length. Each cuvette contained 2.0 mL of N,N-dimethylformamide solution that was 10.0 μM in **Re(BB2)(CO₃)Cl**, **Re(BB3)(CO₃)Cl**, **Re(BB4)(CO₃)Cl**, or **[Ru(bpy)₃](PF₆)₂** (used as a reference, $\Phi = 0.57$)⁴⁸ and 1.0 μM in DPBF. An additional cuvette containing only N,N-dimethylformamide and 1.0 μM DPBF was used as a control. Consumption of DPBF was monitored by observing the change in its integrated emission intensity following irradiation with light from an Intralux 9000 light source (Volpi) fitted with a 10 nm (fwhm) bandpass filter centered at 500 nm (Thor Labs, FB500–10). During the studies of **Re(BB2)(CO₃)Cl** and **Re(BB3)(CO₃)Cl**, the cuvettes were irradiated for 10 second intervals for a total of 40 seconds. Due to its greater $^1\text{O}_2$ generating efficiency, **Re(BB4)(CO₃)Cl** was studied separately and irradiated for 5 second intervals for a total of 20 seconds. DPBF emission spectra were obtained by exciting at $\lambda_{ex} = 405$ nm and scanning from $\lambda_{em} = 400 - 600$ nm using a step size of 1 nm and an integration time of 0.25 seconds.

As has been reported previously,^{49–52} calibration curves of the integrated emission intensity versus the concentration of unreacted DPBF remaining in solution were generated to correct for absorption of the photosensitizers and standard between 400 – 600 nm. Emission spectra were collected from 10.0 μM solutions of **Re(BB2)(CO₃)Cl**, **Re(BB3)(CO₃)Cl**, **Re(BB4)(CO₃)Cl**, and **[Ru(bpy)₃](PF₆)₂** containing DPBF concentrations of 0, 0.25, 0.50, 0.75, 1.00, 1.25, or 1.50 μM . Linear regression lines fit to the calibration data from each solution enabled the integrated emission intensity values obtained from the $^1\text{O}_2$ sensitization experiments to be converted into the corresponding concentrations of unreacted DPBF. A final plot of the concentration of unreacted DPBF versus irradiation time formed a straight line of slope m , which allowed for calculation of the $^1\text{O}_2$ quantum yields via the following expression:

$$\Phi_s = \Phi_{ref} \left(\frac{m_s}{m_{ref}} \right) \left(\frac{\epsilon_{ref}}{\epsilon_s} \right)$$

Where Φ_s and Φ_{ref} are the $^1O_2$ sensitization quantum yields for the sample and [Ru(bpy)₃](PF₆)₂ reference, respectively, m_s and m_{ref} are the slopes of the concentration of unreacted DPBF vs. irradiation time plots for the sample and reference, and ϵ_s and ϵ_{ref} are the extinction coefficients at the wavelength of irradiation (500 nm) for the sample and reference, respectively. All reported $^1O_2$ quantum yields were obtained from an average of three trials.

RESULTS

BB2, **BB3** and **BB4** were synthesized according to published procedures,³⁹ and metalation of these derivatives was achieved by heating solutions of the ligands with Re(CO)₅Cl in 1,2-dichloroethane at reflux for 20 hours (Scheme 1). As reported previously for **Re(BB2)(CO)₃Cl**,³⁷ both **Re(BB3)(CO)₃Cl**, and **Re(BB4)(CO)₃Cl** can be obtained in four steps from commercially available materials, as illustrated by Scheme S1 in the Supporting Information.

Slow evaporation of concentrated solutions of **Re(BB3)(CO)₃Cl**, and **Re(BB4)(CO)₃Cl** dissolved in 1,2-dichloroethane yielded single crystals of both complexes. Figure 1 displays the solid-state structures obtained for **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** along with a reproduction of the structure obtained previously for **Re(BB2)(CO)₃Cl**.³⁷ Crystallographic parameters for **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** are summarized in Table 1, and fully labeled thermal ellipsoid plots for both complexes are available in the Supporting Information (Figures S1 and S2). The crystal structures of all three isomeric Re complexes reveal that the BODIPY units are nearly orthogonal to the bipyridine backbone. Table 2 summarizes the dihedral angles between each BODIPY moiety and the pyridine ring to which it is directly appended for each Re complex. For all three complexes, the bpy—BODIPY dihedral angles range from ~74°–88°. Additionally, the ¹H NMR spectra of **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** show a desymmetrization of the aliphatic resonances attributable to the BODIPY alkyl substituents that is not seen in the ¹H NMR spectra of the free **BB3** or **BB4** ligands (Supporting Information Figures S3 and S4). Similar observations from the ¹H NMR spectra of **BB2** and **Re(BB2)(CO)₃Cl** were attributed to a lack of free rotation about the indacene-bipyridine bonds, which places opposite sides of each BODIPY group in distinct environments as a result of being in closer proximity to either the axial chloride or the axial carbonyl ligand on each complex's Re center (Figure 1).³⁷ Given that dimethyl substitution at the 1- and 7-positions of the BODIPY framework has been reported to hinder rotation of *meso* substituents,^{53–55} the rigid, nearly perpendicular orientation of the BODIPY groups with respect to the 2,2'-bipyridine backbone observed in all three rhenium complexes is not surprising.

The solid-state structures of **Re(BB2)(CO)₃Cl**, **Re(BB3)(CO)₃Cl**, and **Re(BB4)(CO)₃Cl** also allow for the comparison of the proximity of the BODIPY luminophores to each of the complexes rhenium centers to be assessed across the series of isomeric compounds. The average through-space distance from the boron atoms of the two appended BODIPY groups to the central rhenium atom was determined to be 9.43 Å for **Re(BB2)(CO)₃Cl**, 8.39 Å for **Re(BB3)(CO)₃Cl** and 5.56 Å for **Re(BB4)(CO)₃Cl**, as summarized in Table 2. Accordingly, the BODIPY substituents are positioned increasingly closer to the heavy metal as their

points of attachment are changed from the 4,4'- to the 5,5'- to the 6,6'-positions on the 2,2'-bipyridine ligand.

The steady state photophysical properties of **Re(BB2)(CO)₃Cl**, **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** were studied and compared to those of the corresponding bpyBODIPY ligands in DMF. As shown in Figure 2 and summarized in Table 3, the absorption spectrum of each of the three rhenium complexes is dominated by a strong feature in the visible region with a maximum extinction coefficient of 79,400 M⁻¹cm⁻¹ at 530 nm for **Re(BB2)(CO)₃Cl**, 52,700 M⁻¹cm⁻¹ at 533 nm for **Re(BB3)(CO)₃Cl** and 88,500 M⁻¹cm⁻¹ at 542 nm for **Re(BB4)(CO)₃Cl**. These absorption bands are attributed to the BODIPY-based S₀→S₁ (π→π*) transition,³⁸ and similar features with maximum extinction coefficients of 86,000 M⁻¹ cm⁻¹ at 526 nm, 95,100 M⁻¹cm⁻¹ at 528 nm, and 100,800 M⁻¹cm⁻¹ at 526 nm are evident in the absorption profiles of the **BB2**, **BB3** and **BB4** ligands, respectively (Figure 2). While metalation weakly attenuates the characteristic BODIPY absorption features and shifts them towards slightly longer wavelengths, overall the electronic spectra of the metal complexes are very similar to those of the corresponding bpyBODIPY ligands. This observation is notable, as Re(bpy)(CO)₃Cl complexes lacking additional pendant chromophores usually absorb poorly at wavelengths longer than ~450 nm.^{56,57} We note that the BODIPY based adsorption observed for **Re(BB3)(CO)₃Cl** is reduced as compared to the other two isomeric complexes considered. This difference in absorptivity is presumably due to the differences in coupling between the BODIPY units when attached to the 5,5' positions of the bipyridine ligand.

The room temperature emission spectra of **BB2**, **BB3**, **BB4**, **Re(BB2)(CO)₃Cl**, **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** were collected from DMF solutions under a nitrogen atmosphere with λ_{exc} = 500 nm, and are overlaid with the corresponding absorption spectra in Figure 2. In contrast to the minor differences observed in the absorption spectra between each free bpyBODIPY ligand (*i.e.*, **BB2**–**BB4**) and the corresponding Re complexes, incorporation of the BODIPY-based bpy ligands onto the rhenium tricarbonylchloride center results in dramatic changes to the indacenes' emission properties. For example, **BB2** shows a single, intense emission feature centered at λ_{fl} = 550 nm, and similar features appear in the emission spectra of **BB3** and **BB4** centered at λ_{fl} = 548 nm. Table 3 lists the emission quantum yields determined for the bpyBODIPY ligands; **BB2** and **BB4** display Φ_{fl} values close to unity, and the quantum yield for **BB3** (Φ_{fl} = 3.5 × 10⁻¹) is also relatively large. Accordingly, the three ligands exhibit the small Stokes shifts and high fluorescence quantum yields that are hallmarks of traditional BODIPY photophysics.^{2,12} By contrast, each of the isomeric rhenium complexes emit at slightly longer wavelengths (λ_{fl} = 561 nm for **Re(BB2)(CO)₃Cl**; λ_{fl} = 553 nm for **Re(BB3)(CO)₃Cl**; and λ_{fl} = 564 nm for **Re(BB4)(CO)₃Cl**), and with considerably diminished intensity. The fluorescence quantum yields of **Re(BB2)(CO)₃Cl** and **Re(BB3)(CO)₃Cl** are Φ_{fl} = 1.4 × 10⁻² and Φ_{fl} = 1.3 × 10⁻², respectively, representing ~10–100 fold reductions in magnitude relative to the corresponding free bpyBODIPY ligands (*i.e.*, **BB2** and **BB3**). The fluorescence from **Re(BB4)(CO)₃Cl**, in which the BODIPY groups are held closest to the rhenium center, is attenuated even further with Φ_{fl} = 1.3 × 10⁻³. The decrease in fluorescence quantum yield observed when **BB2** and **BB3** are complexed to rhenium, which places the BODIPY fluorophores ~9.5 Å and 8.4 Å,

respectively from the metal center, coupled with the even lower Φ_{fl} obtained when the BODIPY units are brought within ~ 5.5 Å of the rhenium atom in **Re(BB4)(CO)₃Cl**, suggest that enhanced ISC to the triplet excited state is being promoted by the heavy rhenium atom. The rhenium mediated ISC to the triplet provides a competing process to the facile radiative relaxation of the BODIPY singlet excited states in each of the isomeric Re complexes.

Additional evidence of triplet state population in the isomeric rhenium complexes of Scheme 1 is garnered by observation of a second red-shifted feature in the emission spectrum recorded for **Re(BB4)(CO)₃Cl**. This long-wavelength band is centered at $\lambda_{\text{em}} = 828$ nm and has a quantum yield of $\Phi_{\text{ph}} = 1.6 \times 10^{-4}$ (see Figure 2c, inset and Table 3). This long-wavelength signal was conspicuously absent when the emission spectrum of **Re(BB4)(CO)₃Cl** was measured under air, while the shorter-wavelength features observed for **Re(BB2)(CO)₃Cl**, **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** at around 555–565 nm remain unperturbed (see Figure S5). Molecular oxygen in its triplet ground state ($^3\text{O}_2$) can engage in energy transfer with chromophores and metal complexes in triplet excited states to return the excited light-absorber to its ground state, and is converted to singlet oxygen ($^1\text{O}_2$) in the process. Consequently, emission quenching as a result of exposure to air is often diagnostic of phosphorescence. Rhenium bipyridyltricarbonylchloride complexes are often phosphorescent, however, emission from the $^3\text{MLCT}$ state is normally observed between 600–650 nm.^{57,58} The spectral position of the emission feature of **Re(BB4)(CO)₃Cl** (~ 830 nm) is more consistent with emission from the BODIPY $^3\pi-\pi^*$ manifold, which has been previously observed at wavelengths longer than 700 nm.¹⁵ Excitation spectra recorded for **Re(BB4)(CO)₃Cl** (Figure S6) while monitoring either the fluorescence (575 nm) or phosphorescence (828 nm) emission are consistent with the BODIPY absorbance profile (Figure 2), demonstrating that both the singlet and triplet emission of **Re(BB4)(CO)₃Cl** arises from the BODIPY moieties.

To confirm that the air-sensitivity of the long-wavelength emission feature of **Re(BB4)(CO)₃Cl** is a result of energy transfer to molecular oxygen, and to investigate whether **Re(BB2)(CO)₃Cl** and **Re(BB3)(CO)₃Cl** also support triplet photochemistries and photosensitize the formation of $^1\text{O}_2$, the three isomeric complexes of Scheme 1 were probed using the singlet oxygen trapping agent, 1,3-diphenylisobenzofuran (DPBF)^{46,47} in DMF and irradiated with $\lambda_{\text{ex}} = 500$ nm. **Re(BB2)(CO)₃Cl**, **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** all demonstrate an ability to produce $^1\text{O}_2$ indicating that even the 4,4'- and 5,5'-substituted isomers undergo at least some ISC to the triplet excited state upon irradiation with 500 nm light. Furthermore, because most traditional *fac*-Re(bpy)(CO)₃Cl derivatives fail to absorb light longer than ~ 450 nm, **Re(BB2)(CO)₃Cl**, **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** join a small group of such complexes capable of sensitizing the formation of $^1\text{O}_2$ upon irradiation with light of $\lambda \geq 500$ nm.^{11,33,40,41}

[Ru(bpy)₃]²⁺ was used as an actinometer ($\Phi = 0.57$ in DMF)⁴⁸ to determine the $^1\text{O}_2$ quantum yields (Φ) for each of the isomeric Re complexes. The values of Φ for **Re(BB2)(CO)₃Cl** and **Re(BB3)(CO)₃Cl** ($\lambda_{\text{ex}} = 500$ nm) were found to be low and essentially equivalent ($\Phi = 1.7\%$ and 1.6% , respectively). **Re(BB4)(CO)₃Cl** displayed a significantly higher quantum yield for $^1\text{O}_2$ production of $\Phi = 18.6\%$. The enhanced efficiency with which **Re(BB4)(CO)₃Cl** acts as a $^1\text{O}_2$ photosensitizer compared to the other two complexes

of Scheme 1 is consistent with the spectroscopic observations/rationale presented above (vide supra); although the rhenium center promotes ISC due to enhanced spin-orbit coupling in all three Re complexes, the especially close proximities of the BODIPY moieties (~5.5 Å) to the heavy metal center as a result of their attachment at the 6,6'-positions of the bipyridyl backbone leads to much more effective population or longer lifetime of the triplet excited state for **Re(BB4)(CO)₃Cl**. As such, of the three isomeric Re complexes considered in this work, **Re(BB4)(CO)₃Cl** represents a platform that is most suitable for ¹O₂ sensitization and related triplet photochemical applications using visible excitation sources.

We emphasize that metal...BODIPY distance may not be the only factor that ultimately controls ¹O₂ sensitization efficiency. While the series of isomeric rhenium complexes we present demonstrates how distal separation between the BODIPY light absorber and metal center can be correlated with intersystem crossing efficiency and ¹O₂ quantum yield, many other factors ultimately control the efficacy of these two processes. In particular, the strength of spin-orbit coupling, which promotes ISC to the triplet excited state is largely dependent on identity of the metal center.³¹ Similarly, the manner by which the visible light absorber is attached to the metal center is also critical, as a rhenium bipyridine complex in which a BODIPY light absorber is tethered to a Re(bpy)(CO)₃Cl core via an alkyne linker shows Φ = 0.88.⁵⁹ Accordingly molecular topology is an important consideration in the construction of new complexes that support triplet photochemistry.

CONCLUSIONS AND FUTURE DIRECTIONS

Typical BODIPY chromophores strongly absorb visible light, and are characterized by high photostability, and large fluorescence quantum yields that can approach unity. Although these traits make them effective as fluorophores, standard BODIPYs are generally not suitable for applications requiring triplet photosensitizers due to their negligible rates of intersystem crossing. Conversely, *fac*-Re(bpy)(CO)₃Cl complexes readily populate the triplet excited state due to the presence of the heavy metal atom, but their utility as triplet photosensitizers is often limited by poor light absorption within the visible region (especially beyond 450 nm). In this study, we have detailed an isomeric set of *fac*-Re(bpy)(CO)₃Cl homologues with pendant BODIPY chromophores attached at either the 4,4'-, 5,5'-, or 6,6'-positions of the 2,2-bipyridine ligand. We anticipated these complexes would combine the favorable photophysical properties of the two individual classes of light absorbers (*i.e.*, BODPY and *fac*-Re(bpy)(CO)₃Cl) to deliver robust and efficient triplet photosensitizers that strongly absorb visible light.

Re(BB2)(CO)₃Cl, **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** form a series of three isomeric complexes in which the position of the BODIPY units is attached to the 2,2-bipyridine ligand at progressively shorter distances from the rhenium center. The **Re(BB2)(CO)₃Cl** complex (4,4'-BODIPY substituents), **Re(BB3)(CO)₃Cl** complex (5,5'-BODIPY substituents), and **Re(BB4)(CO)₃Cl** complex (6,6'-BODIPY substituents), position the light absorbing indacene moieties at ~9.4 Å, 8.4 Å and 5.5 Å from the heavy Re center, respectively. Electronic absorption spectra recorded for each of these rhenium complexes show that the characteristic BODIPY absorption features are preserved, and confirm that all three compounds are strong visible light absorbers. The high emission quantum yields

typical of traditional BODIPY derivatives are not observed for any of the three rhenium complexes, however, and the decrease in fluorescence intensity becomes more extreme as the BODIPY moiety is placed closer to the rhenium center. **Re(BB4)(CO)₃Cl**, which has the shortest average through-space Re•••BODIPY separation (~5.5 Å) of the three complexes studied also exhibits weak phosphorescence. The quenched BODIPY fluorescence observed across the series of rhenium complexes and the phosphorescence of **Re(BB4)(CO)₃Cl** are consistent with the heavy rhenium atom enhancing the rate of ISC to deliver a triplet excited state using visible light excitation.

The ability of **Re(BB2)(CO)₃Cl**, **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl** to undergo ISC following excitation with light of $\lambda = 450\text{--}550\text{ nm}$, makes these systems potentially suitable for triplet photochemistry applications. To this end, evaluation of these complexes' abilities to photosensitize the formation of ¹O₂ upon irradiation with light of $\lambda_{\text{ex}} = 500\text{ nm}$ demonstrates that each system can produce singlet oxygen using visible light. In keeping with the observed trends in singlet and triplet emission properties across the series of complexes, **Re(BB4)(CO)₃Cl** displays the highest quantum yield for ¹O₂ production ($\Phi = 18.6\%$), consistent with this isomer presenting the shortest distal separation between the heavy Re center and ancillary BODIPY light absorbers.

When viewed in whole, the results presented herein demonstrate that appending chromophores at the 6,6'-positions of the 2,2'-bipyridine ligand of *fac*-Re(bpy)(CO)₃Cl complexes is an effective strategy to construct triplet photosensitizers that effectively harvest visible light. Additionally, the appreciable ¹O₂ quantum yield realized with **Re(BB4)(CO)₃Cl** suggest that architectures that provide a similar topological arrangement of chromophore and heavy metal center may potentially be of use in photodynamic therapy or as photoCORMs^{60–66} that can be used to administer CO inside cells upon visible light irradiation.^{67,68} Although the absorbance profile of **Re(BB4)(CO)₃Cl** only extends out to about 585 nm, which falls just short of the therapeutic window (~600–900 nm) of red to near IR light that most deeply penetrates biological tissues, synthetic modification of the pendant BODIPY moieties or introduction of alternative chromophores that absorb longer wavelength light at the 6,6'-positions of the bipyridyl ligand of *fac*-Re(bpy)(CO)₃Cl may yield promising candidates for applications in photodynamic therapy.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENT

This work was supported by NSF CAREER Award CHE1352120 and NIH P20GM104316. NMR and other data were acquired at UD using instrumentation obtained with assistance from the NSF and NIH (NSF-MIR 0421224, NSF-MIR 1048367, NSF-CRIF MU CHE-0840401 and CHE-0541775, NIH P20 RR017716).

ABBREVIATIONS

¹ O ₂	singlet oxygen
PDT	photodynamic therapy

NMR	nuclear magnetic resonance
MS	mass spectrometry
LC/MS	liquid chromatography/mass spectrometry
DPBF	1,3-diphenylisobenzofuran

REFERENCES

1. Benniston AC; Copley G Lighting the Way Ahead with Boron Dipyrromethene (BODIPY) Dyes. *Phys. Chem. Chem. Phys.* 2009, 11, 4124–4131. [PubMed: 19458813]
2. Ulrich G; Ziessel R; Harriman A The Chemistry of Fluorescent Bodipy Dyes: Versatility Unsurpassed. *Angew. Chem. Int. Ed.* 2008, 47, 1184–1201.
3. Klifout H; Stewart A; Elkhalfa M; He H BODIPYs for Dye Sensitized Solar Cells. *ACS Appl. Mater. Interfaces.* 2017, 9, 39873–39889.
4. Gómez-Durán CFA; García-Moreno I; Costela A; Martín V; Sastre R; Bañuelos J; Arbeloa FL; Arbeloa IL; Peña-Cabrera E 8-PropargylaminoBODIPY: Unprecedented Blue-emitting Pyrromethene Dye. *Synthesis, Photophysics and Laser Properties.* *Chem. Commun.* 2010, 46, 5103–5105.
5. Rosenthal J; Lippard SJ Direct Detection of Nitroxyl in Aqueous Solution Using a Tripodal Copper(II) BODIPY Complex. *J. Am. Chem. Soc.* 2010, 132, 5536–5537. [PubMed: 20355724]
6. Kowada T; Maeda H; Kikuchi K BODIPY-Based Probes for the Fluorescence Imaging of Biomolecules in Living Cells. *Chem. Soc. Rev.* 2015, 44, 4953–4972. [PubMed: 25801415]
7. Boens N; Leen V; Dehaen W Fluorescent Indicators Based on BODIPY. *Chem. Soc. Rev.* 2012, 41, 1130–1172. [PubMed: 21796324]
8. Singh-Rachford TN; Castellano FN Photon Upconversion Based on Sensitized Triplet-Triplet Annihilation. *Coord. Chem. Rev.* 2010, 254, 2560–2573.
9. McCusker CE; Castellano FN Materials Integrating Photochemical Upconversion. *Top. Curr. Chem.* 2016, 374, 1–25.
10. Juergen B; Francis AJ; El Roz KA; Castellano FN; Weare WW; Sommer RD Light-Driven Hydrogen Evolution by BODIPY-Sensitized Cobaloxime Catalysts. *Inorg. Chem.* 2014, 53, 4527–4534. [PubMed: 24725061]
11. Zhao J; Wu W; Sun J; Guo S Triplet Photosensitizers: From Molecular Design to Applications. *Chem. Soc. Rev.* 2013, 42, 5323–5351. [PubMed: 23450221]
12. Loudett A; Burgess K BODIPY Dyes and Their Derivatives: Syntheses and Spectroscopic Properties. *Chem. Rev.* 2007, 107, 4891–4932. [PubMed: 17924696]
13. Boens N; Verbelen B; Dehaen W Postfunctionalization of the BODIPY Core: Synthesis and Spectroscopy. *Eur. J. Org. Chem.* 2015, 2015, 6577–6595.
14. Lakshmi V; Rao MR; Ravikanth M Halogenated Boron-dipyrromethenes: Synthesis, Properties and Applications. *Org. Biomol. Chem.* 2015, 13, 2501–2517. [PubMed: 25594728]
15. Zhao J; Xu K; Yang W; Wang Z; Zhong F The Triplet Excited State of BODIPY: Formation, Modulation and Application. *Chem. Soc. Rev.* 2015, 44, 8904–8939. [PubMed: 26465741]
16. Yogo T; Urano Y; Ishitsuka Y; Maniwa F; Nagano T Highly Efficient and Photostable Photosensitizer Based on BODIPY Chromophore. *J. Am. Chem. Soc.* 2005, 127, 12162–12163.
17. Wu W; Guo H; Wu W; Ji S; Zhao J Organic Triplet Sensitizer Library Derived from a Single Chromophore (BODIPY) with Long-Lived Triplet Excited State for Triplet-Triplet Annihilation-Based Upconversion. *J. Org. Chem.* 2011, 76, 7056–7064. [PubMed: 21786760]
18. Sabatini RP; McCormick TM; Lazarides T; Wilson KC; Eisenberg R; McCamant DW Intersystem Crossing in Halogenated BODIPY Chromophores Used for Solar Hydrogen Production. *J. Phys. Chem. Lett.* 2011, 2, 223–227.
19. Zhang X-F; Yang X; Niu K; Geng H Phosphorescence of BODIPY Dyes. *J. Photochem. Photobiol. A. Chem.* 2014, 285, 16–20.

20. Ortiz MJ; Agarrabeitia AR; Duran-Sampedro G; Bañuelos Prieto J; Lopez TA; Massad WA; Montejano HA; García NA; Lopez Arbeloa I Synthesis and Functionalization of New Polyhalogenated BODIPY Dyes. Studies of Their Photophysical Properties and Singlet Oxygen Generation. *Tetrahedron*, 2012, 68, 1153–1162.
21. Chen Y; Zhao J; Xie L; Guo H; Li Q Thienyl-Substituted BODIPYs with Strong Visible Light Absorption and Long-Lived Triplet Excited States as Organic Triplet Sensitizers for Triplet–Triplet Annihilation Upconversion. *RSC Adv.* 2012, 2, 3942–3953.
22. Galletta M; Campagna S; Quesada M; Ulrich G; Ziesel R The Elusive Phosphorescence of Pyrromethene-BF₂ Dyes Revealed in New Multicomponent Species Containing Ru(II)-terpyridine Subunits. *Chem. Commun.* 2005, 4222–4224.
23. Galletta M; Puntoriero F; Campagna S; Chiorboli C; Quesada M; Goeb S; Ziesel R Absorption Spectra, Photophysical Properties, and Redox Behavior of Ru(II) Polypyridine Complexes Containing Accessory Dipyrromethene-BF₂ Chromophores. *J. Phys. Chem. A* 2006, 110, 4348–4358. [PubMed: 16571037]
24. Wu W; Sun J; Cui X; Zhao J Observation of the Room Temperature Phosphorescence of BODIPY in Visible Light Harvesting Ru(II) Polyimine Complexes and Application as Triplet Photosensitizers for Triplet–Triplet–Annihilation Upconversion and Photocatalytic Oxidation. *J. Mater. Chem. C* 2013, 1, 4577–4589.
25. Nastasi F; Puntoriero F; Campagna S; Diring S; Ziesel R Photoinduced Intercomponent Processes in Multichromophoric Species Made of Pt(II)-terpyridine-acetylide and Dipyrromethene-BF₂ Subunits. *Phys. Chem. Chem. Phys.* 2008, 10, 3982–3986. [PubMed: 18597011]
26. Wu W; Zhao J; Guo H; Sun J; Ji S; Wang Z Long-Lived Room-Temperature Near-IR Phosphorescence of BODIPY in a Visible-Light-Harvesting N^CN Pt(II)-Acetylide Complex with a Directly Metalated BODIPY Chromophore. *Chem. Eur. J.* 2012, 18, 1961–1968. [PubMed: 22250033]
27. Wu W; Liu L; Cui X; Zhang C; Zhao J Red-Light-Absorbing Diimine Pt(II) Bisacetylide Complexes Showing Near-IR Phosphorescence and Long-Lived 3IL Excited State of BODIPY for Application in Triplet–Triplet Annihilation Upconversion. *Dalton Trans.* 2013, 42, 14374–14379.
28. Lazarides T; McCormick TM; Wilson KC; Lee S; McCamant DW; Eisenberg R Sensitizing the Sensitizer: The Synthesis and Photophysical Study of BODIPY-Pt(II)(diamine)(dithiolate) Conjugates. *J. Am. Chem. Soc.* 2011, 133, 350–364. [PubMed: 21175161]
29. Whited MT; Djurovich PI; Roberts ST; Durrell AC; Schlenker CW; Bradforth SE; Thompson ME Singlet and Triplet Excitation Management in a Bichromophoric Near-Infrared-Phosphorescent BODIPY-Benzoporphyrin Platinum Complex. *J. Am. Chem. Soc.* 2011, 133, 88–96. [PubMed: 21142032]
30. Rachford AA; Ziesel R; Bura T; Retailleau P; Castellano FN Boron Dipyrromethene (BODIPY) Phosphorescence Revealed in [Ir(ppy)₂(bpy-C≡C-BODIPY)]⁺ Inorg. Chem. 2010, 49, 3730–3736.
31. Sun J; Zhong F; Yi X; Zhao J Efficient Enhancement of the Visible Light Absorption of Cyclometalated Ir(III) Complexes Triplet Photosensitizers with BODIPY and Applications in Photooxidation and Triplet–Triplet Annihilation Upconversion. *Inorg. Chem.* 2013, 52, 6299–6310. [PubMed: 23327589]
32. Palao E; Sola-Llano R; Tabero A; Manzano H; Agarrabeitia AR; Villanueva A; López-Arbeloa I; Martínez-Martínez V; Ortiz MJ AcetylacetonateBODIPY-Biscyclometalated Ir(III) Complexes: Effective Strategy Towards Smarter Fluorescent Photosensitizer Agents. *Chem. Eur. J.* 2017, 23, 10139–10147.
33. Yi X; Zhao J; Sun J; Guo S; Zhang H Visible Light-Absorbing Rhenium(I) Tricarbonyl Complexes as Triplet Photosensitizers in Photooxidation and Triplet–Triplet Annihilation Upconversion. *Dalton Trans.* 2013, 42, 2062–2074. [PubMed: 23178459]
34. Madu S; Hinge VK; Ravikanth M Synthesis and Properties of Rhenium(I) Bridged Boron Dipyrromethene Diad. *J. Porphyr. Phthalocyanines.* 2014, 18, 535–541.
35. Zhang S; Yuan W; Qin Y; Zhang J; Lu N; Wang Y; Li H; Wang Y; Li Y Bidentate BODIPY-appended 2-pyridylimidazo[1,2-*a*]pyridine Ligand and Fabrication of Luminescent Transition Metal Complexes. *Polyhedron*, 2018, 148, 22–31.

36. Rosenthal J; Nepomnyashchii AB; Kozhukh J; Bard AJ; Lippard SJ Synthesis, Photophysics, Electrochemistry, and Electrogenenerated Chemiluminescence of a Homologous Set of BODIPY-Appended Bipyridine Derivatives. *J. Phys. Chem. C*. 2011, 115, 17993–18001.
37. Teesdale JJ; Pistner AJ; Yap GPA; Ma Y-Z; Lutterman DA; Rosenthal J Reduction of CO₂ Using a Rhenium Bipyridine Complex Containing Ancillary BODIPY Moieties. *Catal. Today*. 2014, 225, 149–157. [PubMed: 25395735]
38. Ulrich G; Ziesel R Convenient and Efficient Synthesis of Functionalized Oligopyridine Ligands Bearing Accessory Pyrromethene-BF₂ Fluorophores. *J. Org. Chem.* 2004, 69, 2070–2083. [PubMed: 15058955]
39. Qi H; Teesdale JJ; Pupillo RC; Rosenthal J; Bard AJ Synthesis, Electrochemistry, and Electrogenenerated Chemiluminescence of Two BODIPY- Appended Bipyridine Homologues. *J. Am. Chem. Soc.* 2013, 135, 13558–13566.
40. Zhong F; Yuan X; Zhao J; Wang Q Visible Light-Harvesting Tricarbonyl Re(I) Complex: Synthesis and Application in Intracellular Photodynamic Effect and Luminescence Imaging. *Sci. China Chem.* 2016, 59, 70–77.
41. McKenzie LK; Bryant HE; Weinstein JA Transition Metal Complexes as Photosensitizers in One- and Two-Photon Photodynamic Therapy. *Coord. Chem. Rev.* 2019, 379, 2–29.
42. Gottlieb HE; Kotlyar V; Nudelman A NMR Chemical Shifts of Common Laboratory Solvents as Trace Impurities. *J. Org. Chem.* 1997, 62, 7512–7515. [PubMed: 11671879]
43. Sheldrick GM A Short History of SHELX. *Acta. Cryst.* 2008, A64, 112–122.
44. Sheldrick GM Crystal Structure Refinement with SHELXL. *Acta. Cryst.* 2015, C71, 3–8.
45. Brouwer AM Standards for Photoluminescence Quantum Yield Measurements in Solution (IUPAC Technical Report). *Pure Appl. Chem.* 2011, 83, 2213–2228.
46. Liu Y; Hammitt R; Lutterman DA; Joyce LE; Thummel RP; Turro C Ru(II) Complexes of New Tridentate Ligands: Unexpected High Yield of Sensitized ¹O₂. *Inorg. Chem.* 2009, 48, 375–385. [PubMed: 19035764]
47. Young RH; Wehrly K; Martin RL Solvent Effects in Dye-Sensitized Photooxidation Reactions. *J. Am. Chem. Soc.* 1971, 93, 5774–5779.
48. Spiller W; Kliesch H; Wöhrle D; Hackbarth S; Röder B; Schnurpfeil G Singlet Oxygen Quantum Yields of Different Photosensitizers in Polar Solvents and Micellar Solutions. *J. Porphyrins Phthalocyanines*. 1998, 2, 145–158.
49. Pistner AJ; Pupillo RC; Yap GPA; Lutterman DA; Ma Y-Z; Rosenthal J Electrochemical, Spectroscopic, and ¹O₂ Sensitization Characteristics of 10,10-Dimethylbiladiene Complexes of Zinc and Copper. *J. Phys. Chem. A* 2014, 118, 10639–10648.
50. Potocny AM; Pistner AJ; Yap GPA; Rosenthal J Electrochemical, Spectroscopic, and ¹O₂ Sensitization Characteristics of Synthetically Accessible Linear Tetrapyrrole Complexes of Palladium and Platinum. *Inorg. Chem.* 2017, 56, 12703–12711.
51. Potocny AM; Riley RS; O'Sullivan RK; Day ES; Rosenthal J Photochemotherapeutic Properties of a Linear Tetrapyrrole Palladium(II) Complex Displaying an Exceptionally High Phototoxicity Index. *Inorg. Chem.* 2018, 57, 10608–10615.
52. Riley RS; O'Sullivan RK; Potocny AM; Rosenthal J; Day ES Evaluating Nanoshells and a Potent Biladiene Photosensitizer for Dual Photothermal and Photodynamic Therapy of Triple Negative Breast Cancer Cells. *Nanomaterials*. 2018, 8, 658–673.
53. Yamada K; Toyota T; Takakura K; Ishimaru M; Sugawara T Preparation of BODIPY Probes for Multicolor Fluorescence Imaging Studies of Membrane Dynamics. *New J. Chem.* 2001, 25, 667–669.
54. Chen Y; Wan L; Zhang D; Bian Y; Jiang J Modulation of the Spectroscopic Property of BODIPY Derivates Through Tuning the Molecular Configuration. *Photochem. Photobiol. Sci.* 2011, 10, 1030–1038. [PubMed: 21384046]
55. Hu W; Zhang X-F; Lu X; Lan S; Tian D; Li T; Wang L; Zhao S; Feng M; Zhang J Modifying the meso-phenyl with Electron Donating Amino Groups Strongly Enhances BODIPY's Ability as Good Singlet Oxygen Photosensitizer. *Dyes and Pigments*. 2018, 149, 306–314.
56. Rossenaar BD; Stufkens DJ; Vl ek A Jr. Halide-Dependent Change of the Lowest-Excited-State Character from MLCT to XLCT for the Complexes Re(X)(CO)₃(a-diimine) (X= Cl, Br, I; α-

- diimine = bpy, iPr-PyCa, iPr-DAB) Studied by Resonance Raman, Time-Resolved Absorption, and Emission Spectroscopy. *Inorg. Chem.* 1996, 35, 2902–2909.
57. Worl LA; Duesing R; Chen P; Ciana LD; Meyer TJ Photophysical Properties of Polypyridyl Carbonyl Complexes of Rhenium(I). *Dalton Trans.* 1991, S, 849–858.
58. Kalyanasundaram K Luminescence and Redox Reactions of the Metal-to-Ligand Charge-Transfer Excited State of Tricarbonylchloro(polypyridyl)rhenium(I) Complexes. *Faraday Trans. 2* 1986, 82, 2401–2415.
59. (Yi X; Zhao J; Sun J; Guo S; Zhang H Visible Light-Absorbing Rhenium(I) Tricarbonyl Complexes as Triplet Photosensitizers in Photooxidation and Triplet–Triplet Annihilation Upconversion. *Dalton. Trans.* 2013, 42, 2062–2074. [PubMed: 23178459]
60. Schatzschneider U PhotoCORMs: Light-Triggered Release of Carbon Monoxide from the Coordination Sphere of Transition Metal Complexes for Biological Applications. *Inorg. Chim. Acta* 2011, 374, 19–23.
61. Wright MA; Wright JA PhotoCORMs: CO Release Moves into the Visible. *Dalton Trans.* 2016, 45, 6801–6811. [PubMed: 27008479]
62. Pierri AE; Pallaoro A; Wu G; Ford PC A Luminescent and Biocompatible PhotoCORM. *J. Am. Chem. Soc.* 2012, 134, 18197–18200.
63. Chakraborty I; Carrington SJ; Mascharak PK Design Strategies to Improve the Sensitivity of Photoactive Metal Carbonyl Complexes (PhotoCORMs) to Visible Light and Their Potential as CO-Donors to Biological Targets. *Acc. Chem. Res.* 2014, 47, 2603–2611. [PubMed: 25003608]
64. Jimenez J; Chakraborty I; Dominguez A; Martinez-Gonzalez J; Sameera WMC; Mascharak PK A Luminescent Manganese PhotoCORM for CO Delivery to Cellular Targets under the Control of Visible Light. *Inorg. Chem.* 2018, 57, 1766–1773. [PubMed: 29393641]
65. Marker SC; MacMillan SN; Zipfel WR; Li Z; Ford PC; Wilson JJ Photoactivated in Vitro Anticancer Activity of Rhenium(I) Tricarbonyl Complexes Bearing Water-Soluble Phosphines. *Inorg. Chem.* 2018, 57, 1311–1331. [PubMed: 29323880]
66. Pickens RN; Neyhouse BJ; Reed DT; Ashton ST; White JK Visible Light-Activated CO Release and $^1\text{O}_2$ Photosensitizer Formation with Ru(II), Mn(I) Complexes. *Inorg. Chem.* 2018, 57, 11616–11625.
67. Mede R; Hoffmann P; Klein M; Görls H; Schmitt M; Neugebauer U; Gessner G; Heinemann SH; Popp J; Westerhausen M A Water-Soluble Mn(CO) $_3$ -Based and Non-Toxic PhotoCORM for Administration of Carbon Monoxide Inside of Cells. *Zeitschrift für Anorg. und Allg. Chemie* 2017, 643, 2057–2062.
68. Chakraborty I; Jimenez J; Mascharak PK CO-Induced Apoptotic Death of Colorectal Cancer Cells by a Luminescent PhotoCORM Grafted on Biocompatible Carboxymethyl Chitosan. *Chem. Commun.* 2017, 53, 5519–5522.

Figure 1.

Solid state structures of $\text{Re}(\text{BB}2)(\text{CO})_3\text{Cl}$,³⁷ $\text{Re}(\text{BB}3)(\text{CO})_3\text{Cl}$, and $\text{Re}(\text{BB}4)(\text{CO})_3\text{Cl}$. Thermal ellipsoids are shown at the 30% probability level (C, black; O, red; N, blue; Cl, dark green; Re, purple; B, brown; F, light green). Hydrogen atoms and cocrystallized solvent molecules have been omitted for clarity.

Figure 2.

Electronic absorption spectra (solid lines) recorded under air and normalized emission spectra (dashed lines) recorded under a nitrogen atmosphere for (a) **BB2** (black) and **Re(BB2)(CO)₃Cl** (green) in DMF, (b) **BB3** (black) in DMF containing 10% CH₂Cl₂ and **Re(BB3)(Co)₃Cl** (orange) in DMF and (c) **BB4** (black) and **Re(BB4)(CO)₃Cl** (purple) in DMF. All spectra were recorded at 298 K. Inset: Emission spectra of **BB4** and **Re(BB4)(CO)₃Cl** rescaled to illustrate the weak emission features of the Re complex.

Scheme 1.

Metallation of bpyBODIPY ligands to generate an isomeric series of *fac*-Re(bpy)(CO)₃Cl Complexes

Table 1.

Crystallographic parameters for **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl**

	Re(BB3)(CO)₃Cl	Re(BB4)(CO)₃Cl
Empirical Formula	C ₄₉ H ₅₄ B ₂ C ₁₃ F ₄ N ₆ O ₃ Re	C ₄₉ H ₅₄ B ₂ C ₁₃ F ₄ N ₆ O ₃ Re
Formula Weight	1165.15	1165.15
Crystal System	Triclinic	Orthorhombic
Space Group	<i>P</i> -1	<i>Pmn</i> 2(1)
<i>a</i>	13.383(3) Å	27.663(4) Å
<i>b</i>	14.555(3) Å	11.5474(15) Å
<i>c</i>	14.563(3) Å	8.0100(10) Å
α	88.305(4)°	90°
β	81.330(4)°	90°
γ	62.646(4)°	90°
Volume	2488.1(9) Å ³	2558.7(6) Å ³
Z	2	2
Temperature	200(2) K	200(2) K
D _{calcd}	1.555 g/cm ⁻³	1.512 g/cm ⁻³
2 θ range	1.725° – 28.302°	1.91° – 28.49°
μ	2.666 mm ⁻¹ (Mo K α)	2.592 mm ⁻¹ (Mo K α)
Reflections	27813	29089
Unique	12249	6513
<i>R</i> (int)	0.0482	0.0716
<i>R</i> _I	0.0588	0.0390
<i>wR</i> ₂	0.1297	0.0772

Table 2.

Dihedral angles and Average **Re**...**BODIPY** Distal Separation of **Re(BB2)(CO)₃Cl**, **Re(BB3)(CO)₃Cl** and **Re(BB4)(CO)₃Cl**

	^a bpy—BODIPY Dihedral Angles	Average Re ... BODIPY Separation
Re(BB2)(CO)₃Cl	73.8°, 83.2°	9.43 Å
Re(BB3)(CO)₃Cl	83.9°, 87.5°	8.39 Å
Re(BB4)(CO)₃Cl	^b 79.5°	^b 5.56 Å

^aDihedral angles measured between planes defined by each BODIPY unit and the directly appended pyridine ring.

^bDue to the presence of a mirror plane, the dihedral angles and **Re**...**BODIPY** distances measured for each BODIPY unit in **Re(BB4)(CO)₃Cl** are identical.

Table 3.

Photophysical data recorded for **BB2**, **BB3**, **BB4**, **Re(BB2)(CO)₃Cl**, **Re(BB3)(CO)₃Cl**, and **Re(BB4)(CO)₃Cl**.

	$\lambda_{\text{max abs}}/\text{nm}$ ($\epsilon \times 10^3 \text{ M}^{-1}\text{cm}^{-1}$)	$\lambda_{\text{fl}}/\text{nm}$ (Φ_{fl})	$\lambda_{\text{ph}}/\text{nm}$ (Φ_{ph})	Φ
BB2	526 nm (86.0)	550 nm (~1)	————	—
Re(BB2)(CO)₃Cl	530 nm (79.4)	561 nm (1.4×10^{-2})	————	1.7%
BB3	528 nm (95.1)	548 nm (3.5×10^{-1})	————	—
Re(BB3)(CO)₃Cl	533 nm (52.7)	553 nm (1.3×10^{-2})	————	1.6%
BB4	526 nm (100.8)	548 nm (~1)	————	—
Re(BB4)(CO)₃Cl	542 nm (88.5)	564 nm (1.3×10^{-3})	829 (1.6×10^{-4})	18.6%

Spectroscopic data for **BB3** was recorded in 10% CH₂Cl₂ in DMF. All other data was recorded in DMF.