Published in final edited form as:

Curr Opin Cell Biol. 2019 February ; 56: 102–108. doi:10.1016/j.ceb.2018.10.009.

How cells exploit tubulin diversity to build functional cellular microtubule mosaics

Antonina Roll-Mecak

Cell Biology and Biophysics Unit, National Institute of Neurological Disorders and Stroke, Bethesda, MD 20892, U.S.A.; Biophysics Center, National Heart, Lung and Blood Institute, , MD 20892, U.S.A.

Abstract

Cellular microtubules are mosaic polymers assembled from multiple α -tubulin isoforms bearing chemically diverse posttranslational modifications. This tubulin diversity constitutes a combinatorial code that regulates microtubule interactions with cellular effectors and alters their intrinsic dynamic and mechanical properties. Cells generate stereotyped and complex tubulin modification patterns that are important for their specialized functions. Here we give a brief overview of the tubulin genetic and chemical diversity and highlight recent advances in our understanding of how the tubulin code regulates essential biological processes ranging from intracellular cargo transport, to cell division and cardiomyocyte contraction. Finally, we speculate on the molecular mechanisms for the generation and maintenance of the complex stereotyped modification patterns that form cellular microtubule mosaics.

Together with actin and intermediate filaments, microtubules form one of the three cytoskeletal filament systems in eukaryotic cells. They are non-covalently linked cylindrical polymers assembled from $\alpha\beta$ -tubulin heterodimers. Microtubules provide mechanical support for cells, function as highways for the delivery of cargo and segregate chromosomes during cell division. In order to perform these disparate functions, cells build microtubule arrays with diverse morphologies and dynamics. These include the stable bundled microtubule arrays in neurons to facilitate efficient cargo transport by molecular motors, the dynamic bipolar spindle to capture chromosomes and generate the forces needed to separate them, the ring of stable microtubules beneath the membrane of platelets that gives these cells their characteristic discoid shape or the highly stable nine-fold symmetric axonemes that are optimized for high-flux cargo movement and built to withstand the high mechanical forces generated during ciliary or flagellar beating (Figure 1).

Microtubule mosaics are formed by diverse tubulin isoforms.

Microtubule arrays display a remarkable diversity in the chemical composition of their constituent subunits, the $\alpha\beta$ tubulin heterodimers. This diversity is generated through the expression of different tubulin isoforms as well as posttranslational modifications. Most

*Correspondence should be addressed to: Antonina Roll-Mecak, Cell Biology and Biophysics Unit, Porter Neuroscience Research Center, National Institutes of Health, Building 35, Room 3B-203, 35 Convent Drive, MSC 3700, Bethesda, MD 20892-3700, Telephone: 301-814-8119, Antonina@mail.nih.gov.

eukaryotes possess multiple tubulin genes (reviewed in [1]). Humans have eight α - and nine β -tubulin genes. Some cell types have restricted tubulin isotype expression, while others express multiple, thus creating many combinations of $\alpha\beta$ heterodimers [1]. Most microtubules are made of mixtures of isotypes, while some contain predominantly a single isotype. For example, the incorporation of the most divergent β -tubulin isoform, β VI (TUBB1) in platelet microtubules is essential for the formation of the platelet marginal band that gives these cells their discoid shape. Loss of this tubulin isoform results in thrombocytopenia [2]. β III tubulin (TUBB3) is specific for neurons [3] and is expressed in non-neuronal cells only during tumorigenesis [3, 4]. Mutations in β III are linked to polymicrogyria [5] and axon guidance disorders ([6], reviewed in [7]). β IVa and b tubulins (TUBB4A and B) are important for the formation of cilia and flagella [8]. They contain an axonemal specific motif on the C-terminal tail identified as important for the generation of the central pair in the motile axonemes of fly sperm [9]. Remarkably, we do not know the exact stoichiometry of tubulin isoforms in most cell types and we have a poor understanding of how tubulin isoform expression levels are regulated throughout development. We also currently lack an understanding of the basic biophysical properties of most tubulin isoforms: their kinetics of incorporation into microtubules and their effect on microtubule structure, dynamics and mechanical properties. Recent studies using recombinant tubulin isoforms and tubulin isolated from an embryonic kidney cell line using an affinity approach revealed that tubulin isoform composition can proportionally tune microtubule dynamic parameters [10, 11]. Thus, changes observed in tubulin isoform expression during differentiation or tumorigenesis can directly control microtubule dynamics in addition to possibly serving to recruit isoform specific effectors *in trans*. We also do not know whether tubulin isoforms incorporate in precise patterns in microtubule arrays and whether such patterns matter for the recruitment of cellular effectors. A comprehensive understanding of tubulin isoform distributions in cells and tissues has been so far hampered by a scarcity of tubulin isoform specific antibodies and difficulties with systematic tagging of tubulin isoforms in many organisms because of adverse effects on folding or microtubule function. The recent advances in producing recombinant tubulin isoforms [10-12] should catalyze the isolation of high-affinity affimers and antibodies with high specificity for different tubulin isoforms to facilitate such efforts.

Cells generate temporally and spatially regulated mosaic patterns of tubulin posttranslational modifications.

Tubulin is further diversified through a large number of chemically diverse posttranslational modifications (reviewed in [13] and [14]). Together with tubulin isoforms they are thought to constitute an evolutionarily conserved “tubulin code” that regulates intrinsic microtubule properties and their recognition by effectors. These modifications include acetylation of α and β -tubulin, phosphorylation of α and β -tubulin, polyamination, detyrosination/tyrosination (the removal and addition of the genetically encoded C-terminal tyrosine on α -tubulin), 2-tubulin (the removal of the last and penultimate glutamates in α -tubulin), glutamylation (the reversible addition of branched chains of glutamates of variable lengths to α or β -tubulin) and glycylation (the addition of branched chains of glycines of variable lengths to α or β -tubulin) (Figure 2). Mass spectrometric studies of tubulin isolated from

cilia have identified varieties with as many as 21 glutamates [15] and 40 glycines [16], while tubulin isolated from brain tissue is functionalized with an average of 3-6 glutamates, with as many as eleven and seven detected on α - and β -tubulin, respectively [17]. To put these numbers in perspective, the C-terminal tails of tubulin that decorate the microtubule surface and to which these amino acid chains are added, range between 10 to 15 residues in length [18]. Thus, these posttranslationally added amino acid chains can drastically change the electrostatic properties of the microtubule surface and also modify the bulk properties of the cytoplasm around this polymer.

Tubulin posttranslational modifications vary widely between cell types and their patterns are stereotyped, indicative of roles in spatial and temporal control. For example, microtubules oriented towards a wound in a confluent cell monolayer are enriched in detyrosination, microtubules in the growth cone are tyrosinated, microtubules in the axon are predominantly glutamylated, detyrosinated, acetylated and enriched in β -tubulin (reviewed in [13], references within), microtubules in the platelet marginal band are acetylated, detyrosinated and enriched in β -tubulin [19], kinetochore microtubules are detyrosinated, astral microtubules are tyrosinated, midbody microtubules are detyrosinated, glutamylated and acetylated and microtubules in centrioles, cilia and flagella are acetylated as well as heavily glutamylated, glycylation (reviewed in [13], references within; Figure 1). Currently we do not understand how the cell achieves the selective enrichment of modifications on these subsets of microtubules. Early studies revealed a correlation between microtubule lifetimes in cells and posttranslational modifications: dynamic microtubules with lifetimes on the order of tens of seconds to minutes are enriched in tyrosinated tubulin while microtubules that persist for hours and are resistant to cold or drug-induced depolymerization are enriched in acetylated, detyrosinated and glutamylated tubulin [20-22]. These subsets of stable microtubules are used for endoplasmic reticulum sliding and are preferred sites of contacts between the endoplasmic reticulum and mitochondria [23]. They are also used for the selective transport of many viruses (reviewed in [24]). Axonemes offer the most dramatic example of specialization of closely positioned microtubule tracks with different posttranslational modifications. A recent elegant study using correlative light and cryo-electron microscopy of the *Chlamydomonas* flagellum revealed that the A-tubule, enriched in tyrosination, is dedicated to retrograde intraflagellar transport (IFT), while the B-tubule, which is enriched in detyrosination, is used for anterograde transport [25]. A more recent study revealed that rapid depletion of tubulin glutamylation in the cilium through rapamycin induced targeting of the deglutamylase CCP5 preferentially impaired kinesin-2 anterograde IFT [26]. This study also demonstrates the utility of the rapamycin dimerization system to interrogate the effects of acute changes in tubulin modifications in cells. The use of photo-regulated modules [27] will allow in the future the investigation of acute changes in modification levels with higher spatial resolution than those achievable with the rapamycin dimerization system.

***Cis* and *trans* regulation of microtubule functions by tubulin modifications.**

While the effects of tubulin posttranslational modifications on microtubule dynamic parameters have still yet to be quantitatively characterized, recent studies show that these modifications regulate the recruitment and activity of cellular effectors and can affect the

mechanical properties of microtubules directly. Tyrosination regulates the recruitment Clip-170 to microtubule tips [28]. It also recruits the motor dynein through its regulator dynactin which interacts directly with the α -tubulin tail and prefers its tyrosinated form over the detyrosinated one [29, 30]. Tyrosination also enhances the depolymerization activity of kinesin-13 MCAK [31]. Detyrosination was recently shown to guide chromosomes towards the cell equator during mitosis by allowing the chromosome associated CENP-E motor to remain attached to microtubules for longer times when subjected to high loads. Disrupting the detyrosination pattern in the spindle caused the chromosomes to be transported in random directions from the spindle poles [32]. A fascinating recent study reports that asymmetric tyrosination in the mammalian oocyte spindle allows preferential meiotic segregation [33]. The asymmetric tyrosination is dependent on signaling by the small GTPase cdc42 from the oocyte cortex which increases tubulin tyrosination activity (or inhibits detyrosination) in the half-spindle close to the cortex compared to the half-spindle facing the egg cytoplasm. Detyrosination also regulates the interaction between sarcomeres and microtubules in cardiomyocytes and this interaction is mediated through desmin intermediate filaments [34]. Microtubules provide mechanical resistance during cardiomyocyte contraction through their interaction with the sarcomere. Disrupting this interaction by decreasing detyrosination reduces stiffness and allows sarcomeres to contract further and faster. Notably, cardiomyopathy patients with increased tubulin detyrosination levels showed a more pronounced functional decline [34], suggesting therapeutic interventions through the detyrosination/tyrosination cycle. Indeed, a recent study shows that cardiomyocytes from failing hearts show a dense network of detyrosinated microtubules that provide increased viscoelastic resistance and reduce contractility. A decrease in stiffness and partial recovery in contractile function can be achieved by inhibiting detyrosination with parthenolide or destabilizing microtubules with colchicine [35].

Glutamylation is ubiquitous in axons and flagella and *in vitro* studies show that this modification tunes the motility of motors that function in these compartments. For example, glutamylation results in modest increases in the run length of kinesin-1 and 2, but not that of cytoplasmic dynein [36]. Microtubule glutamylation controls the activity of microtubule severing enzymes spastin and katanin [37-39]. *In vitro* studies using microtubules with quantitatively defined levels of glutamylation revealed a graded control of spastin microtubule severing by tubulin glutamylation. This mechanism has parallels to the one recently discovered for the actin severing protein cofilin where stereospecific oxidation of actin filaments by MICAL family proteins enhances the binding of cofilin and promotes rapid filament severing [40]. Notably, the microtubule binding affinity of spastin increases linearly with the number of glutamates added to tubulin [37] indicating that glutamylation of the tubulin tails can function as a linear affinity tuner for cellular effectors. Such a graded modulation of microtubule binding affinity is likely at play for many microtubule associated proteins (MAPs) that depend on the C-terminal tails of tubulin for their binding to the microtubule, as hinted by early qualitative experiments using blot overlays [41].

Recent studies demonstrated for the first time that a tubulin posttranslational modification can affect intrinsic properties of the microtubule polymer: acetylation of α -tubulin on Lys40, a residue situated in the microtubule lumen, was shown to mechanically stabilize long-lived microtubules and prevent their breakage in cells [42]. Complementary *in vitro* experiments

showed that acetylation in fact decreases the flexural rigidity of the microtubule, but increases its resilience to repeated deformation, possibly by weakening inter-protofilament contacts and allowing protofilaments to slide with respect to each other when the microtubule bends [43]. Interestingly, acetylated microtubules are often highly curved [23], suggesting that they are subject to high compressive forces in cells. In the mechanosensory neurons of *C. elegans*, loss of the tubulin acetyltransferase produces microtubule structures that are splayed open [44], presumably because of their decreased resilience to the mechanical stress experienced by the worm during its sinusoidal movement. Intriguingly, molecular dynamics simulations proposed that acetylation at Lys40 favors the formation of a stabilizing salt bridge between α -tubulin subunits from neighboring protofilaments [44], offering a possible molecular explanation for the observed changes in mechanical properties. Since tubulin sequence variation also concentrates at polymerization interfaces (in addition to the tubulin C-terminal tails) and structures of microtubules with different tubulin isoform compositions show subtle structural variations at lateral interfaces between protofilaments [45], it will be interesting to establish whether isoform composition impacts microtubule mechanics directly.

Tubulin posttranslational modifications also act on soluble tubulin, affecting its ability to incorporate into microtubules. For example, acetylation of β -tubulin on Lys 252 by the *San* acetyltransferase slows the kinetics of tubulin incorporation into microtubules by possibly affecting the intradimer interface [46]. Recent work shows that phosphorylation by the minibrain kinase MNB/DYRK1 of Ser172 in β -tubulin inhibits tubulin polymerization and affects dendrite morphology in *Drosophila* [47]. This residue is located at the tubulin interdimer interface. Proteomic studies reveal many other modification sites on the tubulin dimer whose physiological relevance is not yet clear.

Towards understanding and reconstituting complex tubulin modification patterns.

Clearly cells expend a considerable portion of their genetic and energetic capital to regulate tubulin isoform and posttranslational diversity. However, we currently lack a comprehensive and mechanistic understanding of how tubulin isoforms are distributed in cells and how tubulin posttranslational patterns are generated and maintained. The length-scales of the mosaic blocks is still to be determined at high-resolution as well as the fidelity in generating patterns. For function, is only the enrichment of a particular type of modification or isoform important on a microtubule or set of microtubules or does the precise patterning of the mosaic blocks matter? The complex microtubule modification patterns observed in cells result from the interplay between microtubule dynamics and the subcellular localization, substrate specificity and kinetics of the tubulin modification enzymes. Kinetic control i.e. the intersection between the turnover rates of the microtubules themselves and the catalytic rates of the modification enzymes, is likely an important mechanism for generating complex spatial patterns. Interestingly, the catalytic rates of tubulin modification enzymes in isolation span a broad range, from sec^{-1} in the case of the tubulin tyrosine ligase TTL [48], min^{-1} in the case of the glutamylase TTLL7 [49] and the glycolase TTLL3 [50], to hrs^{-1} in the case of the tubulin acetyltransferase [51, 52], and thus can selectively modify microtubules with a

broad range of turnover rates as previously hypothesized [53]. In this scenario, an external factor (a MAP or anchoring by a motor to the plasma membrane or kinetochore, for example) acts as the initial stabilizing cue for the microtubule, thus prolonging its lifetime to match that of a modification enzyme. The introduced posttranslational modification can then further reinforce the initial stabilizing cue by specifically recruiting cellular effectors and/or directly affecting the intrinsic stability of the polymer.

To understand the mechanisms used by cells to selectively modify microtubules will require first their characterization at high spatial and temporal resolution in various cell types to compare and contrast patterns with enzyme expression and localization. *In vivo* perturbation analysis by manipulating modification enzymes and their regulators will have to go hand in hand with *in vitro* reconstitution with physiologically relevant tubulin isoform mixtures. Can complex spatial distributions of modified microtubules arise *ex nihilo* from the specificity, kinetics and dynamics of the modification enzymes and microtubules alone? Or do they depend on a pre-existing scaffold generated through nucleation and motor and MAP-dependent microtubule organization? *In vitro* microtubule assays utilize a nucleating structure, in many cases a stabilized GMPCPP microtubule seed, to grow single microtubules without higher order organization. While these experimental setups have been instrumental in studying the basic properties of microtubule regulators, they do not capture the geometric constraints of microtubule effectors in cells. The continuous advancements in micropatterning makes the creation of more physiologically relevant microtubule geometries in confined environments possible [54]. Wider adoption of these technologies coupled with advances in the biochemical characterization of the modification enzymes and the signaling cascades that regulate them may get us closer to recapitulating *in vitro* the chemical and geometric diversities of cellular microtubule arrays.

Acknowledgments

We thank Erina He (National Institute of Health Medical Arts) for help with illustrations.

Funding

A.R.M. is supported by the intramural programs of the National Institute of Neurological Disorders and Stroke and the National Heart, Lung and Blood Institute of the National Institutes of Health, USA.

References and Recommended Reading

Papers of special interest, published within the last two years, have been highlighted as:

*of interest

**of outstanding interest

1. Fojo AT (2009). The role of microtubules in cell biology, neurobiology, and oncology, (Springer Science & Business Media).
2. Schwer HD, Lecine P, Tiwari S, Italiano JE Jr, Hartwig JH, and Shivdasani RA (2001). A lineage-restricted and divergent β -tubulin isoform is essential for the biogenesis, structure and function of blood platelets. *Current Biology* 11, 579–586. [PubMed: 11369202]
3. Burgoyne R, Cambray-Deakin M, Lewis S, Sarkar S, and Cowan N (1988). Differential distribution of beta-tubulin isotypes in cerebellum. *The EMBO journal* 7, 2311–2319. [PubMed: 2461292]

4. Katsetos CD, Herman MM, and Mörk SJ (2003). Class III β -tubulin in human development and cancer. *Cell motility and the cytoskeleton* 55, 77–96. [PubMed: 12740870]
5. Poirier K, Saillour Y, Bahi-Buisson N, Jaglin XH, Fallet-Bianco C, Nabbout R, Castelnau-Ptakhine L, Roubertie A, Attie-Bitach T, and Desguerre I (2010). Mutations in the neuronal β -tubulin subunit TUBB3 result in malformation of cortical development and neuronal migration defects. *Human molecular genetics* 19, 4462–4473. [PubMed: 20829227]
6. Tischfield MA, Baris HN, Wu C, Rudolph G, Van Maldergem L, He W, Chan W-M, Andrews C, Demer JL, and Robertson RL (2010). Human TUBB3 mutations perturb microtubule dynamics, kinesin interactions, and axon guidance. *Cell* 140, 74–87. [PubMed: 20074521]
7. Breuss MW, Leca I, Gstrein T, Hansen AH, and Keays DA (2017). Tubulins and brain development—The origins of functional specification. *Molecular and Cellular Neuroscience*.
8. Jensen-Smith HC, Ludueña RF, and Hallworth R (2003). Requirement for the β I and β IV tubulin isotypes in mammalian cilia. *Cell motility and the cytoskeleton* 55, 213–220. [PubMed: 12789665]
9. Nielsen MG, Turner FR, Hutchens JA, and Raff EC (2001). Axoneme-specific β -tubulin specialization: a conserved C-terminal motif specifies the central pair. *Current Biology* 11, 529–533. [PubMed: 11413005]
- **10. Pamula MC, Ti S-C, and Kapoor TM (2016). The structured core of human β tubulin confers isotype-specific polymerization properties. *J Cell Biol* 213, 425–433. [PubMed: 27185835] Uses recombinant purified -tubulin isoforms to show that sequence differences in the structured core of human β IIItubulin are responsible for a lower catastrophe frequency compared to neuronal β III.
- **11. Vemu A, Atherton J, Spector JO, Moores CA, and Roll-Mecak A (2017). Tubulin isoform composition tunes microtubule dynamics. *Molecular biology of the cell* 28, 3564–3572. [PubMed: 29021343] Shows that microtubules with different tubulin isoforms compositions have different dynamic parameters and reports the 4.2Å cryo-EM structure of a microtubule composed of α 1B/ β I+ β IVb tubulin. Also shows that the dynamic parameters of non-neuronal α 1B/ β I+ β IVb microtubules can be proportionally tuned by the addition of neuronal α 1A/ β III tubulin.
12. Minoura I, Hachikubo Y, Yamakita Y, Takazaki H, Ayukawa R, Uchimura S, and Muto E (2013). Overexpression, purification, and functional analysis of recombinant human tubulin dimer. *FEBS letters* 587, 3450–3455. [PubMed: 24021646]
13. Yu I, Garnham CP, and Roll-Mecak A (2015). Writing and reading the tubulin code. *Journal of Biological Chemistry*, jbc. R115. 637447.
14. Song Y, and Brady ST (2015). Post-translational modifications of tubulin: pathways to functional diversity of microtubules. *Trends in cell biology* 25, 125–136. [PubMed: 25468068]
15. Schneider A, Plessmann U, Felleisen R, and Weber K (1998). Posttranslational modifications of trichomonad tubulins; identification of multiple glutamylation sites. *FEBS letters* 429, 399–402. [PubMed: 9662457]
16. Wall KP, Pagratis M, Armstrong G, Balsbaugh JL, Verbeke E, Pearson CG, and Hough LE (2016). Molecular determinants of tubulin's C-terminal tail conformational ensemble. *ACS chemical biology* 11, 2981–2990. [PubMed: 27541566]
17. Redeker V (2010). Mass spectrometry analysis of C-terminal posttranslational modifications of tubulins In *Methods in cell biology*, Volume 95 (Elsevier), pp. 77–103. [PubMed: 20466131]
18. Roll-Mecak A (2015). Intrinsically disordered tubulin tails: complex tuners of microtubule functions? In *Seminars in cell & developmental biology*, Volume 37 (Elsevier), pp. 11–19. [PubMed: 25307498]
19. Diagouraga B, Grichine A, Fertin A, Wang J, Khochbin S, and Sadoul K (2014). Motor-driven marginal band coiling promotes cell shape change during platelet activation. *J Cell Biol* 204, 177–185. [PubMed: 24421335]
20. Schulze E, and Kirschner M (1987). Dynamic and stable populations of microtubules in cells. *The Journal of cell biology* 104, 277–288. [PubMed: 3543024]
21. Webster DR, Gunderson GG, Bulinski JC, and Borisy GG (1987). Differential turnover of tyrosinated and detyrosinated microtubules. *Proceedings of the National Academy of Sciences* 84, 9040–9044.
22. Webster DR, and Borisy GG (1989). Microtubules are acetylated in domains that turn over slowly. *Journal of cell science* 92, 57–65. [PubMed: 2674164]

23. Friedman JR, Webster BM, Mastronarde DN, Verhey KJ, and Voeltz GK (2010). ER sliding dynamics and ER-mitochondrial contacts occur on acetylated microtubules. *The Journal of cell biology* 190, 363–375. [PubMed: 20696706]
24. Naghavi MH, and Walsh D (2017). Microtubule regulation and function during virus infection. *Journal of virology*, JVI. 00538–00517.
- **25. Stepanek L, and Pigino G (2016). Microtubule doublets are double-track railways for intraflagellar transport trains. *Science* 352, 721–724. [PubMed: 27151870] Uses light microscopy combined with cryo-EM to show that anterograde and retrograde IFT transport are restricted to the B and A-tubule, respectively.
- *26. Hong S-R, Wang C-L, Huang Y-S, Chang Y-C, Chang Y-C, Pusapati GV, Lin C-Y, Hsu N, Cheng H-C, and Chiang Y-C (2018). Spatiotemporal manipulation of ciliary glutamylation reveals its roles in intraciliary trafficking and Hedgehog signaling. *Nature communications* 9. Introduces an effective method for rapid, spatially controlled manipulation of glutamylation levels in cells. Reports use of this method to rapidly deplete glutamylation in cilia and show that it impairs anterograde IFT.
27. Pudasaini A, El-Arab KK, and Zoltowski BD (2015). LOV-based optogenetic devices: light-driven modules to impart photoregulated control of cellular signaling. *Frontiers in molecular biosciences* 2, 18. [PubMed: 25988185]
28. Bieling P, Kandels-Lewis S, Telley IA, van Dijk J, Janke C, and Surrey T (2008). CLIP-170 tracks growing microtubule ends by dynamically recognizing composite EB1/tubulin-binding sites. *J Cell Biol* 183, 1223–1233. [PubMed: 19103809]
- **29. Nirschl JJ, Magiera MM, Lazarus JE, Janke C, and Holzbaur EL (2016). α -Tubulin tyrosination and CLIP-170 phosphorylation regulate the initiation of dynein-driven transport in neurons. *Cell reports* 14, 2637–2652. [PubMed: 26972003] Shows that the α -tubulin tyrosination gradient in the axon and CLIP-170 phosphorylation initiates retrograde transport at the distal axon, through an elegant combination of *in vitro* reconstitution assays with tyrosinated and detyrosinated microtubules, live-cell imaging of cultured neurons, and computer simulations.
- *30. McKenney RJ, Huynh W, Vale RD, and Sirajuddin M (2016). Tyrosination of α -tubulin controls the initiation of processive dynein-dynactin motility. *The EMBO journal*, e201593071. Shows tyrosinated tubulin recruits dynein/dynactin and initiates processive retrograde dynein/dynactin motility through a specific interaction between the CAP-Gly domain of p150 with the C-terminal tyrosine of α -tubulin using *in vitro* motility assays with engineered *S. cerevisiae* tubulin. Once motility is initiated, the C-terminal tyrosine is not needed for processive movement.
31. Peris L, Wagenbach M, Lafanechère L, Brocard J, Moore AT, Kozielski F, Job D, Wordeman L, and Andrieux A (2009). Motor-dependent microtubule disassembly driven by tubulin tyrosination. *The Journal of cell biology* 185, 1159–1166.
- **32. Barisic M, e Sousa RS, Tripathy SK, Magiera MM, Zaytsev AV, Pereira AL, Janke C, Grishchuk EL, and Maiato H (2015). Microtubule detyrosination guides chromosomes during mitosis. *Science* 348, 799–803. [PubMed: 25908662] Shows that detyrosination increases the dwell times of CENP-E under higher load and that this interaction is important for guiding chromosomes to the cell equator during mitosis.
- **33. Akera T, Chmátal L, Trimm E, Yang K, Aonbangkhen C, Chenoweth DM, Janke C, Schultz RM, and Lampson MA (2017). Spindle asymmetry drives non-Mendelian chromosome segregation. *Science* 358, 668–672. [PubMed: 29097549] Shows that non-Mendelian chromosome segregation is mediated by the asymmetric tyrosination of the spindle microtubules in the mouse oocyte. The asymmetric tyrosination is mediated through cdc42 signaling from the cell cortex.
- **34. Robison P, Caporizzo MA, Ahmadzadeh H, Bogush AI, Chen CY, Margulies KB, Shenoy VB, and Prosser BL (2016). Detyrosinated microtubules buckle and bear load in contracting cardiomyocytes. *Science* 352, aaf0659. [PubMed: 27102488] Shows that microtubule provide mechanical resistance during cardiomyocyte contraction and that this resistance is due to an interaction between microtubules and sarcomers that is mediated by desmin and is dependent on microtubule detyrosination.
35. Chen CY, Caporizzo MA, Bedi K, Vite A, Bogush AI, Robison P, Heffler JG, Salomon AK, Kelly NA, and Babu A (2018). Suppression of detyrosinated microtubules improves cardiomyocyte function in human heart failure. *Nature medicine*, 1.

36. Sirajuddin M, Rice LM, and Vale RD (2014). Regulation of microtubule motors by tubulin isotypes and post-translational modifications. *Nature cell biology* 16, 335. [PubMed: 24633327]
- **37. Valenstein ML, and Roll-Mecak A (2016). Graded control of microtubule severing by tubulin glutamylation. *Cell* 164, 911–921. [PubMed: 26875866] Introduces a biochemical platform to examine the effects of tubulin postranslational modifications and shows that glutamylation is the main regulator of the hereditary spastic paraplegia microtubule severing enzyme spastin and that neither detyrosination nor acetylation regulate spastin activity. Glutamylation tunes microtubule severing as a function of glutamate number added per tubulin, providing evidence for a precise graded response to a tubulin polymodification.
38. Sharma N, Bryant J, Wloga D, Donaldson R, Davis RC, Jerka-Dziadosz M, and Gaertig J (2007). Katanin regulates dynamics of microtubules and biogenesis of motile cilia. *The Journal of cell biology* 178, 1065–1079. [PubMed: 17846175]
39. Lacroix B, Van Dijk J, Gold ND, Guizetti J, Aldrian-Herrada G, Rogowski K, Gerlich DW, and Janke C (2010). Tubulin polyglutamylation stimulates spastin-mediated microtubule severing. *The Journal of cell biology* 189, 945–954. [PubMed: 20530212]
40. Grintsevich EE, Yesilyurt HG, Rich SK, Hung R-J, Terman JR, and Reisler E (2016). F-actin dismantling through a redox-driven synergy between Mical and cofilin. *Nature cell biology* 18, 876. [PubMed: 27454820]
41. Bonnet C, Boucher D, Lazereg S, Pedrotti B, Islam K, Denoulet P, and Larcher J-C (2001). Differential binding regulation of microtubule-associated proteins MAP1A, MAP1B and MAP2 by tubulin polyglutamylation. *Journal of Biological Chemistry*.
- **42. Xu Z, Schaedel L, Portran D, Aguilar A, Gaillard J, Marinkovich MP, Théry M, and Nachury MV (2017). Microtubules acquire resistance from mechanical breakage through intraluminal acetylation. *Science* 356, 328–332. [PubMed: 28428427] Reports that acetylation by TAT is required for mechanical stabilization of long-lived microtubules in fibroblasts. *In vitro* bending assays show that acetylation protects microtubules against mechanical rupture by making them more flexible.
- **43. Portran D, Schaedel L, Xu Z, Théry M, and Nachury MV (2017). Tubulin acetylation protects long-lived microtubules against mechanical ageing. *Nature cell biology* 19, 391. [PubMed: 28250419] Reports that acetylation at Lys40 decreases the strength of lateral interactions in the microtubule lattice and results in reduced nucleation and faster depolymerization rates. Using an *in vitro* bending assay reports that acetylation reduces the microtubule persistence length, a finding also reported in ref. 42. The authors propose that this increase in plasticity with acetylation is facilitated by easier sliding of the protofilaments in microtubules under mechanical stress.
44. Cueva JG, Hsin J, Huang KC, and Goodman MB (2012). Posttranslational acetylation of α -tubulin constrains protofilament number in native microtubules. *Current Biology* 22, 1066–1074. [PubMed: 22658592]
- *45. Vemu A, Atherton J, Spector JO, Szyk A, Moores CA, and Roll-Mecak A (2016). Structure and dynamics of single-isoform recombinant neuronal human tubulin. *Journal of Biological Chemistry*, jbc. C116. 731133. Reports the first purification, dynamic parameters and high-resolution cryoEM structure of recombinant tag-free isotypically-pure neuronal α 1A/ β III microtubules.
46. Chu C-W, Hou F, Zhang J, Phu L, Loktev AV, Kirkpatrick DS, Jackson PK, Zhao Y, and Zou H (2011). A novel acetylation of β -tubulin by San modulates microtubule polymerization via down-regulating tubulin incorporation. *Molecular biology of the cell* 22, 448–456. [PubMed: 21177827]
47. Ori-McKenney KM, McKenney RJ, Huang HH, Li T, Meltzer S, Jan LY, Vale RD, Wiita AP, and Jan YN (2016). Phosphorylation of β -tubulin by the down syndrome kinase, Minibrain/DYRK1a, regulates microtubule dynamics and dendrite morphogenesis. *Neuron* 90, 551–563. [PubMed: 27112495]
48. Szyk A, Deaconescu AM, Piszczek G, and Roll-Mecak A (2011). Tubulin tyrosine ligase structure reveals adaptation of an ancient fold to bind and modify tubulin. *Nature structural & molecular biology* 18, 1250.

49. Garnham CP, Vemu A, Wilson-Kubalek EM, Yu I, Szyk A, Lander GC, Milligan RA, and Roll-Mecak A (2015). Multivalent microtubule recognition by tubulin tyrosine ligase-like family glutamylases. *Cell* 161, 1112–1123. [PubMed: 25959773]
50. Garnham CP, Yu I, Li Y, and Roll-Mecak A (2017). Crystal structure of tubulin tyrosine ligase-like 3 reveals essential architectural elements unique to tubulin monoglycylases. *Proceedings of the National Academy of Sciences*, 201617286.
51. Shida T, Cueva JG, Xu Z, Goodman MB, and Nachury MV (2010). The major α -tubulin K40 acetyltransferase α TAT1 promotes rapid ciliogenesis and efficient mechanosensation. *Proceedings of the National Academy of Sciences*, 201013728.
52. Szyk A, Deaconescu AM, Spector J, Goodman B, Valenstein ML, Ziolkowska NE, Kormendi V, Grigorieff N, and Roll-Mecak A (2014). Molecular basis for age-dependent microtubule acetylation by tubulin acetyltransferase. *Cell* 157, 1405–1415. [PubMed: 24906155]
53. Schulze E, Asai DJ, Bulinski JC, and Kirschner M (1987). Posttranslational modification and microtubule stability. *The Journal of Cell Biology* 105, 2167–2177. [PubMed: 3316248]
54. Ghosh S, Hentrich C, and Surrey T (2013). Micropattern-controlled local microtubule nucleation, transport, and mesoscale organization. *ACS chemical biology* 8, 673–678. [PubMed: 23294267]

Figure 1. Cells have microtubule arrays with diverse geometries and chemical compositions
 Microtubule organization in a neuron, fibroblast, platelet, spindle and the motile axoneme of a ciliated protozoan. Dynamic (tyrosinated) microtubules shown in yellow; stable microtubules shown in red. The magnifying glass shows the enrichment in modifications on microtubule subpopulations: tyrosination (yellow), glutamylation or detyrosination (red), glycylation (cyan) and acetylation (magenta). We note that the abundance of these modifications on microtubules varies. For example, the abundance of glutamylated tubulin is higher in axonemes than in the axon, however comprehensive quantitative measures of modification levels on different microtubule subpopulations have yet to be undertaken.

Figure 2. Microtubule structure and posttranslational modifications

Left, microtubule structure showing tubulin C-terminal tails emanating from the microtubule surface and functionalized with various posttranslational modifications. The cross-section shows acetylated Lys40 in the microtubule lumen. Right, cartoon representation of $\alpha\beta$ -tubulin heterodimers functionalized with chemically diverse posttranslational modifications.