Published in final edited form as:

Nature. 2019 October ; 574(7778): 336–337. doi:10.1038/d41586-019-02975-w.

Brain-to-pancreas Signalling Axis Links Nicotine and Diabetes

GIUSEPPE BRUSCHETTA, SABRINA DIANO

Department of Cellular and Molecular Physiology and the Program of Integrative Cell Signaling and Neurobiology of Metabolism, Yale University School of Medicine, New Haven, Connecticut 06520, USA.

Abstract

The discovery of a signalling axis that connects nicotine responses in the brain with glucose metabolism by the pancreas sheds light on why cigarette smoking increases the risk of diabetes.

One of the many dangers of smoking is an increased risk of diabetes, because nicotine uptake leads to altered glucose metabolism and increased blood sugar levels^{1–4}. A group of neurons activated by nicotine is located in the brain's medial habenula (mHb) region; they are characterized by expression of nicotine acetylcholine receptor (nAChR) proteins⁵. These neurons promote aversive responses to nicotine, but until now, no one knew whether they also control the diabetes-associated effects of smoking — and if so, how. On page 372, Duncan *et al.*⁶ identify a signalling pathway that links nAChR-expressing neurons and blood-glucose regulation by the pancreas. It involves a diabetes-associated transcription factor, TCF7L2.

TCF7L2 is part of a signalling pathway⁷ that regulates the production and secretion of the hormone glucagon-like peptide 1 (GLP-1). GLP-1 promotes insulin release from the pancreas, and insulin promotes glucose uptake from the blood into tissues to be stored as fat. Thus, changes in TCF7L2 expression or activity can influence glucose metabolism. (Of note, the pathway is complex because activation of GLP-1 receptors by GLP-1 in turn leads to TCF7L2 activity, creating a signalling loop⁷.)

Duncan *et al.* discovered that TCF7L2 is expressed in the mHb. Given the role of the mHb in the body's response to nicotine, the authors set out to investigate the possibility that TCF7L2 expression in this region is involved in nicotine-induced alterations in glucose metabolism. The researchers first generated two strains of mice — one in which the function of TCF7L2 was impaired in all tissues, and one in which the protein's expression could be inhibited in the mHb by introducing a virus carrying a short RNA sequence. They then provided the mice with a nicotine solution through a drip, rigged such that the animals could push a lever to receive an injection of nicotine at will. Both mice strains showed greater nicotine intake than did control animals.

Prolonged stimulation by nicotine causes nAChR-expressing neurons to become desensitized and stop responding to the molecule⁸. Further examination of the mutant

animals indicated that TCF7L2 improves the ability of nAChR-expressing neurons to recover from this desensitization, presumably therefore promoting aversive responses to smoking. TCF7L2 seems to exert this effect by promoting signalling through cyclic AMP (ref. 9) — a molecule often involved in intracellular signal transduction.

Duncan *et al.* next demonstrated that TCF7L2 augments the ability of nicotine to increase blood-sugar levels: inhibition of TCF7L2 or GLP-1 receptors abolished this effect of nicotine, whereas a drug that stimulates GLP-1 receptors had the opposite effect. Furthermore, when the authors subjected wild-type mice to nicotine exposure and subsequent withdrawal, they observed increased levels of glucose, glucagon and insulin in the blood — indicative of an inability to properly regulate glucose metabolism. The effects were lessened in the mutant mice. Together, these analyses suggest that TCF7L2 activity decreases dependence on nicotine, but, conversely, hampers the body's ability to counteract the detrimental effect of nicotine on glucose metabolism.

To explore the neural circuits involved, Duncan and colleagues injected wild-type mice with a fluorescently labelled 'retrograde' virus, which travels along any neurons connected to an injection site. After injection in the pancreas, the authors found the virus in several areas of the brain, including the mHb, indicating the existence of a neuronal signalling axis from the mHb to the pancreas, by way of other brain regions. Thus nicotine, by stimulating mHb neurons in a TCF7L2-dependent manner, engages neuronal signalling to the pancreas and so alters blood glucose levels.

A 2011 report indicated that only around 6% of smokers quit successfully each year¹⁰, and data indicate that people with diabetes find it harder to quit than do people without the condition¹¹. In a final set of experiments, Duncan *et al.* gave mice sucrose every day for 6 weeks and showed that the subsequent increase in blood sugar led to reduced TCF7L2 levels and a decrease in nicotine-evoked activity in nAChR-expressing neurons. This observation suggests a feedback mechanism that could explain why it is harder for people with diabetes to give up smoking¹¹ (Fig. 1).

Duncan and colleagues' study sheds new light on the role of nicotine in the regulation of glucose metabolism. However, there are some caveats to consider. TCF7L2 is expressed not only in the mHb, but also in other brain regions¹². The pancreas, fat tissues and intestine — all of which are involved in glucose metabolism — also express TCF7L2. The possible contribution of these structures to nicotine-induced dysregulation of glucose metabolism needs to be considered and evaluated.

Furthermore, nicotine is a strong activator of the hypothalamus–pituitary–adrenal (HPA) axis — a network that promotes the release of stress hormones¹³. The function of the mHb is altered by stress, and stress hormones induce changes in blood glucose¹³. As such, it is conceivable that the some of the effects reported by the authors reflect not only the direct effect of nicotine on the mHb, but also an indirect effect through HPA-axis activation.

An interesting question is whether the effect of nicotine on the mHb–pancreas axis is different in males and females. In support of this idea, more men than women smoke, and the risk of female smokers developing diabetes is much greater than the risk for male

smokers, compared with their non-smoking counterparts⁴. Furthermore, nicotine withdrawal induces greater weight gain in women than in men¹⁴.

More broadly, further investigation is needed to confirm the role of the mHb–pancreas circuit in humans. Tobacco addiction in humans involves the interplay of pharmacological, genetic, social and environmental factors. Therefore, the full picture of nicotine’s role in diabetes is likely to involve much more than a single regulatory circuit. Finally, this work raises the question of whether and how TCF7L2 could be targeted to combat tobacco dependency and diabetes. The feasibility of this tantalizing idea will require much more investigation in both mice and humans.

References

1. Chiolero A, Faeh D, Paccaud F & Cornuz J Am. J. Clin. Nutr 87, 801–809 (2008). [PubMed: 18400700]
2. Chase HP et al. J. Am. Med. Assoc 265, 614–617 (1991).
3. Madsbad S et al. Diabetes Care 3, 41–43 (1980). [PubMed: 6996967]
4. Will JC, Galuska DA, Ford ES, Mokdad A & Calle EE Int. J. Epidemiol 30, 540–546 (2001). [PubMed: 11416080]
5. Benowitz NL N. Engl. J. Med 362, 2295–2303 (2010). [PubMed: 20554984]
6. Duncan A et al. Nature 574, 372–377 (2019). [PubMed: 31619789]
7. Liu Z & Habener JF J. Biol. Chem 283, 8723–8735 (2008). [PubMed: 18216022]
8. Picciotto MR, Addy NA, Mineur YS & Brunzell DH Prog. Neurobiol 84, 329–342 (2008). [PubMed: 18242816]
9. Khiroug L, Sokolova E, Giniatullin R, Afzalov R & Nistri A J. Neurosci 18, 2458–2466 (1998). [PubMed: 9502806]
10. Centers for Disease Control and Prevention. MMWR Morbid. Mortal. Wkly Rep 60, 1513–1519 (2011).
11. Solberg LI, Desai JR, O’Connor PJ, Bishop DB & Devlin HM Ann. Fam. Med 2, 26–32 (2004). [PubMed: 15053280]
12. Lee S, Lee CE, Elias CF & Elmquist JK J. Comp. Neurol 517, 925–939 (2009). [PubMed: 19845015]
13. Tweed JO, Hsia SH, Lutfy K & Friedman TC Trends Endocrinol. Metab 23, 334–342 (2012). [PubMed: 22561025]
14. Pirie PL, Murray DM & Luepker RV Am. J. Public Health 81, 324–327 (1991). [PubMed: 1994740]

Figure 1 |. A link between nicotine, diabetes and addiction.

Nicotine activates nicotine acetylcholine receptor (nAChR) proteins in neurons. Activation of a population of nAChR-expressing neurons in the brain's medial habenula (mHb) region leads to aversive responses to nicotine. **a**, Duncan *et al.*⁶ report that activation of these receptors in mice also leads to increases in the release of glucagon and insulin hormones from the pancreas. This leads to increased blood sugar levels — a change associated with an increased risk of diabetes in humans. **b**, The authors find that increased blood sugar levels in turn inhibit the activity of the nAChR-expressing neurons, establishing a feedback loop. **c**, As a result of this feedback, aversive responses to smoking are no longer triggered, resulting in nicotine dependence. The entire circuit is modulated by the protein TCF7L2 that, acting through a second messenger molecule called cAMP, mediates the sensitivity of nAChR-expressing neurons to nicotine.