Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

an Ankura endograft (Lifetech, Shenzhen, China). He received only 3 packs of red blood cells intraoperatively, and after the procedure, he was transferred immediately to the vascular surgery ward. No ICU was needed. He had an uneventful recovery, with full mobilization and oral feeding from the first postoperative day and discharge on the second postoperative day.

Although the type of treatment of rAAA is still debatable,¹ EVAR is considered the first treatment option in an increasing number of vascular departments worldwide because of the reduced perioperative risk and shorter postoperative in-hospital length of stay.² Definitely, the low number suitable for endovascular repair³ should be taken into consideration. The 2018 guidelines for the treatment of AAA recommend EVAR over open repair in anatomically suitable patients with rAAA.² Performing the procedure under local anesthesia is an additional advantage. In a recent analysis of the Vascular Quality Initiative database, patients with rAAA who were treated with EVAR under local anesthesia compared with EVAR under general anesthesia had decreased intraoperative time, decreased number of intraoperative blood transfusions, decreased ICU length of stay, and fewer postoperative pulmonary complications.⁴

Today, when everyone in the health care system struggles with challenges posed by the coronavirus, every choice should be made with the concept in mind of "doing the right thing for the right reason." Using EVAR, if it is anatomically possible, under local anesthesia rather than open repair or EVAR under general anesthesia seems to be the best solution. This way, we can achieve both goals at the same time, that is, treating patients in danger and saving valuable health care resources.

Christos Verikokos, MD

Andreas M. Lazaris, MD, FACS, FEBVS

George Geroulakos, MD, PhD, FRCS

Vascular Surgery Department
Attikon University Hospital
National and Kapodistrian University of Athens
Athens, Greece

C.V. and A.L. share co-first authorship.

REFERENCES

1. IMPROVE Trial Investigators, Powell JT, Sweeting MJ, Thompson MM, Ashleigh R, Bell R, Gomes M, et al. Endovascular or open repair strategy for ruptured abdominal aortic aneurysm: 30 day outcomes from IMPROVE randomized trial. *BMJ* 2014;348:f7661.
2. Chaikof EL, Dalman RL, Eskandari MK, Jackson BM, Lee WA, Mansour MA, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *J Vasc Surg* 2018;67:2-77.e2.
3. Hoornweg LL, Wisselink W, Vahl A, Balm R; Amsterdam Acute Aneurysm Trial Collaborators. The Amsterdam Acute Aneurysm Trial: suitability and application rate for endovascular

repair of ruptured abdominal aortic aneurysms. *Eur J Vasc Endovasc Surg* 2007;33:679-83.

4. Faizer R, Weinhandl E, El Hag, Le Jeune S, Apostolidou I, Shafii SM, et al. Decreased mortality with local versus general anesthesia in endovascular aneurysm repair for ruptured abdominal aortic aneurysm in the Vascular Quality Initiative database. *J Vasc Surg* 2019;70:92-101.

<https://doi.org/10.1016/j.jvs.2020.04.009>

Time to pause, to think, and to recalibrate after COVID-19

It is an eerie feeling entering the theater complex. All elective surgery has stopped as theaters have been converted to additional intensive treatment unit pods, with emergency work restricted to one designated theater.

The number of emergencies has fallen as people stay away from hospitals and surgeons follow the COVID-19 guidelines to reduce the risk of infection to staff and patients.¹ The recommendation is to avoid surgery where possible, especially if it avoids admission and use of an intensive treatment unit bed.^{2,3} There are, however, differences across the United Kingdom and the United States regarding the specifics in the advice.^{4,5} Some of the recommendations are based not on data but on exercising the most cautious and protective approach possible.⁶ Surgeons' and theater staff's risk of acquiring an infection is higher because of a combination of factors and the ability of the virus to remain viable on surfaces for a long time.^{7,8}

It will be interesting to see what happens when we get back to normality again and start seeing patients face to face, instead of virtual consultations on the phone. What will that normality look like for each specialty and for patients who did not get their urgent surgery? It will be interesting to note how many diabetic feet get septic without their urgent débridement or end up with major amputations without their urgent angioplasties and bypasses.

For stroke and transient ischemic attack patients with significant carotid artery stenosis who do not undergo carotid endarterectomy within 7 days, the COVID-19 era will be a defining moment for those championing the benefit of the modern best medical management. It will be a chance to see how many such patients end up with a major stroke within the next 3 months and if the carotid stenting rate goes up.

Similarly, for renal access, no new arteriovenous fistulas or grafts are being made. We are just dealing with complications like infections and blowouts of the old ones. All new starters, crash landers, and those with problematic arteriovenous fistulas are having tunneled lines inserted, with possibly a greater rate of infections as time may tell?

For patients with abdominal aortic aneurysms >5.5 cm, it will be a time of apprehension and frustration. Having been on surveillance for years and reached the threshold for intervention, suddenly they are being told that risk-benefit ratio does not favor surgery in those with <7 -cm abdominal aortic aneurysms (6.5 cm in the American College of Surgeons recommendations). It will be interesting to see how many of them leak or rupture and need an emergency endovascular or open repair.

The effect on surgical training, too, will be immense as surgical residents are redeployed in critical care. The training has essentially been paused and, once it is all over, will have to be extended.

As we sit back and wait for the COVID-19 blanket to lift, the surgical world could be a very different place. Let us not let COVID-19 become a death note for surgery but maintain data and accurately document outcomes to recalibrate what we do.

Raghvinder Pal Singh Gambhir, MS, DNB, FACS, FRCSEd, FFSTEd, FRCS, FEBS

King's College Hospital
London, United Kingdom

REFERENCES

1. Joint surgical colleges guidance for surgeons working during the COVID-19 pandemic. Updated Intercollegiate Guidance. Available at: <https://www.rcseng.ac.uk/coronavirus/joint-guidance-for-surgeons-v2>. Accessed March 28, 2020.
2. NHS England. Clinical guide for the management of vascular patients during the coronavirus pandemic. Available at: <https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/52/2020/03/specialty-guide-management-of-vascular-surgery-patients-v1-20-march-2020.pdf>. Accessed March 28, 2020.
3. Royal College of Surgeons. Guidance for surgeons working during the COVID-19 pandemic. Available at: <https://www.rcseng.ac.uk/coronavirus/joint-guidance-for-surgeons-v1/>. Accessed March 28, 2020.
4. COVID-19 and vascular surgery. Response statement from the Vascular Society for Great Britain and Ireland. Available at: https://www.vascularsociety.org.uk/_userfiles/pages/files/Newsletters/2020/Presidents%20update%2027_03_20.pdf. Accessed March 28, 2020.
5. American College of Surgeons. COVID-19 guidelines for triage of vascular surgery patients. Available at: <https://www.facs.org/covid-19/clinical-guidance/elective-case/vascular-surgery>. Accessed March 28, 2020.
6. SAGES statement on potential viral transmission during the use of laparoscopy, and strategies to address that potential risk. Available at: <https://www.sages.org/recommendations-surgical-response-covid-19/>. Accessed April 5, 2020.
7. Raphael T. Why surgeons in Europe and America don't want to operate right now. Available at: <https://theprint.in/health/why-surgeons-in-europe-and-america-dont-want-to-operate-right-now/387348/>. Accessed March 27, 2020.
8. van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1 [published online ahead of print April 16, 2020]. *N Engl J Med*. doi: [10.1056/NEJMc2004973](https://doi.org/10.1056/NEJMc2004973).

<https://doi.org/10.1016/j.jvs.2020.04.011>

Vascular surgery department adjustments in the era of the COVID-19 pandemic

In Portugal, a state of emergency was declared on March 18 because of the COVID-19 pandemic. Every nonessential business and activity was ceased, and home confinement was required for everyone not involved in essential activities.

This pandemic has brought significant changes to our personal and professional lives. Patients with vascular diseases are especially prone to development of complicated disease if infected by SARS-CoV-2.¹ It is our responsibility as vascular surgeons to protect our patients while keeping an active practice because most of our patients have a life- or limb-threatening disease.

We work in a tertiary university hospital in Lisbon, the capital of Portugal, which has a universal public health system. Our department is one of largest in the country and accommodates a significant part of all referrals to vascular surgery, including complex aortic diseases, with an annual volume of approximately 1700 vascular procedures. Most of our patients treated are urgent or emergent.

Among many other issues, this pandemic presented us with a considerable dilemma. As our hospital is the largest in Lisbon, we had to accommodate a significant number of the COVID-19 patients, including a high demand for intensive care unit (ICU) beds. Under these circumstances, how could we protect our patients and simultaneously keep up a practice to respond to urgent cases?

First, we postponed and discharged every patient we considered nonurgent: asymptomatic carotid stenosis, non-limb-threatening ischemia, aortic aneurysm <5.5 cm, and vascular malformations. All outpatient clinics were carried out by telemedicine, and only urgent patients were seen in person.

Second, we divided our entire staff into two teams alternating on a week-to-week basis. The staff with some ICU experience started to have specific ICU training to be called on to support COVID-19 ICUs if the hospital needed it. Senior staff was put on leave to be called on if necessary. All of this was done to obtain a maximum reserve of team members. Staff symptom checking was routinely performed to ensure that no one had contracted the disease, and all positive nonprotected contacts were sent home for quarantine.