Published in final edited form as:

Sci Immunol. 2018 May 04; 3(23): . doi:10.1126/sciimmunol.aao1392.

MicroRNA-146a controls functional plasticity in $\gamma\delta$ T cells by targeting NOD1

Nina Schmolka^{1,*}, Pedro H. Papotto^{1,†}, Paula Vargas Romero^{1,†}, Tiago Amado¹, Francisco J. Enguita¹, Ana Amorim¹, Ana F. Rodrigues^{2,3}, Katrina E. Gordon⁴, Ana S. Coroadinha^{2,3}, Mark Boldin⁵, Karine Serre¹, Amy H. Buck⁴, Anita Q. Gomes^{1,6,*}, Bruno Silva-Santos^{1,*}

¹Instituto de Medicina Molecular, Faculdade de Medicina, Universidade de Lisboa, 1649-028 Lisboa, Portugal

²iBET, Instituto de Biologia Experimental e Tecnológica, 2780-157 Oeiras, Portugal

³Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, 2780-157 Oeiras, Portugal

⁴Institute of Immunology and Infection and Centre for Immunity, Infection and Evolution, University of Edinburgh, Edinburgh EH9 3FL, UK

⁵Department of Molecular and Cellular Biology, Beckman Research Institute, City of Hope, Duarte, CA 91010, USA

⁶Escola Superior de Tecnologia da Saúde de Lisboa, 1990-096 Lisboa, Portugal.

Abstract

$\gamma\delta$ T cells are major providers of proinflammatory cytokines. They are preprogrammed in the mouse thymus into distinct subsets producing either interleukin-17 (IL-17) or interferon- γ (IFN- γ), which segregate with CD27 expression. In the periphery, CD27⁻ $\gamma\delta$ ($\gamma\delta 27^{-}$) T cells can be induced under inflammatory conditions to coexpress IL-17 and IFN- γ ; the molecular basis of this functional plasticity remains to be determined. On the basis of differential microRNA (miRNA) expression analysis and modulation in $\gamma\delta$ T cell subsets, we identified miR-146a as a thymically imprinted post-transcriptional brake to limit IFN- γ expression in $\gamma\delta 27^{-}$ T cells in vitro and in vivo. On the basis of biochemical purification of Argonaute 2-bound miR-146a targets, we identified *Nod1* to be a relevant mRNA target that regulates $\gamma\delta$ T cell plasticity. In line with this, *Nod1*-deficient mice lacked multifunctional IL-17⁺ IFN- γ ⁺ $\gamma\delta 27^{-}$ cells and were more susceptible

The Authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original U.S. Government Works

*Corresponding author. nina.schmolka@uzh.ch (N.S.); anitagomes@medicina.ulisboa.pt (A.Q.G.); bssantos@medicina.ulisboa.pt (B.S.-S.).

[†]These authors contributed equally to this work.

[‡]These authors contributed equally to this work.

Author contributions: N.S., A.Q.G., and B.S.-S. designed the research and wrote the paper; N.S., P.H.P., P.V.R., T.A., F.J.E., A.A., K.E.G., K.S., and A.Q.G. performed experiments and analyzed the data; N.S. carried out the statistical analyses of the data; A.F.R., M.B., A.S.C., and A.H.B. provided key research tools; B.S.-S. and A.Q.G. supervised the research.

Competing interests: The authors declare that they have no competing interests.

Data and materials availability: miRNA array data and Ago2 RIP-seq data have been deposited in GEO database with the accession codes GSE110565 and GSE109263, respectively.

to *Listeria monocytogenes* infection. Our studies establish the miR-146a/NOD1 axis as a key determinant of $\gamma\delta$ T cell effector functions and plasticity.

INTRODUCTION

The proinflammatory cytokines interferon- γ (IFN- γ) and interleukin-17A (IL-17) are critical mediators of T cell responses against intracellular bacteria and viruses, or extracellular bacteria and fungi, as illustrated by the high susceptibility of patients with mutations in the IFN- γ or IL-17 pathways to *Mycobacterium tuberculosis* or *Candida albicans* infections, respectively (1). In addition to their protective roles in host defense, it has been shown that deregulated IFN- γ or/ and IL-17 production promotes chronic inflammation and autoimmunity. Thus, exacerbated IL-17 responses underlie inflammatory bowel disease, psoriasis, ankylosing spondylitis, and multiple sclerosis (1); and excessive IFN- γ is linked to type 1 diabetes and Crohn's disease (2).

Although IFN- γ and IL-17 can derive from multiple cellular sources, there is often a major contribution from $\gamma\delta$ T cells at early stages of immune responses (3). Namely, IFN- γ -producing $\gamma\delta$ T cells play protective roles in viral, parasitic, and intracellular bacterial infections, such as *Listeria monocytogenes* (4, 5). On the other hand, IL-17-producing $\gamma\delta$ T cells promote neutrophil mobilization and host defense against *Staphylococcus aureus*, *Escherichia coli*, and *C. albicans*, among other infections (3). IL-17-producing $\gamma\delta$ T cells have gathered much recent attention because of their impact on animal models of inflammatory and autoimmune disorders such as psoriasis, colitis, arthritis, or multiple sclerosis (3).

The prompt and abundant production of IFN- γ and IL-17 by $\gamma\delta$ T cells is associated with their "developmental preprogramming" in the thymus (3, 6–9). Thus, we and others have shown that mouse $\gamma\delta$ thymocytes can acquire the capacity to produce IFN- γ or IL-17, which associates with the expression of CD27 and CCR6, respectively (7, 10). Our study on chromatin-based and transcriptional mechanisms governing $\gamma\delta$ T cell differentiation revealed that CD27⁺ $\gamma\delta$ ($\gamma\delta$ 27⁺) T cells are committed to the expression of IFN- γ (but not of IL-17), whereas CD27⁻ $\gamma\delta$ ($\gamma\delta$ 27⁻) T cells exhibit a plastic behavior and can differentiate into cells that produce both IL-17 and IFN- γ (11). *Irf1* and its transcriptional regulator *Tbx21* displayed abundant active histone H3 marks (methylation of lysine K4 and acetylation) and were transcribed in both $\gamma\delta$ T cell subsets. The coexpression of IFN- γ and IL-17 proteins by $\gamma\delta$ 27⁻ T cells was observed under strong inflammatory conditions seemingly driven by IL-1 β and IL-23. The double-producing IL-17⁺ IFN- γ ⁺ $\gamma\delta$ 27⁻ T cells have been documented in animal models of cancer (11), autoimmunity (12), and infection (5) and also in HIV-1-infected patients (13).

However, $\gamma\delta$ 27⁻ T cell plasticity seems limited when compared with that of CD4⁺ T helper 17 (T_H17) cells, which rapidly acquire the capacity to make IFN- γ when exposed to IL-12 or IL-23 in vitro (14–16) and have been found in colitis, Crohn's disease, arthritis, diabetes, multiple sclerosis, and experimental autoimmune encephalomyelitis (EAE) (14, 15). Furthermore, analysis of an IL-17 fate-mapping reporter mouse showed that, whereas CD4⁺ T cells that had expressed IL-17 promptly converted into IFN- γ producers in the spinal cord

during EAE development, IL-17⁺ $\gamma\delta$ T cells were much more resistant to the acquisition of IFN- γ production (14). Because the *Ifny* locus displays active histone H3 marks and active transcription in $\gamma\delta 27^{-}$ T cells (11), we hypothesized that microRNAs (miRNAs) might control IFN- γ expression at the post-transcriptional level in $\gamma\delta 27^{-}$ T cells.

miRNAs drive their mRNA targets into the RNA-inducing silencing complex (RISC), in which Argonaute (AGO) proteins play a central role in preventing protein expression. Several miRNAs have been shown to affect the differentiation of cytokine-producing CD4⁺ T cell subsets [reviewed in 16)]. To date, only two miRNAs have been characterized in $\gamma\delta$ T cells, miR-133b and miR-206, which are co-regulated with IL-17 but have no impact on cytokine production (17). Here, we have identified miR-146a as highly expressed in $\gamma\delta 27^{-}$ T cells compared with their $\gamma\delta 27^{+}$ T cell counterparts, and through combined gain-of-function and loss-of-function approaches, we demonstrate that miR-146a inhibits IFN- γ production in $\gamma\delta$ T cells and thus restricts the functional plasticity of $\gamma\delta 27^{-}$ T cells in vitro and in vivo.

RESULTS

miR-146a is highly expressed in $\gamma\delta 27^{-}$ T cells

To address the role of miRNAs in the differentiation of proinflammatory $\gamma\delta$ T cell subsets, we compared the ex vivo miRNA expression profile of highly purified CD27⁺ ($\gamma\delta 27^{+}$) and CD27⁻ CCR6⁺ ($\gamma\delta 27^{-}$) $\gamma\delta$ T cells isolated from pooled lymph nodes and spleen of naïve C57BL/6J mice. We identified 35 miRNAs differentially expressed (more than twofold) between $\gamma\delta 27^{+}$ and $\gamma\delta 27^{-}$ CCR6⁺ cells (Fig. 1A). Only two related miRNAs, miR-146a and miR-146b, were expressed at higher levels in $\gamma\delta 27^{-}$ CCR6⁺ T cells when compared with $\gamma\delta 27^{+}$ cells. The differential expression of miR-146a (about fourfold) was fully validated by quantitative reverse transcription polymerase chain reaction (RT-qPCR) analysis in peripheral $\gamma\delta 27^{+}$ and $\gamma\delta 27^{-}$ T cell samples (Fig. 1B). In contrast to miR-146a, qPCR for miR-146b revealed very low levels of expression in all T cell subsets analyzed, including $\gamma\delta 27^{-}$ T cells (Fig. 1B and fig. S1A).

The analysis of thymocyte subsets allowed us to establish that miR-146a is already differentially expressed during $\gamma\delta$ T cell development, with a sharp increase as precursors mature into $\gamma\delta 27^{-}$ (but not $\gamma\delta 27^{+}$) thymocytes (Fig. 1C). Furthermore, miR-146a expression was not modulated by exogenous T cell receptor (TCR) stimulation or inflammatory cytokines (fig. S1B). These data demonstrate that miR-146a expression is tightly regulated during $\gamma\delta$ T cell differentiation, being confined to the subset that is preprogrammed in the thymus to produce IL-17 but no IFN- γ (7–9).

miR-146a inhibits IFN- γ production in $\gamma\delta$ T cells

To gain insight into the function of miR-146a in effector $\gamma\delta$ T cell differentiation, we performed gain-of-function studies either by retroviral overexpression of a miR-146a construct or by electroporation of miR-146a mimics in $\gamma\delta$ T cells. Both strategies require T cell activation and therefore act on fully differentiated $\gamma\delta$ T cells obtained from spleen and lymph nodes. Moreover, TCR/IL-2 stimulation is known to enhance IFN- γ production in $\gamma\delta$

T cells (18). Retroviral transduction of the native stem loop of miR-146a in $\gamma\delta$ T cells (Fig. 2A) led to marked overexpression of mature miR-146a (fig. S1C), and this resulted in a significant reduction of the frequency of IFN- γ^+ $\gamma\delta$ T cells compared with either an unrelated miR or a control vector alone (Fig. 2, B and C), without affecting $\gamma\delta$ T cell proliferation, as evaluated by Ki-67 staining (Fig. 2D). These data were validated by an alternative gain-of-function approach based on the electroporation of synthetic miR-146a oligonucleotides (miR-146a mimics) in $\gamma\delta$ T cells (Fig. 2E), which significantly decreased the frequency of IFN- γ^+ $\gamma\delta$ T cells when compared with control oligonucleotides (Fig. 2, F and G). Overexpression of miR-146a did not affect T_H1 differentiation of CD4⁺ T cells (fig. S1, D and E). These in vitro data suggest that miR-146a inhibits IFN- γ production in $\gamma\delta$ T cells, leading us to hypothesize that the high miR-146a levels in $\gamma\delta 27^-$ T cells in vivo (Fig. 1, B and C) restrict their capacity to express IFN- γ .

miR-146a limits IFN- γ production and functional plasticity of $\gamma\delta 27^-$ T cells

To explore the role of miR-146a in $\gamma\delta$ T cell differentiation, we studied $\gamma\delta$ T cells in miR-146a-deficient mice (19). In vitro, $\gamma\delta$ T cell proliferation upon TCR/CD28 stimulation was not affected by the absence of miR-146a (fig. S2, A and B). The numbers of steady-state $\gamma\delta 27^+$ and $\gamma\delta 27^-$ T cells, as well as their production of IFN- γ or IL-17, respectively, were similar between miR-146a-deficient and wild-type (WT) littermate control mice (fig. S2, C and D). We assessed the functional plasticity of $\gamma\delta 27^-$ T cells, namely, their capacity to differentiate into IL-17⁺ IFN- γ^+ cells upon IL-1 β + IL-23 stimulation (fig. S3A). We observed a significantly increased frequency of double producers among miR-146a-deficient (miR-146a^{-/-}) $\gamma\delta 27^-$ T cells when compared with WT (miR-146a^{+/+}) controls (Fig. 3A). As expected (11), TCR/CD28 activation (without IL-1 β or IL-23) failed to generate double producers (Fig. 3A). Neither T_H1 differentiation of CD4⁺ T cells (fig. S3, B and C) nor IFN- γ production by $\gamma\delta 27^+$ T cells (fig. S3, D and E) was affected by the loss of miR-146a. This is consistent with the very low abundance of miR-146a in $\gamma\delta 27^+$ T cells and thus ascribes a miR-146a function that is specific to the high expressing $\gamma\delta 27^-$ T cell subset.

To test whether miR-146a regulates IFN- γ production in $\gamma\delta 27^-$ T cells in a cell-intrinsic manner, we generated (20) mixed (1:1) bone marrow (BM) and mixed (1:1) neonatal thymus chimeras by transferring of BM and neonatal thymocytes from miR-146a^{-/-} mice (CD45.2⁺), together with BM (CD45.1⁺) and neonatal thymocytes (CD45.1/2⁺) from WT mice into recombination activation gene 2 (RAG2)-deficient hosts (Fig. 3B and fig. S4A). The neonatal thymocytes are a prerequisite to generate IL-17-producing $\gamma\delta$ T cells, because they are exclusively generated during fetal and perinatal life (21). This system allows the direct comparison of the progeny of miR-146a^{-/-} versus miR-146a^{+/+} hematopoietic cells developing in the same host (miR-146a^{+/+} in the nonhematopoietic compartment). $\gamma\delta 27^-$ T cells were isolated and activated with IL-1 β plus IL-23 to drive the differentiation of IL-17⁺ IFN- γ^+ double producers. We found that $\gamma\delta 27^-$ T cells derived from miR-146a^{-/-} were more prone to differentiate into double producers and expressed higher levels of IFN- γ than their WT counterparts (Fig. 3C). We also confirmed that the absence of miR-146a did not affect IFN- γ production by $\gamma\delta 27^+$ T cells (fig. S4B). Together, these results indicate that miR-146a acts as a cell-intrinsic brake to IFN- γ production specifically in $\gamma\delta 27^-$ T cells, thereby limiting their functional plasticity.

miR-146a restricts functional plasticity of $\gamma\delta 27^-$ T cells in *Listeria* infection

To validate miR-146a function in vivo, we analyzed miR-146a^{-/-} mice upon oral infection with *L. monocytogenes*, where a sizeable population of IL-17⁺ IFN- γ ⁺ $\gamma\delta 27^-$ T cells had been previously reported (5). We infected miR-146a^{-/-} and miR-146a^{+/+} (WT) littermate controls with *L. monocytogenes* and analyzed the $\gamma\delta$ T cell response at day 7 post infection. $\gamma\delta 27^-$ T cells increased their numbers by about five-fold in both WT and miR-146a^{-/-} mice (fig. S5A). However, the frequency and absolute number of IL-17⁺ IFN- γ ⁺ $\gamma\delta 27^-$ T cells were significantly higher in miR-146a^{-/-} when compared with WT mice (Fig. 4, A and B). This phenotype was largely accounted by V $\gamma 6^+$ T cells (Fig. 4C), which are known to be the main $\gamma\delta$ T cell population responding to *L. monocytogenes* infection (5). The frequency of single IFN- γ^- or single IL-17-producing $\gamma\delta$ T cells was the same in miR-146a^{-/-} when compared to WT mice (fig. S5B). By using this infection model in the mixed BM and neonatal thymocyte chimeras introduced above (Fig. 3B), we found that the IL-17⁺ IFN- γ ⁺ $\gamma\delta 27^-$ T cells present in mesenteric lymph nodes were mostly from miR-146a^{-/-} origin (Fig. 4D). As before, this phenotype was not because of a proliferative advantage of miR-146a^{-/-} compared with WT-derived $\gamma\delta$ T cells (fig. S5, C and D). These data demonstrate that $\gamma\delta$ T cell plasticity under pathophysiological conditions in vivo is regulated by miR-146a.

miR-146a targets Nod1 mRNA that is depleted in $\gamma\delta 27^-$ T cells

To unravel the molecular mechanism(s) underlying miR-146a function, namely, the relevant mRNA target(s), we undertook an unbiased approach based on differential Argonaute 2 (Ago2) RNA immunoprecipitation followed by deep sequencing (Ago2 RIP-seq). Ago2 is a crucial protein that engages with both miRNA and its mRNA target(s) in the RISC. To overcome the problem of lack of commercially available high affinity antibodies against Ago2, we took advantage of a genetically modified mouse in which endogenous Ago2 is replaced with a N-terminal-tagged version of the protein (3 $\times$ FLAG-6 $\times$ HIS), thus allowing stringent purification (i.e., with low unspecific background) of the RNA molecules bound to Ago2 [upon ultraviolet (UV) cross-linking], followed by RNA sequencing (RNA-seq) (fig. S6, A and C). To enrich for miR-146a targets, we transduced purified CD3⁺ T cells (we could not use $\gamma\delta$ T cells because of insufficient cell numbers for this approach) with a miR-146a-expressing retrovirus (RV) (~12-fold overexpression of miR-146a, as shown in fig. S1C). We thereby identified 225 putative mRNAs after differential Ago2 RIP-seq [data submitted to Gene Expression Omnibus (GEO)], from which 96 (42.7%) had a predicted miR-146a binding site in their 3' untranslated region (3' UTR) (fig. S6B). This included two previously validated miR-146a targets, *Zranb2* and *Gtf2e2* [included in DIANA- TarBase v7.0 (22)]. Many of the other putative target transcripts had not been described in $\gamma\delta$ T cells (table S1), and these data thus constitute a useful resource for future studies. For this study, to narrow down the potential mRNA targets of miR-146a involved in the regulation of IFN- γ production in $\gamma\delta$ T cells, we focused on genes with transcription factor or receptor activity (Table 1) that had been associated with the IFN- γ pathway in other immune cells and, in particular, on two candidate genes, *Atf2* (2.5-fold enriched in the RV-miR-146a sample), a transcription factor that binds to the IFN- γ promoter (23), and *Nod1*, an intracellular pattern recognition receptor recently shown to enhance IFN- γ production in $\alpha\beta$ T cells (24, 25), which showed the highest fold difference (4.2-fold) among genes with receptor activity.

Both putative targets have predicted miR-146a binding sites with canonical and noncanonical Watson-Crick base pairing (Fig. 5A). To examine direct miR-146a targeting of *Atf2* and *Nod1*, we designed reporter constructs in a pmirGLO Dual-luciferase miRNA target expression vector for the 3'UTRs of *Atf2* and *Nod1*. We additionally included a negative control vector containing a 3'UTR without miR-146a binding sites and a positive control containing the 3'UTR of *Notch1*, a previously validated miR-146a target (26, 27). We transiently transfected these constructs into human embryonic kidney (HEK) 293 T cells together with an expression plasmid for either miR-146a, an unrelated miRNA, or green fluorescent protein (GFP) control. Cotransfection of miR-146a (but not an unrelated miRNA) showed a significant repression of luciferase activity compared with GFP control for both mRNA targets (Fig. 5B). Mutations in the miR-146a binding sites of both *Atf2* and *Nod1* led to significant recoveries of luciferase levels (Fig. 5, A and B). We undertook various complementary mutational strategies to validate these findings, from mutating 3 to 11 individual base pairs (bp) to deleting the full binding region, always with consistent results (Fig. 5, A and B). Together, these data demonstrate that miR-146a can target *Atf2* and *Nod1*.

To investigate the potential impact of miR-146a on those mRNAs in $\gamma\delta$ T cells, we analyzed their expression levels in $\gamma\delta 27^+$ and $\gamma\delta 27^-$ T cell subsets. Whereas *Atf2* was similarly expressed in $\gamma\delta 27^+$, $\gamma\delta 27^-CCR6^-$, and $\gamma\delta 27^-CCR6^+$ T cells (Fig. 5C), *Nod1* expression (Fig. 5C) showed a notable negative association with miR-146a levels (Fig. 1, A and B). Namely, *Nod1* mRNA was present in $\gamma\delta 27^+$ but essentially absent in $\gamma\delta 27^-CCR6^+$ T cells (Fig. 5C), thus suggesting that *Nod1* is targeted by miR-146a specifically in this $\gamma\delta$ T cell subset. In addition, *Nod1* showed the lowest expression levels in $V\gamma 1^-V\gamma 4^-$ (mostly $V\gamma 6^+$) $\gamma\delta$ T cells (Fig. 5D). We further extended our expression analysis to ex vivo isolated $V\gamma 4^+$ and $V\gamma 1^-V\gamma 4^-$ $\gamma\delta 27^-$ T cells and observed notable increases in *Nod1* (but not *Atf2*) levels in both miR-146a^{-/-} $\gamma\delta 27^-$ T cell subsets when compared with miR-146a^{+/+} counterparts (Fig. 5E). In sum, these results strongly indicate that *Nod1* is a bona fide target of miR-146a in $\gamma\delta 27^-$ T cells. Although we were unable to perform luciferase assays in primary murine $\gamma\delta 27^-$ T cells, we successfully validated *Nod1* targeting by miR-146a in a human $\gamma\delta$ T cell line (fig. S7) that we previously showed to be a suitable model to study human $\gamma\delta$ T cell development (28).

Nod1 is required for IFN- γ production and functional plasticity of $\gamma\delta$ T cells

A key parameter in miRNA research is the demonstration of inverse phenotypes between an miRNA and its mRNA target. We therefore analyzed the production of IFN- γ (and IL-17) in $\gamma\delta$ T cells from *Nod1*^{-/-} mice (as well as *Atf2*^{-/-} mice as expected “negative controls”). Critically, *Nod1*^{-/-} (but not *Atf2*^{-/-}) $\gamma\delta 27^-$ T cells showed a notable inability to differentiate into IL-17⁺ IFN- γ ⁺ cells when compared with littermate controls (Fig. 6A). Again, this was not due to proliferative defects (Fig. 6A). The loss of *Nod1* did not impair IL-17 production by $\gamma\delta 27^-$ T cells (Fig. 6A).

To functionally test the relevance of *Nod1* as the target of miR-146a, we performed a genetic rescue experiment by crossing miR-146a^{-/-} with *Nod1*^{-/-} mice and analyzing littermate controls with various genotypes. We found that reducing *Nod1* (miR-146a^{-/-} *Nod1*^{+/+} mice)

prevented the accumulation of double producers among $\gamma\delta 27^-$ T cells observed in miR-146^{-/-} Nod1^{+/+} mice (relative to miR-146^{+/+} Nod1^{+/+} controls) (Fig. 6B), while not affecting $\gamma\delta 27^+$ T cell differentiation (fig. S8). These data strongly suggest that Nod1 is a dominant target of miR-146a in regulating $\gamma\delta 27^-$ T cell plasticity.

Last, to address the physiological relevance of Nod1 in vivo, we performed *L. monocytogenes* infection. Nod1^{-/-} mice were highly susceptible to *Listeria* infection, as illustrated by increased bacterial burden, severe weight loss, and death of Nod1^{-/-} (compared with both Nod1^{+/+} and Nod1^{+/-}) mice (Fig. 6C). This was associated with strikingly reduced IL-17⁺ IFN- γ ⁺ $\gamma\delta 27^-$ T cells in Nod1^{-/-} $\gamma\delta$ T cells (Fig. 6, D and E). TCRd^{-/-} mice were extremely susceptible to *Listeria* infection even more than Nod1^{-/-} mice (Fig. 6C). Together, these data establish a critical function of Nod1 in promoting IFN- γ production and functional plasticity in $\gamma\delta 27^-$ T cells, which is inverse to the negative role of miR-146a in this process, and thus support the miR-146a/ Nod1 axis as a critical regulator of $\gamma\delta$ T cell differentiation in vivo.

DISCUSSION

miRNAs are endowed with the capacity to regulate various molecular pathways of differentiation and functions in innate and adaptive immunity (29, 30). In T cells, miRNA repertoires are known to change dramatically upon activation and across distinct functional T cell subsets (31, 32). Building on our previous definition of proinflammatory $\gamma\delta$ T cell subsets (7, 11), we show here that miR-146a is highly expressed selectively on $\gamma\delta 27^-$ T cells, where it targets Nod1 and thus limits IFN- γ production by this IL-17–biased subset. The expression pattern of miR-146a was established during thymic $\gamma\delta$ T cell development and was unaffected by exogenous TCR stimulation or inflammatory cytokine signals, further illustrating the importance of developmental preprogramming in $\gamma\delta$ T cell biology (8, 9). In the periphery, inflammatory signals are major drivers of functional plasticity of $\gamma\delta 27^-$ T cells, which manifests itself in the acquisition of IFN- γ coproduction by previous IL-17 single producers. We found that the genetic loss of miR-146a led to the accumulation of multifunctional IL-17⁺ IFN- γ ⁺ double producers in vitro and in vivo, upon *Listeria* infection, where these cells had been associated with a protective memory response against the bacteria (5).

miR-146a has been previously associated with regulation of $\alpha\beta$ T cell responses via dampening of nuclear factor κ B (NF- κ B) activation (33, 34). In the absence of miR-146a, enhanced TCR-induced NF- κ B activation was linked to higher proliferation and increased production of effector cytokines including not only IFN- γ but also IL-17A and IL-2 by CD4⁺ and CD8⁺ T cells—that is, a general activation phenotype (33). However, more recently, miR-146a was shown to inhibit the production of autocrine IL-6 and IL-21 in autoreactive transgenic CD4⁺ T cells, which rather reduced their T_H17 differentiation (35). By contrast, our results show normal proliferation and IL-17 production by miR-146a^{-/-} $\gamma\delta$ T cells, thus highlighting the selective impact on IFN- γ coproduction by $\gamma\delta 27^-$ T cells, which may be linked to the high levels of expression of miR-146a in this subset. As a caveat of our study, we did not have the technical tools to specifically ablate miR-146a in $\gamma\delta$ T cells, so we had to rely on full miR-146a “knockout” mice. No miRNA has yet been shown

to play a role in T_H17/T_H1 plasticity. Looking globally at $CD4^+$ T cell plasticity, miR-10a is thus far an isolated example of an miRNA implicated in limiting “lineage conversion”; being highly expressed in regulatory T cells, miR-10a prevents the acquisition of T_H17 and follicular helper T cell features (36).

In contrast with the conventional targets (Traf6, Irak1, and Stat1, among others) of miR-146a in $CD4^+$ T cells (33, 34), the key mRNA target we identified for miR-146a in $\gamma\delta$ T cells was *Nod1*. This target was identified by an unbiased Ago2-RIP approach in Ago2-tagged T cells after miR-146a overexpression. Although overexpression or deletion of a given miRNA gene regulates distinct sets of target genes (37), we favored an overexpression approach because, as each mRNA is targeted by multiple miRNAs, critical mRNA targets of miR-146a could be unaffected in RISCs of miR-146a^{-/-} T cells. Instead, by providing an excess of miR-146a, we increased the possibility of dragging its mRNA targets into RISCs. Moreover, because miR-146a is a low expressed miRNA, selectively enriched in a minor ($\gamma\delta27^-$) T cell subset, using an overexpression system increases the signals attributable to this low abundant miRNA. The genome-wide Ago2 RIP-seq analysis allowed us to identify many potential previously unknown participants in the differentiation and activation of T cells. Here, we focused our analysis on *Nod1*, and our data suggest that miR-146a targets *Nod1* not by canonical seed sequence-based pairing but by an alternative pathway already shown for other miRNAs [reviewed in (38)], such as miR-155 that targets more than 40% of its mRNAs by noncanonical interactions (39). This type of binding is very difficult to predict using standard strategies as those used by softwares like Targetscan (40). Thus, in this analysis, we modified the standard parameters used by miRWalk (41) by allowing a minimum seed binding of five nucleotides and further refined the results by calculating the Gibbs' function of the miRNA:mRNA hybrid. Because of 3' compensatory binding, the results obtained for the selected miR-146a-5p target at the 3'UTR of *Nod1* ($\Delta G = -22.5$ kcal/mol) are consistent with a stable miR-146:*Nod1* hybrid.

Nod1 is a crucial pattern recognition receptor initially thought to be restricted to innate myeloid cells. However, *Nod1* has been shown to be expressed by $CD8^+$ T cells, at comparable levels to macrophages, and to functionally coactivate T cell proliferation and IFN- γ production (24). We show here that $\gamma\delta$ T cells, particularly the $\gamma\delta27^+$ subset that makes IFN- γ , express *Nod1*. This is consistent with recent data showing that *Nod1* ligation increases IFN- γ production by isolated $CD8^+$ and $CD4^+$ T cells in vitro, and *Nod1*-deficient T cells exhibited impaired IFN- γ production upon dextran sulfate sodium-induced acute inflammation and tumorigenesis in vivo (25). The other prototypic *Nod*-like receptor, *Nod2*, was also suggested to play a T cell-intrinsic role in IFN- γ production, which affected *Toxoplasma gondii* clearance (42).

As confirmed by our data, *Nod1* is a major determinant of resistance to *L. monocytogenes* infection (43). The same applies to $\gamma\delta$ T cells, but we could not directly link the two phenotypes because of lack of genetic tools to conditionally delete *Nod1* specifically in $\gamma\delta$ T cells. On the other hand, although in the infection setting *Nod1* may be recognizing bacterial ligands such as peptidoglycan (44), the inability of *Nod1*-deficient $\gamma\delta27^-$ T cells to differentiate into IFN- γ producers in vitro suggests an additional mechanism of *Nod1* activation. As another limitation of our study, we have dissected neither this upstream

mechanism nor the downstream regulators of cytokine production. Nod1 (and Nod2) are important mediators of endoplasmic reticulum (ER) stress-induced production of proinflammatory cytokines (45). Future research should therefore address whether ER stress and the underlying unfolded protein response, known to be important drivers of inflammation and also to be induced upon T cell activation (46), play a role in Nod1-dependent $\gamma\delta$ T cell differentiation.

In sum, our data demonstrate that high levels of miR-146a in $\gamma\delta 27^-$ T cells suppress Nod1 expression, which is rescued upon ablation of miR-146a, thus promoting IFN- γ expression in thymically precommitted IL-17 producers. Multifunctional IL-17⁺ IFN- γ ⁺ $\gamma\delta$ T cells have been associated with pathogenic $\gamma\delta$ T cell responses in experimental models of (ovarian) cancer (11) and autoimmunity (EAE) (12), and thus, the manipulation of miR-146a levels may have therapeutic potential.

MATERIALS AND METHODS

Mice

All mice used were adults 6 to 12 weeks of age. C57BL/6J and CD45.1 mice were from the Jackson Laboratory, and Rag2^{-/-} mice were from Taconic Biosciences. *III17a*-GFP knock-in mice were from Biocytogen, LLC. miR-146a^{-/-} mice have been described previously (19). Nod1^{-/-} and Atf2^{-/-} mice were obtained from T. Mak (University Health Network, Toronto) and W. Breitwieser (Cancer Research UK Manchester Institute), respectively. The Ago2-tagged knock-in mice (EIF2c2-tag-KI) were generated by A.H.B.'s laboratory in conjunction with Biocytogen. A targeting vector was designed to insert a 147-bp N-terminal tag into the first exon of the Eif2c2 gene downstream of the ATG start site. The tag consists of 3×FLAG tag followed by a PreScission cleavage site and 6×HIS tag (sequence information available by request). A neomycin-selectable cassette was flanked by FLPe sites and inserted into the nonconserved region of intron 1 and targeted embryonic stem (ES) cells generated by homologous recombination. Selected ES cells were microinjected into C57BL/6 blastocysts and embryo transfer to pseudopregnant females. Chimeras were identified using standard procedures. Germline transmission was confirmed by Southern blotting and breeding program set up to establish the line. Both heterozygous and homozygous mice carrying the FLAG-HIS-tagged Ago2 are viable and fertile.

Mice were bred and maintained in the specific pathogen-free animal facilities of Instituto de Medicina Molecular (Lisbon, Portugal). All experiments involving animals were done in compliance with the relevant laws and institutional guidelines and were approved by the ethics committee of Instituto de Medicina Molecular.

Monoclonal antibodies

The following anti-mouse fluorescently labeled monoclonal antibodies (mAbs) were used (antigens and clones): TCR δ (GL3), V γ 1 (2.11), V γ 4 (UC3-10A6), CD3 (145.2C11), CD4 (GK1.5), CD8 (53-6.7), CD25 (PC61), CD27 (LG.7F9), CD44 (IM7), CD45.1 (A20), CD45.2 (104), Ki-67 (16A8), IFN- γ (XMG1.2), and IL-17A (TC11.18H10.1). Antibodies were purchased from BD Biosciences, eBiosciences, or BioLegend.

Cell preparation, flow cytometry, and cell sorting

Cell suspensions were obtained from spleens, lymph nodes, or thymus. Erythrocytes were osmotically lysed in red blood cell lysis buffer (BioLegend). Cells were filtered through 70- μ m cell strainers (BD Biosciences). For cell surface staining, single-cell suspensions were incubated for 30 min with saturating concentrations of mAbs (see above). For intracellular cytokine staining, cells were stimulated with phorbol 12-myristate 13-acetate (50 ng/ml) and ionomycin (1 μ g/ml) in the presence of brefeldin A (10 μ g/ml) (all from Sigma- Aldrich) for 3 hours at 37°C. Cells were stained for the above identified cell surface markers, fixed 30 min at 4°C, permeabilized with the Foxp3/Transcription Factor Staining Buffer set (eBioscience) in the presence of anti-CD16/CD32 (eBioscience) for 15 min at 4°C, and lastly incubated for 1 hour at 4°C with the above identified antibodies in permeabilization buffer. Cells were sorted on FACSARIA (BD Biosciences), and samples were acquired using FACSFortessa (BD Biosciences). Data were analyzed using FlowJo software (Tree Star).

miRNA array

All experiments were conducted at Exiqon Services. The quality of the total RNA was verified by an Agilent 2100 Bioanalyzer profile. Total RNA (250 ng) from sample and reference was labeled with Hy3 and Hy5 fluorescent label, respectively, using the miRCURY LNA miRNA Hi-Power Labelling Kit, Hy3/Hy5 (Exiqon, Denmark), following the procedure described by the manufacturer. The Hy3- labeled samples and a Hy5- labeled reference RNA sample were mixed pairwise and hybridized to the miRCURY LNA miRNA Array (sixth generation: hsa, mmu, and rno) (Exiqon, Denmark), which contains capture probes targeting all miRNAs for human, mouse, or rat registered in the miRBASE 16.0. The hybridization was performed according to the miRCURY LNA miRNA Array instruction manual using a Tecan HS 4800 hybridization station (Tecan, Austria). After hybridization, the microarray slides were scanned and stored in an ozone-free environment (ozone level below 2.0 parts per billion) to prevent potential bleaching of the fluorescent dyes. The miRCURY LNA array microarray slides were scanned using the Agilent G2565BA Microarray Scanner System (Agilent Technologies Inc.), and the image analysis was carried out using the ImaGene 9.0 software (BioDiscovery Inc.). The quantified signals were background-corrected (Normexp with an offset value of 10) and normalized using the Quantile normalization algorithm.

RNA isolation, complementary DNA production, and real-time PCR

RNA was isolated from fluorescence-activated cell sorting (FACS)–sorted cell populations using a High Pure RNA Isolation kit (Roche) for total RNA and an miRNeasy Mini kit (Qiagen) or a High Pure miRNA Isolation kit (Roche) for small RNA. For total RNA, reverse transcription was performed with random oligonucleotides (Invitrogen) using Moloney murine leukemia virus reverse transcriptase (Promega). For miRNA, reverse transcription was performed with a Universal cDNA Synthesis kit II (Exiqon). For total RNA samples, relative quantification of specific complementary DNA (cDNA) species to endogenous references *HPRT* or *Actinb* was carried out using SYBR on a ViiA7 cycler (Applied Biosystems). Primers were either designed manually or by the Universal ProbeLibrary Assay Design Center (Roche), and their sequences were as follows: *HPRT*,

GACCTCTCGAAGTGTGGAT (forward) and GCTTTGTATTTGGCTTTTCC (reverse); *Actb*, CGTGAAAAGATGACCCAGATCA (forward) and TGGTACGACCAGAGGCATACAG (reverse); *Nod1*, CCTTCGTCCTGCATCACTTC (forward) and CTGGTTGACGCTGAGTCTGA (reverse); and *Atf2*, GAAGAGTCTCGCCCACAGTC (forward) and TGGGTCTGAGGAGTTGTGTG (reverse). For miRNA samples, relative quantification of specific cDNA species to reference small RNA RNU5G or miR-423-3p was carried out using SYBR on a ViiA7 cyclor (Applied Biosystems). The respective miRNA LNA primers were used (Exiqon).

Retroviral overexpression of miR-146a

The retroviral constructs encoding either mmu-miR-146a or an unrelated miRNA (mmu-miR-132) were generated by inserting the respective native pre-miRNA sequences flanked by about 200 bp into a modified pMig.IRES-GFP retroviral vector (Addgene #9044). miR-146a and miR-132 sequences were obtained by PCR from genomic DNA from C57BL/6J mice. The internal ribosomal entry site (IRES)-GFP sequence was removed and replaced by PGK-GFP-WPRE sequence. The resulting vector encodes GFP under the control of the phosphoglycerate kinase (PGK) promoter and miR-146a or miR-132 under control of the retroviral long terminal repeat promoter. A Woodchuck Hepatitis Virus Posttranscriptional Regulatory Element (WPRE) was inserted for increased viral titers. For retroviral transduction, $\gamma\delta$ T cells and CD4⁺ T cells were sorted from lymph node and spleen of C57BL/6J mice and stimulated either with plate-bound anti-CD3 ϵ (145.2C11) plus anti-CD28 mAb (37.51) (both 2.5 μ g/ml) or plate-bound anti-CD3 ϵ (145.2C11) (5 μ g/ml) in the presence of soluble anti-CD28 (37.51) (2 μ g/ml) with IL-12 (5 ng/ml) and anti-IL-4 (11B11) (10 μ g/ml), respectively. The transduction was performed with polybrene (4 μ g/ml; Sigma-Aldrich), and medium was supplemented with IL-2 (5 ng/ml). Forty-eight hours after transduction, live cells using a LIVE/DEAD stain kit (Life Technologies), and GFP⁺ cells were sorted and activated for intracellular staining.

$\gamma\delta$ T cell electroporation

$\gamma\delta$ T cells were sorted from lymph node and spleen of C57BL/6J mice and stimulated with plate-bound anti-CD3 and anti-CD28 (both 2.5 μ g/ml). $\gamma\delta$ T cells (2.5×10^4) were resuspended in T buffer and used per transfection with Neon electroporation transfection system (Invitrogen) with three pulses of 10 ms and 1550 V. miRIDIAN mimic of mouse mmu-miR-146a-5p (Dharmacon) and miRIDIAN mimic negative control #1 (Dharmacon) with an RNA concentration of 500 nM per electroporation were used. After electroporation, cells were cultured in media supplemented with IL-2 (5 ng/ml) and 48 hours later activated for intracellular cytokine staining.

In vitro $\gamma\delta$ T cell stimulation and CD4⁺ T_H cell polarization

$\gamma\delta 27^+$ and $\gamma\delta 27^-$ T cells were FACS-sorted and subjected to various stimulation conditions overnight or for 24 to 72 hours. Cells were incubated on plate-bound anti-CD3 ϵ (145.2C11) plus anti-CD28 mAb (37.51) (both at 2.5 μ g/ml) supplemented with IL-12 (10 ng/ml) or, alternatively, in the presence of murine IL-1 β plus IL-23 (both 50 ng/ml) or IL-12 (10 ng/ml). For T_H1 cell culture conditions, FACS-sorted CD4⁺ T cells were incubated with plate-bound anti-CD3 ϵ (145.2C11) and soluble anti-CD28 mAb (37.51) (both at 2 μ g/ml) in the

presence of IL-12 (5 ng/ml) and neutralizing anti-IL-4 mAb (11B11) (5 µg/ml) for 5 days. All cytokines were from PeproTech, except TGF- β and IL-23, which were from R&D systems.

***L. monocytogenes* infection and bacterial quantification**

We used *L. monocytogenes* strain EGDe, carrying a recombinant InlA with a mutation in S192N and Y369S, as previously described (5). Briefly, all mice were deprived of food overnight before infection, housed individually, and given a ~0.5-cm³ piece of mashed food inoculated with 2×10^9 colony-forming units (CFUs) of *L. monocytogenes* in phosphate-buffered saline. CFUs were calculated from single-cell suspensions of spleens plated in serial dilutions on Brain Heart Infusion agar plates. Individual colonies were counted after 30 hours of incubation at 37°C.

Neonatal thymocytes and BM chimeras

Chimeras were generated as previously described (20). Briefly, Rag2^{-/-} mice were lethally irradiated (9.5 gray) and injected 6 hours after with a 1:1 mix of neonatal thymocytes (5×10^6 to 10×10^6 thymocytes per host) isolated from WT (CD45.1/CD45.2) and miR-146a^{-/-} (CD45.2) donors within 48 hours after birth. After 24 hours, the host received a 1:1 mix of 5×10^6 to 10×10^6 BM cells isolated from WT (CD45.1/CD45.2) and miR-146a^{-/-} (CD45.2) donors. Chimeras were kept on antibiotic-containing water (2% Bactrim; Roche) for the first 4 weeks after irradiation. The hematopoietic compartment was allowed to reconstitute for 8 weeks before the animals were used for experiments.

Differential Argonaute 2 RNA immunoprecipitation followed by deep sequencing

CD3⁺ T cells were FACS-sorted from pooled spleens of Ago2-tagged (3×FLAG-6×HIS tag) mice and stimulated with plate-bound anti-CD3 ϵ (145.2C11) plus anti-CD28 mAb (37.51) (both 2.5 µg/ml) and transduced either with RV-miR-146a vector or RV-control vector. Twenty-four hours after transduction, GFP⁺ live cells [using a live/dead stain kit (Life Technologies)] were FACS-sorted. Cells (7×10^5 to 2×10^6) were UV-cross-linked in Stratalinker 2400 (400 mJ cm⁻²) and lysed. AGO-interacting RNA was immunoprecipitated using anti-Flag (M2) magnetic beads (Sigma-Aldrich). Following, Ago2-bound RNA was partially digested using RNase IT (Agilent Technologies) and, in a second step, purified using the HIS-tag and Ni-nitrilotriacetic acid beads (Qiagen) under high stringent denaturing wash conditions. After proteinase K treatment, RNA was isolated using phenol-chloroform isoamyl extraction (detailed protocol available upon request). The quality of the RNA was verified by an Agilent 2100 Bioanalyzer profile, and libraries were prepared using a Clontech SMARTer Stranded RNA-Seq Kit, starting with 8 µl of the immunoprecipitated RNA. The manufacturer's solid phase reversible immobilization (SPRI) protocol was followed, using 4 min for RNA fragmentation, double bead cleanup after reverse transcription, and 20 PCR cycles for amplification. The resulting libraries were sequenced using Illumina HiSeq 2000 instrument with 50-bp single-read sequencing mode.

Ago2 RIP-seq data analysis

To assess overall RNA-seq data quality, sequence length distribution and the base calling accuracy as indicated by the Phred quality score (Q score) were calculated with high-throughput sequence data quality control software FastQC (version 0.10.1; Babraham Bioinformatics, UK). Adaptor sequences were trimmed using FASTX-toolkit. Sequence reads were mapped to the mouse reference genome (GRCm38) using Bowtie2 software (47). Aligned reads were thereafter sorted and indexed by SAMtools (48), Cuffdiff software, a part of the Cufflinks suite (49), was used to detect significant changes in mRNA enrichment between the experimental groups. Genes with a P value, for statistical significance, and q value, to detect the false discovery rate, of <0.05 were considered differentially enriched in Ago2-RIP experiments. Predicted miRNA targets for mmu-miR-146a were determined by miRWalk2 algorithm (41).

Dual luciferase reporter assay

The 3'UTR of Nod1, Atf2, Notch1 (positive control), and PTPBP2 (negative control) were cloned into pmirGLO vector (Promega) from genomic DNA. Primers are available upon request. Mutations in the predicted target sequences of the 3'UTRs were introduced by gene synthesis (GeneCust Europe). The luciferase reporter vectors carrying either the WT or mutant 3'UTR-sequence were transfected with the miR-146a expression vector, an unrelated miRNA (miR-132), or a control vector into HEK293 T cells (ATCC CRL-3216) using Lipofectamine 2000 (Thermo Fisher Scientific) or electroporated into PEER T-ALL cells [DSMZ (German Collection of Microorganisms and Cell Cultures), Braunschweig, Germany], following the manufacturer's protocol. After 48 hours, firefly and *Renilla* luciferase activity were measured by using the Dual-Glo Luciferase Assay System (Promega). *Renilla* luciferase activity served as the internal control, and results were expressed as relative luciferase activity.

Carboxyfluorescein diacetate succinimidyl ester staining

For analysis of proliferation, FACS-sorted CD4⁺ and $\gamma\delta$ T cells were stained by incubation for 5 min at 37°C with 5 mM CFSE (carboxyfluorescein diacetate succinimidyl ester; Molecular Probes). Reactions were quenched by washing with ice-cold RPMI medium supplemented with 10% (v/v) fetal calf serum. CFSE-labeled cells were stimulated with plate-bound anti-CD3 ϵ (145.2C11) plus anti-CD28 mAb (37.51) (both 2.5 μ g/ml). After 4 days, the proliferation was assessed by flow cytometry.

Bromo-deoxyuridine incorporation

Mice were given intraperitoneal injections of 1.5 mg of bromo-deoxyuridine (BrdU) on the day of *L. monocytogenes* infection, and BrdU at a concentration of 0.8 mg/ml was supplemented in the drinking water during the course of infection. Incorporation of BrdU was determined by a BrdU Flow kit (BD Pharmingen) following manufacturer's protocol.

Statistical analysis

The statistical significance of differences between populations was assessed using a two-tailed nonparametric Mann-Whitney test or Student's *t* test. $P < 0.05$ was considered significant and is indicated on the figures.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank F. Caiado, J. Ribot, S. Mensurado, N. Gonçalves-Sousa [Instituto de Medicina Molecular (iMM), Lisboa, Portugal], A. Helwak (Wellcome Centre for Cell Biology, Edinburgh, UK), and K. M. Ansel (University of California, San Francisco, USA) for discussions and advice; V. Benes and D. Pavlinic (European Molecular Biology Laboratory GeneCore facility) for assistance with library preparation for RNA-seq analyses; D. Denner for help in graphical output; and the staff of the Flow Cytometry and Rodent facilities of iMM Lisboa for valuable technical assistance. Ago2-tagged (3×FLAG-6×HIS tag) mice are available from A.H.B. under a material transfer agreement with the University of Edinburgh.

Funding: This work was funded by the European Research Council (CoG_646701) to B.S.-S., Fundação para a Ciência e Tecnologia (FCT; PTDC/BEX-BCM/3592/2014 to A.Q.G. and SFRH/BPD/108821/2015 to N.S.), Wellcome Trust Research Career Development Fellowships (WT097394A1A) to A.H.B., and NIH/National Institute of Allergy and Infectious Diseases and STOP Cancer Foundation to M.B. This publication was sponsored by LISBOA-01-0145-FEDER-007391, with the project cofunded by Fundo europeu de desenvolvimento regional (FEDER), through POR Lisboa 2020 (Programa Operacional Regional de Lisboa, PORTUGAL 2020) and FCT.

REFERENCES AND NOTES

1. Korn T, Bettelli E, Oukka M, Kuchroo VK, IL-17 and Th17 Cells. *Annu. Rev. Immunol.* 27, 485–517 (2009). [PubMed: 19132915]
2. Neurath MF, Weigmann B, Finotto S, Glickman J, Nieuwenhuis E, Iijima H, Mizoguchi A, Mizoguchi E, Mudter J, Galle PR, Bhan A, Autschbach F, Sullivan BM, Szabo SJ, Glimcher LH, Blumberg RS, The transcription factor T-bet regulates mucosal T cell activation in experimental colitis and Crohn's disease. *J. Exp. Med.* 195, 1129–1143 (2002). [PubMed: 11994418]
3. Papotto PH, Ribot JC, Silva-Santos B, IL-17⁺ $\gamma\delta$ T cells as kick-starters of inflammation. *Nat. Immunol.* 18, 604–611 (2017). [PubMed: 28518154]
4. Hiromatsu K, Yoshikai Y, Matsuzaki G, Ohga S, Muramori K, Matsumoto K, Bluestone JA, Nomoto K, A protective role of gamma/delta T cells in primary infection with *Listeria monocytogenes* in mice. *J. Exp. Med.* 175, 49–56 (1992). [PubMed: 1530961]
5. Sheridan BS, Romagnoli PA, Pham Q-M, Fu H-H, Alonzo F III, Schubert W-D, Freitag NE, Lefrançois L, $\gamma\delta$ T cells exhibit multifunctional and protective memory in intestinal tissues. *Immunity* 39, 184–195 (2013). [PubMed: 23890071]
6. Jensen KDC, Su X, Shin S, Li L, Youssef S, Yamasaki S, Steinman L, Saito T, Locksley RM, Davis MM, Baumgarth N, Chien Y.-h., Thymic selection determines $\gamma\delta$ T cell effector fate: Antigen-naïve cells make interleukin-17 and antigen-experienced cells make interferon γ . *Immunity* 29, 90–100 (2008). [PubMed: 18585064]
7. Ribot JC, deBarros A, Pang DJ, Neves JF, Peperzak V, Roberts SJ, Girardi M, Borst J, Hayday AC, Pennington DJ, Silva-Santos B, CD27 is a thymic determinant of the balance between interferon- γ - and interleukin 17-producing $\gamma\delta$ T cell subsets. *Nat. Immunol.* 10, 427–436 (2009). [PubMed: 19270712]
8. Hayday AC, $\gamma\delta$ T cells and the lymphoid stress-surveillance response. *Immunity* 31, 184–196 (2009). [PubMed: 19699170]
9. Muñoz-Ruiz M, Sumaria N, Pennington DJ, Silva-Santos B, Thymic determinants of $\gamma\delta$ T cell differentiation. *Trends Immunol.* 38, 336–344 (2017). [PubMed: 28285814]

10. Haas JD, González FHM, Schmitz S, Chennupati V, Föhse L, Kremmer E, Förster R, Prinz I, CCR6 and NK1.1 distinguish between IL-17A and IFN- γ -producing $\gamma\delta$ effector T cells. *Eur. J. Immunol.* 39, 3488–3497 (2009). [PubMed: 19830744]
11. Schmolka N, Serre K, Grosso AR, Rei M, Pennington DJ, Gomes AQ, Silva-Santos B, Epigenetic and transcriptional signatures of stable versus plastic differentiation of proinflammatory $\gamma\delta$ T cell subsets. *Nat. Immunol.* 14, 1093–1100 (2013). [PubMed: 23995235]
12. Reynolds JM, Martinez GJ, Chung Y, Dong C, Toll-like receptor 4 signaling in T cells promotes autoimmune inflammation. *Proc. Natl. Acad. Sci. U.S.A.* 109, 13064–13069 (2012). [PubMed: 22826216]
13. Fenoglio D, Poggi A, Catellani S, Battaglia F, Ferrera A, Setti M, Murdaca G, Zocchi MR, V δ 1 T lymphocytes producing IFN- γ and IL-17 are expanded in HIV-1-infected patients and respond to *Candida albicans*. *Blood* 113, 6611–6618 (2009). [PubMed: 19395673]
14. Hirota K, Duarte JH, Veldhoen M, Hornsby E, Li Y, Cua DJ, Ahlfors H, Wilhelm C, Tolaini M, Menzel U, Garefalaki A, Potocnik AJ, Stockinger B, Fate mapping of IL-17-producing T cells in inflammatory responses. *Nat. Immunol.* 12, 255–263 (2011). [PubMed: 21278737]
15. Lee YK, Turner H, Maynard CL, Oliver JR, Chen D, Elson CO, Weaver CT, Late developmental plasticity in the T helper 17 lineage. *Immunity* 30, 92–107 (2009). [PubMed: 19119024]
16. Amado T, Schmolka N, Metwally H, Silva-Santos B, Gomes AQ, Cross-regulation between cytokine and microRNA pathways in T cells. *Eur. J. Immunol.* 45, 1584–1595 (2015). [PubMed: 25865116]
17. Haas JD, Nistala K, Petermann F, Saran N, Chennupati V, Schmitz S, Korn T, Wedderburn LR, Forster R, Krueger A, Prinz I, Expression of miRNAs miR-133b and miR-206 in the *Il17a/f* locus is co-regulated with IL-17 production in $\alpha\beta$ and $\gamma\delta$ T cells. *PLOS ONE* 6, e20171 (2011). [PubMed: 21637854]
18. Barros-Martins J, Schmolka N, Fontinha D, Pires de Miranda M, Simas JP, Brok I, Ferreira C, Veldhoen M, Silva-Santos B, Serre K, Effector $\gamma\delta$ T cell differentiation relies on master but not auxiliary Th cell transcription factors. *J. Immunol.* 196, 3642–3652 (2016). [PubMed: 26994218]
19. Boldin MP, Taganov KD, Rao DS, Yang L, Zhao JL, Kalwani M, Garcia-Flores Y, Luong M, Devrekanli A, Xu J, Sun G, Tay J, Linsley PS, Baltimore D, *miR-146a* is a significant brake on autoimmunity, myeloproliferation, and cancer in mice. *J. Exp. Med.* 208, 1189–1201 (2011). [PubMed: 21555486]
20. Cai Y, Xue F, Fleming C, Yang J, Ding C, Ma Y, Liu M, Zhang H.-g., Zheng J, Xiong N, Yan J, Differential developmental requirement and peripheral regulation for dermal V γ 4 and V γ 6T17 cells in health and inflammation. *Nat. Commun.* 5, 3986 (2014). [PubMed: 24909159]
21. Haas JD, Ravens S, Düber S, Sandrock I, Oberdörfer L, Kashani E, Chennupati V, Föhse L, Naumann R, Weiss S, Krueger A, Förster R, Prinz I, Development of interleukin-17-producing $\gamma\delta$ T cells is restricted to a functional embryonic wave. *Immunity* 37, 48–59 (2012). [PubMed: 22770884]
22. Vlachos IS, Paraskevopoulou MD, Karagkouni D, Georgakilas G, Vergoulis T, Kanellos I, Anastasopoulos I-L, Maniou S, Karathanou K, Kalfakakou D, Fevgas A, Dalamagas T, Hatzigeorgiou AG, DIANA-TarBase v7.0: Indexing more than half a million experimentally supported miRNA:mRNA interactions. *Nucleic Acids Res.* 43, D153–D159 (2015). [PubMed: 25416803]
23. Jones B, Chen J, Inhibition of IFN- γ transcription by site-specific methylation during T helper cell development. *EMBO J.* 25, 2443–2452 (2006). [PubMed: 16724115]
24. Mercier BC, Ventre E, Fogeron M-L, Debaud A-L, Tomkowiak M, Marvel J, Bonnefoy N, NOD1 cooperates with TLR2 to enhance T cell receptor-mediated activation in CD8 T cells. *PLOS ONE* 7, e42170 (2012). [PubMed: 22848741]
25. Zhan Y, Seregin SS, Chen J, Chen GY, Nod1 limits colitis-associated tumorigenesis by regulating IFN- γ production. *J. Immunol.* 196, 5121–5129 (2016). [PubMed: 27183588]
26. He Y, Sun X, Huang C, Long X.-r., Lin X, Zhang L, Lv X.-w., Li J, MiR-146a regulates IL-6 production in lipopolysaccharide-induced RAW264.7 macrophage cells by inhibiting Notch1. *Inflammation* 37, 71–82 (2014). [PubMed: 23974950]

27. Mei J, Bachoo R, Zhang C-L, MicroRNA-146a inhibits glioma development by targeting Notch1. *Mol. Cell. Biol.* 31, 3584–3592 (2011). [PubMed: 21730286]
28. Ribeiro ST, Tesio M, Ribot JC, Macintyre E, Barata JT, Silva-Santos B, Casein kinase 2 controls the survival of normal thymic and leukemic $\gamma\delta$ T cells via promotion of AKT signaling. *Leukemia* 31, 1603–1610 (2017). [PubMed: 27899804]
29. Ebert MS, Sharp PA, Roles for microRNAs in conferring robustness to biological processes. *Cell* 149, 515–524 (2012). [PubMed: 22541426]
30. Bronevetsky Y, Ansel KM, Regulation of miRNA biogenesis and turnover in the immune system. *Immunol. Rev.* 253, 304–316 (2013). [PubMed: 23550654]
31. Monticelli S, Ansel KM, Xiao C, Socci ND, Krichevsky AM, Thai T-H, Rajewsky N, Marks DS, Sander C, Rajewsky K, Rao A, Kosik KS, MicroRNA profiling of the murine hematopoietic system. *Genome Biol.* 6, R71 (2005). [PubMed: 16086853]
32. Rossi RL, Rossetti G, Wenandy L, Curti S, Ripamonti A, Bonnal RJP, Birolo RS, Moro M, Crosti MC, Gruarin P, Maglie S, Marabita F, Mascheroni D, Parente V, Comelli M, Trabucchi E, De Francesco R, Geginat J, Abrignani S, Pagani M, Distinct microRNA signatures in human lymphocyte subsets and enforcement of the naive state in CD4⁺ T cells by the microRNA miR-125b. *Nat. Immunol.* 12, 796–803 (2011). [PubMed: 21706005]
33. Yang L, Boldin MP, Yu Y, Liu CS, Ea C-K, Ramakrishnan P, Taganov KD, Zhao JL, Baltimore D, *miR-146a* controls the resolution of T cell responses in mice. *J. Exp. Med.* 209, 1655–1670 (2012). [PubMed: 22891274]
34. Lu L-F, Boldin MP, Chaudhry A, Lin L-L, Taganov KD, Hanada T, Yoshimura A, Baltimore D, Rudensky AY, Function of miR-146a in controlling Treg cell-mediated regulation of Th1 responses. *Cell* 142, 914–929 (2010). [PubMed: 20850013]
35. Li B, Wang X, Choi IY, Wang Y-C, Liu S, Pham AT, Moon H, Smith DJ, Rao DS, Boldin MP, Yang L, miR-146a modulates autoreactive Th17 cell differentiation and regulates organ-specific autoimmunity. *J. Clin. Invest.* 127, 3702–3716 (2017). [PubMed: 28872459]
36. Takahashi H, Kanno T, Nakayamada S, Hirahara K, Sciumè G, Muljo SA, Kuchen S, Casellas R, Wei L, Kanno Y, O'Shea JJ, TGF- β and retinoic acid induce the microRNA miR-10a, which targets Bcl-6 and constrains the plasticity of helper T cells. *Nat. Immunol.* 13, 587–595 (2012). [PubMed: 22544395]
37. Jin HY, Oda H, Chen P, Yang C, Zhou X, Kang SG, Valentine E, Kefauver JM, Liao L, Zhang Y, Gonzalez-Martin A, Shepherd J, Morgan GJ, Mondala TS, Head SR, Kim P-H, Xiao N, Fu G, Liu W-H, Han J, Williamson JR, Xiao C, Differential sensitivity of target genes to translational repression by miR-17~92. *PLOS Genet.* 13, e1006623 (2017). [PubMed: 28241004]
38. Cloonan N, Re-thinking miRNA-mRNA interactions: Intertwining issues confound target discovery. *Bioessays* 37, 379–388 (2015). [PubMed: 25683051]
39. Loeb GB, Khan AA, Canner D, Hiatt JB, Shendure J, Darnell RB, Leslie CS, Rudensky AY, Transcriptome-wide miR-155 binding map reveals widespread noncanonical microRNA targeting. *Mol. Cell* 48, 760–770 (2012). [PubMed: 23142080]
40. Agarwal V, Bell GW, Nam J-W, Bartel DP, Predicting effective microRNA target sites in mammalian mRNAs. *eLife* 4, e05005 (2015).
41. Dweep H, Gretz N, miRWalk2.0: A comprehensive atlas of microRNA-target interactions. *Nat. Methods* 12, 697 (2015). [PubMed: 26226356]
42. Shaw MH, Reimer T, Sánchez-Valdepeñas C, Warner N, Kim Y-G, Fresno M, Nuñez G, T cell–intrinsic role of Nod2 in promoting type 1 immunity to *Toxoplasma gondii*. *Nat. Immunol.* 10, 1267–1274 (2009). [PubMed: 19881508]
43. Kim Y-G, Park J-H, Shaw MH, Franchi L, Inohara N, Núñez G, The cytosolic sensors Nod1 and Nod2 are critical for bacterial recognition and host defense after exposure to Toll-like receptor ligands. *Immunity* 28, 246–257 (2008). [PubMed: 18261938]
44. Chamaillard M, Hashimoto M, Horie Y, Masumoto J, Qiu S, Saab L, Ogura Y, Kawasaki A, Fukase K, Kusumoto S, Valvano MA, Foster SJ, Mak TW, Nuñez G, Inohara N, An essential role for NOD1 in host recognition of bacterial peptidoglycan containing diaminopimelic acid. *Nat. Immunol.* 4, 702–707 (2003). [PubMed: 12796777]

45. Keestra-Gounder AM, Byndloss MX, Seyffert N, Young BM, Chávez-Arroyo A, Tsai AY, Cevallos SA, Winter MG, Pham OH, Tiffany CR, de Jong MF, Kerrinnes T, Ravindran R, Luciw PA, McSorley SJ, Bäumlér AJ, Tsois RM, NOD1 and NOD2 signalling links ER stress with inflammation. *Nature* 532, 394–397 (2016). [PubMed: 27007849]
46. Zhang K, Kaufman RJ, From endoplasmic-reticulum stress to the inflammatory response. *Nature* 454, 455–462 (2008). [PubMed: 18650916]
47. Langmead B, Trapnell C, Pop M, Salzberg SL, Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* 10, R25 (2009). [PubMed: 19261174]
48. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R; 1000 Genome Project Data Processing Subgroup, The sequence alignment/ map format and SAMtools. *Bioinformatics* 25, 2078–2079 (2009). [PubMed: 19505943]
49. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ, Pachter L, Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* 28, 511–515 (2010). [PubMed: 20436464]

Fig. 1. miR-146a is highly expressed selectively on $\gamma\delta 27^-$ T cells.

(A) Microarray heat map of differentially expressed miRNAs in duplicate samples of $\gamma\delta 27^+$ ($n = 4$ mice per sample) and $\gamma\delta 27^-$ CCR6⁺ T cells ($n = 8$ mice per sample) isolated from pooled lymph nodes and spleen of C57BL/6 mice (more than twofold enrichment). (B) RT-qPCR analysis of miR-146a and miR-146b expression in sorted $\gamma\delta 27^+$ and $\gamma\delta 27^-$ T cells from pooled peripheral organs (lymph node and spleen) of C57BL/6 mice. NS, not significant. (C) RT-qPCR analysis of miR-146a expression in sorted DN2 (CD4⁻CD8⁻CD44⁺CD25⁺), DN3 (CD4⁻CD8⁻CD44⁻CD25⁺), $\gamma\delta 25^+$ (CD25⁺CD27⁺), $\gamma\delta 27^+$, and $\gamma\delta 27^-$ thymocytes of C57BL/6 mice. Results are presented relative to miR-423-3p or RNU5G (reference small RNA) expression. Each symbol in (B) and (C) represents an individual mouse. * $P < 0.05$ and ** $P < 0.01$ (Mann-Whitney two-tailed test).

Fig. 2. miR-146a inhibits IFN- γ production in $\gamma\delta$ T cells.

Workflow and results of (A to D) retroviral (RV) overexpression of miR-146a and (E to G) electroporation of miR-146a mimics in sorted peripheral $\gamma\delta$ T cells. (B) Flow cytometry analysis of intracellular IFN- γ and IL-17 expression; and frequency of (C) IFN- γ + and (D) Ki-67+ cells in sorted GFP+ retrovirally transduced $\gamma\delta$ T cells expressing either an unrelated miR (RV-unrelated miR), a control vector (RV-control), or miR-146a (RV-146a). (F) Flow cytometry analysis of intracellular IFN- γ and IL-17 expression and (G) frequency of IFN- γ + in electroporated $\gamma\delta$ T cells with either RNA control or miR-146a mimic. Numbers in

quadrants of flow cytometry plots indicate percentages of cells. Data are representative of five independent experiments. * $P < 0.05$ and ** $P < 0.01$ (Mann-Whitney two-tailed test).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 3. miR-146a limits IFN-γ production and the functional plasticity of γδ27⁻ T cells.

(A) Flow cytometry analysis of intracellular IFN-γ and IL-17 and frequency of IFN-γ⁺ and IL-17⁺ IFN-γ⁺ in γδ27⁻ T cells isolated from peripheral organs and stimulated in vitro overnight with αCD3/28 or for 72 hours in the presence of IL-1β plus IL-23. (B) Schematic for 1:1 mixed miR-146a^{+/+} and miR-146a^{-/-} adult BM and neonatal thymocyte chimeras in Rag2^{-/-} hosts. (C) Flow cytometry analysis of intracellular IFN-γ and IL-17, mean fluorescence intensity (MFI) of IFN-γ⁺, and fold difference of IFN-γ⁺ IL-17⁺ in γδ27⁻ T cells isolated from peripheral organs of above-introduced chimeras after in vitro stimulation

for 72 hours with IL-1 β plus IL-23. Numbers in quadrants of flow cytometry plots indicate percentages of cells. Each symbol in (A) and (C) represents an individual experiment. * $P < 0.05$ (Mann-Whitney two-tailed test).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 4. miR-146a restricts functional plasticity of $\gamma\delta$ 27⁻ T cells in *Listeria* infection.

(A) Flow cytometry analysis of intracellular IFN- γ and IL-17 expression in $\gamma\delta$ T cells isolated from spleens of miR-146a^{+/+} or miR-146a^{-/-} mice, 7 days after infection with *L. monocytogenes*. (B) Frequency and total cell numbers of IL-17⁺ IFN- γ ⁺ in $\gamma\delta$ T cells isolated from spleens of miR-146a^{+/+} and miR-146a^{-/-} mice, 7 days after infection with *L. monocytogenes*. (C) Representative flow cytometry analysis of V γ 6 chain usage (left), frequency of IFN- γ ⁺ IL-17⁺ (middle), and total cell numbers of IFN- γ ⁺ IL-17⁺ (right) of V γ 1V γ 4⁻ $\gamma\delta$ T cells isolated from spleen of miR-146a^{+/+} or miR-146a^{-/-} mice, 7 days after infection with *L. monocytogenes*. (D) Representative flow cytometry analysis of IL-17⁺ IFN- γ ⁺ $\gamma\delta$ T cells isolated from the chimeras established in Fig. 3B. (Right) frequency within IL-17⁺ IFN- γ ⁺ $\gamma\delta$ T cells of either miR-146a^{+/+} or miR-146a^{-/-} origin. Numbers in quadrants of flow cytometry plots indicate percentages of cells. Each symbol in (B) to (D) represents an individual mouse. **P* < 0.05 and ***P* < 0.01 (Mann-Whitney two-tailed test). WT, wild-type; KO, knockout.

Fig. 5. miR-146a targets Nod1 mRNA that is depleted in $\gamma\delta 27^-$ T cells.

(A) Putative binding sites of miR-146a-5p in the 3'UTR region of Nod1 and Atf2. The line represents canonical Watson-Crick base pairing, whereas the dot represents noncanonical base pairing between miR-146a and Nod1 or Atf2 mRNAs. The red bases above illustrated the mutated sites in the region of Nod1 and Atf2 3'UTRs. (B) Dual luciferase reporter assay was performed to verify binding between miR-146a and Nod1 or Atf2 mRNAs. HEK293 T cells were cotransfected with a pmirGLO Dual-Luciferase miRNA Target Expression Vector (Promega) containing either the WT or mutated 3'UTR target sites plus miR-146a, an unrelated miRNA, or GFP control expression vector. A negative construct (without miR-146a binding sites) and a positive construct (3'UTR of Notch1) were included. Data are from three to four independent experiments with technical replicates. * $P < 0.05$, ** $P < 0.005$, *** $P < 0.001$, Student's t test. RT-qPCR analysis of Nod1 and Atf2 expression in (C) sorted $\gamma\delta 27^+$, $\gamma\delta 27^-CCR6^-$, and $\gamma\delta 27^-CCR6^+$ T cells from pooled peripheral organs (lymph node and spleen) of C57BL/6 mice; (D) sorted $V\gamma 1^+$, $V\gamma 4^+$, and $V\gamma 1^-\gamma 4^-$ $\gamma\delta$ T cells and $CD4^+$ T cells from pooled peripheral organs (lymph node and spleen) of C57BL/6 mice; and (E) sorted $V\gamma 4^+CD27^-$ and $V\gamma 1^-\gamma 4^-CD27^-$ $\gamma\delta$ T cells from pooled lymph nodes of either miR-146a^{+/+} or miR-146a^{-/-} mice. Results are presented relative to β -actin and

HPRT expression. Each symbol in (C) to (E) represents an individual mouse. * $P < 0.05$ and ** $P < 0.01$ (Mann-Whitney two-tailed test).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 6. Nod1 is required for IFN- γ production and functional plasticity in $\gamma\delta$ T cells.

(A) Flow cytometry analysis of intracellular IFN- γ and IL-17 expression and frequency of IFN- γ ⁺ IL-17⁺ (left), IL-17⁺ (middle), and Ki-67⁺ (right) in $\gamma\delta$ T cells isolated from peripheral organs (spleen and lymph nodes) of Nod1^{+/+} and Nod1^{-/-} littermates as well as Atf2^{+/+} and Atf2^{-/-} littermates after overnight stimulation with IL-1 β plus IL-23. (B) Flow cytometry analysis of intracellular IFN- γ and IL-17 expression and frequency of IFN- γ ⁺ IL-17⁺ (left), IL-17⁺ (up) and Ki-67⁺ (bottom) in $\gamma\delta$ T cells isolated from peripheral organs (spleen and lymph nodes) of miR-146a^{+/+} Nod1^{+/+}, miR-146a^{-/-} Nod1^{+/+}, and miR-146a^{-/-} Nod1^{-/-} mice.

miR-146a^{-/-}Nod1^{+/-} littermates stimulated for 72 hours in the presence of IL-1 β plus IL-23. (C) Bacterial CFUs of *L. monocytogenes* bacterial burden was enumerated in the spleen 4 days after infection (left), weight loss (middle), and survival (right) of Nod1^{+/+}, Nod^{+/-}, Nod^{-/-}, and TCRd^{-/-} mice infected with *Listeria* ($n = 7$ to 18 in three independent experiments). (D) Flow cytometry analysis of intracellular IFN- γ and IL-17 expression in $\gamma\delta$ T cells isolated from spleen of Nod1^{+/+}, Nod^{+/-}, and Nod^{-/-} mice, 7 days after infection with *Listeria*. Numbers in quadrants of flow cytometry plots indicate percentages of cells. (E) Frequency of IFN- γ ⁺ IL-17⁺ within $\gamma\delta$ T cells (left) and V γ 1V γ 4⁻ $\gamma\delta$ T cells (right) isolated from spleen of Nod1^{+/+}, Nod^{+/-}, and Nod^{-/-}, 7 days after infection with *Listeria* ($n = 7$ to 18 in three independent experiments). Each symbol in (A) and (D) represents an individual mouse. * $P < 0.05$ and ** $P < 0.01$ (Mann-Whitney two-tailed test).

Table 1.

Putative miR-146a targets based on differential Ago2-RNA immunoprecipitation followed by deep sequencing in T cells.

Gene symbol	Fold change	No. of binding
Receptor activity		
<i>Nod1</i>	4.21	1
<i>Tcp11</i>	3.44	1
<i>Ghr</i>	3.07	1
<i>Paqr7</i>	2.72	1
<i>Unc5a</i>	2.59	1
Transcription factor activity		
<i>Ccnh</i>	6.48	1
<i>Tef</i>	4.49	2
<i>Myf6</i>	3.80	1
<i>Zfp788</i>	3.68	1
<i>Tfap2b</i>	3.67	2
<i>Grhl2</i>	3.34	1
<i>Foxc1</i>	3.21	1
<i>Sp1</i>	2.90	1
<i>Milt10</i>	2.83	1
<i>Gtf2e2</i>	2.83	3
<i>Atf2</i>	2.53	2
<i>Yap1</i>	2.50	2
<i>Phf21a</i>	2.32	3