Published in final edited form as:

Endocrinol Metab Clin North Am. 2016 March ; 45(1): 205–216. doi:10.1016/j.ecl.2015.09.013.

Dyslipidemia in Special Ethnic Populations

Jia Pu, PhD^{a,*}, Robert Romanelli, PhD^a, Beinan Zhao, MS^a, Kristen M.J. Azar, RN, MSN, MPH^a, Katherine G. Hastings, BA^b, Vani Nimbal, MPH^a, Stephen P. Fortmann, MD^c, Latha P. Palaniappan, MD, MS^b

^aPalo Alto Medical Foundation Research Institute, Ames Building, 795 El Camino Real, Palo Alto, CA 94301, USA;

^bStanford University School of Medicine, 1265 Welch Road, Stanford, CA 94305, USA;

^cKaiser Permanente Center for Health Research, 3800 North Interstate Avenue, Portland, OR 97227, USA

SUMMARY

There are significant racial/ethnic differences in dyslipidemia prevalence, dyslipidemia-related mortality rates, and response to lipid-lowering agents. Among all racial/ethnic groups, Asian Indians, Filipinos and Hispanics are at most elevated risk for dyslipidemia, which is consistent with the higher CHD mortality rates in these groups. More attention should be paid to these at-risk groups for screening and treatment purposes. Compared with other racial/ethnic groups, statins may have a higher efficacy for Asians, which may potentially be explained by genetic differences in statin metabolism, but overall the data are mixed. At present it may be wise to start with a lower statin dose in an individual with Asian ancestry until the treatment and adverse effects can be determined. In addition, racial/ethnic differences in health behavior patterns should be taken into consideration when promoting lifestyle modification among individuals with dyslipidemia. In particular, Hispanic subgroups (Mexican, Puerto Rican, etc) and Asian (Chinese, South Asian, etc) subgroups should be disaggregated in lifestyle interventions. Further studies are needed to better understand racial/ethnic-specific risk factors contributing to the observed differences in dyslipidemia, CHD, and stroke. Culturally tailored prevention and intervention should be provided to the minority populations with elevated risk for dyslipidemia and considerably more research is needed to determine the best approaches to helping specific subgroups.

Keywords

Dyslipidemia; Racial/ethnic differences; Prevalence; Mortality; Treatment; Lifestyle modification

INTRODUCTION

Dyslipidemia, including high levels of low-density lipoprotein cholesterol (LDL-C; 130 mg/dL), total cholesterol 200 mg/dL), and triglycerides (TG; 150 mg/dL), or low levels of

*Corresponding author. puj@pamfri.org.

Conflicts of Interest: All authors declare they have no conflicts of interest.

high-density lipoprotein cholesterol (HDL-C; <40 [men] and <50 [women] mg/dL), is among the leading risk factors for coronary heart disease (CHD) and stroke.¹ A report of the National Health and Nutrition Examination Survey (NHANES) from 2003 to 2006 estimated that 53% (105.3 million) US adults have at least one lipid abnormality: 27% (53.5 million) have high LDL-C, 23% (46.4 million) have low HDL-C, and 30% (58.9 million) have high TG. In addition, 21% (42.0 million) of US adults have mixed dyslipidemia, defined as the presence of high LDL-C combined with at least one other lipid abnormality.²

Significant heterogeneity in patterns of dyslipidemia prevalence, its relation to CHD and stroke mortality rates, and response to lipid-lowering agents has been observed across racial/ethnic groups.³ These differences in dyslipidemia provide important information that may in part explain the variation in cardiovascular disease (CVD) burden observed across racial/ethnic subgroups. Better understanding of dyslipidemia in special racial/ethnic populations is needed to guide prevention, screening, and treatment efforts.

PREVALENCE OF DYSLIPIDEMIA SUBTYPES AMONG SPECIAL RACIAL/ETHNIC GROUPS

The NHANES is the primary data source for national prevalence rates of dyslipidemia in the United States, sampling mainly non-Hispanic whites (whites), non-Hispanic blacks (blacks), and Mexican Americans. The NHANES has very limited samples from the Asian subgroups.⁴ Other data sources, such as primary care settings and observational studies, contribute to a comprehensive picture of racial/ethnic differences in dyslipidemia by providing important information about races and ethnicities that are less represented in the NHANES. One should be aware that the prevalence of dyslipidemia varies by data source. The observed differences in the prevalence rates of dyslipidemia between studies can be attributable to factors such as study design, sampling methods, time period, geographic variation, and participants' characteristics.

Low-Density Lipoprotein Cholesterol

NHANES data in 2013 showed that the prevalence rate of high LDL-C was highest among Mexican men (40%) and women (30%), followed by non-Hispanic black men (33%) and women (31%). Non-Hispanic white men (30%) and women (29%) had the lowest prevalence of high LDL-C among the 3 racial/ethnic groups.⁵

Similarly, data from a clinic-based cohort in northern California from 2008 to 2011 showed that 63% of black men and 57% of black women had high LDL-C, which were slightly higher than the prevalence rates among non-Hispanic white men (62%) and women (53%).³ Further, Mexican American men (66%) and women (57%) also had higher prevalence of high LDL-C compared with non-Hispanic whites.³ Filipino men (73%) and women (63%) had the highest prevalence rates of high LDL-C among Asian subgroups, non-Hispanic whites, non-Hispanic blacks, and Hispanics.³

Several other studies provide further estimates for variation in prevalence among race/ethnic minority subgroups. Data from the Hispanic Community Health Study (HCHS)/Study of Latinos (SOL), an observational study in San Diego, Chicago, New York City, and Miami,

showed variations among Hispanic subgroups with particularly high prevalence of dyslipidemia among Central American men (55%) and Puerto Rican women (41%).⁶ The Study of Health Assessment and Health Risk in Ethnic groups (SHARE) investigated the prevalence of CHD risk factors for a multiethnic cohort from 3 Canadian cities. They found that South Asians, mainly Asian Indians, had an increased prevalence of higher total and LDL cholesterol compared with Europeans and Chinese.⁷

High-Density Lipoprotein Cholesterol

NHANES data in 2013 showed that 20% of black men and 10% of black women had low HDL-C, defined as less than 40 mg/dL in both men and women, which were lower than the prevalence rates among non-Hispanic white men (33%) and women (12%).⁵ NHANES data also showed Mexican American men (34%) and women (15%) had higher prevalence of low HDL-C compared with non-Hispanic whites.⁵ According to NHANES data from 2011 to 2012, 25% of Asian American men and 5% of Asian American women had low HDL-C.⁸

Although NHANES data showed Asian Americans had the lowest prevalence of low HDL-C as an aggregated group, data from a clinic-based cohort with disaggregated Asian ethnic groups in northern California between 2008 and 2011 found that Asian Indian men (53%) and women (55%) had the highest prevalence of low HDL-C among Asian American subgroups; their prevalence was also higher than Mexican American men (48%) and women (51%), non-Hispanic black men (34%) and women (40%), and non-Hispanic white men (36%) and women (31%).³ Similarly, data from the SHARE study showed South Asians including Asian Indians had an increased prevalence of low HDL-C compared with Europeans and Chinese.⁷

Triglycerides

NHANES data from 1999 through 2008 showed 35% of Mexican Americans had high TG, followed by 33% among non-Hispanic whites, and 16% among non-Hispanic blacks.⁹ Data from a clinic-based cohort in northern California from 2008 to 2011 found that Filipino men (60%) and Mexican women (45%) had the highest prevalence of high TG, compared with Mexican men (56%) and Filipino women (42%), Asian Indian men (55%) and women (37%), non-Hispanic white men (43%) and women (28%), and non-Hispanic black men (30%), and women (18%).³ Data from the SHARE study showed South Asians had the highest prevalence of high TG among South Asians, Chinese, and Europeans.⁷

Potential explanations for racial/ethnic differences in dyslipidemia prevalence have been explored by several studies. In the Multi-Ethnic Study of Atherosclerosis (MESA), researchers found that ethnic disparities were substantially attenuated by adjusting for access to health care.¹⁰ The racial/ethnic differences could also be related to lifestyle, genetic, and cultural differences associated with total and LDL-C concentrations.¹¹ For example, the predilection of South Asians to have lower HDL-C levels has been attributed to the higher prevalence of insulin resistance and related metabolic abnormalities, which may be the consequence of a combination of genetic predisposition, physical inactivity, and a high carbohydrate diet.^{12–14}

DYSLIPIDEMIA-RELATED MORTALITY IN SPECIAL RACIAL/ETHNIC GROUPS

Dyslipidemia often results in an increased risk of premature atherosclerosis, a major risk factor for CHD.¹⁵ CHD is the leading cause of mortality for both men and women in the United States and worldwide, with increasing evidence of gender and racial/ethnic minority disparities in CHD morbidity and mortality.^{16–19} Despite declines in CHD death rates over the past decades, CHD contributes to more than one-third of all deaths for those over the age of 35 years.²⁰

Differences in mortality rate from CHD have been found across races/ethnicities. In the United States, CHD mortality rates are highest in blacks, intermediate in whites and Hispanics, and lowest in some Asian subgroups.^{15,21,22} These CHD rates are paralleled with observed racial/ethnic disparities in dyslipidemia prevalence, with higher CHD rates seen in Hispanics and blacks, potentially owing to limited health care access and other less favorable behavioral factors seen in these groups.¹⁰

Meanwhile, although traditionally known as the “model minority,” disproportionate mortality burden owing to CHD and stroke has been shown among certain Asian subgroups, such as Asian Indians, Filipinos, and Japanese.^{23,24} Of note, the landmark Ni-Hon-San study showed increased CHD mortality rates and decreased stroke rates among Japanese-American men compared with rates in Japan, suggesting a differential disease impact of acculturation to Westernized lifestyles.²⁵ A recent study found higher proportional mortality owing to CHD among Asian Indians, especially in younger age groups, compared with all other racial/ethnic groups.¹⁷ Increased burden from CHD mortality has been well-documented in both native and immigrant Asian Indian populations.^{7,26,27} This observation is consistent with the higher prevalence rates of dyslipidemia (especially low HDL-C) in Asian Indians compared with other Asian subgroups and non-Hispanic whites in the United States.^{3,7,28}

Environmental and social factors (eg, acculturation, socioeconomic status, diet) have been known to increase CHD mortality risk.^{29,30} Cultural diets high in fat, increasing the risk for dyslipidemia, are also of concern.³¹ Differences in susceptibility to CHD may also have a genetic basis, although this has not been determined adequately yet. Thus, it is critical for clinicians to modify lipid management appropriately among these rapidly growing populations.

TREATMENT OF DYSLIPIDEMIA

Overview

There are several US Food and Drug Administration (FDA)-approved 3-hydroxy-3-methylglutaryl-coenzyme A reductase inhibitors (statins) and a variety of nonstatin therapies available for the treatment of dyslipidemia, including bile acids sequestrants, cholesterol absorption inhibitors, fibrates, niacin, and omega-3 fatty acids. Statins are the most widely prescribed treatment for dyslipidemia, and one of the most commonly prescribed drugs in the United States.³² In the most recent national cholesterol treatment guidelines, published

in November 2013 by the American College of Cardiology (ACC)/American Heart Association (AHA),¹ the Expert Panel determined from robust clinical trial data that statins have the most acceptable CVD risk reduction benefit and side effect profile, and that the addition of nonstatin therapy does not seem to provide further benefit in reducing CVD. Clinical trials on which cholesterol treatment guidelines are based have often underrepresented racial/ethnic minority groups. Accordingly, our knowledge of optimal statin treatment regimens, and the effectiveness, tolerability, and safety of these regimens in clinical practice, is limited across diverse racial/ethnic populations.

Racial/Ethnic Differences in Risk Stratification

The most recent ACC/AHA guidelines expand the criteria for patients who would benefit from statin treatment, and more than 80% of newly eligible patients are expected to have no prior CVD.³³ Therefore, they would be assessed for optimal statin treatment based on the new Atherosclerotic CVD (ASCVD) Risk Estimator.³³ This estimator is intended to improve on previous CVD risk estimators, including the Framingham³⁴ and Adult Treatment Panel III³⁵ risk algorithms. It was derived from several longitudinal epidemiologic cohort studies of non-Hispanic whites and blacks, and was validated externally in 2 cohort studies with similar populations (MESA and REGARDS),³⁶ as well as in contemporary samples of the derivation cohorts. However, studies among other race/ethnic minority groups are lacking. The ACC/AHA Work Group, which developed the algorithm for the ASCVD Risk Estimator, acknowledges that it should be used only in men and women of non-Hispanic white or non-Hispanic black decent, and that it may not accurately predict risk in other racial/ethnic groups.³⁷ Specifically, there is concern of overestimation of risk in Mexican Americans and East Asians and underestimation of risk in other groups, such as Puerto Ricans and South Asians.^{37,38} Other countries with diverse racial/ethnic minority populations, such as the United Kingdom, have validated algorithms predicting risk for more than 9 specific racial/ethnic subpopulations (available at: <http://qrisk.org/>).³⁹

Race/Ethnic Differences in Statin Response

Statin metabolism and drug sensitivity—Much of the data on race/ethnic differences in statin metabolism have shown that some Asian subgroups are slower to metabolize statins compared with non-Hispanic whites, which leads to higher systemic drug concentrations. Pharmacokinetic studies indicate rosuvastatin plasma concentrations are 2-fold higher in Japanese relative to non-Hispanic white individuals. Rosuvastatin plasma concentrations have been shown to be elevated similarly in other Asian subgroups; these included Chinese, Malay, and Asian Indians, whose rosuvastatin concentrations were approximately 2 times higher relative to Caucasians.⁴⁰

There are several genetic variants associated with altered statin metabolism. Such variants include, but are not limited to, single nucleotide polymorphisms in the genes that encode the organic anion-transporting polypeptide (OATP)1B1 (521T> C), which regulates hepatic uptake of statins, and the adenosine triphosphate-binding cassette G2 (ABCG2) transporter (421C> A), which regulates hepatic efflux.^{41,42} Variants of both genes are associated with increased statin plasma concentrations within multiple race/ethnic groups.^{41–44} 421C> A has been found to be a single nucleotide polymorphism candidate to explain the observed racial

differences in statin metabolism: the allelic frequency of 421C> A in ABCG2 is higher in Chinese (~35%) relative to Caucasians (9%–14%).^{43,45}

Higher plasma levels of statins in Asians compared with non-Hispanic whites has led to concern about increased risk for statin-induced side effects in this population. Such concerns have led to a revised package insert recommending lower starting doses of rosuvastatin in this population (5 vs 10 mg for non-Hispanic whites).⁴⁶ Notably, in Japan, starting doses for most statins are one-half of what is recommended in the United States.⁴⁷

There is less information on statin metabolism in other racial/ethnic minority groups. We are aware of 1 study that evaluated the pharmacokinetics of single-dose pravastatin in non-Hispanic blacks versus European Americans.⁴⁴ In this study, the OATP1B1 521T> C polymorphism was associated significantly with higher statin plasma concentrations in subjects of European versus African ancestry.⁴⁴ This may be explained by the low allelic frequency of OATP1B1 521T> C in non-Hispanic blacks (~1%)⁴⁴ relative to non-Hispanic whites. To our knowledge, no studies have been published on the pharmacokinetics of statins in Hispanics.

Myalgia (muscle pain) is one of the most commonly observed side effects associated with statins, but has been reported to occur at varying frequencies (5%–20%) in randomized controlled trials (RCT) and observational studies.^{48–50} More severe but rare muscle side effects have also been reported, including myopathy (0.01%–0.3%) and rhabdomyolysis (0.003%–0.01%).⁵¹ To date, high-dose simvastatin (80 mg) is the only statin to receive a US FDA warning for increased risk of muscle damage.⁵² The use of statins in combination with other drugs has also been shown to increase the risk of adverse events.^{53,54} In HPS2-THRIVE, which included 25,673 adults from Europe and China on simvastatin 40 mg/d, a higher risk of adverse events, including myopathy, were reported in patients randomized to niacin-laropiprant relative to placebo.⁵⁵ Furthermore, the relative risk of musculoskeletal events (mostly myopathy) in the niacin-laropiprant versus placebo group was markedly higher among Chinese participants than European participants.⁵⁵ To our knowledge no safety concerns have been raised for statins alone or in combination with agents in non-Hispanic blacks or Hispanics.

Statin efficacy in Asian populations—Studies conducted in Asian countries have documented that lower statin doses can achieve similar therapeutic effects in Asian populations.

- In an open-label study of Japanese patients receiving simvastatin (initial dose 5 mg/d), LDL-C decreased by 26% over 6 months, an effect that corresponds with simvastatin 20 mg/d in Western studies⁵⁶
- In a multicenter, double-blind, RCT in 6 Asian countries, patients randomized to receive 10 mg simvastatin or atorvastatin daily over 8 weeks had an average LDL-C reduction of 35% and 43%, respectively, with more than 80% of patients achieving a National Cholesterol Education Program LDL-C target.⁵⁷ To see similar effects in non-Hispanic whites would generally require at least double this dose of simvastatin.

- In a prospective RCT of the primary prevention of CVD in Japan, pravastatin 10 to 20 mg/d reduced LDL-C by only 18% but reduced CHD by 33% relative to diet-alone⁵⁸; this level of risk reduction in CVD is similar to trials of predominantly non-Hispanic white populations taking higher daily doses of pravastatin (40 mg)⁵⁹ or a higher potency statin (atorvastatin 10 mg/d).⁶⁰

It is tempting to speculate that the apparently increased effect of statins in Asians relative to non-Hispanic whites is owing to genetic differences in statin metabolism; however, because comparisons in these studies were indirect and were conducted on ethnic populations outside of the United States or Europe, underlying cultural differences in diet and lifestyle cannot be ruled out as a causal factor.

In addition, other studies have failed to demonstrate differences in response to statins in Asian and non-Hispanic white populations.

- In a combined analysis of 2 small multicenter open-label studies, GOALLS (included non-Asian and Asian participants) and STATT (included Asians only), the authors compared cholesterol outcomes among patients with CHD treated with simvastatin for 14 weeks.⁶¹ There were no differences in changes in LDL-C among Asians in the STATT study ($n = 133$; -45.4 mg/dL) relative to Asians ($n = 15$) or non-Asians ($n = 183$) in the GOALLS study (-41.1 and 41.2 mg/dL, respectively)⁶¹
- In an observational, prospective study in Canada, no difference in the magnitude of LDL-C lowering was observed among non-Hispanic whites and South Asians (-41% vs -43% , respectively; $P = .40$), taking atorvastatin or simvastatin (20 mg median dose for each) for more than 3 years in the secondary prevention of CVD⁶²
- Using data from an RCT evaluating atorvastatin versus placebo in a multiethnic population in the UK, Chapman and colleagues⁶³ matched White ($n = 198$) and South Asian ($n = 76$) cohorts receiving atorvastatin to evaluate the effects of statins across race/ethnic groups. The authors again found no difference in the percent reduction in LDL-C among whites and South Asians (-40% vs -39% ; $P = .92$)⁶³

To our best knowledge, no existing studies have shown differences in statin efficacy in non-Hispanic blacks, Hispanics, and non-Hispanic whites.^{63–65}

Lifestyle Modification in Special Racial/Ethnic Groups

Lifestyle modification has been recommended to treat dyslipidemia and to reduce ASCVD risk, both before and in addition to the use of lipid-lowering agents.¹ These lifestyle risk factors include unhealthy diet, overweight and obesity, and physical inactivity. The AHA has the following specific recommendations for individuals with dyslipidemia⁶⁶:

- Diet: increase the intake of vegetables, fruits, and whole grains and limit intake of sweets; reduce the intake of saturated fat and trans fat.

- Physical activity: engage in aerobic physical activity; three to four 40-minute sessions per week; moderate-to-vigorous intensity physical activity.

Racial/ethnic disparities also exist in these lifestyle risk factors. According to data from the Behavioral Risk Factor Surveillance System in 2002, non-Hispanic whites had more fruits and vegetables in their diet compared with Hispanics and non-Hispanic blacks.^{6,67} According to the National Health Interview Survey for 2008 through 2010, 71% of Hispanics were overweight or obese, followed by non-Hispanic blacks (70%), non-Hispanic whites (62%), and Asian Americans (42%). In addition, National Health Interview Survey for 2008 through 2010 also found that, compared with non-Hispanic whites (20%), non-Hispanic black (17%), Asian American (16%), and Hispanic (13%) adults were less likely to meet the 2008 guidelines for aerobic and muscle strengthening through leisure time activity.⁶⁸ All these data indicate that both non-Hispanic blacks and Hispanics are at higher risk for having both diet and lifestyle risk factors, and Asian Americans are more likely to be physically inactive.

A recent systematic review included nutrition and physical activity intervention studies in adult African Americans, from 2000 to 2011.⁶⁹ This study found both diet and physical activity interventions for weight loss improved cholesterol clinical outcomes among African Americans.⁶⁹ In particular, it provided evidence to support interventions in community-based settings among African Americans. A 2011 systematic review included intervention studies that promote physical activity in Hispanic adults published between 1988 and 2011.⁷⁰ This study concluded that physical activity interventions in Hispanics should include community-based settings, social support strategy, culturally sensitive intervention design, and staff from the same ethnic group. A systemic review of RCTs of lifestyle interventions for Asian Americans published between 1995 and 2013 concluded that lifestyle interventions improved physical activity, healthy diet, and weight control in Asian Americans.⁷¹ However, the studies included in this review were limited in cultural appropriateness. In particular, Asian subgroups were aggregated together although they have different cultures and health behavior patterns. Other recommendations include individual tailoring, education and modeling of lifestyle behaviors, and providing support during a maintenance phase.

Immigration and acculturation have a profound impact on lifestyle in both Hispanics and Asians in the United States. For example, traditional Hispanic diets contain high levels of fiber. However, studies found US-born Hispanics had a hard time retaining traditional diets and consumed more fat and sugar compared with their counterparts in their home countries.^{72,73} They have less access to high nutritional quality foods and are at risk for overweight/obesity.^{72,74} Similarly, traditional East Asian diets contain less total and saturated fat but more sodium intake compared with Western diets and Asian Indian diets.^{75,76} However, Chinese immigrants who have lived in the United States for longer than 10 years had a more unhealthy diet and less physical activity compared with recent immigrants.³⁰ In contrast, higher acculturation levels were associated with more physical activity among Korean Americans.⁷⁷ Future studies should consider providing culturally tailored, ethnic-specific interventions to these diverse immigrant populations.

REFERENCES

1. Stone NJ, Robinson J, Lichtenstein AH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association task force on practice guidelines. *Circulation* 2014;129:S1–45. [PubMed: 24222016]
2. Toth PP, Potter D, Ming EE. Prevalence of lipid abnormalities in the United States: the National Health and Nutrition Examination Survey 2003–2006. *J Clin Lipidol* 2012;6(4):325–30. [PubMed: 22836069]
3. Frank AT, Zhao B, Jose PO, et al. Racial/ethnic differences in dyslipidemia patterns. *Circulation* 2014;129(5):570–9. [PubMed: 24192801]
4. Holland AT, Palaniappan LP. Problems with the collection and interpretation of Asian-American health data: omission, aggregation, and extrapolation. *Ann Epidemiol* 2012;22(6):397–405. [PubMed: 22625997]
5. Statistical fact sheet 2013 update: high blood cholesterol & other lipids. 2013 Available at: http://www.heart.org/idc/groups/heart-public/@wcm/@sop/@smd/documents/downloadable/ucm_319586.pdf. Accessed November 15, 2014.
6. Rodriguez CJ, Allison M, Daviglus ML, et al. Status of cardiovascular disease and stroke in Hispanics/Latinos in the United States: a science advisory from the American Heart Association. *Circulation* 2014;130:593–625 (1524–4539 [Electronic]). [PubMed: 25098323]
7. Anand SS, Yusuf S, Vuksan V, et al. Differences in risk factors, atherosclerosis and cardiovascular disease between ethnic groups in Canada: the Study of Health Assessment and Risk in Ethnic groups (SHARE). *Indian Heart J* 2000; 52(7 Suppl):S35–43. [PubMed: 11339439]
8. Aoki Y, Yoon SS, Chong Y, et al. Hypertension, abnormal cholesterol, and high body mass index among non-Hispanic Asian adults: United States, 2011–2012. *NCHS Data Brief* 2014;(140):1–8.
9. Miller M, Stone NJ, Ballantyne C, et al. Triglycerides and cardiovascular disease: a scientific statement from the American Heart Association. *Circulation* 2011; 123(20):2292–333. [PubMed: 21502576]
10. Goff DC Jr, Bertoni AG, Kramer H, et al. Dyslipidemia prevalence, treatment, and control in the Multi-Ethnic Study of Atherosclerosis (MESA): gender, ethnicity, and coronary artery calcium. *Circulation* 2006;113(5):647–56. [PubMed: 16461837]
11. Huang MH, Schocken M, Block G, et al. Variation in nutrient intakes by ethnicity: results from the Study of Women's Health Across the Nation (SWAN). *Menopause* 2002;9(5):309–19. [PubMed: 12218719]
12. McKeigue PM, Miller GJ, Marmot MG. Coronary heart disease in South Asians overseas: a review. *J Clin Epidemiol* 1989;42(7):597–609. [PubMed: 2668448]
13. Chambers JC, Kooner JS. Diabetes, insulin resistance and vascular disease among Indian Asians and Europeans. *Semin Vasc Med* 2002;2(2):199–214. [PubMed: 16222611]
14. Radhika G, Ganesan A, Sathya RM, et al. Dietary carbohydrates, glycemic load and serum high-density lipoprotein cholesterol concentrations among South Indian adults. *Eur J Clin Nutr* 2009;63(3):413–20. [PubMed: 17987051]
15. Enas EA. Clinical implications: dyslipidemia in the Asian Indian population. 2002 Available at: <https://southasianheartcenter.org/docs/AAPImonograph.pdf>. Accessed November 15, 2014.
16. Roger VL, Go AS, Lloyd-Jones DM, et al. Heart disease and stroke statistics— 2011 update: a report from the American Heart Association. *Circulation* 2011; 123(4):e18–209. [PubMed: 21160056]
17. Jose PO, Frank AT, Kapphahn KI, et al. Cardiovascular disease mortality in Asian Americans (2003 to 2010). *J Am Coll Cardiol* 2014;64(23):2486–94. [PubMed: 25500233]
18. Palaniappan L, Wang Y, Fortmann SP. Coronary heart disease mortality for six ethnic groups in California, 1990–2000. *Ann Epidemiol* 2004;14(7):499–506. [PubMed: 15310526]
19. Keppel KG, Percy JN, Heron MP. Is there progress toward eliminating racial/ethnic disparities in the leading causes of death? *Public Health Rep* 2010;125(5):689–97. [PubMed: 20873285]
20. Farnier M, Chen E, Johnson-Levonas AO, et al. Effects of extended-release niacin/laropiprant, simvastatin, and the combination on correlations between apolipoprotein B, LDL cholesterol, and

non-HDL cholesterol in patients with dyslipidemia. *Vasc Health Risk Manag* 2014;10:279–90. [PubMed: 24855368]

21. Enas E, Yusuf S, Mehta J. Meeting of international working group on coronary artery disease in South Asians. *Indian Heart J* 1996;48:727–32. [PubMed: 9062031]
22. Hoyert DL. 75 years of mortality in the United States, 1935–2010. *NCHS Data Brief* 2012;(88):1–8.
23. Ye J, Rust G, Baltrus P, et al. Cardiovascular risk factors among Asian Americans: results from a national health survey. *Ann Epidemiol* 2009;19(10):718–23. [PubMed: 19560369]
24. Klatsky AL, Tekawa IS, Armstrong MA. Cardiovascular risk factors among Asian Americans. *Public Health Rep* 1996;111(Suppl 2):62–4. [PubMed: 8898779]
25. Marmot MG, Syme SL, Kagan A, et al. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii and California: prevalence of coronary and hypertensive heart disease and associated risk factors. *Am J Epidemiol* 1975;102(6):514–25. [PubMed: 1202953]
26. Balarajan R Ethnic differences in mortality from ischaemic heart disease and cerebrovascular disease in England and Wales. *BMJ* 1991;302(6776):560–4. [PubMed: 2021718]
27. Enas EA, Garg A, Davidson MA, et al. Coronary heart disease and its risk factors in first-generation immigrant Asian Indians to the United States of America. *Indian Heart J* 1996;48(4):343–53. [PubMed: 8908818]
28. Karthikeyan G, Teo KK, Islam S, et al. Lipid profile, plasma apolipoproteins, and risk of a first myocardial infarction among Asians: an analysis from the INTER-HEART Study. *J Am Coll Cardiol* 2009;53(3):244–53. [PubMed: 19147041]
29. Marmot MG, Syme SL. Acculturation and coronary heart disease in Japanese-Americans. *Am J Epidemiol* 1976;104(3):225–47. [PubMed: 961690]
30. Taylor VM, Yasui Y, Tu SP, et al. Heart disease prevention among Chinese immigrants. *J Community Health* 2007;32(5):299–310. [PubMed: 17922202]
31. Gupta M, Brister S. Is South Asian ethnicity an independent cardiovascular risk factor? *Can J Cardiol* 2006;22(3):193–7. [PubMed: 16520847]
32. Bartholow M Rx focus: top 200 drugs of 2012. *Pharm Times* 2012.
33. Pencina MJ, Navar-Boggan AM, D'Agostino RB Sr, et al. Application of new cholesterol guidelines to a population-based sample. *N Engl J Med* 2014; 370(15):1422–31. [PubMed: 24645848]
34. D'Agostino RB Sr, Vasan RS, Pencina MJ, et al. General cardiovascular risk profile for use in primary care: the Framingham heart study. *Circulation* 2008;117(6): 743–53. [PubMed: 18212285]
35. National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third report of the national cholesterol education program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult treatment panel III) final report. *Circulation* 2002;106(25):3143–421. [PubMed: 12485966]
36. Goff DC Jr, Lloyd-Jones DM, Bennett G, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014;63(25 Pt B):2935–59. [PubMed: 24239921]
37. Goff DC Jr, Lloyd-Jones DM. 2013 Report on the assessment of cardiovascular risk: full work group report supplement. 2013 Available at: http://jaccjacc.cardiosource.com/acc_documents/2013_FPR_S5_Risk_Assessment.pdf. Accessed May 22, 2014.
38. Stone NJ, Robinson JG, Lichtenstein AH. 2013 Report on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular disease in adults: full panel report supplement. 2013 Available at: http://www.kcumb.edu/uploadedFiles/Content/Academics/_Assets/CME_Presentations/Moriarty12_5_full.pdf. Accessed May 23, 2014.
39. Hippisley-Cox J, Coupland C, Vinogradova Y, et al. Predicting cardiovascular risk in England and Wales: prospective derivation and validation of QRISK2. *BMJ* 2008;336(7659):1475–82. [PubMed: 18573856]

40. Lee E, Ryan S, Birmingham B, et al. Rosuvastatin pharmacokinetics and pharmacogenetics in white and Asian subjects residing in the same environment. *Clin Pharmacol Ther* 2005;78(4):330–41. [PubMed: 16198652]
41. Nishizato Y, Ieiri I, Suzuki H, et al. Polymorphisms of OATP-C (SLC21A6) and OAT3 (SLC22A8) genes: consequences for pravastatin pharmacokinetics. *Clin Pharmacol Ther* 2003;73(6):554–65. [PubMed: 12811365]
42. Niemi M Transporter pharmacogenetics and statin toxicity. *Clin Pharmacol Ther* 2010;87(1):130–3. [PubMed: 19890253]
43. Zhang W, Yu BN, He YJ, et al. Role of BCRP 421C>A polymorphism on rosuvastatin pharmacokinetics in healthy Chinese males. *Clin Chim Acta* 2006;373(1–2): 99–103. [PubMed: 16784736]
44. Ho RH, Choi L, Lee W, et al. Effect of drug transporter genotypes on pravastatin disposition in European- and African-American participants. *Pharmacogenet Genomics* 2007;17(8):647–56. [PubMed: 17622941]
45. Keskitalo JE, Zolk O, Fromm MF, et al. ABCG2 polymorphism markedly affects the pharmacokinetics of atorvastatin and rosuvastatin. *Clin Pharmacol Ther* 2009; 86(2):197–203. [PubMed: 19474787]
46. AstraZeneca. CRESTOR (rosuvastatin) prescribing information. 2014 Available at: <http://www1.astrazeneca-us.com/pi/crestor.pdf>. Accessed November 12, 2014.
47. Liao JK. Safety and efficacy of statins in Asians. *Am J Cardiol* 2007;99(3):410–4. [PubMed: 17261409]
48. Bruckert E, Hayem G, Dejager S, et al. Mild to moderate muscular symptoms with high-dosage statin therapy in hyperlipidemic patients—the PRIMO study. *Cardiovasc Drugs Ther* 2005;19(6):403–14. [PubMed: 16453090]
49. Stewart A SLCO1B1 polymorphisms and statin-induced myopathy PLOS Currents Evidence on Genomic Tests December 4, 2013 Edition 1 10.1371/currents.eogt.d21e7f0c58463571bb0d9d3a19b82203.
50. Fung EC, Crook MA. Statin myopathy: a lipid clinic experience on the tolerability of statin rechallenge. *Cardiovasc Ther* 2012;30(5):e212–8. [PubMed: 21884002]
51. Law M, Rudnicka AR. Statin safety: a systematic review. *Am J Cardiol* 2006; 97(8A):52C–60C.
52. FDA drug safety communication: new restrictions, contraindications, and dose limitations for Zocor (simvastatin) to reduce the risk of muscle injury. 2011 Available at: <http://www.fda.gov/Drugs/DrugSafety/ucm256581.htm>. Accessed November 15, 2014.
53. Thompson PD, Clarkson P, Karas RH. Statin-associated myopathy. *JAMA* 2003; 289(13):1681–90. [PubMed: 12672737]
54. Pasternak RC, Smith SC Jr, Bairey-Merz CN, et al. ACC/AHA/NHLBI clinical advisory on the use and safety of statins. *J Am Coll Cardiol* 2002;40(3):567–72. [PubMed: 12142128]
55. Group HT, Landray MJ, Haynes R, et al. Effects of extended-release niacin with laropiprant in high-risk patients. *N Engl J Med* 2014;371(3):203–12. [PubMed: 25014686]
56. Matsuzawa Y, Kita T, Mabuchi H, et al. Sustained reduction of serum cholesterol in low-dose 6-year simvastatin treatment with minimum side effects in 51,321 Japanese hypercholesterolemic patients. *Circ J* 2003;67(4): 287–94. [PubMed: 12655157]
57. Wu CC, Sy R, Tanphaichitr V, et al. Comparing the efficacy and safety of atorvastatin and simvastatin in Asians with elevated low-density lipoprotein-cholesterol—a multinational, multicenter, double-blind study. *J Formos Med Assoc* 2002;101(7): 478–87. [PubMed: 12353340]
58. Nakamura H, Arakawa K, Itakura H, et al. Primary prevention of cardiovascular disease with pravastatin in Japan (MEGA Study): a prospective randomised controlled trial. *Lancet* 2006;368(9542):1155–63. [PubMed: 17011942]
59. Shepherd J, Cobbe SM, Ford I, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. West of Scotland coronary prevention study group. *N Engl J Med* 1995;333(20):1301–7. [PubMed: 7566020]
60. Colhoun HM, Betteridge DJ, Durrington PN, et al. Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS):

multicentre randomised placebo-controlled trial. *Lancet* 2004;364(9435):685–96. [PubMed: 15325833]

61. Morales D, Chung N, Zhu JR, et al. Efficacy and safety of simvastatin in Asian and non-Asian coronary heart disease patients: a comparison of the GOALLS and STATT studies. *Curr Med Res Opin* 2004;20(8):1235–43. [PubMed: 15324526]
62. Gupta M, Braga MF, Teoh H, et al. Statin effects on LDL and HDL cholesterol in South Asian and white populations. *J Clin Pharmacol* 2009;49(7):831–7. [PubMed: 19398601]
63. Chapman N, Chang CL, Caulfield M, et al. Ethnic variations in lipid-lowering in response to a statin (EVIREST): a substudy of the Anglo-Scandinavian Cardiac Outcomes Trial (ASCOT). *Ethn Dis* 2011;21(2):150–7. [PubMed: 21749017]
64. Albert MA, Glynn RJ, Fonseca FA, et al. Race, ethnicity, and the efficacy of rosuvastatin in primary prevention: the justification for the use of statins in prevention: an intervention trial evaluating rosuvastatin (JUPITER) trial. *Am Heart J* 2011; 162(1):106–14.e102. [PubMed: 21742096]
65. Lipworth L, Fazio S, Kabagambe EK, et al. A prospective study of statin use and mortality among 67,385 blacks and whites in the Southeastern United States. *Clin Epidemiol* 2014;6:15–25. [PubMed: 24379700]
66. Eckel RH, Jakicic JM, Ard JD, et al. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/AMERICAN HEART ASSOCIATION Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014;63:2960–84 (1524–4539 [Electronic]). [PubMed: 24239922]
67. Behavioral risk factor surveillance system: BRFSS 2002 survey data and documentation. Available at: http://www.cdc.gov/brfss/annual_data/annual_2002.htm. Accessed November 15, 2014.
68. Schoenborn CA, Adams PF, Peregoy JA. Health behaviors of adults: United States, 2008–2010. *Vital Health Stat* 10 2013;(257):1–184.
69. Lemacks J, Wells BA, Ilich JZ, et al. Interventions for improving nutrition and physical activity behaviors in adult African American populations: a systematic review, January 2000 through December 2011. *Prev Chronic Dis* 2013;10:E99. [PubMed: 23786910]
70. Ickes MJ, Sharma M. A systematic review of physical activity interventions in Hispanic adults. *J Environ Public Health* 2012;2012:156435. [PubMed: 22496702]
71. Bender MS, Choi J, Won GY, et al. Randomized controlled trial lifestyle interventions for Asian Americans: a systematic review. *Prev Med* 2014;67:171–81. [PubMed: 25086326]
72. Dhokarh R, Himmelgreen DA, Peng YK, et al. Food insecurity is associated with acculturation and social networks in Puerto Rican households. *J Nutr Educ Behav* 2011;43:288–94 (1878–2620 [Electronic]). [PubMed: 20952259]
73. Mainous AG 3rd, Diaz VA, Geesey ME. Acculturation and healthy lifestyle among Latinos with diabetes. *Ann Fam Med* 2008;6:131–7 (1544–1717 [Electronic]). [PubMed: 18332405]
74. Lin H, Bermudez OI, Tucker KL. Dietary patterns of Hispanic elders are associated with acculturation and obesity. *J Nutr* 2003;133:3651–7 (0022–3166 [Print]). [PubMed: 14608089]
75. Palaniappan LP, Araneta MR, Assimes TL, et al. Call to action: cardiovascular disease in Asian Americans: a science advisory from the American Heart Association. *Circulation* 2010;122(12):1242–52. [PubMed: 20733105]
76. Zhou BF, Stamler J, Dennis B, et al. Nutrient intakes of middle-aged men and women in China, Japan, United Kingdom, and United States in the late 1990s: the INTERMAP study. *J Hum Hypertens* 2003;17:623–30 (0950–9240 [Print]). [PubMed: 13679952]
77. Song YJ, Hofstetter CR, Hovell MF, et al. Acculturation and health risk behaviors among Californians of Korean descent. *Prev Med* 2004;39:147–56 (0091–7435 [Print]). [PubMed: 15207996]

KEY POINTS

- Among racial/ethnic groups, Asian Indians, Filipinos and Hispanics are at greater risk for dyslipidemia, which is consistent with the higher coronary heart disease (CHD) mortality rates.
- Compared with other racial/ethnic groups, statins may have a higher efficacy for Asians. Studies suggest lower starting dosage in Asians, but the data are mixed.
- Genetic differences in statin metabolism can in part explain this racial/ethnic difference in statin sensitivity and adverse effects.
- Lifestyle modification is recommended as part of dyslipidemia control and management; African Americans and Hispanics have more sedentary behavior and a less favorable diet profile.